

Hypokalemia in Patients with COVID-19

Alfano Gaetano^{1,2}, Ferrari Annachiara¹, Fontana Francesco², Rossella Perrone¹, Giacomo Mori², Ascione Elisabetta², Magistrone Riccardo^{1,2}, Venturi Giulia³, Simone Pederzoli³, Gianluca Margiotta³, Marilina Romeo³, Francesca Piccinini³, Franceschi Giacomo³, Volpi Sara³, Faltoni Matteo³, Ciusa Giacomo³, Bacca Erica³, Tutone Marco³, Raimondi Alessandro³, Menozzi Marianna³, Franceschini Erica³, Cuomo Gianluca³, Orlando Gabriella³, Santoro Antonella, Di Gaetano Margherita³, Puzzolante Cinzia³, Carli Federica³, Bedini Andrea³, Milic Jovana³, Meschiari Marianna³, Mussini Cristina³, Cappelli Gianni^{1,2}, Guaraldi Giovanni³

¹Surgical, Medical and Dental Department of Morphological Sciences, Section of Nephrology, University of Modena and Reggio Emilia, Modena, Italy; electronic address: gaetano.alfano@unimore.it

²Nephrology Dialysis and Transplant Unit, University Hospital of Modena, Modena, Italy

³Clinic of Infectious Diseases, University Hospital of Modena, Modena, Italy

#Modena Covid-19 Working Group (MoCo19) includes

Cristina Mussini, Giovanni Guaraldi, Erica Bacca, Andrea Bedini, Vanni Borghi, Giulia Burastero, Federica Carli, Giacomo Ciusa, Luca Corradi, Gianluca Cuomo, Margherita Digaetano, Giovanni Dolci, Matteo Faltoni, Riccardo Fantini, Giacomo Franceschi, Erica Franceschini, Vittorio Iadisernia, Damiano Larné, Marianna Menozzi, Marianna Meschiari, Jovana Milic, Gabriella Orlando, Francesco Pellegrino, Alessandro Raimondi, Carlotta Rogati, Antonella Santoro, Roberto Tonelli, Marco Tutone, Sara Volpi, and Dina Yaacoub (Infectious Diseases Clinics, University Hospital, via del Pozzo 71, 41124 Modena, Italy); Gianni Cappelli, Riccardo Magistrone, Gaetano Alfano, Francesco Fontana, Ballestri Marco, Giacomo Mori, Roberto Pulizzi, Elisabetta Ascione, Marco Leonelli, Francesca Facchini, Francesca Damiano, (Nephrology Dialysis and Transplant Unit, University Hospital of Modena, Modena, Italy); Massimo Girardis, Alberto Andreotti, Emanuela Biagioni, Filippo Bondi, Stefano Busani, Giovanni Chierigo, Marzia Scotti, and Lucia Serio (Department of Anesthesia and Intensive Care, University Hospital, via del Pozzo 71, 41124 Modena, Italy); Andrea Cossarizza, Caterina Bellinazzi, Rebecca Borella, Sara De Biasi, Anna De Gaetano, Lucia Fidanza, Lara Gibellini, Anna Iannone, Domenico Lo Tartaro, Marco Mattioli, Milena Nasi, Annamaria Paolini, and Marcello Pinti (Chair of Pathology and Immunology, University of Modena and Reggio Emilia, Via Campi, 287, 41125 Modena, Italy).

Abstract

Patients with COVID-19 may experience multiple conditions (e.g., fever, hyperventilation, anorexia, gastroenteritis, acid-base disorder) that may cause electrolyte imbalances. Hypokalemia is a concerning electrolyte disorder that may increase the susceptibility to various kinds of arrhythmia. This study aimed to estimate prevalence, risk factors and outcome of hypokalemia in a cohort of non-critically ill patients.

A retrospective analysis was conducted on 290 hospitalized patients with confirmed COVID-19 infection at the tertiary teaching hospital of Modena, Italy.

Hypokalemia (<3.5 mEq/L) was detected in 119 patients (41%). The decrease of serum potassium level was of mild entity (3-3.4 mEq/L) and occurred in association with hypocalcemia (P=0.001) and lower level of serum magnesium (P=0.028) compared to normokaliemic patients. Urine K: creatinine ratio, measured in a small subset of patients (n=45; 36.1%), showed an increase of urinary potassium excretion in the majority of the cases (95.5%). Causes of kaliuria were diuretic therapy (53.4%) and corticosteroids (23.3%). In the remaining patients, urinary potassium loss was associated with normal serum magnesium, low sodium excretion (FENa< 1%) and metabolic alkalosis. Risk factors for hypokalemia were female gender (P=0.002; HR 0.41, 95%CI 0.23-0.73) and diuretic therapy (P=0.027; HR 1.94, 95%CI 1.08-3.48). Hypokalemia, adjusted for sex, age and SOFA score, resulted not associated with ICU admission (P=0.131, 95% CI 0.228-1.212) and in-hospital mortality (P=0.474; 95% CI 0,170-1,324) in our cohort of patients.

Hypokalemia is a frequent disorder in COVID-19 patients and urinary potassium loss may be the main cause of hypokalemia. The disorder was mild in the majority of the patients and was unrelated to poor outcomes. Nevertheless, hypokalemic patients required potassium supplements to dampen the risk of arrhythmias.

Introduction

COVID-19 is an emerging pandemic spreading worldwide from December 2019. On through 14 June, the death of more than 430.000 people and the massive increase in the number of newly infected people are posing a growing threat to global health[1].

A Chinese epidemiological study documented that the majority of reported COVID-19 cases had a mild presentation (81%) but the development of severe respiratory symptoms (14%) such as dyspnea, tachypnea and hypoxia requires hospital admission.[2] The lung is the principal target of the novel coronavirus and atypical pneumonia is the most common clinical presentation [3]. Multiple organs such as the heart, brain, liver and kidneys can be involved during this infection[4]. Systemic release of cytokines is thought to be the cause of organ dysfunction and, therefore, the major determinant of morbidity in these patients[5]. It provokes a severe pro-inflammatory state leading to hypoxemia and sepsis requiring intensification of supportive therapy. Generally, hospitalized patients belong to a vulnerable subset of the population because they are older adults and have several underlying medical conditions[6]. Patients with COVID-19 can experience a long hospital stay and undergo multiple treatments varying from delivering of ventilatory support to the administration of experimental agents for SARS-CoV-2 infection. In this setting, fever, hyperventilation, sweating, medication-related side effects, change of eating habits may favorite concerning electrolyte imbalance.

Maturing experience in the care of COVID-19 patients showed that hypokalemia is a frequent lab abnormality. There is particular concern about this disorder as it may increase the susceptibility to potential fatal arrhythmia in COVID patients[7,8]. Hypokalemia has been described in a few case-series during the SARS-CoV1 outbreak in 2003, but there is a lack of data on the exact mechanisms underlying this disorder[9,10]. Possible causes of hypokalemia in the context of SARS-CoV2 infection may result from activation of the renin-angiotensin-aldosterone system, gastrointestinal losses, anorexia secondary to concurrent illness and tubular damage caused by ischemic or nephrotoxic kidney damage. Regarding this latter hypothesis, tubular damage may be linked to the direct cytotoxic effect of SARS-CoV-2, since the virus has been associate to diffuse tubular damage[11].

In light of these unanswered clinical questions, we evaluated a cohort of hospitalized patients with the aim to describe the entity of hypokalemia, impact on clinical outcome, and to achieve a better understanding of the

underlying pathophysiological mechanism leading this condition.

Methods

Study design

A single-center, retrospective, observational study was conducted at the University Hospital of Modena. We retrospectively reviewed all the electronic records of 320 non-critically ill patients admitted to our hospital from February 16, 2020 to April 14, 2020, who were diagnosed with SARS-CoV-2 infection, according to WHO interim guidelines[12].

The study was approved by the regional ethical committee of Emilia Romagna.

Data collection

We collected data on age, sex, laboratory values during admission (hemoglobin concentration, lymphocyte count, platelet count, arterial blood gas analysis, markers of inflammation, serum and urine creatinine, urea, serum and serum and urine electrolyte), comorbidities, sign and symptoms at presentation, vital signs, administered treatment (diuretics, electrolyte supplement), as well as the admission on ICU and living status at the end of the observation period.

Outcomes measures

The primary objective of our study was to evaluate the prevalence of hypokalemia. Secondary outcomes included measurement of urinary potassium excretion, the proportion of patients requiring supplemental K, and the impact of hypokalemia on ICU admission and in-hospital mortality compared to normokalemia.

Variables and definition

Hypokalemia was defined as a serum potassium level < 3.5 mEq/l. The normal level of serum potassium ranges from 3.5 to 5.3 mEq/L. The diagnosis of hypokalemia was performed on a single value of serum potassium < 3.5 meq/L at any time during hospitalization. We excluded all serum values of potassium measured on plasma and by blood gas analyzer.

Severity was classified as mild when the serum potassium level was 3 to 3.4 mmol/L, moderate when the serum potassium level was 2.5 to 3 mmol/L, and severe when the serum potassium level was less than 2.5 mmol/L

K urinary excretion was measured as urine K⁺-to-creatinine ratio. A value of K⁺ urinary excretion > 1.5 indicated inappropriate renal potassium loss. K urinary excretion was not measured in patients with chronic and acute kidney disease.

Na urinary excretion was measured with the fractional excretion of sodium (FeNa); a value < 1% declared a reduced excretion of sodium, whereas it resulted augmented for value ≥1%.

Sequential Organ Failure Assessment (SOFA) score was calculated at hospital admission.

Statistical analysis

Baseline characteristics were analyzed using descriptive statistics and reported as proportions and mean (standard deviation [SD]) when appropriate. Categorical variables were analyzed using χ^2 test or Fisher's test as appropriate. Analyses of continuous variables were compared using an unpaired t-test and Mann-Whitney test, as appropriate. Logistic regression analysis was used to identify the predictors of hypokalemia and to test the risk of in-hospital mortality, adjusted for sex, age and SOFA score at ward admission. A P value of less than 0.05 was considered statistically significant. SPSS 23® was used for statistical analysis.

Results

Patients

Serum K⁺ level was measured in 290 (1671 samples) patients with confirmed COVID-19. Three patients with hyperkalemia were excluded for further analysis whereas 171 patients with normokalemia were selected as control group. Hypokalemia (K <3.5 mEq/L) was detected in 119 patients (41%) and serum potassium ranged from 2.4 mEq/l to 3.5 mEq/l with a mean average of 3.1 (SD±0.18) mEq/l.

Hypokalemia was mild (3.4-3 mEq/l) in the majority of the cases (90.7%). It was associated with hypocalcemia (P=0.001) and a lower level of serum magnesium (P=0.028) compared to normokaliemic patients. Patients with hypokalemia had a longer follow-up (P=0.002) compared to normokaliemic patients likely for a severe course of COVID-19, as showed by a statistically significant higher SOFA score (P=0.014) at the admission.

Mean serum creatinine of hypokalemic patients was 0.9 ± 0.9 mg/dl. Demographics and clinical manifestations of patients are reported in Table 1 and 2.

Potassium excretion

Urine K-to-creatinine ratio was measured only in a minority of hypokalemic patients (n=45;36.1%). Urine K-to-creatinine ratio >1.5 mEq/mmol, expression of high urinary excretion of a potassium[13], was documented in 43 (95.5%) patients. Twenty-three patients (53.4%) with high urinary potassium excretion were on diuretic therapy (furosemide) and 10 (23.3%) were on corticosteroids at the time of serum K measurement. The remaining 10 patients (23.3%) with augmented kaliuresis had normal serum level of magnesium and 90% had low sodium excretion ($FENa < 1\%$) (Fig.1). Acid-base status of these ten patients showed that 70% had metabolic alkalosis, 10% respiratory alkalosis, 10% respiratory acidosis and the remaining 10% a normal acid-base status

Therapy in hypokalemic patients

Potassium supplements were administrated to correct hypokalemia (Table 2). Salts of potassium (KCl, 600mg) were administered via the oral route in 28 (23.5%) patients and IV in 7 (5.8%) patients, both oral route and iv were used in one (0.8%) patient.

Intravenous magnesium sulfate was administered in 22 (18.4%) patients for hypomagnesemia

Potassium-sparing diuretics were largely administered among hypokalemic patients (26 %) compared to normokalemia (7%), likely for its pharmacological action targeted to maintain a normal kalemia and prevent hypokalemia (Table 2). Forty-five (37.5%) patients were on diuretic therapy with furosemide and two (1.6%) with thiazide at the time of diagnosis with hypokalemia. Hypertension (57.4%), cardiovascular disease (38.2%) and renal impairment (31.9%) were the potential causes of diuretic use in these patients.

Risk factor and Outcome

Univariate analysis showed that female gender, corticosteroids and diuretics were significantly associated with hypokalemia in our cohort of patients. Multivariate regression model analysis showed that diuretic therapy ($P=0.027$; HR 1.94; 95% CI 1.08-3.48) and female sex ($P=0.003$; HR 2.44; 95% CI 1.36-4.37) were significant risk factors for hypokalemia. Survival regression analysis showed that hypokalemia adjusted for

sex, age and SOFA score was not associated with intensive care admission (P=0.131, 95% CI 0.228-1.212) and in-hospital mortality (P=0.474; 95% CI 0.170-1.324).

Discussion

The results of our study showed that hypokalemia is a common electrolytic abnormality among patients with COVID-19. Hypokalemia has been found in 41% of patients during hospitalization for severe symptoms of SARS-CoV-2 infection. The prevalence and causes of hypokalemia are unknown until now, as this phenomenon has been only mentioned as a potential manifestation of COVID-19[7]. However, there is a general concern about the risk of hypokalemia to trigger life-threatening arrhythmia in patients affected by SARS-CoV-2[7,8].

Diuretic therapy along with female sex was significantly associated with hypokalemia in our study-cohort. Diuretics are pharmacological agents largely used in the general population to treat hypertension and prevent fluid overload. The slightly higher prevalence of hypertension, cardiovascular disease and renal impairment in hypokaliemic than normokaliemic patients may be a plausible explanation of the higher use of diuretics in this group of patients. Diuretics are widely known to cause hypokalemia and other electrolyte imbalances with long-term administration [14], as they act by inhibiting reabsorption of electrolytes in the renal tubules. This side effect was likely the major determinant of other electrolyte disturbances in our patients. Hypokalemia was indeed associated with hypocalcemia and a lower level of serum magnesium compared to normokaliemic patients. Although the female sex appears poorly interlinked with hypokalemia, the first experimental studies conducted in the 50s [15] and then confirmed in the 90s [11] showed that women, especially aged ones, have less exchangeable body potassium than other subset of the population. The women are, therefore, at high risk to develop hypokalemia because they have depleted deposits of potassium due to their different body composition, characterized by less amount of extracellular water compared to men.

Hypokalemia is a frequent electrolyte disturbance in hospitalized patients. Multiple mechanisms underlying this condition include diuretic therapy, gastrointestinal loss, anorexia or alkalosis[22]. More rarely, congenital or acquired tubular defect, side effects of drugs or tubular ischemia induce tubular potassium losses due to the disruption of renal electrolytes handling[23,24]. Hyperaldosteronism, primitive or as a consequence of

activation of the renin-angiotensin system, is also known to stimulate urinary potassium excretion. In the setting of COVID-19 infection, several adjunctive conditions can disrupt potassium homeostasis during hospitalization. Respiratory alkalosis due to hypoxia-driven hyperventilation may provoke transcellular shifts with increased intracellular uptake, and anorexia, as a consequence of continuative use of face mask or ventilation helmet or status of severe illness, may lead to a decrease in potassium intake; diarrhea due to medication (e.g., lopinavir/ritonavir) or the cytopathic effect of the virus on the gastrointestinal cells may be a cause of potassium losses.

Moderate and severe hypokalemia has widely been associated with potentially life-threatening complications such as cardiac dysrhythmias, paralysis and rhabdomyolysis [16]. Underlying cardiovascular disease [17,18] and chronic kidney disease [19,20] are widely recognized as a poor prognostic factor for mortality among hypokalemic patients.

The findings of our study show that patients with hypokalemia had a statistically longer hospitalization than controls. Likely the higher rate of respiratory symptoms (dyspnea and cough) and the worse SOFA score (2.7 vs 2.2) at admission reflected a more severe systemic inflammatory response leading to a prolonged hospital stay than normokaliemic patients. Nevertheless, we did not notice that hypokalemia was not a poor prognostic factor for intensive care admission ($P=0.131$, 95% CI 0.228-1.212) and hospital mortality ($P=0.474$; 95% CI 0,170-1,324). Likely the mild severity of hypokalemia (90.7% had serum levels of potassium between 3.4 and 3 mEq/L) may explain the null effect of hypokalemia on major outcomes in our cohort of patients.

Despite these reassuring data, serum potassium should be carefully monitored in this vulnerable subset of the population for the risk of potentially lethal arrhythmia. Hypokalemia may further prolong QT interval in COVID-19 patients with abnormal QT interval due to the large use of off-label agents for SARS-CoV-2 infection such as azithromycin and hydroxychloroquine.

To date, few reports document an interplay between hypokalemia and coronavirus infection. Hypokalemia has been reported during the SARS-CoV-1 outbreak in Canada. Booth et al[9] described that 62 out of 144 (43%) patients had mild hypokalemia (mean, 3.2 mEq/l) during hospitalization. Hypokalemia was not isolated and occurred with other electrolyte abnormalities such as hypomagnesemia and hypophosphatemia. The authors were unable to establish clear evidence for electrolyte imbalance and suggested that it may be a clinical

manifestation of the infection or a nephrotoxic effect of ribavirin or other medicaments on renal electrolytes handling. The abstract of a Chinese article, reported hypokalemia as a side effect of glucocorticoid administration in SARS-CoV-1 infected patients[10]. These agents through activation of the mineralocorticoid receptor on the renal tubular cells could have increased urinary potassium excretion[25].

In an attempt to clarify tubular potassium handling in COVID-19 we investigated the magnitude of potassium secretion in a minority of hypokalemic patients. Urine K-to-creatinine ratio was augmented in 95.5% of patients who underwent urinary examination (n=45), expression of pathological tubular potassium handling. After the exclusion of about three-fourths of urinary samples because of intercurrent diuretic and corticosteroid therapy, we found that in the small proportion of remaining patients (n=10, 23.3%), urinary examination was characterized by an augmented urinary potassium excretion in the absence of concurrent causes (normal serum magnesemia). Interestingly, most of them (70%) had metabolic alkalosis on arterial blood gas analysis. Based on these data, the etiology of hypokalemia of COVID patients seems multifactorial and a side effect of diuretic and corticosteroid therapy. However, we cannot overlook the inappropriate loss of urinary potassium occurring in a small number of patients in the absence of apparent pharmacological interferences. Two hypotheses should be enunciated regarding this issue. First, SARS-COV-2 may mediate a direct viral cytopathic effect on kidney tubular proximal cells with loss of renal tubular cell integrity and function. In support of this hypothesis, the virus has been identified in urine samples in patients with severe COVID-19[26], as ACE 2, virus entry receptor, is largely expressed within the kidney, especially in tubular cells[27]. The virus may cause diffuse acute proximal tubular injury characterized by loss of apical brush border and vacuolation[11]. The second hypothesis relies on hyperaldosteronism due to the abnormal activation of the renin-angiotensin system as a result of ACE2 disruption following cell infection[28]. The findings of metabolic alkalosis, low sodium excretion corroborate this theory, but the lack of data on serum renin and aldosterone levels cannot corroborate our hypothesis.

Several other limitations of this study must be enunciated. First, we were unable to trace hypokalemia-related clinical manifestations despite the completeness of the electronic chart. Second, the restricted group of patients who underwent urinary examination has reduced the accuracy of analysis to identify the causes of

hypokalemia. However, despite the small sample, we have collected comprehensive data on magnesium level, urine sodium and acid-base status.

This study is the first report describing in detail the clinical characteristic of hypokalemic patients affected by SARS-CoV-2. Although urine examination was available only in a small number of patients, we traced the potential causes of hyperkalemia, including diuretic therapy, acid-base disorder and corticosteroid therapy, in a cohort of hospitalized patients. Our results support the need to frequently measure the level of serum potassium along with urine K-to-creatinine ratio or 24h potassium excretion in COVID-19 patients in order to improve the management of these patients. Based on the magnitude of the urinary loss, potassium should be supplemented per os or IV, and a careful assessment of the ECG should be performed, especially in concomitance of potentially arrhythmogenic drugs. Further studies are required to investigate the hypothesized cytopathic effect of SARS-CoV-2 on renal tubular cells and other secondary injury mechanisms such as hypoxia and cytokine storm.

Conclusions

Hypokalemia was a frequent electrolyte disorder in our hospitalized patients with COVID-19. It was highly associated with urinary loss due principally to iatrogenic causes. In a small proportion of patients (22.2%), the etiology of tubular potassium loss remained elusive. Generally, hypokalemia was mild, not associated with poor outcomes and was corrected with oral supplementation of potassium. Although hypokalemia was not associated with mortality in our cohort of patients, it may be a life-threatening condition (e.g., long QT) if it remains untreated.

Bibliography

- 1 COVID-19 Map; Coronavirus COVID-19 Global Cases by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU) [Internet]. Johns Hopkins Coronavirus Resour Cent. [cited 2020 June 14]. Available from: <https://coronavirus.jhu.edu/map.html>
- 2 Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020 Apr;323(13):1239–42.
- 3 Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020 Feb;0(0):null.
- 4 Wang T, Du Z, Zhu F, Cao Z, An Y, Gao Y, et al. Comorbidities and multi-organ injuries in the treatment of COVID-19. *The Lancet*. 2020 Mar;395(10228):e52.
- 5 Moore BJB, June CH. Cytokine release syndrome in severe COVID-19. *Science*. 2020 Apr DOI: 10.1126/science.abb8925
- 6 Garg S. Hospitalization Rates and Characteristics of Patients Hospitalized with Laboratory-Confirmed Coronavirus Disease 2019 — COVID-NET, 14 States, March 1–30, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69. DOI: 10.15585/mmwr.mm6915e3
- 7 Bansal M. Cardiovascular disease and COVID-19. *Diabetes Metab Syndr Clin Res Rev*. 2020 May;14(3):247–50.
- 8 Basu-Ray I, Soos MP. Cardiac Manifestations Of Coronavirus (COVID-19). *StatPearls*. Treasure Island (FL): StatPearls Publishing; 2020; [cited 2020 May 30]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK556152/>
- 9 Booth CM, Matukas LM, Tomlinson GA, Rachlis AR, Rose DB, Dwosh HA, et al. Clinical features and short-term outcomes of 144 patients with SARS in the greater Toronto area. *JAMA*. 2003 Jun;289(21):2801–9.
- 10 Li N, Wang G, Wu Y, Xie G, Xiao F, Chen B, et al. [Side effects of glucocorticosteroids in the management of 1291 patients of SARS]. *Beijing Da Xue Xue Bao*. 2004 Oct;36(5):519–24.
- 11 Su H, Yang M, Wan C, Yi L-X, Tang F, Zhu H-Y, et al. Renal histopathological analysis of 26 postmortem findings of patients with COVID-19 in China. *Kidney Int*. 2020 Apr;0(0). DOI: 10.1016/j.kint.2020.04.003

- 12 WHO/COVID-19/laboratory/2020.5. Laboratory testing for 2019 novel coronavirus (2019-nCoV) in suspected human cases [Internet]. [cited 2020 Apr 12]. Available from: <https://www.who.int/publications-detail/laboratory-testing-for-2019-novel-coronavirus-in-suspected-human-cases-20200117>
- 13 Jędrusik P, Symonides B, Wojciechowska E, Gryglas A, Gaciong Z. Diagnostic value of potassium level in a spot urine sample as an index of 24-hour urinary potassium excretion in unselected patients hospitalized in a hypertension unit. *PloS One*. 2017;12(6):e0180117.
- 14 Arampatzis S, Funk G-C, Leichtle AB, Fiedler G-M, Schwarz C, Zimmermann H, et al. Impact of diuretic therapy-associated electrolyte disorders present on admission to the emergency department: a cross-sectional analysis. *BMC Med*. 2013 Mar;11:83.
- 15 Sagild U. Total exchangeable potassium in normal subjects with special reference to changes with age. *Scand J Clin Lab Invest*. 1956;8(1):44–50.
- 16 Gennari FJ. Disorders of potassium homeostasis. Hypokalemia and hyperkalemia. *Crit Care Clin*. 2002 Apr;18(2):273–88, vi.
- 17 Bowling CB, Pitt B, Ahmed MI, Aban IB, Sanders PW, Mujib M, et al. Hypokalemia and outcomes in patients with chronic heart failure and chronic kidney disease: findings from propensity-matched studies. *Circ Heart Fail*. 2010 Mar;3(2):253–60.
- 18 Goyal A, Spertus JA, Gosch K, Venkitachalam L, Jones PG, Berghe GV den, et al. Serum Potassium Levels and Mortality in Acute Myocardial Infarction. *JAMA*. 2012 Jan;307(2):157–64.
- 19 Gilligan S, Raphael KL. Hyperkalemia and Hypokalemia in CKD: Prevalence, Risk Factors, and Clinical Outcomes. *Adv Chronic Kidney Dis*. 2017 Sep;24(5):315–8.
- 20 Lee S, Kang E, Yoo KD, Choi Y, Kim DK, Joo KW, et al. Lower serum potassium associated with increased mortality in dialysis patients: A nationwide prospective observational cohort study in Korea. *PLoS ONE*. 2017 Mar;12(3). DOI: 10.1371/journal.pone.0171842
- 21 Jensen HK, Brabrand M, Vinholt PJ, Hallas J, Lassen AT. Hypokalemia in acute medical patients: risk factors and prognosis. *Am J Med*. 2015 Jan;128(1):60-67.e1.
- 22 Paice BJ, Paterson KR, Onyanga-Omara F, Donnelly T, Gray JM, Lawson DH. Record linkage study of hypokalaemia in hospitalized patients. *Postgrad Med J*. 1986 Mar;62(725):187–91.
- 23 Asmar A, Mohandas R, Wingo CS. A Physiologic-Based Approach to the Treatment of a Patient With Hypokalemia. *Am J Kidney Dis Off J Natl Kidney Found*. 2012 Sep;60(3):492–7.

- 24 Unwin RJ, Luft FC, Shirley DG. Pathophysiology and management of hypokalemia: a clinical perspective. *Nat Rev Nephrol*. 2011 Feb;7(2):75–84.
- 25 Palmer BF. Regulation of Potassium Homeostasis. *Clin J Am Soc Nephrol CJASN*. 2015 Jun;10(6):1050–60.
- 26 Guan W-J, Zhong N-S. Clinical Characteristics of Covid-19 in China. Reply. *N Engl J Med*. 2020 Mar;382. DOI: 10.1056/NEJMc2005203
- 27 Soler MJ, Wysocki J, Batlle D. ACE2 alterations in kidney disease. *Nephrol Dial Transplant Off Publ Eur Dial Transpl Assoc - Eur Ren Assoc*. 2013 Nov;28(11):2687–97.
- 28 Alfano G, Guaraldi G, Fontana F, Ferrari A, Magistroni R, Mussini C, et al. The Role of the Renin-Angiotensin System in Severe Acute Respiratory Syndrome-CoV-2 Infection. *Blood Purif*. 2020 Apr;1–5.

Table 1. Demographic and clinical characteristics of hypokalemic (K<3.5; mEq/L) and normokaliemic (K 3.5-5.3 mEq/L) COVID-19 patients.

Characteristic	Hypokalemic N= 119	Normokaliemic N=171	P
<i>Age, years</i>			
Mean (\pm SD)	65,8 (12,9)	64.1 (14.4)	0.258
<i>Gender, n(%)</i>			
Male	73 (61,3)	138 (80.7)	0.001
<i>Comorbidities, n(%)*</i>			
Diabetes	26 (27.9)	24 (16.3)	0.113
Hypertension	54 (83)	74(76.2)	0.81
Cardiovascular Disease	33 (38)	32 (23)	0.085
Chronic Renal Insufficiency	21 (21.4)	29 (20.4)	0.875
<i>Sign and symptoms, n(%)*</i>			
Fever, mean (\pm SD)	37,1 (1)	36.9 (1)	0.287
Dyspnea	61(56)	64 (50)	0.022
Diarrhea	17(15,7)	15 (11.7)	0.182
Cough	66 (61.1%)	68 (53.1)	0.009
Fatigue	25(23,1)	39 (30.5)	0.774
Myalgia	12(11,1)	8 (6.3)	0.098
Headache	9(8,3)	15 (11.7)	0.83
<i>Clinical characteristic</i>			
Systolic pressure, mmHg mean (\pm SD)	125 (18,4)	122.1 (20)	0.245
Diastolic pressure, mmHg (\pm SD)	73,3 (11,7)	72.8 (11.3)	0.911
Respiratory rate	24,5 (8,3)	23 (6.3)	0.428
Baseline PaO ₂ /FiO ₂	217.9 (101.5)	235.6 (108.8)	0.215
<i>Acid-base status n(%)*</i>			
Normal	44 (40.3)	68 (45.6)	0.713
Metabolic alkalosis	49 (44.9)	55 (36.9)	0.135
Respiratory alkalosis	12 (11)	8 (5.3)	0.098
Respiratory acidosis	4 (3.6)	9 (6)	0.569
SOFA Score	2,7 (2,1)	2.2 (2.1)	0.014
<i>Treatment, n(%)*</i>			
Furosemide	45 (37.8)	46 (26.3)	0.066
Tiazide	2 (1,6)	2 (1,1)	0.713

Potassium-sparing diuretics	31 (26)	12 (7)	0.0001
Corticosteroid	46 (38.6)	53 (30.9)	0.219
<i>Status at the end of F-U</i>			
Follow-up, days	14,1 (7,5)	11.4 (7.1)	0.002
ICU admission (%)	11(9,2)	29(16,9)	0.083
Death	17 (14,2)	28(16.3)	0.742

ICU denotes intensive care unit; SD, standard deviation; SOFA, sequential organ failure assessment

* T-test or Mann Whitney test as appropriate; χ^2 test for categorical variable

Table 2. Main lab examinations of hypokalemic and normokaliemic patients

<i>Lab test, Mean (±SD)</i>	Hypokalemic	Normokalemic	<i>P value*</i>
Hemoglobin, g/l	12,1 (1,6)	12.8 (1.793)	0.001
White cells, mm ³	7393 (5044)	8156.6 (8228.1)	0.750
Platelets, 10 ⁹ /l	252,3 (119,3)	243.2 (116.1)	0.482
Glycemia	105,2 (35,9)	110.6 (54.4)	0.656
Potassium, mmol/l	3,1 (0,1)	4.0 (0.3)	< 0.0001
Sodium, mmol/l	137.4 (3,9)	137.2 (3.6)	0.818
Calcium, mg/l	8,3 (0,5)	8.7 (0.6)	0.001
Magnesium	2,1 (0,3)	2.6 (0.3)	0.028
Chloride, mmol/l	99.6 (4,5)	100.2 (4.4)	0.509
Creatinine, mg/l	0,9 (0,9)	1.1 (1.3)	< 0.0001
Urea, mg/dl	55,31 (47,8)	52 (30.3)	0.564
Albumin, gr/dl	3,2 (0,5)	3.1 (05)	0.519
Urine K ⁺ (spot)	30,1 (16,7)	38.1 (21.8)	0.229
Urine Creatinine (spot)	144,4 (72,4)	110.1 (62.1)	0.079
Urine protein-to-creatinine	0,5 (0,5)	0.4(0.31)	0.300
D-dimer, mg/l	2478 (4070.8)	3045.7 (5718.3)	0.826
Alanine amino-transferase, U/l	45,9(37,2)	109.65 (761.7)	0.371
Lactate dehydrogenase, U/l	687,2 (310)	812.2 (1914.1)	0.348
Ferritin, mg/dl	867 (416.7)	856.4 (824.1)	0.373
Bilirubin, mg/l	0.7 (0.5, 0.3)	0.7 (0.7)	0.301
C-reactive protein, mg/l	10,2 (9,9)	9.8 (7.6)	0.524

*T-test or Mann Whitney test as appropriate

Table 3A. Predictive factors of hypokalemia revealed by univariate logistic regression analysis

	HR	CI(95%)		<i>p</i> value
Age	1,01	0,99	1,03	0,31
PO₂/FiO₂	1,00	0,97	1,00	0,30
Sex				
Female	2,64	1,55	4,48	<0,001
SOFA	1,00	1,00	1,00	0,23
Serum magnesium	0,31	0,09	1,01	0,05
Alkalosis	1,55	0,94	2,57	0,09
Corticosteroid				
Yes/No	1,66	1,01	2,74	0,05
Diuretic Therapy				
Yes/No	2,00	1,22	3,27	0,01

Table 3B. Multivariate regression model of predictive factors of hypokalemia

	HR	CI(95%)		<i>p</i> value
Alkalosis	1,50	0,89	2,53	0,128
Diuretic Therapy	1,94	1,08	3,48	0,027
Sex				
Female	2,44	1,36	4,37	0,003
Corticosteroid	1,25	0,69	2,27	0,467

Figure1. Flow chart of patients' selection

