

Could masks curtail the post-lockdown resurgence of COVID-19 in the US?

Calistus N. Ngonghala^{†,††,*}, Enahoro Iboi[‡], and Abba B. Gumel^{†,††}

[†] *Department of Mathematics, University of Florida, Gainesville, FL 32611, USA.*

^{††} *Emerging Pathogens Institute, University of Florida, Gainesville, FL 32610, USA.*

[‡] *School of Mathematical and Statistical Sciences, Arizona State University, Tempe, Arizona, 85287, USA.*

^{††} *Other affiliation: Department of Mathematics and Applied Mathematics, University of Pretoria, Pretoria 0002, South Africa.*

July 6, 2020

Abstract

The community lockdown measures implemented in the United States, during late March to end of May of 2020, resulted in a significant reduction in the community transmission of the COVID-19 pandemic throughout the country. However, a number of US states are currently experiencing an alarming post-lockdown resurgence of the pandemic, triggering the fear for a possible severe second wave of the pandemic in some US jurisdictions. We designed a mathematical model for addressing the key question of whether or not the universal use of face masks can halt or curtail such resurgence (and possibly avert a second wave, without having to undergo another cycle of major community lockdown) in the states of Arizona, Florida, New York and the entire US. The model was parametrized and fitted using cumulative mortality data from the four jurisdictions. Our study highlights the importance of early implementation of the community lockdown measures. In particular, a sizable reduction in the burden of the pandemic would have been recorded in each of the four jurisdictions if the community lockdown measures were implemented a week or two earlier. These reductions are greatly augmented if the early implementation of the lockdown measures is complemented with a public face masks use strategy. It is shown that the pandemic would have been almost completely suppressed from significantly taking off if the lockdown measures were implemented two weeks earlier, and if a sizable percentage of the residents of the four jurisdictions wore face masks during the respective lockdown periods. If the level of lifting of community lockdown is high (which entails allowing for greater community contacts and re-opening of businesses and social activities, in comparison to what was allowed during the community lockdown period), the states of Arizona and Florida will record a devastating second wave of the pandemic by the end of 2020, while the state of New York and the entire US will record milder second waves. If the level of lifting for the community lockdown was mild (i.e., only allowing very limited community contacts and business activities, in comparison to the lockdown period), only the state of Florida will experience a second wave. The severity of the projected second wave depends on the level of lifting of the community lockdown. For instance, the projected second wave for Arizona and Florida, associated with moderate and high level of lifting of lockdown, will be more severe than their respective first wave. For high level of lifting of lockdown measures, the increased use of face masks after the lockdown period greatly reduces the burden of the pandemic. In particular, for this high lifting scenario, none of the four jurisdictions will experience a second wave if half of their residents wear face masks consistently after their respective lockdown period). A testing strategy that increases the maximum detection rate of asymptomatic infected individuals (followed by contact tracing and self-isolation of the detected cases) greatly reduces the burden of the pandemic in all four jurisdictions, particularly if also combined with universal face mask use strategy. Universal use of face masks in public, with at least moderate level of compliance, could halt the post-lockdown resurgence of COVID-19, in addition to averting the potential for (or severity of) a second wave of the pandemic in each of the four jurisdictions.

Keywords: *Mathematical model; COVID-19; community lockdown; face masks; compliance; lifting; detection; testing; contact tracing; self-isolation.*

*Corresponding author: Email: calistusnn@ufl.edu

1 Introduction

The novel coronavirus (COVID-19) pandemic that emerged in December 2019 is, undoubtedly, the most important public health challenge facing mankind since the 1918 pandemic of influenza. The rampaging pandemic (which spread to all countries on earth), caused by a new Severe Acute Respiratory syndrome (SARS-CoV-2), continues to cause devastating public health and socio-economic impact in many parts of the world, including the US [1, 2]. The pandemic accounted for over 10.5 million confirmed cases and 513,000 deaths by the end of June 2020 [3, 4]. Although the pandemic emerged from China, the US and Brazil bore the brunt of its public health burden (with over 2.7 million cases and 130,122 deaths for the US; and over 1.4 million cases and 59,656 deaths for Brazil, as of June 30, 2020). In particular, the US state of New York (alone) recorded about 417,836 confirmed cases and 32,129 COVID-induced mortality as of June 30, 2020. Owing to the effective and sustained control measures implemented in New York state, the state (which was once the global epicenter of the pandemic) is now recording low daily new cases and COVID-related mortality (in fact, the mortality numbers now recorded in New York state is among the lowest in the US). Although many US states have been witnessing declines in disease incidence (since the lifting of lockdown measures), a number of states (notably Arizona, California, Florida and Texas) have recently started experiencing a dramatic increase in the number of cases and hospitalizations. In particular, Arizona and Florida recorded 4,877 (on July 1, 2020) and 10,109 (on July 2, 2020), respectively (these staggering numbers are the highest daily case count for both states since the pandemic started) [5]. Furthermore, ICU admissions at the Texas Medical Center in Houston City was at 97% bed capacity as of June 24, 2020 [6]. The current trend suggests that Arizona and Florida may be the next epicenters of the pandemic in the US. As of July 2, 2020, the two states accounted for about 79,215 and 152,434 confirmed cases, and 1,632 and 3,505 cumulative deaths, respectively). A total of 57,232 new confirmed cases was reported nationwide on the same day.

The main COVID-19 transmission pathways are person-to-person transmission through respiratory droplets, and transmission through contaminated surfaces [7]. Studies have shown that some individuals infected with the novel coronavirus can be asymptomatic or symptomatic with mild, moderate, severe, or critical symptoms [8–15]. Asymptomatic individuals exhibit no COVID-19 symptoms, although they contribute in disease transmission [8]. They include pre-symptomatic individuals who start shedding the virus before the onset of symptoms [16] and infectious individuals who do not show clinical disease symptoms after the incubation period. Those with mild clinical symptoms suffer from light fever, sneezing, cough, discomfort, etc., but no pneumonia, acute respiratory distress syndrome (ARDS). These individuals do not require Intensive Care Unit (ICU) admission [8, 9]. Some of the individuals with moderate symptoms, particularly the elderly (those aged 65+) and those with pre-existing health conditions, might experience a mild form of pneumonia that require self-isolation or hospitalization, but not ARDS or ICU admission [10–13]. Clinically severe cases of COVID-19 develop symptoms that include acute respiratory distress and failure, which might damage the lungs, as well as complications requiring hospitalization and possible ventilation [9, 14]. Critical cases require ICU admission and ARDS ventilators for survival. They are at high mortality risk and generally include the elderly (those above 65 years) and people with underlying health conditions [11]. It is known that about 81% of COVID-19 confirmed cases show mild to moderate or no symptoms, 14% show severe clinical symptoms, and approximately 5% of the cases are clinically ill [8, 14]. In addition to transmission by the clinically symptomatic, pre-symptomatic and asymptomatic individuals contribute significantly to disease transmission [17–19]. This complicates control efforts, especially when they are focused only on the severely or critically ill cases. When infectious individuals show only mild to moderate or no symptoms, it is likely that they have no urge to seek medical aid or adhere to preventive measures, thereby causing negative impact on disease mitigation or containment efforts. Hence, distinguishing between various forms of disease severity is important not only in reducing community transmission (and socio-economic burden), but also contribute in lowering the risk of infection health care workers face.

Although concerted global efforts are exerted towards developing a safe and effective vaccine against COVID-19 [20–22], no such vaccine is expected to be ready for use in humans within the next three months. Furthermore, there is currently no safe and effective antiviral against the virus. Consequently, control and mitigation efforts against COVID-19 are restricted to the implementation of non-pharmaceutical interventions (NPIs). These interventions include community lockdowns, maintaining social (or physical)-distancing in public, wearing face masks

in public, quarantine of suspected cases, isolation of confirmed cases, contact-tracing and testing. Community lockdowns and social-distancing have been widely implemented in most US states (they entailed asking people to shelter at home, closing large gathering places, such as schools, malls and non-essential businesses). The use of face masks in public is also encouraged by many state and local governments [23]. The success of any of these measures relies on a number of factors. For example, the success of lockdown, social-distancing and face mask use measures relies on the willingness of the members of the general population to comply with the prescribed measures and the effectiveness of the measures (e.g., face mask effectiveness to prevent the transmission or acquisition of infection) [23–26]. Diagnostic (surveillance) testing relies on the ability to test as many people as possible and how sensitive the test is in identifying cases, especially people at the early stage of the COVID-19 infection.

By April 7, 2020, mandatory lockdown measures were put in place in over 42 US states [27]. While these lockdown measures and other NPIs (such as social-distancing, the use of face masks in public, personal hygiene, testing, etc.) have been effective in curtailing the spread of COVID-19 in many US states, especially the state of New York, the late implementation and early relaxation of these lockdown measures seem to be having negative effects on the effort to effectively curtail the pandemic in some US states. It should be recalled that the US White House Coronavirus Task Force announced a four-phase guidelines, on April 16, 2020, for states to meet before considering relaxing the community lockdowns they implemented [28]. Specifically, Phase 1 of the guidelines requires states to consider lifting lockdown measures if they (i) achieve two weeks of continuous decline in the number of COVID-19 cases, (ii) have enough testing capacity for at-risk health care workers, and (iii) have enough hospital capacity. Relaxation of strict lockdown measures should rely heavily on the ability of states in particular, and the nation in general, to test and contact trace as many people as possible, encourage a sizable population to use face masks consistently in public and socially (or physically)-distance (stay 6 feet apart). Although no US state has met all four phases as of May 20, 2020 (in fact, it is doubtful if any state has met even the first phase [29, 30]), almost all the US states started relaxing the community lockdown measures by this time. This premature relaxation or lifting of the community lockdown measures (in an effort to re-open the economy) is clearly responsible for the post-lockdown resurgence of COVID-19 in a number of states (associated with spikes in confirmed cases and hospitalizations), including in the aforementioned states of Arizona, California, Florida and Texas [29, 30]. A natural question to ask, based on the current trends, is whether or not the post-lockdown resurgence would trigger a severe second wave of the pandemic in some US states, or in the entire country (and how severe the second wave might be, in comparison to the first). It is notable that some US states have started taking important measures to help halt the post-lockdown resurgence, such as pausing the re-opening phases, re-locking down some businesses, and enacting mandatory mask usage ordinances. In fact, some counties have implemented, or are contemplating, re-imposing stay-at-home orders [31–34].

In this study, we develop and use a mathematical model to address the important question of whether or not masks could curtail the post-lockdown resurgence of COVID-19 in the US. Specifically, we use the model to assess the impact of (1) early implementation of lockdown measures, combined with increased mask usage; (2) varying levels of lifting of community lockdown measures (and how increases in face mask compliance affect the impact) and (3) detection (including tracing, testing and self-isolation) of asymptomatic infected individuals on control and mitigating the burden of the pandemic in the chosen jurisdictions. The model will also be used to predict the likelihood (and severity) of a second wave of the pandemic in the US. The model is designed and calibrated, using cumulative mortality data from the four jurisdictions, in Section 2. Theoretical analyses of the model, with respect to the asymptotic stability of its associated continuum of disease-free equilibria, is carried out in Section 3. An expression for the final size of the pandemic is also derived analytically. Numerical simulations of the model are reported in Section 4.

2 Materials and Methods

2.1 Model Formulation

The objectives of this study will be achieved *via* the design, analysis, parametrization and simulations of a Kermack-McKendrick-type SEIR (susceptible-exposed-infectious-recovered) epidemic (no human demography) model for the

transmission dynamics and control of COVID-19 in a population. In addition to incorporating pertinent aspects of the epidemiology of the disease, the model to be developed will allow for the assessment of the non-pharmaceutical interventions being implemented in the US, notably social-distancing, face mask usage and contact-tracing and testing. The model to be developed is based on stratifying the total human population at time t , denoted by $N(t)$, into mutually exclusive compartments based on disease status. Specifically, we split $N(t)$ into the sub-populations of individuals who are susceptible ($S(t)$; i.e., uninfected individuals who may contract the disease at a later time), exposed ($E(t)$; i.e., individuals who are newly-infected but are not yet infectious), pre-symptomatic exposed ($E_p(t)$; i.e., newly-infected individuals who start shedding the virus before the end of the incubation period), asymptotically-infectious ($I_a(t)$; i.e., who do not show clinical symptoms of the disease at the end of the incubation period), infectious with mild symptoms ($I_m(t)$), infectious with severe symptoms ($I_s(t)$), infectious in self-isolation ($I_i(t)$), hospitalized or isolated at a health care facility ($I_h(t)$), in intensive care units ($I_c(t)$), recovered but not tested ($R_u(t)$) and tested recovered ($R_t(t)$). Further serologic (antibody) testing of recovered humans is important for determining the level of immunity to COVID-19. Thus,

$$N(t) = S(t) + E(t) + E_p(t) + I_a(t) + I_m(t) + I_s(t) + I_h(t) + I_c(t) + R_u(t) + R_t(t).$$

The model is given by the following deterministic system of nonlinear differential equations (where a dot represents differentiation with respect to time t):

$$\begin{aligned} \dot{S} &= -\lambda S, \\ \dot{E} &= \lambda S - (\sigma_e + \tau_d)E, \\ \dot{E}_p &= \sigma_e E - (\sigma_p + \tau_d)E_p, \\ \dot{I}_a &= r\sigma_p E_p - (\gamma_a + \tau_d)I_a, \\ \dot{I}_m &= (1-r)g\sigma_p E_p - \rho_m I_m, \\ \dot{I}_s &= (1-r)(1-g)\sigma_p E_p - (\delta_s + \gamma_s + \rho_s)I_s, \\ \dot{I}_i &= \tau_d(E + E_p + I_a) + \rho_m I_m + (1-f)\rho_s I_s - (\gamma_i + \xi_i + \delta_i)I_i, \\ \dot{I}_h &= f\rho_s I_s + \xi_i I_i - (\gamma_h + \psi_h + \delta_h)I_h, \\ \dot{I}_c &= \psi_h I_h - (\gamma_c + \delta_c)I_c, \\ \dot{R}_u &= \gamma_a I_a + \gamma_s I_s + \gamma_i I_i - \tau_s R_u, \\ \dot{R}_t &= \gamma_h I_h + \gamma_c I_c + \tau_s R_u, \end{aligned} \tag{2.1}$$

where the *force of infection* λ is defined as:

$$\lambda = (1 - \varepsilon_m c_m) \frac{\beta_p E_p + \beta_a I_a + \beta_m I_m + \beta_s I_s}{N - \theta(I_i + I_h + I_c)}. \tag{2.2}$$

In the model (2.1), susceptible individuals acquire COVID-19 following effective contact with individuals in the pre-symptomatic (E_p), asymptotically-infectious (I_a), mildly-symptomatic (I_m) and severely-symptomatic (I_s) classes, at a rate λ (defined in (2.2) below). Newly-infected individuals (in the E class) progress to the pre-symptomatic class at a rate σ_e . At the end of the incubation period, a proportion, $0 < r \leq 1$, of humans in the E_p class progress to the asymptotically-infectious class (I_a), at a rate $r\sigma_p$ (where $0 < r < 1$ is the proportion of

pre-symptomatic humans who do not show clinical symptoms of COVID-19 at the end of the incubation period). The remaining proportion, $1 - r$, who show clinical symptoms of COVID-19 at the end of the incubation period, are split into those who show mild symptoms (at a rate $g(1 - r)$, where $0 < g \leq 1$ is the proportion of the fraction), $1 - r$, that shows mild symptoms of COVID-19) and those who show severe symptoms (at a rate $(1 - g)(1 - r)$).

Asymptomatic individuals in the E , E_p and I_a classes are detected, *via* random diagnostic/surveillance testing, at a rate τ_d . Similarly, serology testing is administered to untested recovered individuals (i.e., those in R_u class) at a rate τ_s . Recovery in I_k ($k = a, i, c$) class occurs at a rate γ_k . Individuals with mild and severe symptoms are self-isolated at a rate ρ_m and $(1 - f)\rho_s$ (where $0 < f \leq 1$ is the proportion of individuals with severe symptoms that are hospitalized), respectively. Self-isolated individuals are hospitalized at a rate ξ_i . COVID-induced mortality occurs in I_k ($k = s, i, h, c$) at a rate δ_k . Hospitalized individuals are admitted into ICU at a rate ψ_h .

In (2.2), the term $1 - \varepsilon_m c_m$ represents a measure of the reduction in community contacts due to face mask use in the community. In particular, $0 \leq \varepsilon_m \leq 1$ is face mask efficacy in preventing the spread (i.e., outward efficacy) or acquisition (i.e., inward efficacy) of the disease and $0 \leq c_m \leq 1$ is the community-wide compliance in face mask usage. The parameters β_p , β_a , β_m and β_s represent, respectively, the effective contact rates of infected individuals in the pre-symptomatic (E_p), asymptotically-infectious (I_a), symptomatically-infectious with mild symptoms (I_m) and symptomatically-infectious with severe symptoms (I_s) class. The parameter $0 \leq \theta \leq 1$ is a major of the effectiveness of hospitalization, self-isolation and ICU admission to prevent COVID-19 transmission by individuals in the I_i , I_h and I_c classes.

Diagnostic and serology testing are universally considered to be highly crucial to slowing community transmission of COVID-19. In particular, diagnostic testing allows for the detection, rapid isolation and contact tracing of cases, thereby breaking the chain of community transmission that would have otherwise ensued. In the model (2.1), diagnostic testing is accounted for by way of detecting asymptomatic infected individuals, in the E , E_p and I_a classes (at the rate τ_d). We use the following functional form for τ_d :

$$\tau_d = \tau_{dmax} \left(\frac{T_n}{1 + T_n} \right), \quad (2.3)$$

where τ_{dmax} is the maximum detection rate *via* diagnostic testing and T_n is the average total number of tests administered in the community *per* day. It should be noted that T_n can also be interpreted as a function of the number of available test kits in the community. During the 7-day period from April 30 to May 6, 2020, the average number of tests *per* day for the US was 264, 249. The maximum number of tests for the States of Arizona, Florida and New York, for the same time period, were 3, 275 (with 10.9% of these positive), 12, 223 (with 4.5% of these positive) and 22, 345 (with 13.8% of these positive), respectively [35]. These numbers are below the minimum levels recommended by the World Health Organization (WHO) (10, 743 for Arizona; 23,937 for Florida; 112,802 for New York and 917, 450 for the whole of US). The positive test ratio benchmark recommended by the WHO is 10% or less [35]. Similarly, we use the following definition for τ_s :

$$\tau_s = \tau_{smax} \left(\frac{T_n}{1 + T_n} \right), \quad (2.4)$$

where τ_{smax} is the maximum detection rate for the serology (antibody) testing in the community.

The model (2.1) monitors human population. Hence, all its state variables are non-negative for all time t (further, all parameters of the model are non-negative). For housekeeping purposes, we introduce an equation for the rate of change of the population of COVID-deceased individuals (denoted by $D(t)$), given by:

$$\dot{D} = \delta_s I_s + \delta_i I_i + \delta_h I_h + \delta_c I_c. \quad (2.5)$$

A flow diagram of the model (2.1) is depicted in Figure 1, and the state variables and parameters of the model are described in Tables A.1 and A.2 (in Appendix A), respectively.




Figure 2: Data fitting of the model (2.1), for the pre-lockdown and lockdown periods for the states of Arizona, Florida, New York and the entire US. (a)-(d): cumulative mortality data and model fit for the pre-lockdown period for the state of Arizona, Florida, New York, and the entire US, respectively. (e)-(h): cumulative mortality data and model fit for the lockdown period for the state of Arizona, Florida, New York and the entire US, respectively.

3 Mathematical Analysis

3.1 Computation of Final Epidemic Size and Reproduction Number

The model (2.1) has a continuum (or family) of disease-free equilibria given by:

$$\mathcal{E}_0 : (S^*, E^*, E_p^*, I_a^*, I_m^*, I_s^*, I_i^*, I_h^*, I_c^*, R_u^*, R_t^*) = (S(0), 0, 0, 0, 0, 0, 0, 0, 0, 0, 0),$$

where $S(0)$ is the initial number of susceptible individuals in the community. The asymptotic stability analysis of this family of equilibria will be explored below to gain qualitative insight into the behavior of initial solutions of the model, particularly with respect to the control of the burden of the pandemic. For an epidemic model (i.e., a model without demographic dynamics), such as (2.1), a useful feature to compute is the *final size* of the disease [42–44]. It provides a measure of the number of individuals who remain susceptible at the end of the epidemic. The approach in [42] will be used to compute the final size relation for the model (2.1). To apply this method, it is convenient to let $\mathbf{x} \in \mathbb{R}_+^8$, $y \in \mathbb{R}_+$, and $\mathbf{z} \in \mathbb{R}_+^2$ be vectors representing the compartments of infected, susceptible and recovered individuals in the model (2.1), respectively. That is, $\mathbf{x} = (E, E_p, I_a, I_m, I_s, I_i, I_h, I_c)^T$, $y = S$ and $\mathbf{z} = (R_u, R_t)^T$. Furthermore, suppose, Π is an 8×1 matrix in which the (i, j) entry is the fraction of the j^{th} susceptible compartment that goes into the i^{th} infected class upon becoming infected, and \mathbf{b} is an 8–dimensional row vector of relative horizontal transmissions. It then follows from Arino *et al.* [42] that the model (2.1) reduces to the following three-dimensional system of differential equations:

$$\dot{\mathbf{x}} = \Pi D y \lambda(\mathbf{x}, y, \mathbf{z}) \mathbf{b} \mathbf{x} - V \mathbf{x},$$

$$\dot{y} = -D y \beta(\mathbf{x}, y, \mathbf{z}) \mathbf{b} \mathbf{x},$$

$$\dot{\mathbf{z}} = W \mathbf{x},$$

where λ is the *force of infection* of the model (2.1) defined in (2.2), W is a 2×8 matrix in which the (i, j) entry is the rate at which individuals of the j^{th} infected compartment transition into the recovered (i^{th} \mathbf{z}) compartment upon recovery, and V is the standard M-matrix of the linear transition terms between the infected compartments of

the model (2.1) [42] (see also the application of next generation operator method to analyse the asymptotic stability of disease-free equilibria of disease transmission models [45, 46]). It can be seen that, in the context of model (2.1),

$$D = 1, \mathbf{b} = (0, 1, 1, 1, 1, 0, 0, 0), \Pi = (1, 0, 0, 0, 0, 0, 0, 0)^T,$$

and,

$$V = \begin{pmatrix} A_e & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\sigma_e & A_p & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -r\sigma_p & A_a & 0 & 0 & 0 & 0 & 0 \\ 0 & -(1-r)g\sigma_p & 0 & \rho_m & 0 & 0 & 0 & 0 \\ 0 & -(1-r)(1-g)\sigma_p & 0 & 0 & A_s & 0 & 0 & 0 \\ -\tau_d & -\tau_d & -\tau_d & -\rho_m & -(1-f)\rho_s & A_i & 0 & 0 \\ 0 & 0 & 0 & 0 & -f\rho_s & -\xi_i & A_h & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -\varphi_h & A_c \end{pmatrix}.$$

Let $y(0) = S(0), \mathbf{z}(\mathbf{0}) = (R_u(0), R_t(0))^T$. It then follows from Theorem 2.1 in [42] that the reproduction number of the model (2.1), denoted by \mathcal{R} , is given by:

$$\mathcal{R}_c = \lambda(0, y(0), \mathbf{z}(\mathbf{0}))\mathbf{b}V^{-1}\Pi y(0) = \mathcal{R}_{E_p} + \mathcal{R}_{I_a} + \mathcal{R}_{I_m} + \mathcal{R}_{I_s}, \quad (3.1)$$

where,

$$\mathcal{R}_{E_p} = \frac{(1 - \varepsilon_m c_m)\beta_p \sigma_e}{A_e A_p}, \quad \mathcal{R}_{I_a} = \frac{(1 - \varepsilon_m c_m)\beta_a r \sigma_e \sigma_p}{A_a A_e A_p}, \quad \mathcal{R}_{I_m} = \frac{(1 - \varepsilon_m c_m)\beta_m \sigma_e \sigma_p (1 - r)g}{\rho_m A_e A_p},$$

$$\mathcal{R}_{I_s} = \frac{(1 - \varepsilon_m c_m)\beta_s \sigma_e \sigma_p (1 - r)(1 - g)}{A_e A_p A_s},$$

with,

$$A_e = \sigma_e + \tau_d, \quad A_p = \sigma_p + \tau_d, \quad A_a = \gamma_a + \tau_d, \quad A_s = \delta_s + \gamma_s + \rho_s,$$

$$A_i = \delta_i + \gamma_i + \xi_i, \quad A_h = \delta_h + \gamma_h + \psi_h, \quad A_c = \delta_c + \gamma_c.$$

The reproduction number \mathcal{R}_c is an epidemiological quantity that measures the average number of new COVID-19 cases generated by a typical infected individual introduced into a community where some anti-COVID-19 interventions (such as social-distancing, face mask usage and self-isolation) are implemented. The result below follows from Theorem 2.1 of [42].

Theorem 3.1. *The continuum of disease-free equilibria (\mathcal{E}_0) of the model (2.1) is locally-asymptotically stable whenever $\mathcal{R}_c < 1$.*

The epidemiological implication of Theorem 3.1 is that community transmission of COVID-19 can be effectively suppressed in the community if the control and mitigation strategies implemented can bring the reproduction number (\mathcal{R}_c) to a value less than unity.

Remark 3.1. *For epidemic models (i.e., models with no birth and death demographic processes), such as (2.1), having the associated reproduction (\mathcal{R}_c) reduced to a value less than unity is sufficient, but not necessary, for eliminating the disease. That is, even if \mathcal{R}_c exceeds unity, the disease eventually dies out over time. There are many reasons to explain this. First, for $\mathcal{R}_c > 1$, a sizable number of the population will get infected particularly if no control or mitigation interventions are implemented (or are delayed in being implemented), some of whom would*

survive the infection (i.e., they would recover). Since the model assumes that recovery confers perfect and permanent natural immunity against future infections, this process leads to, or contributes to, building community-wide natural herd immunity (so that many of the remaining susceptible members of the community can be protected). This, together with the implementation of control and mitigation measures, contribute in significantly suppressing community transmission, and the disease eventually dies out (even if $\mathcal{R}_c > 1$).

Finally, it follows from Theorem 5.1 of [42] that the final epidemic size relation for the model (2.1) is given by

$$\ln\left(\frac{S(0)}{S(\infty)}\right) \geq \frac{\mathcal{R}_c}{S(0)} \left(S(0) - S_\infty + E(0) + \frac{A_e E_p(0)}{\sigma_e} \right) + \frac{1 - \varepsilon_m c_m}{S(0)} \left[\frac{\beta_a}{A_a} I_a(0) + \left(\frac{\beta_m}{\rho_m} I_m(0) + \frac{\beta_s}{A_s} I_s(0) \right) \right],$$

where $S(\infty) = \lim_{t \rightarrow \infty} S(t)$. This final size relation can be solved recursively for $S(\infty)$ using, for instance, the Newton-Raphson's method, to obtain the number of susceptible humans who remain uninfected after the epidemic.

3.2 Sensitivity of Reproduction Number With Respect to Case Detection and Face Mask Usage

In this section, the effect of case detection and self-isolation of asymptomatic individuals in the E , E_p and I_a classes (using diagnostic testing) and face mask usage on the reproduction number (\mathcal{R}_c) of the model (2.1) will be assessed, for the states of Arizona, Florida, New York and the entire US. We first simulate the effect of maximum case detection (τ_{dmax}) for two scenarios, where face masks of low quality (such as home-made cloth masks with estimated efficacy of 25%) or of moderate quality (such as medical/surgical masks, with estimated efficacy of 50%) are used in the community. A contour plot (heat map) of \mathcal{R}_c , as a function of percentage increase in the maximum case detection rate (i.e., increase in the baseline value of τ_{dmax}) and face mask compliance (c_m), is depicted in Figure 3. This figure shows that the reproduction number (\mathcal{R}_c) decreases with increasing values of the percentage increase in the maximum case detection rate and face mask compliance, for the states of Arizona (Figure 3 (a) and (e)), Florida (Figure 3 (b) and (f)), New York (Figure 3 (c) and (g)) and the entire US (Figure 3 (d) and (h)), regardless of the type (or efficacy) of the face mask used in the community. In particular, it is evident from the contour plots that the reproduction number can be brought to a value less than unity (thereby resulting in effectively curtailing community transmission of COVID-19) in Arizona, Florida and New York if the baseline maximum case detection rate is increased by at least 10%, the face mask efficacy is 25%, and face mask compliance exceeds 73%, 95%, and 35%, respectively. If the face mask efficacy is increased to 50%, then for the same maximum detection rate of 10%, face mask use compliance 37%, 48%, and 18% is required to reduce the reproduction number below one. Greater reductions in \mathcal{R}_c are recorded if the moderately-effective face mask (Figure 3 (e)-(h)) are used in the community, in comparison to when low effective face masks are used (Figure 3 (a)-(d)). These plots clearly show that widespread (random) testing (which then implies rapid detection, tracing and isolation of confirmed cases), combined with face mask usage, significantly contributes in reducing community transmission (by breaking the chain of transmission) regardless of the efficacy of face masks.

The sensitivity of the reproduction number with respect to the singular effect of face masks is monitored by generating contour plots of \mathcal{R}_c , as a function of face mask efficacy (ε_m) and compliance (c_m). The results obtained, depicted in Figure 3 (i)-(l), for Arizona, Florida, New York and the entire US, show decreasing values of the reproduction number with increasing values of face mask efficacy and compliance, as expected. It is shown that, with the assumed baseline face mask efficacy of 50% (i.e., $\varepsilon_m = 0.5$), a minimum compliance of 53% will be needed to effectively curtail community transmission in the state of Arizona (Figure 3 (i)). The corresponding minimum compliance percentages needed to achieve some control in Florida, New York and the whole of US are, 59%, 38% and 46%, respectively. The contours in Figure 3 (i)-(l) also show that using low effective face masks (e.g., the cloth mask with estimated efficacy of 25%) with high compliance can lead to significant reduction in disease burden. However, using such (cloth) masks will fail to reduce the reproduction number to a value below unity, even if 100% compliance is achieved, in the states of Arizona and Florida. On the other hand, such masks can bring the reproduction number to a value less than unity for the state of New York and the entire nation if the compliance in its usage is at least 76% and 92%, respectively.




Figure 3: **Sensitivity of the control reproduction number (\mathcal{R}_c) to important control parameters.** Heat maps of the control reproduction number, as a function of cloth face mask use compliance and percentage increases in the maximum detection rate of asymptomatic infected individuals (τ_{dmax}). (a)-(d): impact of cloth masks (with efficacy $\varepsilon_m = 0.25$) on disease dynamics for the state of Arizona, Florida, New York and all of US, respectively. (e)-(h): impact of surgical/medical masks (with efficacy $\varepsilon_m = 0.50$) on disease dynamics for the state of Arizona, Florida, New York and all of US, respectively. (h)-(k): impact of face masks efficacy (ε_m) and compliance (c_m) on disease dynamics for the state of Arizona, Florida, New York and all of US, respectively. Parameter values used in the simulations are as given in Tables A.3-A.4 in Appendix A.

4 Numerical Simulations

In this section, the model (2.1) will be simulated, using the baseline parameter values in Tables A.3-A.4 in Appendix A (unless otherwise stated), to assess the community-wide impact of early implementation of strict lockdown measures (aimed at minimizing community transmission of COVID-19) in the states of Arizona, Florida, New York and the entire US. The impact of lifting of community lockdown, as well as the effect of the use of non-pharmaceutical interventions after the lifting of the lockdown, will also be assessed. Parameter values for the pre-lockdown (sub-tables (a) of Tables A.3-A.4 in Appendix A) and the lockdown period (sub-tables (b) of Tables A.3-A.4 in Appendix A) are used to simulate the COVID-19 dynamics in the four jurisdictions during the pre-lockdown and the lockdown periods, respectively. It should be mentioned that, in all of the simulations to be carried out, the control-related parameters (e.g., the face masks compliance and efficacy, c_m and ε_m , and the diagnostic and serology detection rate parameters, τ_d , τ_s and τ_{dmax}) are kept at their baseline values, for both the pre-lockdown and lockdown periods in Tables A.3-A.4 (unless otherwise stated).

4.1 Impact of Early Implementation of Lockdown Measures and Mask use Compliance

It should be recalled that lockdown measures were implemented in the states of Arizona, Florida and New York on March 31, 2020, April 3, 2020 and March 22, 2020, respectively, while the entire US was on lockdown by April 7, 2020. Furthermore, partial lifting of the lockdown measures were announced in the three states by May 15, 2020,

May 4, 2020 and May 28, 2020, respectively. Although some US states delayed announcing some form of lifting of lockdown measures until June 2020, the overwhelming majority of the US states implemented some form of lifting by May 28, 2020. Consequently, we assume that the entire US started partial lifting of lockdown on May 28, 2020, as was the case for the state of New York. We now run simulations of the model (2.1) for the three scenarios: (a) the precise period when community lockdown was actually implemented, (b) lockdown measures were implemented a week earlier than the actual time it was implemented, and (c) lockdown measures were implemented two weeks earlier than the actual time they were implemented, for each of the three states and the entire US.

The simulation results obtained for scenario in which lockdown was implemented one week earlier are depicted in Figure 4, showing daily and cumulative mortality for the three states and the US. This figure shows a marked decrease in the peak daily deaths in each of the four locations (Figures 4 (a)-(d), magenta curves). In other words, the pandemic curves in the three states and in the entire US would have been flattened earlier (characterized by very low numbers at the pandemic peak). The decrease in daily mortality is even more dramatic if lockdown was implemented two weeks earlier (Figures 4 (a)-(d), green curves). Furthermore, our simulations show that if the lockdown measures were implemented a week earlier in Arizona, up to 75% of the cumulative number of COVID-19 deaths in the state could have been averted (Figure 4 (e), magenta curve). Similarly, 76% of the cumulative deaths in the state of Florida (Figure 4 (f), magenta curve) and 99% of the cumulative mortality in the state of New York (Figure 4 (f), magenta curve) would have been prevented. At least 77% of the cumulative deaths recorded in the entire US would have been averted if lockdown measures were implemented a week earlier (Figure 4 (h), magenta curve). More dramatic reductions in cumulative mortality would have been recorded if the lockdown measures were implemented two weeks earlier (Figure 4 (e)-(h), green curves).




Figure 4: Impact of early lockdown: Simulations of the model (2.1), showing the daily and cumulative mortality, as a function of time, for various durations of the onset of lockdown measures for the states of Arizona, Florida, New York and the whole of US. The model is ran from the date of the index case, for each of the four jurisdictions, and extended one month beyond the first day of the relaxation (or partial lifting) of the lockdown measures. (a)-(d): daily deaths for the state of Arizona, Florida, New York and all of US, respectively. (e)-(h): cumulative deaths for the state of Arizona, Florida, New York and all of US, respectively. The predictions of the model, for the case when community lockdown measures were implemented one week earlier, two weeks earlier, or on the actual date the lockdown measures were implemented, are represented by magenta, green, and blue curves, respectively. Red dots represent the actual observed data, while blue dashed vertical lines depict the start and end of the actual lockdown period (shaded in cyan). Parameter used in the simulations are given in Tables A.3-A.4 in Appendix A, with various start dates for the implementation of community lockdown measures.

Simulations were also carried out to assess the community-wide impact of increased face mask usage during the lockdown period in each of the four jurisdictions considered in this study. For these simulations, four levels of

face mask compliance (in relation to the baseline value of the face mask compliance during lockdown, tabulated in Tables A.3-A.4 in Appendix A), namely 25%, 50%, 75% and 100% compliance, are considered. The results obtained, depicted in Figure 5, show a marked decrease in cumulative mortality with increasing compliance level of mask usage in each of the four jurisdictions. In particular, it can be seen that, if half of the residents of the state of Arizona consistently wear face mask in public during the lockdown period, up to 47% of the cumulative mortality recorded by the day of the partial lifting of the lockdown measures in the state (May 14, 2020) would have been averted (Figure 5 (a)). Similarly, up to 30%, 50% and 26% of the cumulative deaths recorded by the day of the lifting of the lockdown in the states of Florida (Figure 5 (b)), New York (Figure 5 (c)) and in the entire US (Figure 5 (d)) would have been averted if half their residents wear face masks consistently during their respective lockdown periods.

A dramatic increase in the cumulative mortality averted in the four jurisdictions is recorded if 75% of the residents of the four jurisdictions wear face masks during the lockdown period. Specifically, while up to 60% and 42% of mortality recorded by the day of lifting of lockdown measures in Arizona and Florida would be averted, the state of New York and the entire US would prevent up to 65% and 37% of the deaths they recorded by the day of the lifting (if 75% of their respective residents wear face masks during the lockdown). If everyone in the four jurisdictions wear face mask during the lockdown, the percentage reduction of the cumulative mortality on the day of lifting of lockdown in the states of Arizona, Florida, New York and the whole of US further increase to 66%, 50%, 74% and 44%, respectively (this corresponds to 468, 714, 21, 928, and 44, 380 deaths being avoided in the respective jurisdictions, under this 100% face mask compliance scenario). The actual cumulative mortality recorded in the four jurisdictions by their respective day of the partial lifting of the lockdown (based on using the baseline values of the face mask compliance, given in Tables A.3-A.4 in Appendix A), together with the cumulative mortality that would have been recorded on the day of the lifting for various face mask compliance levels, are tabulated in Table 4.1.

In summary, it can be concluded from Figures 4 and 5 that early implementation of lockdown (i.e., if they were implemented a week or two earlier than the day they were implemented), combined with increased level of face masks use during the lockdown periods, will result in a dramatic reduction in the burden of the pandemic in each of the four jurisdictions considered in this study. In fact, these figures show that it is quite possible that the pandemic might not have even (significantly) taken off in any of the four jurisdictions if the lockdown measures were implemented two weeks earlier, and if most of the residents of the four jurisdictions wore face masks consistently in public.

Table 4.1: Actual cumulative mortality on the day lockdown was lifted (Row 2), and cumulative mortality as a function of mask use compliance that would have been recorded on the day lockdown was lifted? for the states of Arizona, Florida, New York, and entire US. Baseline mask compliance corresponds to the estimated mask-use compliance during the lockdown period in Tables A.3-A.4 in Appendix A.

Mask compliance level	Arizona	Florida	New York	USA
Baseline mask compliance during lockdown	651	1, 399	30, 140	105, 896
25% mask compliance	558	1, 282	23, 540	93, 410
50% mask compliance	374	996	1, 4780	74, 030
75% mask compliance	285	819	10, 320	62, 540
100% mask compliance	236	705	7, 682	55, 300




Figure 5: **Impact of face masks compliance during lockdown:** Simulations of the model (2.1), showing cumulative mortality, as a function of time, for the US state of Arizona, Florida, New York and the entire US. The model is simulated from the date of the index case for each of the four jurisdictions, for different levels of face masks compliance (c_m). (a)-(d): cumulative deaths for the states of Arizona, Florida, New York and all of US, respectively. The blue curve represents the case for the baseline value of the face masks compliance, and the red dots represent the actual data. The blue dashed vertical lines represent the onset and termination dates for the lockdown measures (this region is shaded in cyan color). Parameter values used in the simulations are as given in Tables A.3-A.4 in Appendix A, with various levels of face masks compliance (c_m).

4.2 Impact of Varying Levels of Lifting of Community Lockdown Measures

The model (2.1) is now simulated, using the parameter values in Tables A.3-A.4 in Appendix A, to assess the community-wide impact of varying levels of lifting of the community lockdown measures implemented in the states of Arizona, Florida, New York and the entire US. Various states and jurisdictions within the US (with the exception of the states of New York, New Jersey, New Hampshire, District of Columbia, Illinois, and Delaware) implemented various levels of lifting of the lockdown measures (by May 20, 2020) [27], in an effort to re-open socio-economic activities. We consider three levels of community lockdown lifting, namely mild, moderate and high, as described below (adapted from [47]):

Mild re-opening: The main components of mild lifting or re-opening of community lockdown include lifting of stay-at-home (lockdown) measures (except for vulnerable individuals, who should shelter-at-home) and maintaining six-foot social-distancing in public places (e.g., parks, outdoor recreation areas, shopping areas, etc). For this re-opening level, no socializing in groups with more than 10 people and non-essential travels are allowed. Workers work from home (telework) and nonessential businesses (such as construction sites, manufacturers, nonessential retail stores for delivery, curbside and in-store pickup, wholesalers, outdoor business, such as drive-in movies, landscaping and gardening) are allowed to reopen. Furthermore, restaurants and bars are allowed to offer take-out, while schools and organized youth activities (e.g., daycare, camp) remain closed. Visitation to senior care facilities and hospitals are prohibited.

Moderate re-opening: Under this moderate level of community re-opening, indoor businesses (such as indoor dining at restaurants, hair salons and barber shops, but with waiting areas closed, offices, etc.) are allowed to reopen (but with limits on capacity, strict cleaning requirements and mandatory social distancing). Furthermore, real estate firms, in-store retails (excluding Malls, but individual stores can provide curbside pickup), vehicle sales, leases and rentals, commercial building management, etc., are allowed to reopen. Nonessential travels and gatherings of up to 25 people are allowed. The use of face coverings in public is encouraged.

High re-opening: For high level of lifting of community lockdown, gatherings of up to 50 people are allowed. Additionally, nonessential businesses (such as indoor dining at restaurants with up to 50% capacity and tables spaced at least six feet away from each other; seating in bar areas allowed, but only if six feet of distance can be maintained between parties) are allowed to re-open. Furthermore, personal care services, such as nail salons,

massage parlors, spa services and tanning salons can reopen. Some “low-risk” youth sports are allowed. Similarly, indoor religious gatherings can operate at 33% of maximum capacity. Nonessential businesses, such as, low-risk outdoor arts and entertainment activities, including zoos, botanical gardens, historical sites, outdoor museums and parks, low-risk indoor arts and entertainment activities, including museums, aquariums and historical sites; film, movie and music production; higher education institutions; professional sports, without fans in the stands, are allowed. Places that often draw large crowds (e.g., movie theaters, gyms and fitness centers, shopping malls, etc.) are allowed to reopen.

We model lifting of community lockdown based on increases in community contacts, as measured in terms of increases in the baseline values of the community contact rate parameters, β_p , β_a , β_i and β_s (tabulated in Tables A.4-A.7 (b)). In particular, we assume that mild lifting of lockdown measures corresponds to a 5% increase in the baseline values of the contact rate parameters. Similarly, we assume that moderate and high lifting correspond to a 10% and 20% increase in the baseline values of these parameters, respectively. While numerous US states adopted mild or moderate levels of reopening, none, to the authors’ knowledge, adopted the high reopening level. In other words, high reopening represents a worst-case scenario that is not, at the current moment, realistically plausible. We do not expect any US state to adopt the high reopening level (until community transmission of the pandemic is greatly curtailed and/or a safe and effective vaccine or antiviral is available).

It should further be stated that, for the simulations to be carried out in this section, all control-related parameters of the model (e.g., parameters related to the use of face masks in public (c_m and ε_m and case detection (τ_d , τ_s and τ_{dmax} etc.) are kept at their baseline values. That is, for the simulations in this section, no additional improvements in the baseline values of the control-related parameters of the model will be allowed (i.e., all control measures are implemented at their baseline levels given in Tables A.4-A.7).

The simulation results obtained, depicted in Figure 6, show an increase in both the daily and cumulative COVID-19-induced mortality with increasing lifting levels of the community lockdown. This is expected, since increasing the level of lifting of the community lockdown measures implies increased community contacts, thereby resulting in increased number of COVID-19 infections, hospitalizations and deaths. For the worst-case scenario, where the lifting level is high (i.e., the baseline values of the community contact rate parameters are increased by 20%), the states of Arizona, Florida and the entire US will experience a devastating second wave of the pandemic, occurring five to ten months after the lockdown measures were lifted, while the state of New York will only experience a mild second wave (Figure 6, red curves). In particular, under this high lifting scenario (and with control measures at their baseline values), the second wave for the states of Arizona and Florida will peak in mid and late October, 2020, respectively, (with about 1, 148 and 2, 733 COVID-19 related deaths in Arizona and Florida, respectively on the day the pandemic peaks (Figures 6 (a)-(b), red curves)). Similarly, the second wave for the entire US will peak early in February 2021 (recording about 2, 788 deaths on the day of the second wave peak), and no pandemic peak will be experienced at the state of New York (Figures 6 (c)-(d), red curves).

On the other hand, if the level of lifting of the community lockdown was moderate, our results show a marked decrease in the size of the pandemic peaks, as well as a shift in the timing of the peaks (Figures 6 (a)-(d), magenta curves). Specifically, for this scenario, the peak of the second wave for Arizona and Florida will occur in mid to late April and February, 2021, respectively, while the US state of New York and the entire US do not experience a second wave. Furthermore, the moderate lifting scenario will avert over 92% and 84% of the deaths that would have occurred at the peak of the pandemic in Arizona and Florida, respectively, if high level of lifting of the control measures is implemented (i.e., compare red and magenta curves in Figures 6 (a)-(d)).

If mild lifting of the community lockdown was implemented, our simulations show that while the states of Florida, New York and the whole of US will not experience a second wave, Florida will experience a relatively mild second wave that will peak in late November, 2020 (Figures 6 (a)-(d), green curves). In this (mild lifting) scenario, Florida will record about 46 deaths at the peak of the second wave. This represents a 98% reduction of the mortality that would have been recorded at the peak of the second wave in Florida if high lifting of lockdown measures was implemented. Furthermore, for this mild lifting of lockdown scenario, the states of Arizona, Florida, New York, and the entire US will record about 2, 274, 12, 06032, 380, and 155, 900 cumulative deaths, respectively, by the end of December 2020. These cumulative death numbers represent increases of about 438, 4, 252, 30, and 5, 300, respectively, in the cumulative mortality that would have been recorded by the end of September 2020. Additionally,

the projected cumulative mortality numbers by the end of the year represent increases of about 1,623, 10,661, 2,240, and 50,004, respectively, from the actual cumulative mortality numbers when each of the four jurisdiction started re-opening (Figures 6 (e)-(h)). Table 4.2 summarizes the cumulative COVID-19 related deaths that would have been averted in the states of Arizona, Florida, New York, and the entire US, by the end of September, October, November, and December of 2020, if the mild level of lifting of community lockdown was implemented in the four locations, in comparison to implementing the moderate level of lifting of community lockdown (showing greater reductions in the states of Arizona and Florida, in comparison to the mild or moderate reductions in the state of New York and the whole of US).

It is worth emphasizing that the size of the second wave depends on the level of lifting of the lockdown measures implemented and the location. For instance, while the state of Florida will always record a second wave, regardless of the level of the lifting of the community lockdown measures (mild, moderate, or high), the states of Arizona and New York, as well as the entire US, will not record a (significant) second wave if the level of lifting of the community lockdown was mild. In all four locations, the size (or public health burden of the second wave) decreases with decreasing level of the lifting of the community lockdown measures. Further, moderate and mild lifting not only decreases the peak daily mortality, they shift the time of the second wave. It is also notable from Figure 6 that the second wave in Arizona and Florida is far more severe than the first wave, regardless of the level of lifting of the community lockdown implemented in the two states.




Figure 6: **Impact of re-opening of community lockdown.** Simulations of the model (2.1), showing daily and cumulative mortality, as a function of time, for various levels of re-opening of community lockdown measures in the states of Arizona, Florida, New York and the entire US. Mild, moderate, and high re-opening of community lockdown corresponds to a 5%, 10%, and 20% increase in the baseline value of the community contact rate parameters (β_p , β_a , β_i , and β_s), respectively. (a)-(d): Daily deaths for the state of Arizona, Florida, New York and all of US, respectively. (e)-(h): Cumulative deaths for the state of Arizona, Florida, New York and all of US, respectively. Other parameter values used in the simulations are as given in Tables A.3-A.4 in Appendix A.

Table 4.2: Percentage reduction in cumulative COVID-19 mortality averted in the states of Arizona, Florida, New York and the entire US (by the end of September, October, November, and December of 2020) if mild levels of lifting of community lockdown was implemented, in comparison to the case where moderate level of lifting of the community lockdown was implemented. Mild and moderate lifting of community lockdown correspond, respectively, to a 5% and 10% increase in the baseline values of the community contact rate parameters (β_p , β_a , β_i and β_s).

Location	September 30, 2020	October 31, 2020	November 30, 2020	December 31, 2020
Arizona	48 %	58 %	66 %	73 %
Florida	52 %	60 %	67 %	72 %
New York	2.3 %	2.5 %	2.6 %	2.7 %
US	10 %	13 %	15 %	17 %

4.3 Effect of Mask Usage After High Lifting of Community Lockdown Measures

In this section, the model (2.1) is simulated, using the parameter values in Tables A.3-A.4 in Appendix A, to assess the community-wide impact of mask usage during the post-lockdown period in the states of Arizona, Florida, New York and the whole of US. For these simulations, we consider high level of lifting of the community lockdown measures in all four jurisdictions (i.e., we consider the scenario where the baseline values of the community contact rate parameters, β_p , β_a , β_i and β_s , are increased by 20% in all four locations) and various values of face mask compliance (c_m). Note that the baseline efficacy of the face mask to protect the wearer from acquiring infection is assumed to be 50% (i.e., $\varepsilon_m = 0.5$). The simulation results obtained are depicted in Figure 7, from which it follows that the cumulative number of cases, hospitalizations and COVID-induced mortality decrease with increasing face mask compliance. It was observed that if two in every five people (i.e., 40%) in each of the jurisdictions considered in this study wear face masks in public after the lockdown measures were lifted, only the state of Florida will experience a (very mild) second wave of the pandemic. In this scenario, the states of Arizona and New York, as well as the entire US, will not have a second wave of the pandemic. Furthermore, no second wave of the pandemic will be experienced in any of the four jurisdictions if half of the residents wear face masks after the lockdown (this is obtained from plotting the corresponding daily mortality; the plot is not included to save space).

If half of Arizonans wear face masks after the lockdown period, under this high lifting scenario, Figure 7 (a) shows that the state will record a cumulative mortality of 1, 180 by the end of December 2020 (and this corresponds to a 98% reduction in the cumulative mortality that would have been recorded in the state if the face mask compliance was at the estimated value of 14% during the lockdown period, tabulated in Table A.4)). Similarly, the states of Florida, New York, and the entire US will record 4, 465, 31, 770, and 138, 800 cumulative deaths, respectively (over the same time period), if half their populations wear face masks in public after the lockdown period. This represents a reduction of 96%, 26% and 67% in the cumulative mortality, respectively, that would have been recorded if the respective estimated face mask compliance during the lockdown period was used (Figures 7 (b)-(d)). These numbers decrease dramatically if 75% of the residents of the four jurisdictions wear face masks immediately after the lifting of the lockdown measures. For example, the states of Arizona, Florida, New York and the whole US will record cumulative mortality of 951, 2, 516, 31, 270 and 124, 200, respectively (over the same time period), if 75% of their respective residents wear face masks after the lockdown period (representing a 99%, 99%, 28% and 74% reduction in the respective cumulative mortality in comparison to when the estimated lockdown face mask compliance value is used). Higher reductions are observed when everyone wears face mask after the lockdown (*albeit* the reductions are far more pronounced in the states of Arizona and Florida, than in the state of New York and the entire US). Figure 7 shows similar results for the reductions in the cumulative hospitalizations (Figures 7 (e)-(h)) and the cumulative new COVID cases (Figures 7 (i)-(l)).




Figure 7: **Impact of face mask usage on COVID-19 burden after mhigh lifting of community lockdown.** Simulations of the model (2.1), for high lifting of community lockdown in the states of Arizona, Florida, New York and the entire US, and various levels of face mask compliance (c_m). (a)-(d): Cumulative mortality for the state of Arizona, Florida, New York and all of US, respectively. (e)-(h): Cumulative hospitalizations for the state of Arizona, Florida, New York and all of US, respectively. (i)-(l): Cumulative cases for the states of Arizona, Florida, New York and all of US, respectively. Parameter values used are as given in Tables A.3-A.4 in Appendix A, and various levels of face mask compliance (c_m).

4.4 Effect of Mask Usage After High Lifting of Community Lockdown Measures

When high level of lifting is implemented (i.e., the associated community contact rate parameters (β_p , β_a , β_i and β_s) are increased by over 20% from their baseline values), our simulations show that a second wave in all four jurisdictions would be inevitable, depending on the level of the increase of the high lifting scenario. We consider a hypothetical scenario where the level of lifting of the community lockdown measures is further increased, such as to a level that entails increasing the baseline values of the community contact rate parameters (β_p , β_a , β_i and β_s) by 50%. This higher level may be akin to essentially returning to business as usual, where no significant restrictions are imposed (except, perhaps, no large gatherings such as major sporting, social and political campaign events). The simulation results obtained, for this hypothetical scenario in the four jurisdictions, are depicted in Figure 8. This figure shows a dramatic decrease in daily COVID-induced mortality with increasing face mask compliance. For this higher community lockdown lifting scenario, the states of Arizona and Florida will experience a major second wave peaking on October 19, 2020 (with 1,294 deaths on this day) and October 3, 2020 (with 3,599 deaths on this day), respectively, even if half their residents wear face masks after the lockdown has been lifted (Figures 8 (a)-(b), red curves). The state of New York and the entire US will have milder second waves peaking on March 24, 2021 (with 123 deaths at the peak) and January 13, 2021 (with 5,896 deaths at the peak), respectively. Furthermore, no second wave will occur in any of the four jurisdictions if 75% of their respective residents wear face masks after the lockdown period (Figures 8 (a)-(d), purple curves). In fact, under this very high lifting scenario, all four jurisdictions will have very mild or no significant outbreaks of COVID-19 if the face masks compliance is at least

75%. In other words, high masks compliance after community lockdown lifting (even if the level of lifting was very high) greatly reduces the community transmission of COVID-19 in all four jurisdictions (resulting in very mild or no outbreaks in all four jurisdictions). If everyone in the jurisdictions wear face masks after the lockdown (i.e., $c_m = 1$), our simulations show that, for this very high level of lifting of community lockdown (with baseline community contact rates increased by 50%), the pandemic will be effectively curtailed (as measured in terms of major suppression of community transmission) in all four locations within two to three months after the community lockdown measures have been lifted.




Figure 8: **Impact of higher lifting of community lockdown measures** (as measured in terms of 50% increase in the baseline values of the community contact rate parameters, β_e , β_i , β_a and β_s) and various compliance levels of face mask usage in the four jurisdictions. Simulations for the model (2.1) showing the daily mortality, as a function of time. (a)-(d): daily deaths for the state of Arizona, Florida, New York and all of US, respectively. Parameter values used are as given in Tables A.3-A.4 in Appendix A, with various levels of face masks compliance (c_m).

4.5 Impact of Case Detection and High Lifting of Community Lockdown Measures

In this section, the model (2.1) is simulated to assess the impact of detection of exposed (E), pre-symptomatic (E_p), and asymptotically-infectious (I_a) individuals, *via* the implementation of a COVID-19 diagnostic/surveillance testing strategy (as measured in terms of increase in the baseline values of the maximum detection rate parameter, τ_{dmax}), and high level of lifting of community lockdown measures, on the burden of COVID-19 in the four jurisdictions considered in this study. The results obtained, depicted in Figure 9, show a dramatic decrease in the daily and cumulative mortality with increasing values of the maximum detection rate. This figure shows that, while Arizona, Florida, and the entire US will suffer a major second wave if the baseline value of the maximum detection rate (τ_{dmax}) is used (as depicted in Figure 6), much milder second waves will be recorded in the states of Arizona and Florida if the baseline value of the maximum detection rate is increased by 10% (Figures 9 (a)-(d), red curves). It should be mentioned that the milder second waves will still be larger than the first waves recorded in the two jurisdictions. On the other hand, it can be seen from this figure that (for this scenario) the state of New York and the entire US will not suffer a second wave of the pandemic. If the baseline value of the maximum detection rate can be increased by 15%, our simulations show that none of the four jurisdictions will suffer a second wave. However, the decline in the daily deaths for the states of Arizona and Florida is slower than that for the state of New York and the entire US (Figures 9 (a)-(d), magenta curves). If the baseline value of the maximum detection rate is increased by 20% from its baseline value, our simulations show that none of the four jurisdictions will experience a second wave (Figures 9 (a)-(d), green curves).

Furthermore, a significant reduction in the cumulative mortality is achieved in all four jurisdictions, with increasing levels of the baseline maximum detection rate (Figures 9 (e)-(h)). For example, if the maximum detection rate is increased by 10% from its baseline value (noting that, for these simulations, high level of lifting of lockdown is used), our simulations for the state of Arizona show that up to 96% of cumulative mortality that would have been recorded by the end of December 2020, under the high lifting scenario with baseline value of maximum detection rate (depicted in Figure 6) will be averted in the state if τ_{dmax} is increased by 10% from its baseline value (i.e., compare Figures Figure 6 and Figure 9). Similarly, the states of Florida, New York and the entire US will avert 91%, 25%, and 60% of the respective cumulative deaths that would have been recorded by the end of December, 2020, under this scenario. Our simulations show even more dramatic decrease in the cumulative number of deaths

if the baseline maximum detection rate is increased by 20% (Figure 9, green curves). It can be concluded from Figure 9 that widespread (random) diagnostic testing (i.e., the detection, tracing and rapid self-isolation of infected individuals with no symptoms of COVID-19) plays a major role in curtailing community transmission in each of the four jurisdictions. Consequently, widespread testing greatly contributes in suppressing community transmission of COVID-19. In particular, ramping up testing to a level that increases the baseline value of the maximum detection rate (τ_{dmax}), by as low as 20%, could greatly diminish community transmission, to the extent that Arizona, Florida, New York state, and the entire US may not experience a second wave. In summary, widespread random testing contributes in detecting, tracing and isolating asymptomatic cases (hence, breaking their transmission chains) that would otherwise be spreading the virus in the community. More testing clearly does not mean more new cases. More testing means more detection (and rapid isolation) of asymptomatic cases, thereby reducing community transmission. It should be emphasized that face masks compliance is maintained at the baseline value depicted in Tables A.3-A.4.




Figure 9: **Impact of percentage increase in maximum detection rate of asymptomatic infected individuals**, for the case where high level of lifting of community lockdown (i.e., 20% increase in the community contact rates) was implemented in each of the four jurisdictions. Simulations of the model (2.1), showing daily and cumulative mortality, as a function of time. (a)-(d): daily deaths for the state of Arizona, Florida, New York and all of US, respectively. (e)-(h): cumulative deaths for the state of Arizona, Florida, New York and all of US, respectively. Parameter values used are as given in Tables A.3-A.4 in Appendix A, with various percentage increase in the value of τ_{dmax} .

Additional simulations were carried out to assess the combined impact of percentage increases in the maximum case detection rate (τ_{dmax}) of asymptomatic infected individuals (i.e., infected individuals in the E , E_p and I_a classes) and increased mask use compliance (c_m), from their baseline values, on the public health burden of the pandemic in the four jurisdictions. The simulations were carried out for the case where a high level of lifting of the community lockdown (i.e., 20% increase in the baseline values of the community contact rate parameters) was implemented in each of the four jurisdictions. The results obtained are depicted in Figure 10, from which it follows that the daily and cumulative mortality significantly decreases with increasing levels of the maximum case detection rate (from its baseline value). For instance, even a 10% increase in the baseline value of the maximum case detection rate could lead to a sizable reduction in the cumulative mortality in all four jurisdictions. Furthermore, a 20% increase in the baseline level of the maximum case detection rate can dramatically decrease the burden of the pandemic in each of the four jurisdictions even if combined with low level of face mask compliance, such as 25% face mask compliance (Figure 10, green curves). In fact, this combination (of 20% increase in baseline value of τ_{dmax} and 25% face mask compliance) can avert a second wave in each of the four jurisdictions. For this particular case (with 20% increase in baseline value of τ_{dmax} and 25% mask compliance), the states of Arizona, Florida, New York

and the whole of US will record, respectively, about 1, 075, 3, 394, 31, 620, and 138, 000 cumulative deaths by the first week of December 2020. This corresponds to 99%, 99%, 28% and 71% decrease in the cumulative deaths if the maximum case detection rate (τ_{dmax}) was combined only with baseline face mask compliance (i.e., if only baseline face mask compliance is used). Larger increases in the maximum case detection rate and increased face mask compliance will lead to even more significant decrease in the burden of the pandemic in all four jurisdictions.




Figure 10: **Impact of increase in maximum detection rate of asymptomatic infected individuals (τ_{dmax}) and face masks use.** Simulations of the model (2.1), showing daily and cumulative mortality, as a function of time, for high level of lockdown lifting in the states of Arizona, Florida, New York and the entire US. (a)-(d): daily deaths for the state of Arizona, Florida, New York and all of US, respectively. (e)-(h): cumulative deaths for the state of Arizona, Florida, New York and all of US, respectively. Parameter values used are as given in Tables A.3-A.4 in Appendix A, with various values of percentage increases in τ_{dmax} and face masks compliance fixed at 25% (i.e., $c_m = 0.25$).

Discussion and Conclusions

In this study, we developed a new Kermack-Mckendrick epidemic model (i.e., a disease transmission model with no demographic processes) for the transmission dynamics and control of COVID-19 in the states of Arizona, Florida, New York, and the entire US. Some of the notable features of the compartmental model, which takes the form of a deterministic system of nonlinear differential equations, include accounting for the dynamics of pre-symptomatic and asymptotically-infectious individuals (who contribute to disease transmission) and allowing for the assessment of the community-wide impact of some non-pharmaceutical interventions, particularly the use of face masks in public. The model, which was parameterized using cumulative mortality data from the aforementioned four jurisdictions, was used to address the important question of whether or not the widespread use of face masks could halt the post-lockdown resurgence of COVID-19 in the US, without having to undergo another cycle of community lockdowns. Specifically, the model was used to assess the community-wide impact of early lockdown, various levels of lockdown lifting, case detection of asymptomatic individuals, and the use of face masks on the dynamics and control of COVID-19 in each of the four jurisdictions considered in this study.

Rigorous qualitative analysis of the model reveal that it has a continuum of disease-free equilibria, which is asymptotically-stable whenever a certain epidemiological threshold, known as the *control reproduction number* is less than unity. The epidemiological implication of this result, which represents a sufficient condition for the effective control of the disease, is that community transmission of COVID-19 can be significantly suppressed in the four jurisdictions if the control and mitigation interventions can bring, and maintain, the reproduction threshold to a value less than unity. A relation for the final size of the pandemic was also derived analytically. We explored

the sensitivity of the reproduction number with respect to face mask compliance in the four jurisdictions. In particular, we showed that community transmission of COVID-19 can be significantly reduced, using a face mask with efficacy of 50%, if at least 53% of Arizonans wear a face mask (from the beginning of the index case in Arizona). Similar face mask compliance for the states of Florida (59%), New York (38%) and the entire US (46%) were also obtained. In other words, our study shows that COVID-19 could have been effectively controlled if a public face mask strategy (particularly using surgical/medical masks, with protective efficacy of at least 50%) is implemented in each of the four locations, and the aforementioned minimum compliance attained (and maintained) from the beginning of the pandemic. This result is consistent with what actually happened in other parts of the world (particularly in some Asian countries, such as China, Japan, Singapore and South Korea), where the universal use of face masks greatly curtailed community transmission of COVID-19 (and brought the pandemic under very effective control, essentially at scales that can be, for all intents and purposes, considered as pandemic elimination).

Community lockdown was one of the major non-pharmaceutical intervention implemented in the US (and around the world), in an effort to curtail the community spread of COVID-19. Many states of the US were in community lockdown generally from mid March to the end of May, 2020 (we consider, for simulation purposes, that the entire US was (generally) in community lockdown between April 7, 2020 to May 28, 2020). In particular, the state of New York, once the global epicenter of COVID-19, was among the first of the US states to implement a community lockdown (on March 22, 2020) and among the last to begin to partially-reopen (on May 28, 2020). As a consequence, the state of New York is currently recording very low COVID-19 case numbers and one of the lowest mortality numbers in the US [48]. On the other hand, the states of Arizona and Florida that were among the last to lockdown and among the first to partially lift the lockdown measures, are now experiencing an alarming resurgence of COVID-19 (with record case and hospitalization numbers [4, 5, 49]). It should be emphasized that, although the states of Arizona and Florida succeeded in bending their epidemic curves during the lockdown period, they were not in community lockdown long enough to achieve the two weeks of continuous decline in confirmed cases stipulated in the re-opening guidelines by the White House Coronavirus Task Force [28]. Our study highlights the importance of early implementation of control measures (community lockdown in this case) and sustaining these measures until a time is reached when it is safe to (responsibly) begin to relax these measures (i.e., begin to partially lift the lockdown measures). Our simulations showed that if the lockdown measures were implemented a week earlier (than the actual days they were implemented in the four locations), all four jurisdictions would have recorded very significant reductions in their respective daily and cumulative mortality numbers by the day the lockdown measures were partially lifted. For example, our simulation results showed that about 77% of the cumulative mortality number for the US (as of the day lockdown measures were partially lifted) would have been averted.

Our simulation results also highlight the importance of combining early implementation of the lockdown measures with increased face mask usage during the community lockdown. In particular, we showed that combining the early implementation of lockdown (i.e., implementing the lockdown measures 1-2 weeks earlier) with a public face mask strategy during the lockdown (and with reasonably high face mask compliance level) resulted in dramatic reductions in disease burden in each of the four jurisdictions. In fact, our simulations showed that the pandemic might not have even taken off significantly in any of the the four jurisdictions if the community lockdown measures were implemented two weeks earlier and most people in the jurisdictions wear face masks during the lockdown.

The lifting of community lockdown in many communities in the US was done in multiple phases, and at varying levels. For this reason, we used our model to assess the community-wide impact of the varying levels of lifting of community lockdown implemented in the four jurisdictions. We considered three levels of lifting of community lockdown, namely mild, moderate and high (based on the level of community interactions allowed. For example, based on whether or not restaurants, gyms, salons, malls, etc., are opened or closed). Mechanistically, the heterogeneity in the lifting levels is incorporated into our model by increasing the baseline values of the community contact rate parameters ($\beta_p, \beta_a, \beta_i$ and β_s). Our results showed an increase in daily and cumulative mortality in each of the four jurisdictions with increasing lifting levels of the community lockdown implemented in the respective jurisdictions. Furthermore, we showed that if a high level of lifting of the community lockdown is implemented (and all control measures are fixed at their baseline levels), the states of Arizona and Florida will have a devastating second wave of the COVID-19 pandemic that would occur about five months after their respective lifting of the

community lockdown, while the state of New York and the entire US will record much milder second waves. However, if the level of lifting was moderate, our results showed a sizable decrease in the COVID-19 burden in all four jurisdictions. If mild lifting was implemented, only the state of Florida will experience a second wave of the pandemic. In other words, our study showed that the size (or severity) of the second wave depends on the level of lifting of the community lockdown. Furthermore, the severity (or size) of the second wave decreases with decreasing levels of the lockdown lifting. The second waves recorded in Arizona and Florida during moderate and high lifting of the community lockdown measures are drastically more severe than their respective first waves.

We also assessed the impact of universal use of face masks after the lockdown. In particular, we showed that for high lifting, the disease burden in each of the four jurisdictions decreases with increasing face mask compliance. If two in every five residents of each of the four jurisdictions (i.e., 40%) wear face masks after the lockdown, our study showed that only the state of Florida will have a second wave (which will be relatively mild). In fact, no second wave will be experienced in any of the four locations if half of their residents wear face masks. Higher reductions of pandemic burden will be recorded if face mask compliance is 75%. Thus, this study showed that, for mild moderate, or high level of lifting of community lockdown, face masks are extremely useful, and can greatly suppress COVID-19 (in addition to obviating the likelihood of a devastating second wave). On the other hand, if the level of lifting is unreasonably high, characterized by a 50% increase in the community contact rate parameters (this hypothetical scenario is essentially equivalent to returning to almost a “business as usual” scenario), then all four jurisdictions will experience a second wave (*albeit* the second wave for New York state and the entire US will be mild) if half the residents of the respective jurisdictions wear face masks after the lockdown. There will, however, be no second wave in any of the four jurisdictions if the post-lockdown face mask compliance is 75%. Thus, our study showed that high face masks compliance after lockdown will greatly curtail COVID-19 even if the lifting of the lockdown was high.

The community-wide impact of early detection and self-isolation of asymptomatic cases (as measured by increase in the maximum detection rate parameter, τ_{dmax}) was also assessed in our study. Our simulation results showed a significant reduction in the daily and cumulative mortality with increasing values of the maximum detection rate, in all four jurisdictions. Although the states of Arizona and Florida, as well as the entire US, will suffer devastating second waves when their respective baseline value of the maximum detection rate parameter are used, ramping up testing (and contact tracing, followed by self-isolation of cases) to a level that increases the respective baseline value of the maximum detection rate by just 10% will result in a much milder second wave in Arizona and Florida, and no second wave in New York state and the whole of the US. Further increasing the baseline maximum detection rate to 20% of its baseline value will guarantee no second wave as well as trigger a substantial reduction in the number of deaths in any of the four jurisdictions. More dramatic reductions in disease burden are achieved if the testing strategy is combined with a universal face masks use strategy.

In summary, this study emphasizes the importance of early implementation of effective control strategies. In the absence of a safe and effective vaccine or antiviral, the control and mitigation of COVID-19 rely solely on the implementation of non-pharmaceutical interventions. While the lockdown measures implemented in the US have greatly curtailed community transmission of COVID-19 during the lockdown period, numerous states within the US federation rushed to pre-maturely re-open or lift the lockdown measures, triggering an alarming resurgence of COVID-19 in numerous states. We showed that the implementation of an effective public face mask strategy will greatly control community transmission in the states of Arizona, Florida, New York, and the entire US if the compliance level in each of the four jurisdictions is high enough (regardless of the level of lifting of lockdown implemented). We showed that such a face mask use strategy can avert the projected devastating second expected to hit these jurisdictions by the end of the year 2020. The effectiveness of face masks to curtail the burden of COVID-19 is enhanced if it is combined with an effective testing strategy that can increase the maximum detection rate (of asymptomatic infected individuals) in the community. This study showed that the prospect for the effective control of the post-lockdown resurging COVID-19 in the states of Arizona, Florida, New York and the entire US is very promising using a face mask strategy, if the mask strategy is universally adopted in the country (and compliance is at least moderate). Before a safe and effective vaccine and/or antiviral is developed and approved for use in humans, the use of face masks in public is, undoubtedly, the best and most effective way to curtail community transmission of the COVID-19 pandemic in the four jurisdictions we considered. Of course, our study is not

advocating that face masks must be worn in public until the vaccine or antiviral is developed. They should be worn until community transmission is greatly suppressed. Once this is achieved, the use of face masks in public can be relaxed, as long as diagnostic testing and contact tracing can be ramped up to quickly identify and suppress any new post-mask outbreak in the community. Countries in Asia, particularly (and some in Europe) adopted this approach to great effect. Once their mask use strategy (and other NPIs, such as social-distancing) succeeded in suppressing community transmission, they relaxed the use of face masks in public but continue to be very vigilant (*vis a vis* the rapid identification and suppression of any future outbreaks). It should be mentioned that contact tracing is more feasible (and effective) when community transmission is already suppressed to a low (and manageable) level. It is logistically-easier (or feasible) to effectively trace a small number of cases, than a large number. The US is currently recording about 56,000 confirmed cases daily, and the number of confirmed cases is expected to rise (up to 100,000 cases daily, or more). Tracing the contacts of these cases will be almost an impossible undertaking. Our study emphasize the urgent need to decrease community transmission (*via*, primarily, the universal use of face masks, complemented by social-distancing, avoiding large gatherings and other NPIs). Once the reduction is achieved, the face masks use strategy can be relaxed, and tracing and testing can now become the main strategy to rapidly detect and contain any future COVID-19 outbreaks in the community.

Finally, this study highlights the fact that widespread random testing contributes in detecting, tracing and isolating asymptomatic cases (hence, breaking their transmission chains) that would otherwise be spreading the virus in the community. More testing clearly does not mean more new cases. More testing means more detection (and rapid isolation) of asymptomatic cases, thereby reducing community transmission. Hence, more testing reduces number of new cases, hospitalizations and deaths. More dramatic reduction in COVID-19 burden (measured in terms of reduction in the number of new cases, hospitalizations and deaths) is achieved when the public face masks use strategy (with at least moderate compliance) is combined with a robust and effective random testing (and subsequent tracing and rapid isolation of cases) strategy in the community.

Acknowledgements

ABG acknowledges the support, in part, of the Simons Foundation (Awards #585022) and the National Science Foundation (Award #1917512). CNN acknowledges the support of the Simons Foundation (Award #627346).

References

- [1] World Health Organization, Emergencies, preparedness, response. Pneumonia of unknown origin – China, Disease Outbreak News (5 January, Accessed on March 5, 2020).
- [2] Q. Li, X. Guan, P. Wu, X. Wang, L. Zhou, Y. Tong, R. Ren, K. S. Leung, E. H. Lau, J. Y. Wong, et al., Early transmission dynamics in Wuhan, China, of novel coronavirus–infected pneumonia, *New England Journal of Medicine* (2020).
- [3] World Health Organization, Coronavirus disease (COVID-2019) situation reports, WHO (Accessed on April 26, 2020).
- [4] E. Dong, H. Du, L. Gardner, An interactive web-based dashboard to track COVID-19 in real time, *The Lancet Infectious Diseases* (2020).
- [5] Worldometer, COVID-19 coronavirus pandemic: reported cases and deaths by country, territory, or conveyance, Worldometer (Assessed on May 2, 2020).
- [6] E. Czachor, Houston ICUs at 97 percent capacity as texas coronavirus cases break records, *News Week* (Assessed on June 25, 2020).

- [7] N. van Doremalen, T. Bushmaker, D. H. Morris, M. G. Holbrook, A. Gamble, B. N. Williamson, A. Tamin, J. L. Harcourt, N. J. Thornburg, S. I. Gerber, et al., Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1, *New England Journal of Medicine* (2020).
- [8] World Health Organization, Coronavirus disease 2019 (COVID-19): situation report, 46, WHO (2020).
- [9] Y. Liu, L.-M. Yan, L. Wan, T.-X. Xiang, A. Le, J.-M. Liu, M. Peiris, L. L. Poon, W. Zhang, Viral dynamics in mild and severe cases of covid-19, *The Lancet Infectious Diseases* (2020).
- [10] W. K. Silverstein, L. Stroud, G. E. Cleghorn, J. A. Leis, First imported case of 2019 novel coronavirus in Canada, presenting as mild pneumonia, *The Lancet* 395 (2020) 734.
- [11] X. Yang, Y. Yu, J. Xu, H. Shu, H. Liu, Y. Wu, L. Zhang, Z. Yu, M. Fang, T. Yu, et al., Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study, *The Lancet Respiratory Medicine* (2020).
- [12] X.-W. Xu, X.-X. Wu, X.-G. Jiang, K.-J. Xu, L.-J. Ying, C.-L. Ma, S.-B. Li, H.-Y. Wang, S. Zhang, H.-N. Gao, et al., Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series, *British Medical Journal* 368 (2020).
- [13] B. E. Young, S. W. X. Ong, S. Kalimuddin, J. G. Low, S. Y. Tan, J. Loh, O.-T. Ng, K. Marimuthu, L. W. Ang, T. M. Mak, et al., Epidemiologic features and clinical course of patients infected with SARS-CoV-2 in Singapore, *JAMA* (2020).
- [14] Z. Wu, J. M. McGoogan, Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention, *JAMA* (2020).
- [15] T. G. Rajesh, B. L. John, C. del Rio, Mild or moderate Covid-19, *The New England Journal of Medicine (Clinical Practice)* (2020).
- [16] X. He, E. H. Y. Lau, P. Wu, X. Deng, J. Wang, X. Hao, Y. C. Lau, J. Y. Wong, Y. Guan, X. Tan, X. Mo, Y. Chen, B. Liao, W. Chen, F. Hu, Q. Zhang, M. Zhong, Y. Wu, L. Zhao, F. Zhang, B. J. Cowling, F. Li, G. M. Leung, Temporal dynamics in viral shedding and transmissibility of COVID-19, *Nature Medicine* (2020).
- [17] L. Zou, F. Ruan, M. Huang, L. Liang, H. Huang, Z. Hong, J. Yu, M. Kang, Y. Song, J. Xia, et al., SARS-CoV-2 viral load in upper respiratory specimens of infected patients, *New England Journal of Medicine* 382 (2020) 1177–1179.
- [18] H. Nishiura, T. Kobayashi, T. Miyama, A. Suzuki, S. Jung, K. Hayashi, R. Kinoshita, Y. Yang, B. Yuan, A. R. Akhmetzhanov, et al., Estimation of the asymptomatic ratio of novel coronavirus infections (COVID-19), *medRxiv* (2020).
- [19] K. Mizumoto, K. Kagaya, A. Zarebski, G. Chowell, Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020, *Euro-surveillance* 25 (2020) 2000180.
- [20] E. Callaway, Scores of coronavirus vaccines are in competition — how will scientists choose the best?, a *nature* research journal (Assessed on April 30, 2020).
- [21] D. Kirkpatrick, In race for a coronavirus vaccine, an oxford group leaps ahead, <https://www.nytimes.com/2020/04/27/world/europe/coronavirus-vaccine-update-oxford.html> (Assessed on May 2, 2020).
- [22] S. Knvul, T. Katie, More coronavirus vaccines and treatments move toward human trials, <https://www.nytimes.com/2020/04/08/health/coronavirus-vaccines.html> (Assessed on May 2, 2020).

- [23] J. Cohen, Not wearing masks to protect against coronavirus is a ‘big mistake’, top Chinese scientist says’, *Science Mag* 27 (2020).
- [24] L. J. Radonovich, M. S. Simberkoff, M. T. Bessesen, A. C. Brown, D. A. Cummings, C. A. Gaydos, J. G. Los, A. E. Krosche, C. L. Gibert, G. J. Gorse, et al., N95 respirators vs medical masks for preventing influenza among health care personnel: a randomized clinical trial, *JAMA* 322 (2019) 824–833.
- [25] V. Offeddu, C. F. Yung, M. S. F. Low, C. C. Tam, Effectiveness of masks and respirators against respiratory infections in healthcare workers: a systematic review and meta-analysis, *Clinical Infectious Diseases* 65 (2017) 1934–1942.
- [26] S. Feng, C. Shen, N. Xia, W. Song, M. Fan, B. J. Cowling, Rational use of face masks in the COVID-19 pandemic, *The Lancet Respiratory Medicine* (2020).
- [27] S. Mervosh, J. C. Lee, L. Gamio, N. Popovich, See how all 50 states are reopening, *The New York Times* (Assessed on June 25, 2020).
- [28] White House Coronavirus Task Force, Opening up America again, White House (Assessed on July 2, 2020).
- [29] K. Collins, L. Leatherby, Most states that are reopening fail to meet White House guidelines, *The New York Times* (Assessed on July 2, 2020).
- [30] A. Bendix, Nearly half of states reopened before meeting the White House safety criteria. Their new surges were entirely predictable., *Business Insider (Science News)* (Assessed on July 2, 2020).
- [31] J. Bote, G. Hauck, At least 21 states pause reopening or take new steps to limit the spread of COVID-19, *USA Today* (Assessed on July 3, 2020).
- [32] S. R. Kelleher, These U.S. cities and states have hit the pause button on reopening, *Forbes* (Assessed on July 3, 2020).
- [33] S. A. Allen Kim, J. Froio, These are the states requiring people to wear masks when out in public, *CNN* (Assessed on July 3, 2020).
- [34] L. Mendelson, Facing your face mask duties – a list of statewide orders, as of July 2, 2020, *Littler* (Assessed on July 3, 2020).
- [35] R. Stein, C. Wroth, A. Hurt, US Coronavirus testing still falls short. How’s your state doing?, *Health News from National Public Radio* ((Assessed on May 11, 2020)).
- [36] World Health Organization, Coronavirus disease (COVID-2019) situation dashboard, WHO (Accessed on May 2, 2020).
- [37] World Health Organization, Novel coronavirus (2019-nCoV) situation reports, WHO (Accessed on May 2, 2020).
- [38] E. Dong, H. Du, L. Gardner, Coronavirus COVID-19 Global Cases by Johns Hopkins CSSE, *The Lancet Infectious Diseases* (2020).
- [39] H. Lau, T. Khosrawipour, P. Kocbach, H. Ichii, J. Bania, V. Khosrawipour, Evaluating the massive underreporting and undertesting of COVID-19 cases in multiple global epicenters, *Pulmonology* (2020).
- [40] L. H. Sun, J. Achenbach, Cdc chief says coronavirus cases may be 10 times higher than reported, *The Washington Post (Health)* (Assessed on July 3, 2020).

- [41] C. N. Ngonghala, E. Iboi, S. Eikenberry, M. Scotch, C. R. MacIntyre, M. H. Bonds, A. B. Gumel, Mathematical assessment of the impact of non-pharmaceutical interventions on curtailing the 2019 novel coronavirus, *Mathematical Biosciences* (2020) 108364.
- [42] J. Arino, F. Brauer, P. Van Den Driessche, J. Watmough, J. Wu, A final size relation for epidemic models, *Mathematical Biosciences & Engineering* 4 (2007) 159.
- [43] F. Brauer, A final size relation for epidemic models of vector-transmitted diseases, *Infectious Disease Modelling* 2 (2017) 12–20.
- [44] F. Brauer, The final size of a serious epidemic, *Bulletin of mathematical biology* 81 (2019) 869–877.
- [45] O. Diekmann, J. A. P. Heesterbeek, J. A. Metz, On the definition and the computation of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous populations, *Journal of Mathematical Biology* 28 (1990) 365–382.
- [46] P. van den Driessche, J. Watmough, Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission, *Mathematical Biosciences* 180 (2002) 29–48.
- [47] New York Times, What Are the Phases of New York’s Reopening Plan?, *The New York Times* (Assessed on July 4, 2020).
- [48] A. Cuomo, Governor Cuomo announces lowest covid-19 deaths and hospitalizations in new york since the pandemic began, *Information on Coronavirus* (Assessed on July 2, 2020).
- [49] J. Croft, C. Maxouris, D. Wood, Record increases of new Covid-19 cases in 3 US states with the most population, *CNN* (Assessed on June 25, 2020).
- [50] A. Davies, K.-A. Thompson, K. Giri, G. Kafatos, J. Walker, A. Bennett, Testing the efficacy of homemade masks: would they protect in an influenza pandemic?, *Disaster Medicine and Public Health Preparedness* 7 (2013) 413–418.
- [51] L. Peng, W. Yang, D. Zhang, C. Zhuge, L. Hong, Epidemic analysis of COVID-19 in China by dynamical modeling, *arXiv preprint arXiv:2002.06563* (2020).
- [52] C. Zhou, Evaluating new evidence in the early dynamics of the novel coronavirus COVID-19 outbreak in Wuhan, China with real time domestic traffic and potential asymptomatic transmissions, *medRxiv* (2020).
- [53] N. M. Linton, T. Kobayashi, Y. Yang, K. Hayashi, A. R. Akhmetzhanov, S.-m. Jung, B. Yuan, R. Kinoshita, H. Nishiura, Incubation period and other epidemiological characteristics of 2019 novel coronavirus infections with right truncation: a statistical analysis of publicly available case data, *Journal of Clinical Medicine* 9 (2020) 538.
- [54] H. Sun, Y. Qiu, H. Yan, Y. Huang, Y. Zhu, S. X. Chen, Tracking and Predicting COVID-19 Epidemic in China Mainland, *medRxiv* (2020).
- [55] C. You, Y. Deng, W. Hu, J. Sun, Q. Lin, F. Zhou, C. H. Pang, Y. Zhang, Z. Chen, X.-H. Zhou, Estimation of the time-varying reproduction number of COVID-19 outbreak in China, Available at SSRN 3539694 (2020).
- [56] M. Arentz, E. Yim, L. Klaff, S. Lokhandwala, F. X. Riedo, M. Chong, M. Lee, Characteristics and outcomes of 21 critically ill patients with COVID-19 in Washington State, *Jama* (2020).
- [57] S. Kissler, C. Tedijanto, E. Goldstein, Y. Grad, M. Lipsitch, Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period, *Science* (2020).

- [58] N. M. Ferguson, D. Laydon, G. Nedjati-Gilani, N. Imai, K. Ainslie, M. Baguelin, S. Bhatia, A. Boonyasiri, Z. Cucunubá, G. Cuomo-Dannenburg, et al., Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand, London: Imperial College COVID-19 Response Team, March 16 (2020).
- [59] Centers for Disease Control and Prevention, Coronavirus disease 2019 (COVID-19): Caring for someone sick at home, CDC (Assessed on April 29, 2020).

Appendices

A Tables of variable descriptions, parameter descriptions, and parameter values

Table A.1: Description of the state variables of the model (2.1)

State variable	Description
S	Population of susceptible individuals
E	Population of exposed (newly-infected but not infectious) individuals
E_p	Population of pre-symptomatic (infectious) individuals
I_a	Population of asymptotically-Infectious individuals
I_m	Population of infectious individuals with mild or moderate clinical symptoms of COVID-19
I_s	Population of infectious individuals with severe clinical symptoms of COVID-19
I_i	Population of infectious individuals in self-isolation
I_h	Population of hospitalized individuals
I_c	Population of individuals in ICU
R_u	Population of untested recovered individuals
R_t	Population of recovered individuals who received serology (antibody) test

Table A.2: Description of parameters of the model (2.1)

Parameter	Description
$\beta_p(\beta_a)(\beta_m)(\beta_s)$	Effective contact rate for individuals in the $E_p(I_a)(I_m)(I_s)$ class
ε_m	Efficacy of face masks to prevent transmission and acquisition of infection ($0 < \varepsilon_m \leq 1$)
c_m	Compliance in face mask usage in the community ($0 < c_m \leq 1$)
τ_d	Detection rate of asymptotically-infected individuals <i>via</i> random diagnostic/surveillance testing
τ_s	Detection rate of untested recovered individuals <i>via</i> serology (antibody) testing
$\tau_{dmax}(\tau_{smax})$	Maximum diagnostic (serology) detection rate
T_n	Average number of testing conducted <i>per</i> day
σ_e	Progression rate from E to E_p class
σ_p	Progression rate of pre-symptomatic exposed individuals to I_a, I_m or I_s class
r	Proportion of individuals in the E_p class who show no clinical symptoms of COVID-19 at the end of the incubation period (and move to the I_a class)
$1 - r$	Proportion of individuals in the E_p class who show clinical symptoms of COVID-19 at the end of the incubation period (and move to the I_m class, at a rate $g(1 - r)\sigma$, or to the I_s class, at a rate $(1 - g)(1 - r)\sigma$)
g	Proportion of individuals in the E_p class who develop mild symptoms of COVID-19 at the end of the incubation period
$1 - g$	Proportion of individuals in the E_p class who develop severe symptoms of COVID-19 at the end of the incubation period
f	Proportion of individuals in the I_s class who are hospitalized
$1 - f$	Proportion of individuals in the I_s class who are self-isolated
ρ_m	Self-isolation rate for individuals in the I_m class
$(1 - f)\rho_s$	Self-isolation rate for individuals in the I_s class
$f\rho_s$	Hospitalization rate for individuals in the I_s class
ξ_i	Hospitalization rate of self-isolated individuals
ψ_h	ICU admission rate for hospitalized individuals
$\gamma_a(\gamma_s)(\gamma_h)(\gamma_i)(\gamma_c)$	Recovery rate for individuals in the $I_a(I_s)(I_h)(I_i)(I_c)$ class
$\delta_s(\delta_h)(\delta_i)(\delta_c)$	Disease-induced death rate for individuals in the $I_s(I_h)(I_i)(I_c)$ class

Table A.3: Baseline parameter values for the model (2.1) drawn from the literature.

Parameter	value	Reference
ε_m	0.5	Estimated from [50]
σ_e	1/2.5	[17, 51, 52, 52–54]
σ_p	1/2.5	[17, 53–55]
r	0.324	Estimated from [8, 14]
g	0.719	Estimated from [8, 14]
ρ_s	1/3.5	[56]
ψ_h	1/6	[57]
γ_a	1/5	[57]
γ_s	1/10	[17, 58]
γ_h	1/8	[58]
γ_i	1/7	[59]
γ_c	1/10	[17, 58]

Table A.4: Calibrated parameter values for the model (2.1) using cumulative mortality data for the state of Arizona. (a) Pre-lockdown period (i.e., from March 6 to March 31, 2020). (b) Lockdown period (i.e., from March 31 to May 15, 2020). The lower 95% confidence interval bound is denoted by CI (low), while the upper 95% confidence interval bound is denoted by CI (up).

(a) Estimated parameters for the pre-lockdown period.

(b) Estimated parameters for the lockdown period.

Parameter	Value	95% CI (low)	95% CI (up)	Parameter	Value	95% CI (low)	95% CI (up)
β_p	1.4491	0.1988	1.8275	β_p	1.3149	1.0225	1.4491
β_a	0.5746	0.1000	0.9678	β_a	0.5746	0.4512	0.5746
β_m	0.6456	0.1000	3.1649	β_m	0.4372	0.1000	0.6456
β_s	0.4233	0.1000	0.6055	β_s	0.3893	0.2265	0.4233
c_m	0.0435	0.0357	0.0500	c_m	0.1392	0.1100	0.1715
τ_{dmax}	0.2388	0.1537	0.4171	τ_{dmax}	0.4525	0.3257	0.5721
ρ_m	0.2857	0.0571	0.2972	ρ_m	0.2552	0.0982	0.2857
ξ_i	0.2857	0.0857	0.2972	ξ_i	0.2601	0.1287	0.2857
δ_h	0.0155	0.0050	0.0380	δ_h	0.0081	0.0073	0.0103
δ_c	0.0304	0.0300	0.0800	δ_c	0.0108	0.0097	0.0137
δ_s	0.0228	0.0225	0.0600	δ_s	0.0054	0.0073	0.0069
δ_i	0.0228	0.0225	0.0600	δ_i	0.0054	0.0049	0.0069
\mathcal{R}_c	2.0510	0.7623	2.5078	\mathcal{R}_c	0.9415	0.0049	0.9877

Table A.5: Calibrated parameter values for the model (2.1) using cumulative mortality data for the state of Florida. (a) Pre-lockdown period (i.e., from March 1 to April 3, 2020). (b) Lockdown period (i.e., from April 3 to May 4, 2020). The lower 95% confidence interval bound is denoted by CI (low), while the upper 95% confidence interval bound is denoted by CI (up).

(a) Estimated parameters for the pre-lockdown period.

(b) Estimated parameters for the lockdown period.

Parameter	Value	95% CI (low)	95% CI (up)	Parameter	Value	95% CI (low)	95% CI (up)
β_p	1.3523	0.8804	1.6706	β_p	1.1505	0.6004	2.3827
β_a	0.9232	0.4013	1.926	β_a	0.9232	0.1160	0.9832
β_m	0.6800	0.1000	0.8675	β_m	0.5064	0.1154	0.6800
β_s	0.2062	0.0100	0.2479	β_s	0.1642	0.1021	0.2062
c_m	0.0396	0.0374	0.05	c_m	0.1647	0.0800	0.2056
τ_{dmax}	0.2309	0.1065	0.3011	τ_{dmax}	0.4681	0.3242	0.8105
ρ_m	0.2857	0.0571	0.2956	ρ_m	0.1612	0.0857	0.1714
ξ_i	0.2561	0.0973	0.2857	ξ_i	0.2568	0.1143	0.2571
δ_h	0.0051	0.0050	0.0320	δ_h	0.0067	0.0040	0.0150
δ_c	0.0308	0.0300	0.0702	δ_c	0.0075	0.0014	0.0135
δ_s	0.0231	0.0225	0.0527	δ_s	0.0056	0.0011	0.0101
δ_i	0.0231	0.0225	0.0527	δ_i	0.0056	0.0011	0.0101
\mathcal{R}_c	2.0997	1.9118	2.2214	\mathcal{R}_c	0.9772	0.6638	0.9859

Table A.6: Calibrated parameter values for the model (2.1) using cumulative mortality data for the state of New York. (a) Pre-lockdown period (i.e., from March 1 to March 22, 2020). (b) Lockdown period (i.e., from March 22 to May 28, 2020). The lower 95% confidence interval bound is denoted by CI (low), while the upper 95% confidence interval bound is denoted by CI (up).

(a) Estimated parameters for the pre-lockdown period.

(b) Estimated parameters for the lockdown period.

Parameter	Value	95% CI (low)	95% CI (up)	Parameter	Value	95% CI (low)	95% CI (up)
β_p	2.7450	1.3772	3.3895	β_p	1.0363	0.4207	1.8901
β_a	0.8989	0.1000	2.0134	β_a	0.8989	0.2228	0.9889
β_m	0.1254	0.1000	3.1228	β_m	0.4046	0.1735	0.5978
β_s	0.4246	0.1000	0.8043	β_s	0.5943	0.1000	0.7425
c_m	0.0190	0.0000	0.0500	c_m	0.1500	0.1222	0.2162
τ_{dmax}	0.3010	0.3002	0.4940	τ_{dmax}	0.4485	0.3010	0.6142
ρ_m	0.2857	0.0571	0.2896	ρ_m	0.2466	0.1716	0.2857
ξ_i	0.2822	0.0857	0.2857	ξ_i	0.2426	0.1714	0.2571
δ_h	0.0371	0.0050	0.0400	δ_h	0.0097	0.0067	0.0221
δ_c	0.0434	0.0300	0.0800	δ_c	0.0129	0.0089	0.0221
δ_s	0.0326	0.0225	0.0600	δ_s	0.0065	0.0044	0.0111
δ_i	0.0326	0.0225	0.0600	δ_i	0.0065	0.0044	0.0111
\mathcal{R}_c	2.5321	1.5460	3.2335	\mathcal{R}_c	0.8480	0.5307	0.8955

Table A.7: Calibrated parameter values for the model (2.1) using cumulative mortality data for the entire US. (a) Pre-lockdown period (i.e., from January 22 to April 7, 2020). (b) Lockdown period (i.e., from April 7 to May 28, 2020). The lower 95% confidence interval bound is denoted by CI (low), while the upper 95% confidence interval bound is denoted by CI (up).

(a) Estimated parameters for the pre-lockdown period.

(b) Estimated parameters for the lockdown period.

Parameter	Value	95% CI (low)	95% CI (up)	Parameter	Value	95% CI (low)	95% CI (up)
β_p	0.7290	0.6677	0.7751	β_p	0.4075	0.3327	0.5454
β_a	0.8969	0.7745	0.9764	β_a	0.6036	0.1008	0.8614
β_m	0.8086	0.7530	0.9224	β_m	0.4592	0.2531	0.7336
β_s	0.4010	0.1870	0.4642	β_s	0.1344	0.1053	0.2635
c_m	0.0278	0.0276	0.0294	c_m	0.1835	0.1312	0.2300
τ_{dmax}	0.1797	0.1722	0.1960	τ_{dmax}	0.2294	0.1799	0.3812
ρ_m	0.1801	0.1584	0.1853	ρ_m	0.2571	0.1045	0.2684
ξ_i	0.1765	0.1316	0.2417	ξ_i	0.0795	0.0571	0.1557
δ_h	0.0073	0.0052	0.0386	δ_h	0.0048	0.0043	0.0050
δ_c	0.0304	0.0300	0.0800	δ_c	0.0064	0.0058	0.0067
δ_s	0.0228	0.0225	0.0600	δ_s	0.0032	0.0029	0.0034
δ_i	0.0228	0.0225	0.0600	δ_i	0.0032	0.0029	0.0034
\mathcal{R}_c	2.3277	2.2854	2.3758	\mathcal{R}_c	0.8833	0.6363	0.8929