## Genomic epidemiology reveals multiple introductions and spread of SARS-CoV-2 in the Indian state of Karnataka

Chitra Pattabiraman<sup>1</sup>, Farhat Habib<sup>2</sup>, Harsha PK<sup>1</sup>, Risha Rasheed<sup>1</sup>, Vijayalakshmi Reddy<sup>1</sup>, Prameela Dinesh<sup>3</sup>, Tina Damodar<sup>1</sup>, Kiran Hosallimath<sup>1</sup>, Anson K George<sup>1</sup>, Nakka Vijay Kiran Reddy<sup>1</sup>, Banerjee John<sup>1</sup>, Amrita Pattanaik<sup>1</sup>, Narendra Kumar<sup>1</sup>, Reeta S Mani<sup>1</sup>, Manjunatha M Venkataswamy<sup>1</sup>, Shafeeq K Shahul Hameed<sup>1</sup>, Prakash Kumar B.G<sup>3</sup>, Anita Desai<sup>1</sup>, Ravi Vasanthapuram<sup>1</sup>

An earlier version of this manuscript was removed owing to inclusion of confidential information relating to an individual

<sup>&</sup>lt;sup>1</sup> Department of Neurovirology, National Institute of Mental Health and Neurosciences, Bengaluru-560029

<sup>&</sup>lt;sup>2</sup> TruFactor-InMobi Group, Bengaluru-560103

<sup>&</sup>lt;sup>3</sup> Directorate of Health and Family Welfare Services, Government of Karnataka, Bengaluru-560012

<sup>\$</sup>Corresponding author. Email – virusravi@gmail.com