

1 **Original Article**

2
3 **The utility of Resting Pulse Rate in Defining High Blood Pressure among Adolescents in Mbarara Municipality,**
4 **Uganda**

5
6 **Godfrey Katamba,^{1,*} David Collins Agaba,² Rosemary Namayanja², Agnes Namaganda¹, Abdul Musasizi³,**
7 **Mivule Abdul Kinene⁴, Richard Migisha²**

8
9 ¹Department of Physiology, School of Medicine, King Ceasor University, Kampala, Uganda

10 ²Department of Physiology, Faculty of Medicine, Mbarara University of Science and Technology, Mbarara Uganda

11 ³Department of Biochemistry, School of Medicine, King Ceasor University, Kampala, Uganda

12 ⁴Department of Human Anatomy, School of Medicine, King Ceasor University, Kampala, Uganda

13
14 Corresponding author; Godfrey Katamba, King Ceasor University, Kampala Uganda; email; cgmkats13@gmail.com

15
16 **Abstract**

17
18 High resting pulse rate (RPR) is associated with adverse cardiovascular events, and could be
19 used as a marker of cardiovascular health. We determined the correlation between RPR and
20 blood pressure (BP); and its accuracy in defining high blood pressure among adolescents
21 attending secondary schools in Mbarara municipality, south-western Uganda. We conducted a
22 cross-sectional study among secondary school adolescents aged 12-19 years in Mbarara
23 municipality, Uganda. We captured demographic characteristics using a structured questionnaire;
24 and measured anthropometric indices and BP. We performed a linear regression analysis to
25 determine the relationship between RPR and blood pressure and plotted receiver operating
26 characteristics curves (ROC) to assess the accuracy of RPR in defining high BP. We enrolled
27 616 adolescents with a mean age of 15.6±2.0 years and 65.6% (404/616) were female. The RPR
28 was significantly correlated with diastolic blood pressure (DBP) in both boys (Beta = 0.22
29 [95%CI: 0.10; 0.36]), p<0.001 and girls (Beta = 0.51 [95%CI: 0.43; 0.60]), p<0.001. RPR was
30 significantly correlated with systolic blood pressure (SBP) only in the girls (Beta = 0.23 [95%CI:
31 0.15; 0.30]), p<0.001. The optimal threshold for RPR in defining prehypertension was
32 RPR≥76bpm with an area under the curve (AUC) of 0.653[95%CI: 0.583-0.722], the sensitivity
33 of 0.737 and specificity of 0.577. In defining hypertension, the optimal threshold was RPR ≥
34 79bpm at a sensitivity of 0.737 and specificity of 0.719, with an AUC of 0.728[95%CI: 0.624-
35 0.831]. Resting pulse rate was positively correlated with BP and was more accurate in defining
36 hypertension compared to prehypertension in the study.
37
38

39 **Introduction**

40 Hypertension (HTN) was previously considered a rare condition among adolescents¹. In contrast, it has become more
41 common and is increasing unexpectedly in this age group^{2, 3}. The prevalence of HTN among adolescents has been
42 reported in several countries and varies from 3.6% to 30%³⁻¹¹. High blood pressure (BP) in young subjects is
43 associated with accelerated vascular aging and increased risk for adult HTN and cardiovascular disease^{12, 13}. The
44 diagnostic algorithm for HTN in children and adolescents does not rely on single BP cut-offs as is the case for adults.
45 This is believed to be the cause of its under-diagnosis in young populations, because the normal and abnormal BP
46 values vary with sex, height, and age, making it difficult for clinicians to remember¹⁴. Furthermore, the algorithm
47 involves repeated BP measurements and follow up visits^{15, 16}. This is laborious and may not be suitable for resource-
48 limited settings, with a low clinician to patient ratio. Thus, easier to use diagnostic approaches are needed in resource-
49 limited settings, to facilitate the identification of adolescents with high BP and those who need interventions to prevent
50 or control HTN to improve long-term outcomes.

51 Resting pulse rate (RPR) has recently attracted attention as an important predictor of high BP^{17, 18} and a risk factor for
52 cardiovascular morbidity and mortality in general populations¹⁹⁻²². Additionally, various studies among children and
53 adolescents, have reported a significant correlation of elevated RPR with both systolic (SBP) and diastolic blood
54 pressure (DBP)²³⁻²⁶, highlighting its possible role in screening for high BP. Hypertension results from alterations in
55 the overall hemodynamic load on the cardiovascular system²⁷, which is an interplay between the cardiac output (CO)
56 and total peripheral vascular resistance (TPR)²⁸. The CO is affected by changes in the heart rate, which in turn depends
57 on the autonomic tone of the cardiovascular system²⁹. The relationship between RPR and BP, among adolescents, has
58 not been well studied in sub-Saharan Africa. RPR is considered a simple and cheap measurement, which does not
59 necessitate the use of special instruments and patients can conveniently measure it by themselves.

60 In this cross-sectional study, we aimed to assess the utility of RPR in defining high blood pressure (SBP and DBP)
61 among adolescents from selected secondary schools in Mbarara municipality. South-western Uganda.
62

63 **Methods**

64 *Study sample*

65 The participants were part of a cross-sectional study ((IRB No. 18/03-18), which involved three secondary schools in
66 Mbarara municipality, southwestern Uganda between May and November 2018. The sample size was estimated using
67 the Kish Leslie method of 1965³⁰ assuming a prevalence of hypertension of 10.7% among secondary school
68 adolescents³¹, and 95% confidence interval within a 3% error margin. The final sample size was adjusted for an
69 anticipated participant non-response rate of 10%; thus 449 participants were required. In this analysis, a total sample
70 of 616 adolescents aged 12-19 years was used.

71 *Socio-demographic information*

72 The age of the participant was self-reported and was the number of complete years since birth while the sex of the
73 participant was a binary variable (male or female) and based on the sexual characteristics as was observed by the
74 researcher.

75 *Determination of anthropometric indices*

76 The methods used to determine the height, weight, neck circumference, waist circumference, and their respective
77 ratios, have been described in detail elsewhere³². Briefly, height was measured using a wall mount height board in
78 centimeters with the participant having no shoes⁶. A standard Seca scale was used to determine the weight to the
79 nearest 0.5kg³³. Body mass index (BMI) was calculated as the ratio of weight in kilograms to height in square meters.
80 Waist circumference (WC) was measured at the midpoint between the lowest border of the rib cage and the top of the
81 lateral border of the iliac crest while hip circumference was measured at the greatest horizontal circumference below
82 the iliac crest at the level of the greater trochanter using a non-elastic measuring tape (Seca 203 Ergonomic
83 circumference measuring tape, Hamburg, Germany). Waist to hip ratio (WHR) was the ratio of WC to HC³⁴ while
84 waist to height ratio (WHtR) was the ratio of WC divided by height³⁵. The Neck circumference (NC) assessed as a
85 surrogate measure for upper body adipose tissue distribution. It was measured at the level of the laryngeal prominence
86 using an inelastic flexible measuring tape (Seca 203 Ergonomic circumference measuring tape, Hamburg, Germany),
87 with the subjects in the standing position and the head held erect and eyes facing forward to the nearest 0.1cm³⁶.

88 *Measurement of blood pressure and resting pulse rate*

89 Blood pressure and resting pulse rate were measured using a digital blood pressure machine (Scian SP-582 Digital
90 Blood Pressure Monitor) as previously described by Katamba et al³². Each participant was allowed to seat on a chair
91 with back supported, feet on the floor, arm supported and cubital fossa at heart level after 5 minutes of sitting rest
92 without talking^{37, 38}. The cuff of appropriate size (ranged from 12x22cm to 16x30cm) was placed at the bare upper
93 arm, one inch above the bend of the participant's elbow. It was ensured that the tubing fell over the front center of the
94 arm so that the sensor was correctly placed. The end of the cuff was pulled so that it was evenly tight around the arm.
95 The cuff was placed tight enough so that only two fingertips could be slipped under the top edge of the cuff. It was
96 made sure that the skin did not pinch when the cuff inflated. The participant was asked to remain and quiet as the
97 machine begins measuring. The cuff inflated automatically after placing the start button, and then slowly deflated so
98 that the machine took the measurement. When the reading was complete, the monitor displayed the BP and pulse rate
99 on the digital panel³⁹. Three readings were recorded per participant at 5 minutes interval. The average of the 2nd and
100 3rd SBP and DBP measurements was used as the subject's BP respectively³⁸. Those adolescents who had elevated
101 BP in the first session were identified. A re-measurement, using the same procedure was done after one week to
102 confirm that BP is truly and constantly elevated.
103

104 *Data management and analysis*

105 Data were analyzed using the Stata software version 13.0 (College Station, Texas, USA). Continuous variables such
106 were described as mean±SD while categorical variables such as sex were described as percentages and frequencies.
107 The outcome variable was BP (both SBP and DBP). Pearson correlation analysis was done to determine the strength
108 of the relationship between BP, RPR, and anthropometric indices. The Pearson correlation coefficients and their 95%
109 confidence intervals were reported. Multivariate linear regression analysis was used to control for anthropometric
110 indices in the relationship between RPR and BPs. A p-value of less than 0.05 was considered for assessing the
111 statistical significance and 95% CI of the changes in BP values associated with a unit change in RPR was calculated.
112 Receiver operating characteristic (ROC) curve analyses were performed to determine the accuracy of RPR in defining
113 high BP (prehypertension and hypertension) among adolescents and identify optimal thresholds of RPR for identifying
114 high arterial BP. Optimal thresholds were selected as the values corresponding to the maximum of Youden's index
115 on the ROC curve. Prehypertension and hypertension were then redefined by the determined optimal thresholds of
116 RPR. These were compared with the gold standard blood pressure cut-offs in adolescents as stated by the Joint
117 National Committee on hypertension guidelines⁴⁰. The sensitivities and specificities and their respective 95%
118 confidence intervals (CI) were obtained to assess the performance of the determined RPR optimal thresholds. All
119 analyses were stratified by sex because of the physiological difference in resting pulse rates between the two groups.
120

121 **Results**

122 *Characteristics of study participants by body mass index*

123 A sample of 616 had complete data and was included in the final analysis as was obtained from a larger prevalence
124 study. These were aged between 12 to 19 years; with a mean age of 15.6±2.0 years. The prevalence of hypertension
125 was found at 3.1% while prehypertension was estimated at 7.1%⁴¹. In table 1, the characteristics of the participants
126 are described according to BMI categories.
127

128 *Correlation analysis*

129 RPR was positively correlated with both SBP and DBP among adolescents in our study. While the overall linear
130 correlation coefficients of 0.22 [95% CI: 0.14 – 0.29] for SBP and 0.44 [95% CI: 0.37 – 0.50] for DBP were
131 statistically significant, they are weak. The correlation of RPR with both SBP and DBP was still significant and
132 positive even after controlling for the sex of participants as shown in figure 1. Most of the indices showed positive
133 linear relations with both DBP and SBP among the girls, except for the WHR, which exhibited a negative correlation
134 with SBP as seen in table 2.

135 *Linear regression analysis*

136 A bivariate linear regression of RPR and BP (SBP and DBP) indicated a significant positive linear relationship
137 between two variables. However, the variability (using the linear regression R squared value) in DBP, which could be
138 explained by RPR, was only 19% compared to 5% for SBP. For the entire sample, a multivariate linear regression
139 showed that a unit increase in RPR was significantly associated with increases in BP (SBP and DBP). The change in
140 SBP reduced after adjusting for anthropometric indices as shown in table figures 2a and 2b and figures 3a and 3b.

141 *Optimal thresholds of RPR for identifying prehypertension and hypertension among adolescents*

142 The optimal threshold for identifying prehypertension was determined as $RPR \geq 76$ bpm with a sensitivity of 0.737
143 and specificity of 0.577. The performance of this threshold based on the AUC was 0.652 [CI; 0.583-0.722] as shown
144 in figure 4. This optimal threshold classified 58.8% of the participants correctly as prehypertensive. The positive
145 predictive value (PPV) of 0.118 and a negative predictive value (NPV) of 0.882. The optimal threshold for
146 hypertension in our study was determined as $RPR \geq 79$ bpm, with a sensitivity of 0.737, the specificity of 0.719 and
147 an AUC of 0.728 [CI; 0.624-0.831] as shown in figure 5. This threshold classified 71.92% of the participants
148 correctly, with a PPV of 0.077 and NPV of 0.988. Both sensitivity and specificity of RPR were higher for the
149 identification of HTN than prehypertension among our study participants as shown in table 3.
150

151 **Discussion**

152 Our study demonstrated a positive linear relationship between RPR and blood pressure (SBP and DBP), even after
153 adjusting for anthropometric indices and sex. This is in agreement with findings from several studies done among
154 children and adolescents in Brazil^{42, 43}, Nigeria³¹, India⁴⁴, USA⁴⁵, and China^{25, 46, 47}. The association between RPR
155 and elevated BP is believed to be due to alterations in the cardiovascular autonomic tone, including a reduction in
156 cardiac vagal stimulation, increased sympathetic activation and imbalance in the sympathovagal system⁴⁸. As
157 previously reported^{49, 50}, these alterations are associated with the recognized increased risk for HTN among individuals
158 with elevated resting heart rates. Moreover, it has been previously postulated that the effect of elevated heart rate on
159 cardiovascular mortality might be mediated through high BP²⁶.

160
161 Our study also found RPR to be more accurate in defining HTN as opposed to defining prehypertension among
162 adolescents. The AUC of 0.609 to 0.854 suggested a high and robust discriminatory accuracy of RPR in identifying
163 HTN among secondary school adolescents in our study setting. In defining prehypertension, the AUC was lower but
164 well above 0.6 which is the recommended minimum. Besides, the much higher NPVs of the cut off for HTN
165 indicated that our test is unlikely to omit adolescents with HTN and prehypertension. However, due to the low
166 prevalence of HTN in our study, the PPVs of our optimal cut-offs were also low. This showed that many adolescents
167 with both HTN and PreHTN would be misclassified as not having prehypertension or hypertension. This makes
168 RPR not the best substitute for the commonly used age, gender and height BP percentiles for diagnosing
169 hypertension. However, it can be used for screening adolescents at high risk of high BP in populations with a high
170 prevalence of high BP and in resource-limited settings. The study findings are in agreement with those as reported
171 by Tjugen *et al*⁵¹, which involved elderly subjects to determine whether the heart rate itself was a risk factor for the
172 development of hypertension or just a marker of sympathetic overactivation. It was found that a high heart rate was
173 a strong predictor of cardiovascular disease and a reliable predictor for the development of hypertension⁵¹. The
174 study also addresses recommendations for further research on the use of RPR in defining HTN among adolescents
175 from China⁴⁶, Hong Kong²⁵ among others. Thus, this study responds to such recommendations with fair news,
176 showing that RPR may be useful in discriminating people who are at risk of developing elevated BP; however, it
177 may not be used as the basis for the diagnosis of the condition. Nonetheless, our findings contribute to the existing
178 body of evidence that proposes the use of RPR as a useful clinical measurement and as a risk factor for
179 cardiovascular disease as was reported from the WHO Cardiovascular Disease and Alimentary Comparison Study²³.

180 Our study has some limitations as outlined; First, the participants were not objectively evaluated secondary causes of
181 high blood pressure and elevated heart rate. Nevertheless, we did not exclude adolescents with a self-reported
182 history of other endocrine disorders and those with acute febrile conditions. Additionally, we assessed adolescents
183 with high BP and elevated RPR on more than one occasion before confirming the diagnosis. Second, we conducted
184 the study in a few selected secondary schools, because of logistical concerns. This limits the generalization of our

185 findings to adolescent populations outside the peri-urban secondary schools of Mbarara municipality, south-western
186 Uganda. We thus recommend larger surveys in a wider population to corroborate our findings. Finally, we cannot
187 assess the temporal relationship between RPR and BP in our study population, because of the cross-sectional nature
188 of our study design. Future longitudinal studies in larger adolescent populations are needed to explore this
189 relationship.

190 **Conclusions**

191 Our study demonstrated a positive linear relationship between RPR and BP among adolescents after controlling for
192 anthropometric indices and sex. RPR was found to be more accurate in defining hypertension than in defining
193 prehypertension in our study population.

194 **List of abbreviations**

195 AUC: area under the curve
196 BMI: Body mass index
197 BP: blood pressure
198 CI: Confidence interval
199 CO: cardiac output
200 CVD: Cardiovascular disease
201 DBP: diastolic blood pressure
202 HTN: Hypertension
203 NC: Neck circumference
204 NPV: negative predictive value
205 PPV: positive predictive value
206 PreHTN: Prehypertension
207 ROC: receiver operating characteristic
208 RPR: resting pulse rate
209 SBP: systolic blood pressure
210 TPR: total peripheral resistance
211 WHR: Waist hip ratio
212 WHtR: Waist height ratio

213 **Declarations**

214 **Ethical approval and consent to participate**

215 The study was approved by the research ethics committee of Mbarara University of Science and Technology (IRB
216 No. 18/03-18) We also obtained permission to collect data from the school headteachers. The class teachers were
217 informed about the purpose of the study and all potential participants were first sensitized about study procedures,
218 possible benefits, and risks. Adolescents of 12 – 17 years freely assented, and consent for their participation was
219 obtained through their teachers. We obtained written informed consent from adolescents aged 18-19 years.

220 **Consent for publication**

221 Not applicable

222 **Competing interests**

223 There is no conflict of interest in this work.

224 **Funding**

225 This research received no external funding

226 **Availability of data and materials**

227 The dataset is available on request from the corresponding other

228 **Authors' contributions**

229 GK, RM, and DCA: Conceptualization of work & its realization, wrote the manuscript, checked the references,
230 compiled the literature sources, data collection, statistical analysis, and interpretation of data, and wrote the
231 manuscript and is the corresponding author.

232 RN, DCA, RM: mentored the conceptualization of work & its realization, compiling literature sources and statistical
233 analysis, helped in data interpretation, guided manuscript writing, checked the references.

234 AN, AM, MAK: Assent and consent form administration, data collection, data entry, and analysis.

235 All authors read and approved the study manuscript

236 **Acknowledgments**

237 I am thankful to my family, friends and the participants from the various data collection sites.

238 **References**

- 239 1. Chiolero A, Bovet P, Paradis G. Screening for elevated blood pressure in children and
240 adolescents: a critical appraisal. *JAMA pediatrics* 2013; **167**(3): 266-273.
241
- 242 2. Chen X, Wang Y. Tracking of blood pressure from childhood to adulthood. *Circulation*
243 2008; **117**(25): 3171-3180.
244
- 245 3. Abdul-Jalil I. Obesity and Hypertension among Adolescents in Senior High Schools in the
246 Asante Akim North District, Ghana. University of Ghana, 2018.
247
- 248 4. Gonçalves VSS, Galvão TF, Andrade KRCd, Dutra ES, Bertolin MNT, Carvalho KMBd
249 *et al.* Prevalence of hypertension among adolescents: systematic review and meta-analysis.
250 *Revista de saude publica* 2016; **50**.
251
- 252 5. Leung LC, Sung RY, So H-K, Wong SN, Lee KW, Lee KP *et al.* Prevalence and risk
253 factors for hypertension in Hong Kong Chinese adolescents: waist circumference predicts
254 hypertension, exercise decreases risk. *Archives of disease in childhood* 2011; **96**(9): 804-
255 809.
256
- 257 6. McNiece KL, Poffenbarger TS, Turner JL, Franco KD, Sorof JM, Portman RJ *et al.*
258 Prevalence of hypertension and pre-hypertension among adolescents. 2007; **150**(6): 640-
259 644. e1.
260
- 261 7. Noubiap JJ, Essouma M, Bigna JJ, Jingi AM, Aminde LN, Nansseu JR. Prevalence of
262 elevated blood pressure in children and adolescents in Africa: a systematic review and
263 meta-analysis. *The Lancet Public Health* 2017; **2**(8): e375-e386.
264
- 265 8. Nkeh-Chungag BN, Sekokotla AM, Sewani-Rusike C, Namugowa A, Iputo JE. Prevalence
266 of hypertension and pre-hypertension in 13-17 year old adolescents living in Mthatha-
267 South Africa: A cross-sectional study. *Central European journal of public health* 2015;
268 **23**(1): 59.
269

- 270 9. Luma GB, Spiotta RT. Hypertension in children and adolescents. *Am Fam Physician* 2006;
271 **73(9)**: 1558-68.
272
- 273 10. Uwaezuoke S, Okoli C, Ubesie A, Ikefuna AJNjocp. Primary hypertension among a
274 population of Nigerian secondary school adolescents: Prevalence and correlation with
275 anthropometric indices: A cross-sectional study. 2016; **19(5)**: 649-654.
276
- 277 11. Emmanuel EE, Dada SA, Amu EO, Aduayi VA, Atoyebi OA, Marcus O *et al.*
278 Hypertension and its correlates among in-school adolescents in Ekiti State, South-west,
279 Nigeria. *Asian Journal of Medical Sciences* 2017; **8(4)**: 1-5.
280
- 281 12. Chen X, Wang Y. Tracking of blood pressure from childhood to adulthood: a systematic
282 review and meta-regression analysis. *Circulation* 2008; **117(25)**: 3171.
283
- 284 13. Harvey A, Montezano AC, Touyz RM. Vascular biology of ageing—Implications in
285 hypertension. *Journal of molecular and cellular cardiology* 2015; **83**: 112-121.
286
- 287 14. Hansen ML, Gunn PW, Kaelber DCJJ. Underdiagnosis of hypertension in children and
288 adolescents. *JAMA* 2007; **298(8)**: 874-879.
289
- 290 15. Program NHBPE. *The fourth report on the diagnosis, evaluation, and treatment of high*
291 *blood pressure in children and adolescents*, US Department of Health and Human
292 Services, National Institutes of Health ..., 2005.
293
- 294 16. Spagnolo A, Giussani M, Ambrozzi AM, Bianchetti M, Maringhini S, Matteucci MC *et al.*
295 Focus on prevention, diagnosis and treatment of hypertension in children and adolescents.
296 *Italian journal of pediatrics* 2013; **39(1)**: 20.
297
- 298 17. Koskela JK, Tahvanainen A, Haring A, Tikkakoski AJ, Ilveskoski E, Viitala J *et al.*
299 Association of resting heart rate with cardiovascular function: a cross-sectional study in
300 522 Finnish subjects. *BMC cardiovascular disorders* 2013; **13(1)**: 102.
301
- 302 18. Yang HI, Kim HC, Jeon JY. The association of resting heart rate with diabetes,
303 hypertension, and metabolic syndrome in the Korean adult population: The fifth Korea
304 National Health and Nutrition Examination Survey. *Clinica Chimica Acta* 2016; **455**: 195-
305 200.
306
- 307 19. Zhang M, Han C, Wang C, Wang J, Li L, Zhang L *et al.* Association of resting heart rate
308 and cardiovascular disease mortality in hypertensive and normotensive rural Chinese.
309 *Journal of cardiology* 2017; **69(5)**: 779-784.
310
- 311 20. Cooney MT, Vartiainen E, Laakitainen T, Juolevi A, Dudina A, Graham IM. Elevated
312 resting heart rate is an independent risk factor for cardiovascular disease in healthy men
313 and women. *American heart journal* 2010; **159(4)**: 612-619. e3.
314

- 315 21. Jouven X, Empana J-P, Schwartz PJ, Desnos M, Courbon D, Ducimetière P. Heart-rate
316 profile during exercise as a predictor of sudden death. *New England Journal of Medicine*
317 2005; **352**(19): 1951-1958.
318
- 319 22. Reunanen A, Karjalainen J, Ristola P, Heliövaara M, Knekt P, Aromaa A. Heart rate and
320 mortality. *Journal of internal medicine* 2000; **247**(2): 231-239.
321
- 322 23. Liu L, Mizushima S, Ikeda K, Nara Y, Yamori Y, Group CS. Resting heart rate in relation
323 to blood pressure: Results from the World Health Organization—Cardiovascular Disease
324 and Alimentary Comparison Study. *International journal of cardiology* 2010; **145**(1): 73-
325 74.
326
- 327 24. Christofaro DGD, Casonatto J, Vanderlei LCM, Cucato GG, Dias RMR. Relationship
328 between Resting Heart Rate, Blood Pressure and Pulse Pressure in Adolescents. *Arquivos*
329 *brasileiros de cardiologia* 2017; (AHEAD): 0-0.
330
- 331 25. Kwok S-Y, So H-K, Choi K-C, Lo AF, Li AM, Sung RY *et al.* Resting heart rate in children
332 and adolescents: association with blood pressure, exercise and obesity. *Archives of disease*
333 *in childhood* 2013: archdischild-2012-302794.
334
- 335 26. Fernandes RA, Júnior IFF, Codogno JS, Christofaro DGD, Monteiro HL, Lopes DMR.
336 Resting heart rate is associated with blood pressure in male children and adolescents. *The*
337 *Journal of pediatrics* 2011; **158**(4): 634-637.
338
- 339 27. Ventura HO, Taler SJ, Strobeck JE. Hypertension as a hemodynamic disease: the role of
340 impedance cardiography in diagnostic, prognostic, and therapeutic decision making.
341 *American journal of hypertension* 2005; **18**(S2): 26S-43S.
342
- 343 28. Mayet J, Hughes A. Cardiac and vascular pathophysiology in hypertension. *Heart* 2003;
344 **89**(9): 1104-1109.
345
- 346 29. Grassi G, Mark A, Esler M. The sympathetic nervous system alterations in human
347 hypertension. *Circulation research* 2015; **116**(6): 976-990.
348
- 349 30. Kish LJA. Sampling organizations and groups of unequal sizes. 1965: 564-572.
350
- 351 31. Uwaezuoke S, Okoli C, Ubesie A, Ikefuna A. Primary hypertension among a population
352 of Nigerian secondary school adolescents: Prevalence and correlation with anthropometric
353 indices: A cross-sectional study. *Nigerian journal of clinical practice* 2016; **19**(5): 649-
354 654.
355
- 356 32. Katamba G, Collins Agaba D, Migisha R, Namaganda A, Namayanja R, Turyakira E.
357 Using blood pressure height index to define hypertension among secondary school
358 adolescents in southwestern Uganda. *Journal of Human Hypertension* 2019.
359

- 360 33. Al-Isa AJEJoCN. Body mass index, overweight and obesity among Kuwaiti intermediate
361 school adolescents aged 10–14 years. *European Journal of Clinical Nutrition* 2004; **58**(9):
362 1273.
363
- 364 34. WHO. Waist circumference and waist-hip ratio: Report of a WHO expert consultation,
365 Geneva, 8-11 December 2008. 2011.
366
- 367 35. Ashwell M, Gunn P, Gibson SJOr. Waist-to-height ratio is a better screening tool than
368 waist circumference and BMI for adult cardiometabolic risk factors: systematic review and
369 meta-analysis. *Obesity reviews* 2012; **13**(3): 275-286.
370
- 371 36. Kuciene R, Dulskiene V, Medzioniene JJBp. Association of neck circumference and high
372 blood pressure in children and adolescents: a case–control study. *BMC pediatrics* 2015;
373 **15**(1): 127.
374
- 375 37. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN *et al.* Recommendations
376 for blood pressure measurement in humans and experimental animals: part 1: blood
377 pressure measurement in humans: a statement for professionals from the Subcommittee of
378 Professional and Public Education of the American Heart Association Council on High
379 Blood Pressure Research. *Circulation* 2005; **111**(5): 697-716.
380
- 381 38. WHO. A global brief on hypertension: silent killer, global public health crisis: World
382 Health Day 2013. 2013.
383
- 384 39. O'brien E, Waeber B, Parati G, Staessen J, Myers MGJBBMJ. Blood pressure measuring
385 devices: recommendations of the European Society of Hypertension. *British Medical*
386 *Journal* 2001; **322**(7285): 531.
387
- 388 40. Rodriguez-Cruz E. Pediatric Hypertension. In: Medscape, 2017.
389
- 390 41. Katamba G, Agaba DC, Migisha R, Namaganda A, Namayanja R, Turyakira E. Prevalence
391 of hypertension in relation to anthropometric indices among secondary adolescents in
392 Mbarara, Southwestern Uganda. *Italian Journal of Pediatrics* 2020; **46**(1): 76.
393
- 394 42. Christofaro DGD, Casonatto J, Vanderlei LCM, Cucato GG, Dias RMR. Relationship
395 between resting heart rate, blood pressure and pulse pressure in adolescents. *Arquivos*
396 *brasileiros de cardiologia* 2017; **108**(5): 405-410.
397
- 398 43. Farah BQ, Christofaro DGD, Balagopal PB, Cavalcante BR, de Barros MVG, Ritti-Dias
399 RM. Association between resting heart rate and cardiovascular risk factors in adolescents.
400 *European journal of pediatrics* 2015; **174**(12): 1621-1628.
401
- 402 44. Chitrapu RV, Thakkallapalli ZM. Prehypertension among medical students and its
403 association with cardiovascular risk factors. *Journal of Dr. NTR University of Health*
404 *Sciences* 2015; **4**(1): 8.
405

- 406 45. Yilmazel G. PREHYPERTENSION PREVALENCE AND ITS ASSOCIATION WITH
 407 NECK AND ABDOMINAL OBESITY IN DISEASE-FREE YOUNG ADULTS. *ACTA*
 408 *MEDICA MEDITERRANEA* 2017; **33**(2): 329-334.
 409
- 410 46. Hu J, Shen H, Chu G-P, Fu H, Huang F-F, Zheng Y-M *et al.* Association of elevated resting
 411 pulse rate with increased risk of hypertension development in children: A prospective study
 412 in Suzhou, China. *Medicine* 2017; **96**(32).
 413
- 414 47. Kwok S-Y, So H-K, Choi K-C, Lo AF, Li AM, Sung RY *et al.* Resting heart rate in children
 415 and adolescents: association with blood pressure, exercise and obesity. *Archives of disease*
 416 *in childhood* 2013; **98**(4): 287-291.
 417
- 418 48. Palatini P, Julius S. The physiological determinants and risk correlations of elevated heart
 419 rate. *American journal of hypertension* 1999; **12**(S1): 3S-8S.
 420
- 421 49. Ji C, Zheng X, Chen S, Dong Y, Yang G, Gao X *et al.* Impact of resting heart rate on the
 422 progression to hypertension in prehypertension patients. *Zhonghua xin xue guan bing za*
 423 *zhi* 2014; **42**(10): 860-865.
 424
- 425 50. Aladin AI, Al Rifai M, Rasool SH, Keteyian SJ, Brawner CA, Michos ED *et al.* The
 426 association of resting heart rate and incident hypertension: the Henry Ford Hospital
 427 Exercise Testing (FIT) Project. *American journal of hypertension* 2016; **29**(2): 251-257.
 428
- 429 51. Tjugen TB, Flaa A, Kjeldsen SE. High Heart Rate as Predictor of Essential Hypertension::
 430 The Hyperkinetic State, Evidence of Prediction of Hypertension, and Hemodynamic
 431 Transition to Full Hypertension. *Progress in cardiovascular diseases* 2009; **52**(1): 20-25.
 432
 433
 434
 435
 436

Table 1: Showing characteristics of participants by BMI categories

Variable	Obese (n=21)	Overweight (n=188)	Normal weight (n=387)	Underweight (n=20)
Female	(n=18)	(n=166)	(n=215)	(n=5)
Male	(n=3)	(n=22)	(n=172)	(n=15)
	<i>Mean (SD)</i>	<i>Mean (SD)</i>	<i>Mean (SD)</i>	<i>Mean (SD)</i>
Age (years)	17.7(1.6)	15.6(1.9)	15.5(2.0)	15.9(1.9)
NC (cm)	31.9(1.8)	30.3(1.7)	29.9(2.2)	28.8(1.6)
WHR	0.84(0.05)	0.76(0.07)	0.8(0.08)	0.79(0.07)
RPR (bpm)	77.9(6.8)	74.9(7.0)	75.6(8.7)	76.4(13.5)
SBP (mmHg)	117.3(14.6)	114.3(7.3)	112.9(8.9)	107(14.4)
DBP (mmHg)	75.3(7.9)	65.8(7.4)	66.4(8.1)	66.4(9.7)

437 SBP=Systolic blood pressure, DBP=Diastolic blood pressure, NC=neck circumference, WHR=waist height ratio,
 438 RPR=resting pulse rate
 439

Table 2: Showing the correlation between blood pressure and anthropometric indices

Variables	r (Pearson correlation) [95% confidence interval]	p-value
------------------	---	----------------

Systolic blood pressure		
BMI (kgm ⁻²)	0.10 [0.02 - 0.18]	0.011
NC (cm)	0.50 [0.44 - 0.56]	<0.001
WHR	-0.14[-0.21-0.39]	<0.001
WHtR	0.24 [0.17 - 0.31]	<0.001
Diastolic blood pressure		
BMI (kgm ⁻²)	0.05 [0.03 - 0.12]	0.265
NC (cm)	0.32 [0.25 - 0.39]	<0.001
WHR	-0.14[-0.22-0.07]	<0.001
WHtR	0.15 [0.07 - 0.23]	<0.001


441 *BMI=Body mass index, NC= Neck circumference, WHR= Waist hip ratio, WHtR=Waist height ratio.*

442
443 Table 3: Optimal thresholds of RPR for identifying PreHTN and HTN among adolescents

Category	Threshold	PPV	NPV	Se	Sp	AUC [95% CI]
Overall PreHTN	≥76bpm	0.117	0.965	0.727	0.577	0.653[0.583-0.722]
	HTN	≥79bpm	0.077	0.989	0.737	0.719
Boys PreHTN	≥75bpm	0.167	0.975	0.722	0.608 0.629	0.685[0.558-0.811]
	HTN	≥76bpm	0.154	0.993		0.700
Girls PreHTN	≥76bpm	0.141	0.966	0.769 0.100	0.545 0.620	0.623[0.515-0.731]
	HTN	≥78bpm	0.053			0.100


444 *PreHTN: Prehypertension, HTN: Hypertension, Se: sensitivity, Sp: Specificity, PPV: positive predictive value,*
445 *NPV: Negative predictive value, AUC: area under the curve.*

446




447
448
449
450
451

Figure 1: Showing the Pearson correlation of resting pulse rate with blood pressure of the participants




452
453
454
455


Figure 2a: Showing linear regression of resting pulse rate with systolic blood pressure by sex of the participants




456
457 Figure 2b: Adjusted linear regression of resting pulse rate with systolic blood pressure by sex of the participants



458
459 Figure 3a: Showing linear regression of resting pulse rate with systolic blood pressure by sex of the participants
460




461
462 Figure 3b: Adjusted linear regression of resting pulse rate with diastolic blood pressure by sex of the participant.



463
464
465
466
467
468
469

prehtn: *prehypertension*, rprprehtn: *resting pulse rate in defining prehypertension*
Figure 4: ROC curve showing the overall prehypertension definition by resting pulse rate among adolescents



470
471
472
473
474

htn: *hypertension*, rprhtn: *resting pulse rate in defining hypertension*

Figure 5: ROC curve showing the overall AUC of resting pulse rate in defining hypertension among adolescents