

# Prevalence of mask wearing in northern Vermont in response to SARS-CoV-2

Brian Beckage<sup>1,2,3\*</sup>, Thomas Buckley<sup>4</sup>, Maegan Beckage<sup>5</sup>

**1** Department of Plant Biology, University of Vermont, Burlington, Vermont, USA

**2** Department of Computer Science, University of Vermont, Burlington, Vermont, USA

**3** Gund Institute for Environment, University of Vermont, Burlington, Vermont, USA

**4** Colchester High School, Colchester, Vermont, USA

**5** Essex High School, Essex Junction, Vermont, USA

These authors contributed equally to this work.

\* [brian.beckage@uvm.edu](mailto:brian.beckage@uvm.edu)

## Abstract

Mask wearing is integral to reducing the spread of SARS-CoV-2. Information on prevalence of face mask usage is required both to model disease spread and to improve compliance with mask usage through targeted messaging. We sought to (1) estimate the prevalence of mask usage in the most populous county of Vermont (Chittenden County; > 25% of state population) and to (2) assess the effect of age and sex on mask use. We monitored the entrances to eight different business types and visually assessed individuals' age, gender, and mask use from a distance. We collected 1004 observations from 16 May through 30 May 2020 as businesses began to reopen following an extended state-wide lock down. We analyzed these data using a Bayesian random effects logistic regression model. We found that overall 75.5% of individuals used a mask with significant effects of age and gender on mask usage. Females were more likely to wear masks than males (83.8%, n=488 vs. 67.6%, n=516 mask usage, respectively); the odds of a male wearing a mask was 53% that of the female odds. Across age groups, the elderly were most likely to wear a mask (91.4%, n=209) followed by young adults (74.8%, n=246), middle-aged adults (70.7%, n=519) and children (53.3%, n=30). The odds of an elderly person wearing a mask were 16.7 times that of a child, while the odds for young adults and middle-aged adults were ~3 times greater than a child. Highest mask usage was in elder females (96.3%, n=109) and lowest mask usage was in male children (43.8%, n=16).

## Introduction

The use of face masks can be effective at reducing the transmission of SARS-CoV-2 [1] [2]. Theoretical studies show that the extent of disease outbreaks is dependent on population compliance with mask usage and effectiveness of the masks at reducing viral transmission [3]. Simulation studies also suggest that there is a rapid phase transition from low to high prevalence of infection in populations as a function of compliance with social mitigation strategies such as mask usage. Mask usage below a threshold percentage of individuals results in widespread infection of the population [4]. Information on population compliance with mask usage is thus important to understanding the continuing dynamics of SARS-CoV-2.

Our objective in this study was i) to estimate the prevalence of mask usage in public places of business and ii) to assess the effect of gender and age on mask usage in northern Vermont. We visually assessed mask usage and estimated age and gender of individuals from a distance without any interaction with the patrons. We collected 1004 observations from outside 8 businesses and analyzed these data using a logistic regression models.

## Materials and methods

### Regional context

Vermont reported its first case of SARS-CoV-2 on 7 March 2020. The governor declared a state of emergency on 13 Mar 2020, followed by a series of orders to mitigate spread of SARS-CoV-2 [5]. These included a “Stay Home, Stay Safe” order on 24 March that directed the closure of in-person operations for all non-essential businesses and mandated that the public remain at home unless leaving for reasons critical to health or safety. On 15 May 2020, the order was updated to “Be Smart, Stay Safe”, relaxing the previous restrictions and beginning the process of reopening businesses, while urging Vermonters to socially distance and use masks in public.

Our study was carried out in Chittenden County, located in northwestern Vermont, USA. Chittenden County is the most urbanized and densely populated county in Vermont, containing 26.2% of the state population; population estimates and demographics for Chittenden County are available from the US Census Bureau [6].

### Data collection

We assessed the prevalence of mask wearing by passively observing the entrances to businesses from 16 May through 30 May 2020. that included grocery, hardware, and convenience stores, and a golf course. We recorded individuals’ mask usage as well as their estimated age and gender. Age and gender were assessed visually without interactions with the individuals. We assigned age into the following categories: child ( $\leq 14$  years old), young adult ( $>14$  and  $\leq 25$  years old), middle age adult ( $>25$  and  $\leq 60$  years old) and elderly adult ( $>60$  years old). Mask use was determined as individuals were entering the establishment since they sometimes carried the mask in their hand in the parking lot and only put on masks when entering the store. The observers were unobtrusively stationed in a vehicle near the store entrance or outside the businesses and did not interact with the observed individuals.

### Statistical analysis

We analyzed these data using logistic regression. We first selected the best-supported model of mask usage from a set of fixed-effect logistic regression models using Akaike’s Information Criterion [7]. These models were fit using the GLM function in R [8]. We then added business type as a random effect to the selected model and fit this Bayesian random-effects logistic regression using the NIMBLE library within R [9]. We did not treat business locations as the random effect but rather the business identity, e.g., all supermarkets of a given chain were treated as a single random effect. This implies that any effect of a given business is associated with the demographic utilizing that business brand (e.g., grocery store chain 1 vs. grocery store chain 2) as opposed to an effect of any particular store location. We ran 4 MCMC chains, each of length 2 million, used a burn-in length of 10,000, and thinned our chains by a factor of 100. We used Gelman and Rubin’s Convergence Diagnostic to confirm convergence of our chains [10]. We report parameter estimates and odds ratios.

## Results

Mask usage was 75.5% overall, but varied widely with gender and age group (Table 1). Mask usage for males was 67.6% compared to 83.8% for females. Mask use was highest for the elders age group with 91.4% mask use. Children had the lowest rate of compliance at 53.3%, though our sample size of children was low compared to other groups. Highest overall use of face masks was elder females with 96.3% compliance, while lowest compliance was with male children (43.8% mask use), followed by middle age males (62.5% mask use).

**Table 1. Number of people wearing mask/not wearing mask (and % mask use) by age and gender for a total of 1004 observations.**

	Male	Female	Total
Children	7/9 (43.8 )	9/5 (64.3)	16/14 (53.3)
Young adults	78/37 (67.8)	106/25 (80.9)	184/62 (74.8)
middle-aged	178/107 (62.5)	189/45 (80.8)	367/152 (70.7)
Elders	86/14 (86.0)	105/4 (96.3)	191/18 (91.4)
Total	349/167 (67.6)	409/79 (83.8)	758/246 (75.5)

The most parsimonious, fixed effects model of mask usage included age and gender but not their interaction (Table 2). The gender:age interaction term was, however, included in the second 'best' fixed effects model. Our final model, which included age and gender as fixed effects, and business type as a random effect, estimated that the odds of a male using a mask were only 53% of those of a female (Table 3). The odds of an elder wearing a mask were >16 times greater than a child and ~5 times (5.58 and 5.12, respectively) that of young or middle-aged adults, while the odds for young adults and middle-aged adults were ~3 times (2.99 and 3.26, respectively) greater than for a child.

**Table 2. Comparison of fixed effects models using AIC.**

Model	AIC	$\Delta$ AIC
1. Intercept only	1120.05	74.63
2. Gender	1085.89	40.47
3. Age	1077.57	32.15
4. Gender + Age	1045.42	0
5. Gender + Age + Gender:Age	1049.98	4.56

**Table 3. Parameter estimates of fixed gender and age effects for final model.**

	Mean	2.5%	50%	97.5%	Odds
Intercept	-0.24	-2.73	-0.22	2.11	–
Male	-0.65	-0.99	-0.65	-0.31	0.53
Young	1.00	0.15	1.00	1.85	2.99
Middle Age	1.10	0.28	1.10	1.91	3.26
Elder	2.69	1.72	2.69	3.68	16.69


Table notes: The male odds ratio is relative to females and the young, middle age, and elder age categories are relative to children. The final model was model 4 from Table 2 with a random business effect added.

There was large variation in mask usage across businesses identities (Fig 1). The

lowest mask usage was at a golf course (LF), likely because the individuals were outdoors and thus perceived less need to wear a mask, followed by a gas station convenience store (MF) (Table 4). The highest mask usage was associated with a grocery store (SH).

75  
76  
77  
78

**Fig 1. Mask usage as a function of gender and business type. Business types are defined in Table 4.**



**Table 4. Estimates of random effects of eight business types in final model.**

	Mean	2.5%	50%	97.5%	Odds
AH	0.89	-1.44	0.85	3.39	2.43
HF	0.73	-1.51	0.71	3.10	2.08
LF	-4.44	-9.95	-4.00	-1.56	0.012
MF	-1.19	-3.46	-1.21	1.16	0.30
PC	1.06	-1.20	1.04	3.45	2.90
SC	0.98	-3.06	0.63	7.22	2.67
SH	1.40	-0.86	1.36	3.80	4.04
WM	0.60	-1.66	0.58	2.99	1.83

Table notes: AH~hardware store; HF,PC,SC,SH~grocery store chains; LF~golf course; MF~convenience store with gas station; WM~department store.

## Discussion

79  
80  
81  
82

The overall mask usage of 75.5% found in our study was high compared to the 41.2% mask usage observed in grocery stores in Wisconsin over the same period [11]. Similar trends to what we observed with respect to gender and age were also found in their

study: Mask usage was higher in older adults (59.5%) and females compared to males (44.8% vs 36.9%, respectively), and lowest mask usage was among minors (26.2%). We do note that infections have continued to increase in Wisconsin [12] compared to Vermont [13], although this is just an observation rather than a claim that these divergent trends are related to differences in mask usage.

## Conclusion

Social mitigation strategies such as wearing masks in public are important to reducing spread of SARS-CoV-2. Information on compliance with use of face masks is required both for constructing models of disease spread and also for targeted messaging to increase compliance with mask usage. Studies such as this provide baseline information for assessing both demographic effects on mask usage, and also provide baseline information for regional comparisons or to assess trends in mask usage across time within a region.

## Acknowledgments

We acknowledge Prof. Jane Molofsky for reading and commenting on an earlier draft of this manuscript.

## References

1. Leung NHL, Chu DKW, Shiu EYC, Chan KH, McDevitt JJ, Hau BJP, et al. Respiratory virus shedding in exhaled breath and efficacy of face masks. *Nature Medicine*. 2020;26(5):676–680. doi:10.1038/s41591-020-0843-2.
2. Chan JFW, Yuan S, Zhang AJ, Poon VKM, Chan CCS, Lee ACY, et al. Surgical mask partition reduces the risk of non-contact transmission in a golden Syrian hamster model for Coronavirus Disease 2019 (COVID-19). *Clinical Infectious Diseases*. 2020;doi:10.1093/cid/ciaa644.
3. Kot AD. Critical levels of mask efficiency and of mask adoption that theoretically extinguish respiratory virus epidemics. *medRxiv*. 2020; p. 2020.05.09.20096644. doi:10.1101/2020.05.09.20096644.
4. Braun B, Taraktaş B, Beckage B, Molofsky J. Phase transitions and control measures for network epidemics caused by infections with presymptomatic, asymptomatic, and symptomatic stages. *arXiv:200509751 [physics]*. 2020;.
5. Novel Coronavirus (COVID-19): Vermont State Response & Resources | Office of Governor Phil Scott;. Available from: <https://governor.vermont.gov/covid19response>.
6. U S Census Bureau. QuickFacts; 2019. Available from: <https://www.census.gov/quickfacts/fact/table/VT,chittendencountyvermont/PST045219>.
7. Akaike H. A new look at the statistical model identification. *IEEE Transactions on Automatic Control*. 1974;19(6):716–723. doi:10.1109/TAC.1974.1100705.
8. R Core Team. R: A language and environment for statistical computing.; 2020. Available from: <https://www.R-project.org/>.

9. Valpine Pd, Turek D, Paciorek CJ, Anderson-Bergman C, Lang DT, Bodik R. Programming With Models: Writing Statistical Algorithms for General Model Structures With NIMBLE. *Journal of Computational and Graphical Statistics*. 2017;26(2):403–413. doi:10.1080/10618600.2016.1172487.
10. Plummer M, Best N, Cowles K, Vines K. CODA: convergence diagnosis and output analysis for MCMC. *R News*. 2006;6(1):7–11.
11. Arp NL, Nguyen TH, Graham Linck EJ, Feeney AK, Schrope JH, Ruedinger KL, et al. Use of face coverings by the public during the COVID-19 pandemic: an observational study. *Public and Global Health*; 2020. Available from: <http://medrxiv.org/lookup/doi/10.1101/2020.06.09.20126946>.
12. Timeline of COVID-19 policies, cases, and deaths in Wisconsin;. Available from: <https://coronavirus.jhu.edu/data/state-timeline/new-confirmed-cases/wisconsin>.
13. Timeline of COVID-19 policies, cases, and deaths in Vermont;. Available from: <https://coronavirus.jhu.edu/data/state-timeline/new-confirmed-cases/vermont>.