

1 Body mass index multiple regression formula testable by
2 all nine Bradford Hill causality criteria: Artificial
3 intelligence analytics applied to worldwide ecological
4 BMI and risk factor data to model obesity

5
6 David K Cundiff, MD (0000-0002-3206-9665), independent researcher,^{1 3} Chunyi Wu, PhD
7 (0000-0002-2186-3433), Research Epidemiologist/Statistician^{2 3}

8 ¹ Long Beach, California, USA

9 ² Area Specialist Lead in Epidemiology and Statistics, Michigan Medicine, Ann Arbor,
10 Michigan, USA

11 ³ Volunteer collaborators with the Institute of Health Metrics and Evaluation, Seattle,
12 Washington, USA

13 Correspondence to: David K Cundiff davidkcundiff@gmail.com Phone: 1-562-438-880

14 Word count: abstract=350, text=2984

15

16

Key Points

17 **Question:** Can worldwide global burden of disease (GBD) data analysis derive a body mass
18 index (BMI) multiple regression formula testable by Bradford Hill causality criteria?

19 **Findings:** □ A multiple regression derived formula including population attributable fractions
20 (PAFs) of 20 dietary risk factors, physical activity, childhood severe underweight, and sex
21 satisfied all nine Bradford Hill causality criteria. The BMI formula also plausibly predicted the
22 long-term BMI outcomes related to various dietary and physical activity scenarios.

23 **Meaning:** GBD data analysis of BMI and associated risk factors may infer causality with
24 overweight/obesity and possibly for health outcomes in nutritional epidemiology generally.

25

26 **Abstract**

27 **Importance** Artificial intelligence analytics may be applied to global burden of disease (GBD)

28 data on body mass index (BMI) and associated risk factors.

29 **Objective** Rigorously quantify the interactions of dietary and other risk factors that result in

30 adult BMI

31 **Design** We formatted global burden of disease data relevant to body mass index and associated

32 risk factors. We empirically explored the univariate and multiple regression correlations of BMI

33 risk factors with worldwide BMI to derive a BMI multiple regression formula (BMI formula).

34 **Setting** Worldwide

35 **Participants** Institute of Health Metrics and Evaluation staff and volunteer collaborators

36 analyzed over 12,000 GBD risk factor surveys of people from 195 countries and synthesized the

37 data into representative cohort BMI and risk factor values.

38 **Main outcome measures** the performances of the BMI formula when tested with all nine

39 Bradford Hill causality criteria each scored on a 0-5 scale: 0=negative to 5=very strong support

40 **Results** In the BMI formula derived, all foods were expressed in kilocalories/day (kcal/day). We

41 adjusted BMI formula risk factor coefficients to equate with their population attributable

42 fractions (PAFs in percent's) relating to worldwide mean BMI. BMI increasing foods had “+”

43 signs and BMI decreasing foods had “-“ signs. Total PAF of BMI formula=80.96%. BMI

44 formula=(0.37%*processed meat + 4.23%*red meat + 0.02%*fish + 2.24%*milk +

45 5.67%*poultry + 1.77%*eggs + 0.34%*alcohol + 0.99%*sugary beverages + 0.04%*corn +

46 0.72%*potatoes + 8.48%*saturated fatty acids + 3.89%*polyunsaturated fatty acids +

47 0.27%*trans fatty acids - 2.99%*fruit - 4.07%*vegetables - 0.37%*nuts and seeds -

48 0.45%*whole grains - 1.49%*legumes - 8.62%*rice - 0.10%*sweet potatoes - 7.45% physical

49 activity (METs/week) - 20.38%*child underweight + 6.02%*sex (male=1, female=2))*0.05012
50 + 21.77. (BMI formula versus BMI: $r=0.907$ (95% CI: 0.903 to 0.911) $p<0.0001$, Bradford Hill
51 causality criteria test scores (0-5): (1) strength=5, (2) experimentation=5, (3) consistency=5, (4)
52 dose-response=5, (5) temporality=5, (6) analogy=4, (7), plausibility=5, (8) specificity=5, and (9)
53 coherence=5. Total score=44/45.

54 **Conclusions and relevance** Nine Bradford Hill causality criteria strongly supported a causal
55 relationship between the BMI formula derived and mean BMIs of worldwide cohorts. The
56 artificial intelligence methodology introduced could inform individual, clinical, and public health
57 strategies regarding overweight/obesity prevention/treatment and other health outcomes.
58

59 **Introduction**

60 According to IBM Cloud Education, “At its simplest form, artificial intelligence is a
61 field, which combines computer science and robust datasets, to enable problem-
62 solving.”¹ The analysis dataset here formatted from 1.4 gigabytes of global burden of
63 disease (GBD) worldwide data and the BMI multiple regression derived risk factor
64 formula (BMI formula) satisfy this definition of artificial intelligence.

65
66 In an editorial in the BMJ titled, “Implausible results in human nutrition research”, influential
67 Stanford University meta researcher, Dr. John Ioannidis detailed why, in his words, “definitive
68 solutions won’t come from another million observational papers or small randomized trials.”² Dr.
69 Ioannidis called for radical reform of nutritional epidemiology methodologies used to influence
70 food/agricultural policies and to produce dietary guidelines for clinicians and the public.³ This
71 study analysing GBD worldwide data of dietary and other risk factors for BMI attempts to
72 answer Dr. Ioannidis’ call.

73
74 This paper will use formatted, population weighted GBD data to derive a multiple regression
75 BMI formula (BMI formula). The BMI formula modeled will be comprised of dietary and other
76 risk factors with coefficients equated to their population attributable fractions (PAFs) related to
77 the mean BMIs of worldwide cohorts.

78
79 We used Bradford Hill causality criteria to test the BMI formula. Satisfying Bradford Hill
80 causality criteria is considered validating in epidemiological research.⁴ We hypothesized that AI
81 analytics would significantly contribute to understanding the risks for obesity.

82 **Methods**

83 As volunteer collaborators with the Institute of Health Metrics and Evaluation (IHME), we
84 received raw GBD ecological data (≈ 1.4 Gigabytes) on mean BMIs of male and female cohorts
85 15-49 years old and 50-69 years old from each year 1990-2017 from 195 countries and 365
86 subnational locations ($n=1120$ cohorts). We also utilized GBD data on exposures to 32 risk
87 factors and covariates potentially related to BMI.

88
89 Food risk factors came from surveys of individuals as g/day. IHME dietary covariate data
90 originally came from Food and Agriculture Organization data on animal and plant food
91 commodities available percapita in countries worldwide—as opposed to consumption data from
92 participant interviews.⁵ Supplementary Table 1 lists the relevant GBD risk factors, covariates,
93 and other available variables with definitions of those risk factor exposures.⁶

94
95 GBD worldwide citations of over 12,000 surveys constituting ecological data inputs for this
96 analysis are available from IHME.⁵ The main characteristics of IHME GBD data sources for
97 BMI, the protocol for the GBD study, and all risk factor values have been published by IHME
98 GBD data researchers and discussed elsewhere.⁷⁻¹⁰ These included detailed descriptions of
99 categories of input data, potentially important biases, and methodologies of analysis. We did not
100 clean or pre-process any of the GBD data. GBD cohort risk factor and BMI data from the IHME
101 had no missing records. The updated 2019 raw data with variables we used for this analysis may
102 be obtained by volunteer researchers collaborating with IHME.¹¹

103

104 To maximally utilize the available data, we averaged the values for ages 15-49 years old together
105 with 50-69 years old for BMI and for each risk factor exposure for each male and female cohort
106 for each year. Finally, for each male and female cohort, data from all 28 years (1990-2017) on
107 mean BMI and on each of the risk factor exposures were averaged using the computer software
108 program R.

109
110 To weigh the country and subnational data according to population, internet searches (mostly
111 Wikipedia) yielded the most recent population estimates for countries and subnational states,
112 provinces, and regions. World population data from the World Bank and the Organisation for
113 Economic Co-operation and Development could not be used because they did not include all 195
114 countries or any subnational data.

115
116 Using the above-described formatted dataset of risk factors, covariates, and BMIs, a software
117 program in R generated a population-weighted analysis dataset. Each male or female cohort in
118 the population-weighted analysis dataset represented approximately 1 million people (range: <
119 100,000 to 1.5 million). The analysis dataset had n=7846 cohorts (rows of data), half male and
120 half female, representative of over seven billion people.

121
122 Supplementary Table 2 details how we converted omega-3 fatty acid g/day to fish g/day using
123 data on the omega-3 fatty acid content of frequently eaten fish from the National Institutes of
124 Health Office of Dietary Supplements (USA).¹² As shown in Supplementary Table 3, we
125 converted all of the animal and plant food data, including alcohol and sugary beverage
126 consumption, from g/day to kcal/day. For the g to kcal conversions, we used the Nutritionix track

127 app,¹³ which tracks types and quantities of foods consumed. Saturated fatty acids (SFA) risk
128 factor (0-1 portion of the entire diet) was not available with GBD data from 2017, so GBD SFA
129 risk factor data from 2016 was used. Polyunsaturated fatty acid (PUFA) and trans fatty acid
130 (TFA) GBD risk factor data from 2017 (0-1 portion of the entire diet) were also utilized. These
131 fatty acid data were converted to kcal/day by multiplying the fatty acid 0-1 portion of the entire
132 diet by the kcal/day available for each cohort.

133

134 **Statistical methods**

135 To determine the strengths of the risk factor correlations with mean BMIs of population
136 weighted worldwide cohorts (7846 cohorts) or subgroups of cohorts (e.g., continents, socio-
137 demographic quartiles, etc.), we utilized Pearson correlation coefficients: r , 95% confidence
138 intervals (95% CIs), and p values.

139

140 In seeking to derive the optimal BMI multiple regression formula from worldwide data, we
141 determined our methodology as we proceeded by experimenting with strategies to optimize the
142 functioning of candidate BMI formulas derived by

143

- 144 1. including as many as possible of the available dietary variables,
- 145 2. combining dietary variables if appropriate, and
- 146 3. including physical activity and other plausibly informative variables.

147

148 Appendix 1 explains the use of Bradford Hill causality criteria to assess whether the risk factors
149 in the BMI formula derived accurately modeled the worldwide mean BMI. Briefly, we tested the

150 BMI formula output with the Bradford Hill causality criteria (1) strength, (2) experimentation,
151 (3) consistency, (4) dose response, (5) temporality, (6) analogy, (7) plausibility, (8) specificity,
152 and (9) coherence. For each criterion, we used a 0-5 scale to assess the magnitude of support of
153 the BMI formula output being causally related to the BMIs worldwide (0=no support of causality
154 to 5=very strong support of causality, total possible points=45).

155
156 In determining the variables to include and exclude in worldwide BMI formula, we set the
157 statistical threshold for a variable to enter and to remain in the formula at $p < 0.25$. We used SAS
158 and SAS Studio statistical software 9.4 (SAS Institute, Cary, NC) for the data analysis.

159

160

161 **Results**

162 Table 1 shows the basic statistics and univariate correlations of mean cohort BMI worldwide
163 with mean dietary and other risk factor values. Whole grains, legumes, rice, and sweet potatoes
164 negatively correlated with BMI. We designated them, “BMI decreasing foods.” The six animal
165 foods (processed meat, red meat, fish, milk, poultry, and eggs), alcohol, sugary beverages, corn
166 availability, potato availability, added SFA, added PUFA, and added TFA all positively
167 correlated with BMI. We designated them, “BMI increasing foods.”

168
169 In Supplementary Table 4, which shows BMI risk factor to risk factor correlations, corn
170 availability (kcal/day percapita, a covariate) correlated moderately strongly with sugary
171 beverages ($r=0.419$, 95% CI 0.400 to 0.437, $p<0.0001$), suggesting that high fructose corn syrup
172 may have accounted for the unexpected positive correlation of corn with BMI.

173
174 According to the International Potato Center, potato availability (a covariate), which positively
175 correlated with BMI, included $\geq 50\%$ highly processed potato products worldwide,¹⁴ likely
176 accounting for the positive correlation of potatoes with BMI.

177
178 The strong positive correlations of fruits, vegetables, and nuts and seeds with BMI (Table 1)
179 were unexpected. These findings suggested likely multicollinearity of fruits, vegetables, and nuts
180 and seeds with BMI increasing foods (when an independent variable is highly correlated with
181 one or more known or unknown other independent variables). Supplementary Table 4 shows the
182 univariate correlations of each of the BMI formula risk factors with each other risk factor. Fruits,
183 vegetables and nuts and seeds strongly positively correlated with 10 of the 13 dietary variables

184 that positively correlated with BMI (six animal-based foods, alcohol, SFA, PUFA, and TFA).
185 Fruits, vegetables and nuts and seeds negatively correlated with the four other plant-based foods
186 (whole grains, legumes, rice, and sweet potatoes), except fruits was not significantly correlated
187 with whole grains. Together, these findings strongly suggested that multicollinearities accounted
188 for the positive correlations of fruits, vegetables, and nuts and seeds with BMI.

189
190 Supplementary Table 5 shows the Excel spreadsheet calculations, which were coordinated with
191 SAS multiple regressions involved in the derivation of the worldwide BMI formula. Appendix 2
192 detailed the steps in the BMI (dependent variable) versus BMI risk factors (independent
193 variables) multiple regression formula derivation. As seen in Supplementary Table 5, described
194 in Appendix 2, and contrasted with the current method of deriving PAFs,¹⁵⁻¹⁷ the BMI formula
195 versus BMI variance ($R^2=0.8096$) * 100 determined the total PAF (80.96%) accounted for by the
196 BMI formula. The 23 risk factor coefficients could then be equated to the PAFs of those risk
197 factors. The resulting BMI formula is shown below (all foods in kcal/day):

198
199 BMI formula = (0.37% * processed meat + 4.23% * red meat + 0.020% * fish + 2.24% *
200 milk + 5.67% * poultry + 1.77% * eggs + 0.338% * alcohol + 0.99% * sugary beverages
201 + 0.04% * corn + 0.72% * potatoes + 8.48% * saturated fatty acids + 3.89% *
202 polyunsaturated fatty acids + 0.27% * trans fatty acids - 2.99% * fruit - 4.07% *
203 vegetables - 0.37% * nuts and seeds - 0.45 * whole grains - 1.49% * legumes - 8.62% *
204 rice - 0.10% * sweet potatoes - 7.45% * physical activity (METs/week) - 20.38% * child
205 underweight (≥ 2 SD below mean BMI) + 6.02% * sex (male=1, female=2)) * 0.05012 +
206 21.77.

207

208 **BMI formula output analyzed by Bradford Hill causality criteria**

209 As mentioned in the methods and detailed in Appendix 1, the nine Bradford Hill causality
210 criteria tested the functionality of the BMI formula:

- 211 1. Strength score=5 for $r > 0.500$, $p < 0.0001$ —The BMI formula's output regressed with
212 mean BMI of worldwide cohorts gave $r = 0.907$ (95% CI: 0.903 to 0.911) $p < 0.0001$,
213 $R^2 = 0.8096$, total percent weight=80.96%.
- 214 2. Experiment score=5 for all 20 bootstrap trial BMI formulas with $r > 0.500$, $p < 0.0001$ —
215 Table 2 shows bootstrapping the BMI formula related to mean worldwide BMI with 20
216 trials (repeated sampling from the worldwide analysis dataset with replacements¹⁸). Each
217 trial had 100 randomly selected cohorts to generate a unique BMI formula. As Table 2
218 shows, the mean values for BMI and the risk factor PAFs were all quite close to the mean
219 values for BMI and BMI risk factors in the worldwide BMI formula (last column).
- 220 3. Consistency score=5 for the mean of absolute differences of BMI formula outputs and
221 mean BMI $< 0.300 \text{ kg/M}^2$ —In Table 3, we used 37 subsets of worldwide BMI and risk
222 factor data to test the consistency of BMI formula outputs, utilizing the 20 bootstrap trials
223 (#2 Experiment) to generate 80% confidence intervals for each subset. Table 3 shows the
224 average absolute difference between the 37 subgroups mean BMIs and BMI formula
225 outputs was 0.252 kg/M^2 .
- 226 4. Dose-response (Biological gradient) score=5 for the mean of absolute differences of BMI
227 formula outputs and mean BMI for the dose-response quartiles $< 0.300 \text{ kg/M}^2$ —Table 3
228 also shows that the mean of the BMI absolute differences between the BMI formula
229 estimates and mean BMIs in the four dose-response quartiles was 0.220 kg/M^2 .

230 **5.** Temporality score=5 for the BMI trend formula versus BMI trend $r>0.500$,
231 $p<0.0001$ —As is shown in Supplementary Table 6 and detailed in Appendix 1, a
232 multiple regression analysis with the worldwide BMI trend 1990-2007 (dependent
233 variable) versus 22 of the 23 risk factor trends (independent variables excluding sex)
234 generated a BMI trend formula (All foods are in Kcal/day.):

235
236 BMI trend formula= $-0.00 * \text{processed meat} + 2.52 * \text{red meat} - 0.15 * \text{fish} +$
237 $0.22 * \text{milk} + 5.46 * \text{poultry} + 1.89 * \text{eggs} + 0.07 * \text{alcohol} + 0.12 * \text{sugary}$
238 $\text{beverages} - 0.02 * \text{corn} + 0.49 * \text{potatoes} - 0.01 * \text{saturated fatty acids} + 1.01 * \text{polyunsaturated fatty acids}$
239 $- 0.01 * \text{trans fatty acids} + 3.89 * \text{fruit} + 5.92 * \text{vegetables} + 0.11 * \text{nuts and seeds} + 0.05 * \text{whole grains} - 0.88 * \text{legumes} + 3.28$
240 $* \text{rice} + 3.09 * \text{sweet potatoes} + 4.91 * \text{physical activity (METs/week)} - 0.93 * \text{child underweight} (\geq 2 \text{ SD below mean BMI}).$

241
242
243
244 As with the BMI formula, the coefficients of the BMI risk factor trends formula were
245 equated to their PAFs. BMI trend formula $r=0.592$, 95% CI: 0.577 to 0.606,
246 $p<0.0001$, and total PAF=35.02%.

247 **6.** Analogy score=4—The three metabolic risk factors correlated with BMI to test
248 analogy were systolic blood pressure (SBP), low density lipoprotein cholesterol
249 (LDL-C), and fasting plasma glucose (FPG).

- 250 • BMI correlated with SBP: $r=0.102$, 95% CI: 0.080 to 0.124, $p<0.0001$.
- 251 • BMI correlated with FPG: $r=0.558$, 95% CI: 0.542 to 0.573, $p<0.0001$.
- 252 • BMI correlated with LDL-C: $r=0.756$, 95% CI: 0.746 to 0.765, $p<0.0001$.

253 Since BMI versus SBP $r<0.500$, Bradford Hill causality score=4.

254 **7. Plausibility: Score=5**—Based on systematic medical literature reviews, physical
255 activity inversely correlated with BMI,¹⁹ and BMI directly correlated with intakes
256 of sugary beverages,²⁰ alcohol,²¹ and animal foods.²² The relationship of adult
257 BMI with early childhood severe underweight has not been reported worldwide.
258 Since people in poor countries with high infant/childhood malnutrition have fewer
259 resources to obtain animal foods, fatty acids, and alcohol and more need for
260 physical exercise than in developed countries, it is highly plausible that childhood
261 severe underweight negatively correlated with lower BMI in adulthood.

262 **8. Specificity: Score=5** for the BMI formula being unique.—The BMI formula was
263 specific to worldwide BMI and would have been different from risk factor
264 formulas modeling SBP, FPG, LDL-C, or any health outcome.

265 **9. Coherence: Score=5**—As evidenced by the near perfect score 39/40 on the first eight
266 criteria, the BMI formula accurately modeled worldwide mean BMI—total causality
267 criteria score: 44/45.

268
269 Table 5 shows BMI formula estimates for various patterns of diet and/or other BMI formula risk
270 factors. For instance, increasing physical activity by adding a run for 1 hour/day at six miles/hour
271 on average along with decreasing BMI increasing foods by isocalorically shifting 25% of BMI
272 increasing foods (Kcal/day) to BMI decreasing foods was projected to reduce the mean cohort
273 BMI from 26.66 kg/M² to mean BMI=22.26 kg/M².

274

275

276 **Discussion**

277 Appendix 2 discusses the derivation of PAFs with comparable risk assessment (CRA)
278 methodology by the World Health Organization¹⁵ and the IHME¹⁶ based on systemic literature
279 reviews. Appendix 2 details the steps in the derivation of the GBD data based BMI formula
280 (Supplemental Table 5), which precisely calculate 23 risk factor PAFs, totaling the BMI formula
281 PAF. Appendix 1 provides the metrics for validation by the nine Bradford Hill causality criteria.

282
283 A review of the literature on food costs relative to nutrient quality found that the median costs of
284 starches (€0.14/100 kcal) was quite low relative to fruits and vegetables (€0.82/100 kcal),
285 meat/eggs/fish (€0.64/100 kcal), fresh dairy (€0.32/100 kcal), and nuts (€0.25/100 kcal).²³ This
286 suggests that people who can afford fruits, vegetables and nuts and seeds are people who also
287 can afford high quantities of the BMI increasing foods—animal foods, sugary beverages, high
288 fructose corn syrup, alcohol, and highly processed foods (e.g., potato chips and fries). The
289 economics of food, including the high cost of fruits, vegetables, and nuts and seeds, may
290 contribute to overweight and obesity in the USA and in other affluent countries.

291
292 The Supplemental Nutrition Assistance Program (SNAP—formerly Food Stamps) spent an
293 estimated 22.6% of its \$73 billion/year budget²⁴ on payments to low-income Americans for
294 “sweetened beverages, prepared desserts, salty snacks, candy, and sugar.”²⁵ Additionally, the US
295 Department of Agriculture (USDA) has subsidized crops that go primarily for animal feed or that
296 are processed into sugars while not subsidizing fruits and vegetables.²⁶ While the USDA
297 recognizes the relatively low intake of fruits and vegetables in the USA and sponsors a publicity

298 campaign to increase fruits and vegetables consumption,²⁷ USDA expenditures should promote
299 reduced prices of BMI decreasing foods and increased prices of BMI increasing foods.

300
301 Following a low-carbohydrate, high-fat diet has been demonstrated to cause modest short term
302 weight loss in obese people,²⁸ However, the BMI formula projects that the long-term effect of a
303 low-carb, high fat diet would be a mean cohort BMI of 29.05 kg/M² (Table 5).

304

305 **Limitations**

306 The GBD data on animal foods, plant foods, alcohol, and fatty acids were not comprehensive and
307 comprised only 1191.4 Kcal/day on average worldwide. Subnational data were available on only
308 four countries. Because the data formatting and statistical methodology were new, this was
309 necessarily a post hoc analysis and no pre-analysis protocol or Bradford Hill causality criteria
310 scoring system was possible. As detailed in the Foresight Report on obesity,²⁹ obesity is affected
311 by a complex system of interacting factors besides diet, physical activity, and childhood feeding
312 patterns, and sex. So genes,³⁰ gut microbiome,³¹ ultra-processing of food,^{32,33} and other
313 influences on BMI were outside of the purview of this analysis.

314

315 **Generalizability**

316 Given the strength and consistency of the relationship of the mean cohort BMI with the BMI
317 formula estimates (Tables 1- 4), the findings should be generalizable to people all over the
318 world. The results could be further refined for relatively high SDI countries by deriving a BMI
319 formula from the four countries that have subnational data (UK, USA, Mexico, and Japan).

320 **Conclusion**

321 Nine Bradford Hill causality criteria strongly supported that the worldwide obesity epidemic is
322 causally related, in large part but not exclusively to 23 risk factors in proportion to their PAFs in
323 the BMI formula. The findings in this study should be considered by health policymakers
324 drafting dietary guidelines for healthy weight management. While this study dealt only with
325 dietary and other available risk factors for BMI (overweight/obesity), the AI methodology
326 introduced could easily apply to estimating PAFs of multiple dietary and other risk factors that
327 pertain to dozens of non-communicable diseases, for which the IHME have GBD data.

328

329

330

331 **Table 1. BMI risk factor basic statistics** □ (n=7846)

BMI versus BMI risk factors worldwide n=7846 □	Mean	SD	Min	Max	r	95% CI low	95% CI high	p	R ²
BMI kg/M ²	21.77	2.290	17.95	29.39					
Processed meat Kcal/day	5.334	9.720	0.20	68.77	0.5939	0.579	0.608	<.0001	0.3527
Red meat Kcal/day	50.27	45.13	3.21	235.95	0.6561	0.643	0.668	<.0001	0.4304
Fish Kcal/day	9.985	36.52	0.40	370.36	0.1021	0.080	0.124	<.0001	0.0104
Milk Kcal/day	25.04	27.05	1.06	146.8	0.6762	0.664	0.688	<.0001	0.4573
Poultry Kcal/day available	44.32	50.08	1.06	411.9	0.8086	0.801	0.816	<.0001	0.6539
Eggs Kcal/day available	19.36	14.71	0.79	69.64	0.6835	0.672	0.695	<.0001	0.4672
Alcohol Kcal/day	81.03	57.33	4.25	429.8	0.1461	0.124	0.168	<.0001	0.0213
Sugary beverages Kcal/day	298.4	152.4	72.91	1472	0.1305	0.109	0.152	<.0001	0.0170
Corn Kcal/day available	34.72	48.28	0.16	305.2	0.0760	0.054	0.098	<.0001	0.0058
Potatoes Kcal/day available	84.04	74.60	3.07	533.9	0.2095	0.188	0.231	<.0001	0.0439
Saturated fatty acids Kcal/day	190.6	63.94	70.79	481.1	0.7033	0.692	0.714	<.0001	0.4946
PUFAs Kcal/day	81.10	73.44	2.93	381.3	0.7301	0.720	0.740	<.0001	0.5330
Trans fatty acids Kcal/day	13.23	13.65	1.99	77.76	0.4749	0.458	0.492	<.0001	0.2256
Fruits Kcal/day	40.21	22.50	3.58	161.4	0.6163	0.602	0.630	<.0001	0.3798
Vegetables Kcal/day	79.76	43.12	9.48	304.2	0.5111	0.495	0.527	<.0001	0.2612
Nuts and seeds Kcal/day	8.414	8.357	0.05	103.0	0.4741	0.457	0.491	<.0001	0.2248
Whole grains Kcal/day	55.65	30.93	1.14	235.1	-0.2029	-0.224	-0.182	<.0001	0.0412
Legumes Kcal/day	51.74	32.23	0.51	194.7	-0.3836	-0.402	-0.364	<.0001	0.1471
Rice Kcal/day available	141.9	116.3	1.42	461.8	-0.5572	-0.572	-0.542	<.0001	0.3104
Sweet potatoes Kcal/day available	22.76	35.95	0.02	364.7	-0.1476	-0.169	-0.126	<.0001	0.0218
Physical activity METs	4714	1368	1609	7669	-0.4431	-0.461	-0.425	<.0001	0.1964
Child underweight by >2SD	0.1862	0.1707	0.0039	0.5300	-0.7941	-0.802	-0.786	<.0001	0.6306
Sex male 1 and female 2	1.50	0.500	1.00	2.00	0.1236	0.102	0.145	<.0001	0.0153
Total Kcal/day available	2574	418	1579	3898	0.8402	0.834	0.847	<.0001	0.7059
Stop breast feeding <6 months	0.1193	0.0555	0.0159	0.2400	0.7989	0.791	0.807	<.0001	0.6383
Sodium g/d	4.45	2.34	1.33	9.21	-0.0170	-0.039	0.005	0.1268	0.0000
Calcium g/d	0.301	0.179	0.081	1.04	0.7650	0.756	0.774	<.0001	0.5850
Dietary fiber g/d	9.21	3.15	2.72	22.68	0.3090	0.289	0.329	<.0001	0.0960
Fasting plasma glucose mmol/L	4.30	0.350	3.32	5.58	0.5570	0.542	0.572	<.0001	0.3110
LDL cholesterol mmol/L	2.35	0.400	1.27	3.25	0.7560	0.746	0.765	<.0001	0.5710
Systolic BP mm Hg	133.9	4.320	123.4	147.9	0.1020	0.080	0.124	<.0001	0.0100
Socio-demographic index (0-1)	0.543	0.174	0.112	0.900	0.7330	0.722	0.743	<.0001	0.5370

332
333 □ See Supplementary Table 1 for definitions

334 **Table 2. Bootstrap validation experiment: 20 BMI formulas each with 100 random cohorts**

Worldwide BMI and BMI risk factors	RCT1	RCT2	RCT3	RCT4	RCT5	RCT6	RCT7	RCT8	RCT9	RCT 10	RCT 11	RCT 12	RCT 13
BMI kg/M ²	21.71	22.04	21.68	21.92	21.57	21.42	21.79	22.18	21.75	21.85	21.49	21.88	21.71
Processed meat	0.33	0.32	0.34	0.34	0.28	0.31	0.32	0.39	0.57	0.36	0.21	0.26	0.28
Red meat	3.95	3.73	4.92	3.47	3.80	3.70	3.45	4.89	5.61	3.69	3.56	3.20	3.16
Fish	0.03	0.02	0.03	0.01	0.64	0.03	0.03	0.03	0.04	0.02	0.02	0.03	0.49
Milk	2.13	2.29	2.40	2.19	2.04	1.95	1.89	2.39	2.66	2.14	1.68	1.83	1.43
Poultry	6.62	6.24	6.60	4.82	5.36	4.33	5.92	6.72	5.41	4.84	4.48	6.55	4.15
Eggs	1.81	1.68	1.84	1.45	2.17	1.70	1.57	1.85	2.10	1.65	1.59	1.46	1.39
Alcohol	0.09	0.74	0.03	0.28	0.05	0.04	0.77	0.59	1.65	0.38	0.04	0.00	0.01
Sugary beverages	1.90	2.25	6.36	0.78	4.79	2.44	0.43	3.96	0.30	1.55	2.04	3.43	2.90
Corn	0.04	0.00	0.09	0.01	0.26	0.02	0.14	0.27	0.00	0.01	0.32	0.28	0.32
Potatoes	0.23	0.10	0.60	0.67	0.69	0.15	0.11	1.77	1.26	0.68	1.39	0.02	1.34
SFA	8.56	7.14	8.55	6.96	9.12	9.81	7.44	8.52	8.80	7.75	6.66	7.10	5.97
PUFA	4.30	3.97	4.31	3.16	4.09	3.59	4.09	4.12	3.96	3.66	2.84	3.30	2.84
TFA	0.33	0.29	0.23	0.23	0.26	0.14	0.29	0.30	0.38	0.26	0.19	0.35	0.14
Fruits	3.18	2.55	3.99	3.25	3.88	2.39	1.95	3.38	2.44	2.00	2.31	2.82	2.65
Vegetables	3.22	3.39	4.70	5.83	3.67	2.95	4.38	3.79	5.06	4.04	5.51	4.46	1.87
Nuts and seeds	0.22	0.32	0.51	0.30	0.55	0.24	0.36	0.59	0.60	0.25	0.07	0.13	0.13
Whole grains	0.14	0.40	0.24	0.85	0.26	0.68	0.18	0.01	0.12	0.29	1.32	0.53	0.17
Legumes	2.86	1.41	0.71	1.38	1.44	1.82	1.44	1.06	1.65	1.48	1.12	1.71	1.33
Rice	7.24	6.91	7.76	9.80	10.26	8.25	6.35	9.19	5.84	4.31	12.17	8.20	7.82
Sweet potatoes	0.05	0.29	0.12	0.20	0.04	0.07	0.22	0.12	0.17	0.17	0.02	0.30	0.05
Physical activity	6.65	10.11	3.69	9.29	0.66	5.71	14.65	2.63	3.81	10.17	7.80	7.37	6.38
Child underweight	8.86	8.41	5.17	5.47	9.68	21.13	23.33	15.63	20.69	27.27	22.28	15.50	23.20
Sex M=1, F=2	21.20	19.02	13.78	18.92	15.67	9.49	1.65	7.23	5.06	4.36	4.29	10.32	9.07
Total % weights	83.94	81.57	76.96	79.67	79.66	80.95	80.97	79.45	78.17	81.35	81.91	79.14	77.11

Worldwide BMI and BMI risk factors	RCT 14	RCT 15	RCT 16	RCT 17	RCT 18	RCT 19	RCT 20	Mean RCTs 1-20	SD	80% CI low	80% CI high	World mean n=7846
BMI kg/M ²	21.79	21.65	21.81	21.81	22.01	21.76	21.79	21.78	0.13	21.49	22.04	21.77
Processed meat	0.30	0.53	0.37	0.41	0.28	0.31	0.55	0.36	0.09	0.26	0.55	0.37
Red meat	2.65	4.77	3.57	4.55	2.47	3.49	5.71	3.95	0.84	2.65	5.61	4.23
Fish	0.02	0.02	0.01	0.03	0.02	0.56	0.70	0.12	0.23	0.01	0.64	0.02
Milk	1.84	2.29	1.79	2.51	1.58	1.80	2.50	2.10	0.33	1.58	2.51	2.24
Poultry	4.55	4.91	4.87	5.66	4.66	4.52	6.87	5.43	0.87	4.33	6.72	5.67
Eggs	1.47	1.51	1.33	2.01	1.32	1.61	2.08	1.69	0.25	1.33	2.10	1.77
Alcohol	0.01	2.10	0.66	0.67	0.01	0.19	1.32	0.46	0.56	0.01	1.65	0.34
Sugary beverages	2.58	1.31	0.79	1.11	0.99	0.61	0.29	1.90	1.55	0.30	4.79	0.99
Corn	0.37	0.00	0.54	0.11	0.22	0.35	0.00	0.15	0.16	0.00	0.37	0.04
Potatoes	0.49	0.06	1.49	0.26	0.16	1.71	0.21	0.68	0.56	0.06	1.71	0.72
SFA	6.28	7.85	6.26	9.46	5.28	5.56	10.62	7.77	1.41	5.56	9.81	8.48
PUFA	3.09	3.93	3.53	4.87	2.99	2.81	5.02	3.73	0.62	2.84	4.87	3.89
TFA	0.36	0.29	0.46	0.31	0.34	0.28	0.29	0.28	0.07	0.14	0.38	0.27
Fruits	2.93	3.55	3.77	2.45	2.58	2.01	2.41	2.88	0.61	2.00	3.88	2.99
Vegetables	2.89	3.97	2.99	2.68	3.26	4.29	1.73	3.88	1.15	1.87	5.51	4.08
Nuts and seeds	0.32	0.39	0.38	0.32	0.26	0.27	0.33	0.33	0.14	0.13	0.59	0.37
Whole grains	0.35	0.74	0.06	0.01	0.11	0.36	0.27	0.39	0.33	0.01	0.85	0.45
Legumes	0.68	1.18	0.55	2.30	0.95	0.78	1.61	1.39	0.52	0.68	2.30	1.49
Rice	10.38	6.31	11.98	3.53	5.90	10.30	9.13	8.26	2.25	4.31	11.98	8.62
Sweet potatoes	0.12	0.11	0.03	0.18	0.12	0.10	0.18	0.13	0.08	0.03	0.29	0.10
Physical activity	8.54	8.15	7.04	8.47	12.49	7.95	5.06	7.48	3.10	2.63	12.49	7.45
Child underweight	19.16	19.70	22.65	22.31	28.86	24.16	12.46	18.18	6.77	5.47	27.27	20.38
Sex M=1, F=2	10.56	6.31	5.80	7.13	6.59	6.62	10.47	9.21	5.20	4.29	19.02	6.02
Total % weights	79.92	80.01	74.68	81.34	81.42	80.63	79.83	80.74	2.17	76.96	81.91	80.96

335

336

337 **Table 3. Consistency measured with 37 subgroups of worldwide GBD data**

Subsets of worldwide data on risk factors related to BMI	Final BMI formula n	Mean BMI	BMI formula output	Absolute differences	RCT 1-20 Mean	80% CI low	80% CI high	BMI output out of 80% CI
SDI quartile 1	1926	24.486	24.496	0.010	24.455	24.138	24.738	
SDI quartile 2	2096	21.827	22.158	0.331	22.185	21.900	22.523	1
SDI quartile 3	2220	20.401	19.931	0.471	19.976	19.584	20.338	1
SDI quartile 4	1604	20.306	20.546	0.240	20.554	20.329	20.903	1
Africa	1682	21.666	21.429	0.237	21.447	21.121	21.772	
Asia	4188	20.478	20.492	0.014	20.515	20.180	20.780	
Europe	880	24.313	24.604	0.292	24.557	24.356	24.730	1
North America	558	25.912	25.494	0.418	25.477	25.016	25.810	1
Oceania	54	24.433	25.205	0.772	25.033	24.721	25.430	1
South America	468	23.463	24.128	0.665	24.222	23.788	24.594	1
Four countries with subnational data	730	24.848	24.795	0.052	24.775	24.296	25.110	
Quartile 1 BMI increasing foods	1967	24.089	24.365	0.275	24.362	24.120	24.584	1
Q2	1834	21.962	21.961	0.000	21.943	21.639	22.290	
Q3	2050	20.607	20.167	0.440	20.188	19.885	20.462	1
Q4	1995	20.494	20.682	0.188	20.738	20.446	21.066	
Quartile 1 BMI decreasing foods	1475	20.612	20.273	0.339	20.312	19.955	20.650	
Q2	2275	21.125	21.049	0.076	21.056	20.731	21.345	
Q3	2171	22.405	22.377	0.028	22.410	22.117	22.772	
Q4	1925	22.695	23.084	0.389	23.067	22.846	23.411	
Quartile 1 physical activity	2371	20.480	20.311	0.169	20.314	20.062	20.555	
Q2	1573	21.663	21.881	0.217	21.912	21.658	22.230	
Q3	1998	21.325	21.324	0.001	21.343	21.026	21.727	
Q4	1904	23.923	23.963	0.039	23.974	23.553	24.369	
Male	3923	21.485	21.420	0.065	21.388	21.138	21.566	
Female	3923	22.051	22.120	0.068	22.180	21.858	22.501	
Bootstrap trial 1	100	21.715	21.741	0.025	21.755	21.503	22.013	
Bootstrap trial 2	100	22.040	21.986	0.055	22.011	21.724	22.254	
Bootstrap trial 3	100	21.676	21.744	0.068	21.769	21.479	22.019	
Bootstrap trial 4	100	21.926	21.944	0.018	21.954	21.659	22.206	
UK	66	24.994	25.245	0.250	25.155	24.739	25.497	
USA	376	26.017	25.809	0.208	25.684	25.177	26.352	
Japan	158	21.892	22.888	0.996	22.898	22.254	23.327	1
Mexico	130	24.985	23.953	1.032	24.222	23.679	24.977	1
Quartile 1 Dose-response: BMI formula versus BMI	1928	24.763	24.949	0.186	24.936	24.617	25.230	
Q2	1469	22.568	22.480	0.088	22.525	22.187	22.805	
Q3	2647	20.872	21.069	0.197	21.069	20.884	21.325	1
Q4	1802	19.227	18.818	0.409	18.859	18.443	19.220	1
Mean of BMI - BMI formula absolute val				0.252				
Mean of BMI-BMI formula dose response				0.220				
Total out of range								13

338
339

340 **Table 4. Trends of BMI and the BMI risk factors (1990-2017)**

BMI trend versus risk factor trends, n=7846 cohorts	Mean	SD	Min	Max	R	95% CI	95% CI	P	R²
BMI trend	0.076	0.031	-0.018	0.216					
Processed meat Kcal/day trend	-0.051	0.156	-0.812	1.680	-0.009	-0.031	0.013	0.44	0.000
Red meat Kcal/day trend	0.556	1.084	-3.291	4.731	0.410	0.392	0.428	<.0001	0.168
Fish Kcal/day trend	-0.224	0.953	-0.130	9.790	-0.158	-0.180	-0.137	<.0001	0.025
Milk Kcal/day trend	0.190	0.299	-2.018	2.503	0.206	0.185	0.227	<.0001	0.043
Poultry Kcal/day trend	1.146	1.042	-0.871	7.655	0.420	0.402	0.439	<.0001	0.177
Eggs Kcal/day trend	0.300	0.492	-0.877	1.275	0.484	0.467	0.501	<.0001	0.234
Alcohol Kcal/day trend	0.031	0.164	-0.914	1.993	0.289	0.268	0.309	<.0001	0.083
Sugary beverages Kcal/day trend	0.289	1.139	-5.401	14.001	0.125	0.103	0.147	<.0001	0.016
Corn Kcal/day available trend	-0.028	0.606	-4.355	2.809	0.141	0.119	0.162	<.0001	0.020
Potatoes Kcal/day available trend	0.246	1.837	-12.59	11.031	0.272	0.251	0.292	<.0001	0.074
SFA Kcal/day trend	0.210	0.925	-3.660	2.095	-0.050	-0.072	-0.028	<.0001	0.002
PUFA Kcal/day trend	0.996	0.959	-3.026	5.473	0.286	0.266	0.306	<.0001	0.082
TFA kcal/day trend	-0.210	0.446	-2.681	0.013	-0.066	-0.088	-0.044	<.0001	0.004
Fruits Kcal/day trend	0.623	0.604	-2.078	4.104	0.481	0.464	0.498	<.0001	0.232
Vegetables Kcal/day trend	1.399	1.614	-7.961	8.465	0.396	0.377	0.415	<.0001	0.157
Nuts and seeds Kcal/day trend	0.276	0.265	-0.338	3.124	0.123	0.101	0.144	<.0001	0.015
Whole grains Kcal/day trend	0.173	0.586	-2.604	8.463	0.104	0.082	0.126	<.0001	0.011
Legumes Kcal/day trend	0.370	0.979	-3.240	5.401	-0.298	-0.318	-0.277	<.0001	0.089
Rice Kcal/day available trend	-0.787	1.292	-5.051	3.496	-0.393	-0.412	-0.374	<.0001	0.155
Sweet potatoes Kcal/day available trend	-0.460	1.339	-4.818	3.905	-0.499	-0.516	-0.482	<.0001	0.249
Physical activity METs trend	4.724	6.826	-33.86	28.629	0.379	0.360	0.398	<.0001	0.144
Child underweight >2SD trend	-0.004	0.004	-0.014	0.001	0.045	0.023	0.067	<.0001	0.002

341

342 **Table 5. BMI formula predictions with different diet or other risk factor scenarios β**

Risk factor scenario	BMI kg/M²	BMI kg/M² predicted	80 % CI low	80 % CI high
World (n=7846 cohorts mean 1990-2017)	21.77	21.77	21.33	22.13
World with all children severely underweight		16.90	15.79	18.05
World with no children severely underweight		22.88	22.35	23.38
USA	26.67	26.08	25.30	26.43
USA 25% kcal shifted to BMI decreasing foods[†]		23.41	22.31	23.93
USA physical activity mean plus 1 hour/day run		24.94	24.05	25.67
USA physical activity mean + 1 hour run and 25% shift to BMI decreasing foods[†]		22.26	21.63	23.00
USA no red or processed meat		25.36	24.67	25.72
USA no sugary beverages		26.03	25.17	26.34
USA vegetarian		24.13	23.44	24.47
USA vegan		23.47	22.67	23.97
EAT-Lancet diet		22.86	21.67	23.53
Low Carb Mediterranean Diet		29.05	27.92	29.54

343 β BMI formula estimates based on 28 years of following dietary and risk factor patterns

344 [†] Kcal/day 13 BMI increasing foods isocalorically shifted to the 7 BMI decreasing foods in the

345 BMI formula, distributed proportionally.

346

347

348 Supplementary Table 1. Definitions of IHME GBD risk factors and covariates related to BMI

Variables	Definition
Alcohol	Any alcohol consumption (g/day)
Body-mass index	Body mass index (BMI) (kg/m ²)—the dependent variable of interest
Child underweight	Proportion of children – 3 SD to – 2 SD of the WHO 2006 standard weight-for-age curve (0-1)
Corn	Corn availability percapita (g/day), a covariate
Discontinued breast feeding	Proportion of children aged 6-23 months who do not receive any breast milk
Eggs	Eggs availability percapita (g/day) a covariate
Fasting plasma glucose	Fasting plasma glucose (mmol/L)
Fish	This variable expressed in g/day was derived by determining the weight of fish in g corresponding to 1 g of omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) by averaging the fish g per 1 g of omega-3 fatty acids 20 species of fish= 117.04 g/day fish/1 g/day omega-3 fatty acids (Supplementary Table 2)
Fruits	Consumption of fruits (includes fresh, frozen, cooked, canned, or dried fruit but excludes fruit juices and salted or pickled fruits) (g/day)
Kilocalories available /day	The mean number of kilocalories percapita available per day to people in each location (kcal/day available), a covariate
LDL cholesterol	Serum low-density lipoprotein cholesterol (mmol/L)
Legumes	Consumption of beans, lentils, pulses (g/day)
Milk	Consumption of milk including non-fat, low-fat, and full-fat milk but excluding soy milk and other plant derivatives (g/day)
Nuts and seeds	Consumption of nuts and seeds (g/day)
Physical activity	Average weekly physical activity at work, home, transport-related and recreational measured by MET min per week. Less than 3000 METs per week constitutes low physical activity.
Poultry	Poultry availability percapita (g/day), a covariate
Potatoes	Potatoes availability percapita (g/day), a covariate
Processed meat	Consumption of any processed meat (includes meat preserved by smoking, curing, salting, or addition of chemical preservatives, including bacon, salami, sausages, or deli or luncheon meats like ham, turkey, and pastrami) (g/day)
Red meat	Consumption of red meat (includes beef, pork, lamb, and goat but excludes poultry, fish, eggs, and all processed meats) (g/day)
Rice	Rice availability percapita (g/day), a covariate
Seafood omega-3 fatty acids	Seafood omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) in tablet or fish form (g/day)
Socio-demographic index	SDI is a composite indicator of development status that was originally constructed for GBD 2015, and is derived from components that correlate strongly with health outcomes. It is the geometric mean for indices of the total fertility rate among women younger than 25 years, mean education for those aged 15 years or older, and lag-distributed income per capita. The resulting metric ranges from 0 to 1, with higher values corresponding to higher levels of development.
Sugar-sweetened beverages	Consumption of any beverage with ≥50 calories of sugar per one-cup serving, including carbonated beverages, sodas, energy drinks, fruit drinks but excluding 100% fruit and vegetable juices (g/day)
Sweet potatoes	Sweet potato availability percapita (g/day), a covariate
Systolic blood pressure	Systolic blood pressure (mm Hg)
Total sugar	Total sugar availability percapita (g/day), a covariate

Vegetables	Consumption of frozen, cooked, canned, or dried vegetables (including legumes but excluding salted or pickled, juices, nuts and seeds, and starchy vegetables such as potatoes or corn) (g/day)
Whole grains	Consumption of whole grains (bran, germ, and endosperm in their natural proportions) from breakfast cereals, bread, rice, pasta, biscuits, muffins, tortillas, pancakes, and others (g/day)

349

350

351 **Supplementary Table 2. Omega-3 Fatty Acid g to fish g calculation¶**

Fish	DHA g/3 ounce fish	EPA g/3 ounce fish	Omega-3 Fatty Acids (DHA + EPA) g/3 ounce fish mean	Fish 3 ounces = 85.02 g	Fish (g) per omega-3 Fatty Acids (g)=columns E/F
Salmon Atlantic farmed	1.24	0.59			
Salmon Atlantic wild	1.22	0.35			
Herring Atlantic	0.94	0.77			
Sardines canned in tomato sauce drained	0.74	0.45			
Mackerel Atlantic	0.59	0.43			
Salmon pink canned drained	0.63	0.28			
Trout rainbow wild	0.44	0.40			
Oysters eastern wild	0.23	0.30			
Sea bass	0.47	0.18			
Shrimp	0.12	0.12			
Lobster	0.07	0.10			
Tuna light canned in water drained	0.17	0.02			
Tilapia	0.11				
Scallops	0.09	0.06			
Cod Pacific	0.1	0.04			
Tuna yellowfin	0.09	0.01			
Mean DHA and EPA Omega-3 Fatty Acids g/3 ounce fish	0.4531	0.2733			
Calculations total Omega-3 FA g to fish g			0.7264	85.02	117.043

352 ¶ Data on omega-3 fatty acid content of varieties of fish came from the National Institutes of
 353 Health Office of Dietary Supplements (USA)

354
 355
 356

357 **Supplementary Table 3. Calculations of kcal/day from g/day of animal and plant foods¶**

Foods	Food sub-categories	kcal/serving	g/serving	kcal/g
Milk (2% fat)		122	244	0.5
Fish		218	170	1.28
Eggs		72	50	1.44
Poultry		187	85	2.91
Red meat		247	85	2.91
Processed meat				
	Salami	222	59	3.76
	Pastrami	104	71	1.46
	Ring baloney	86	28	3.07
	Pepperoni	94	100	0.94
Average processed meat		126.5	64.5	1.96
Fruits		97	162	0.60
Vegetables		59	91	0.65
Legumes		249	179	1.39
Nuts		172	28	6.14
Seeds				
	Flax seeds	55	10	5.5
	Chia seeds	58	12	4.83
	Fennel seeds	34.5	10	3.45
	Hemp seeds	55.3	10	5.53
Average of seeds		50.7	10.5	4.83
Average of nuts and seeds		111.4	19.25	5.78
Corn		99	103	0.96
Potatoes		161	173	0.93
Sweet potatoes		115	151	0.76
Rice		205	158	1.3
Whole grains		120	52	2.31

358 ¶ Source: NutritionIX app

359

360

361

362 **Supplementary Table 4. Pearson correlations between BMI formula risk factors from**
 363 **Table 1**

	Red meat	Fish	Milk	Poultry	Eggs	Alcohol	Sugary beverages	Corn	Potatoes	SFA	PUFA
Correlations between BMI risk factors (r)											
Processed meat KC/d	0.647	0.286	0.756	0.662	0.569	0.576	-0.108	-0.050	0.159	0.624	0.699
Red meat KC/d	1.000	0.118	0.676	0.664	0.784	0.466	0.046	-0.143	0.123	0.704	0.694
Fish KC/d	0.118	1.000	0.109	0.163	0.393	0.294	-0.163	-0.018	-0.058	0.038	0.163
Milk KC/d	0.676	0.109	1.000	0.673	0.525	0.385	0.072	-0.107	0.202	0.766	0.680
Poultry KC/d	0.664	0.163	0.673	1.000	0.628	0.351	0.142	0.011	0.102	0.701	0.837
Eggs KC/d	0.784	0.393	0.525	0.628	1.000	0.341	0.021	-0.099	0.030	0.595	0.631
Alcohol KC/d	0.466	0.294	0.385	0.351	0.341	1.000	-0.090	-0.107	-0.011	0.287	0.379
Sugary beverages KC/d	0.046	-0.163	0.072	0.142	0.021	-0.090	1.000	0.419	-0.025	0.045	0.113
Corn KC/d	-0.143	-0.018	-0.107	0.011	-0.099	-0.107	0.419	1.000	0.158	-0.133	-0.034
Potatoes KC/d	0.123	-0.058	0.202	0.102	0.030	-0.011	-0.025	0.158	1.000	0.211	0.086
SFA KC/d	0.704	0.038	0.766	0.701	0.595	0.287	0.045	-0.133	0.211	1.000	0.635
PUFAs KC/d	0.694	0.163	0.680	0.837	0.631	0.379	0.113	-0.034	0.086	0.635	1.000
TFA KC/d	0.187	-0.008	0.428	0.446	0.203	0.152	0.198	0.269	-0.091	0.233	0.433
Fruits KC/d	0.371	0.098	0.490	0.606	0.366	0.058	0.198	0.102	0.243	0.470	0.589
Vegetables KC/d	0.382	0.272	0.349	0.345	0.513	0.191	-0.167	-0.203	-0.047	0.224	0.385
Nuts and seeds KC/d	0.386	0.072	0.579	0.504	0.333	0.232	-0.146	-0.112	0.187	0.471	0.527
Whole grains KC/d	-0.112	0.091	-0.254	-0.023	-0.077	0.066	0.204	0.428	-0.058	-0.165	-0.063
Legumes KC/d	-0.404	0.061	-0.194	-0.220	-0.416	0.140	0.197	0.269	0.019	-0.389	-0.145
Rice KC/d	-0.351	-0.070	-0.565	-0.389	-0.271	-0.129	-0.148	-0.250	-0.377	-0.378	-0.368
Sweet potatoes KC/d	0.005	-0.074	-0.395	-0.240	0.049	-0.063	-0.218	-0.028	0.239	-0.130	-0.234
Physical activity METs	-0.068	-0.163	-0.430	-0.396	-0.114	0.105	-0.103	0.028	0.022	-0.309	-0.431
Child underweight	-0.736	-0.189	-0.403	-0.618	-0.803	-0.131	-0.065	-0.083	-0.215	-0.584	-0.622

	TFA	Fruits	Vegetables	Nuts-seeds	Whole grains	Legumes	Rice	Sweet potatoes	Physical activity	Child underweight	sex m/f
Correlations between BMI risk factors (r)											
Processed meat KC/d	0.463	0.379	0.290	0.504	-0.023	-0.197	-0.391	-0.194	-0.212	-0.411	-0.086
Red meat KC/d	0.187	0.371	0.382	0.386	-0.112	-0.404	-0.351	0.005	-0.068	-0.736	-0.230
Fish KC/d	-0.008	0.098	0.272	0.072	0.091	0.061	-0.070	-0.074	-0.163	-0.189	-0.023
Milk KC/d	0.428	0.490	0.349	0.579	-0.254	-0.194	-0.565	-0.395	-0.430	-0.403	-0.018
Poultry KC/d	0.446	0.606	0.345	0.504	-0.023	-0.220	-0.389	-0.240	-0.396	-0.618	0.000
Eggs KC/d	0.203	0.366	0.513	0.333	-0.077	-0.416	-0.271	0.049	-0.114	-0.803	0.000
Alcohol KC/d	0.152	0.058	0.191	0.232	0.066	0.140	-0.129	-0.063	0.105	-0.131	-0.432
Sugary beverages KC/d	0.198	0.198	-0.167	-0.146	0.204	0.197	-0.148	-0.218	-0.103	-0.065	-0.311
Corn KC/d	0.269	0.102	-0.203	-0.112	0.428	0.269	-0.250	-0.028	0.028	-0.083	0.000
Potatoes KC/d	-0.091	0.243	-0.047	0.187	-0.058	0.019	-0.377	0.239	0.022	-0.215	0.000
SFA KC/d	0.233	0.470	0.224	0.471	-0.165	-0.389	-0.378	-0.130	-0.309	-0.584	0.007
PUFAs KC/d	0.433	0.589	0.385	0.527	-0.063	-0.145	-0.368	-0.234	-0.431	-0.622	-0.013
TFA KC/d	1.000	0.387	0.284	0.312	0.087	0.044	-0.261	-0.322	-0.412	-0.121	0.096
Fruits KC/d	0.387	1.000	0.386	0.579	0.012	-0.028	-0.319	-0.096	-0.391	-0.493	0.116
Vegetables KC/d	0.284	0.386	1.000	0.547	-0.349	-0.157	-0.230	-0.095	-0.208	-0.453	-0.066
Nuts and seeds KC/d	0.312	0.579	0.547	1.000	-0.211	-0.023	-0.352	-0.074	-0.302	-0.293	-0.030
Whole grains KC/d	0.087	0.012	-0.349	-0.211	1.000	0.100	0.548	-0.039	0.268	0.089	-0.064
Legumes KC/d	0.044	-0.028	-0.157	-0.023	0.100	1.000	-0.050	-0.054	-0.068	0.506	-0.142
Rice KC/d	-0.261	-0.319	-0.230	-0.352	0.548	-0.050	1.000	0.070	0.386	0.374	0.000
Sweet potatoes KC/d	-0.322	-0.096	-0.095	-0.074	-0.039	-0.054	0.070	1.000	0.327	-0.167	0.000
Physical activity	-0.412	-0.391	-0.208	-0.302	0.268	-0.068	0.386	0.327	1.000	0.135	-0.257
Child underweight	-0.121	-0.493	-0.453	-0.293	0.089	0.506	0.374	-0.167	0.135	1.000	-0.017

364 **Red: Fruits, vegetables, and nuts and seeds correlated positively with BMI increasing foods**
 365 **Blue: Fruits, vegetables, and nuts and seeds correlated positively with themselves**
 366 **Green: Fruits, vegetables, and nuts and seeds correlated negatively with BMI decreasing foods**
 367
 368

369 **Supplementary Table 5. Derivation of the BMI formula**

370

		Column A		Column B		Column C		Column D		Column E		Column F
Row		BMI risk factors		Mean		BMI v RF R ²		Adjust for fatty acids		Multiple regression parameter estimates		Column B Column C Column D Column F
1	+	Processed meat	*	5.334	*	0.3527	*	1	*	0.0034	=	0.0064
2	+	Red meat	*	50.27	*	0.4304	*	1	*	0.0034	=	0.0736
3	+	Fish	*	9.985	*	0.0104	*	1	*	0.0034	=	0.0004
4	+	Milk	*	25.04	*	0.4573	*	1	*	0.0034	=	0.0389
5	+	Poultry available	*	44.32	*	0.6539	*	1	*	0.0034	=	0.0985
6	+	Eggs available	*	19.36	*	0.4672	*	1	*	0.0034	=	0.0308
7	+	Alcohol	*	81.03	*	0.0213	*	1	*	0.0034	=	0.0059
8	+	Sugary beverages	*	298.36	*	0.0170	*	1	*	0.0034	=	0.0173
9	+	Corn available	*	34.72	*	0.0058	*	1	*	0.0034	=	0.0007
10	+	Potatoes available	*	84.04	*	0.0439	*	1	*	0.0034	=	0.0125
11	+	SFA	*	190.58	*	0.4946	*	0.46	*	0.0034	=	0.1474
12	+	PUFAs	*	81.096	*	0.5330	*	0.46	*	0.0034	=	0.0676
13	+	TFA	*	13.23	*	0.2256	*	0.46	*	0.0034	=	0.0047
14	+	Fruits	*	40.21	*	0.3798	*	1	*	0.0034	=	0.0519
15	+	Vegetables	*	79.76	*	0.2612	*	1	*	0.0034	=	0.0708
16	+	Nuts and seeds	*	8.414	*	0.2248	*	1	*	0.0034	=	0.0064
17	-	Whole grains	*	55.65	*	0.0412	*	1	*	0.0034	=	0.0078
18	-	Legumes	*	51.74	*	0.1471	*	1	*	0.0034	=	0.0259
19	-	Rice available	*	141.86	*	0.3104	*	1	*	0.0034	=	0.1497
20	-	Sweet potatoes available	*	22.76	*	0.0218	*	1	*	0.0034	=	0.0017
21	-	Physical activity METs							*	0.12949	=	0.1295
23	-	Child underweight by >2SD							*	0.35420		0.3542
22	+	Sex male 1 and female 2							*	0.10455	=	0.1046
24		Sum of parameter estimates										1.4071
25		R ² of BMI formula										0.8585
26		R ² of BMI formula / Sum of parameter estimates										0.6101
27		F1-F23 * F27 * 100 to convert to percent weights										61.01

371

372

373 **Supplementary Table 5. Derivation of the BMI formula continued**

Row		Column G		Column H		Column I		Column J
		BMI risk factors		Un-adjusted Percent weights		BMI risk factors with fruits, vegetables, and nuts and seeds with negative signs		Percent weights (J1- J23) * 0.94299 (J27)
1	+	Processed meat	*	0.3903	+	Processed meat	*	0.3680
2	+	Red meat	*	4.4882	+	Red meat	*	4.2324
3	+	Fish	*	0.0216	+	Fish	*	0.0204
4	+	Milk	*	2.3753	+	Milk	*	2.2399
5	+	Poultry available	*	6.0115	+	Poultry available	*	5.6688
6	+	Eggs available	*	1.8764	+	Eggs available	*	1.7694
7	+	Alcohol	*	0.3585	+	Alcohol	*	0.3381
8	+	Sugary beverages	*	1.0532	+	Sugary beverages	*	0.9932
9	+	Corn available	*	0.0416	+	Corn available	*	0.0392
10	+	Potatoes available	*	0.7654	+	Potatoes available	*	0.7218
11	+	SFA	*	8.9947	+	SFA	*	8.4819
12	+	PUFAs	*	4.1245	+	PUFAs	*	3.8893
13	+	TFA	*	0.2848	+	TFA	*	0.2685
14	+	Fruits	*	3.1683	-	Fruits	*	2.9877
15	+	Vegetables	*	4.3224	-	Vegetables	*	4.0760
16	+	Nuts and seeds	*	0.3923	-	Nuts and seeds	*	0.3700
17	-	Whole grains	*	0.4751	-	Whole grains	*	0.4480
18	-	Legumes	*	1.5789	-	Legumes	*	1.4889
19	-	Rice available	*	9.1354	-	Rice available	*	8.6146
20	-	Sweet potatoes available	*	0.1029	-	Sweet potatoes available	*	0.0970
21	-	Physical activity METs	*	7.9004	-	Physical activity METs	*	7.4500
22	-	Child underweight by >2SD	*	21.6105	-	Child underweight by >2SD	*	20.3784
23	+	Sex male 1 and female 2	*	6.3788	+	Sex male 1 and female 2	*	6.0151
24		Sum of parameter estimates		85.8512		Sum of parameter estimates		80.9568
25		r of unadjusted BMI formula		0.9266		r of adjusted BMI formula		0.8998
26		R ² of unadjusted BMI formula		0.8585		R ² of adjusted BMI formula		0.8096
27						Sum percent weights (J24) / unadjusted sum percent weights (H24)		0.4299
28		H26 * 100 to convert to total percent weights=H24		85.8512		J26 * 100 to convert to percent weights=J24		80.9560

374

375

376 **Supplementary Table 6. BMI trend versus risk factor trends formula derivation (all**
 377 **variables standardized)**

		Column A		Column B		Column C		Column D		Column E		Column F
Row		BMI trend and risk factor trends (change/year)		Mean		BMI trend v RF trends R ²		Adjustment for fatty acids		Multiple regression parameter estimates		Column B * Column C * Column D * Column E
		BMI kg/M ² trend		0.07603								
1	-	Processed meat KC/d trend	*	0.0513	*	0.0001	*	1.00	*	1.1427	=	0.0000
2	+	Red meat KC/d trend	*	0.5556	*	0.1682	*	1.00	*	1.1427	=	0.1068
3	-	Fish KC/d trend	*	0.2244	*	0.0251	*	1.00	*	1.1427	=	0.0064
4	+	Milk KC/d trend	*	0.1901	*	0.0425	*	1.00	*	1.1427	=	0.0092
5	+	Poultry KC/d trend	*	1.1457	*	0.1768	*	1.00	*	1.1427	=	0.2315
6	+	Eggs KC/d trend	*	0.2996	*	0.2340	*	1.00	*	1.1427	=	0.0801
7	+	Alcohol KC/d trend	*	0.0307	*	0.0833	*	1.00	*	1.1427	=	0.0029
8	+	Sugary beverages KC/d trend	*	0.2890	*	0.0156	*	1.00	*	1.1427	=	0.0051
9	-	Corn KC/d trend	*	0.0284	*	0.0198	*	1.00	*	1.1427	=	0.0006
10	+	Potatoes KC/d trend	*	0.2461	*	0.0740	*	1.00	*	1.1427	=	0.0208
11	-	SFA KC/d trend	*	0.2101	*	0.0025	*	0.46	*	1.1427	=	0.0003
12	+	PUFA KC/d trend	*	0.9963	*	0.0820	*	0.46	*	1.1427	=	0.0429
13	+	TFA trend	*	0.2096	*	0.0044	*	0.46	*	1.1427	=	0.0005
14	+	Fruits KC/d trend	*	0.6234	*	0.2317	*	1.00	*	1.1427	=	0.1651
15	+	Vegetables KC/d trend	*	1.3992	*	0.1570	*	1.00	*	1.1427	=	0.2511
16	+	Nuts and seeds KC/d trend	*	0.2762	*	0.0150	*	1.00	*	1.1427	=	0.0047
17	+	Whole grains KC/d trend	*	0.1729	*	0.0108	*	1.00	*	1.1427	=	0.0021
18	-	Legumes KC/d trend	*	0.3696	*	0.0886	*	1.00	*	1.1427	=	0.0374
19	+	Rice KC/d trend	*	0.7869	*	0.0886	*	1.00	*	1.1427	=	0.1389
20	+	Sweet potatoes KC/d trend	*	0.4603	*	0.0886	*	1.00	*	1.1427	=	0.1311
21	+	Physical activity METs trend	*						*	0.20825	=	0.20825
22	-	Child underweight >2SD trend	*						*	0.03926	=	0.03926
23												1.48519

378
379

380 **Supplementary Table 6. BMI trend versus risk factor trends formula derivation continued**

Row		Column G		Column H		Column I		Column J
		BMI trend and risk factor trends		Preliminary risk factor coefficients (Column F repeated)		BMI trend and risk factor trends		Column H: $H_{26} * (H_1 - H_{22}) = \text{BMI trend formula percent weights}$
	+	BMI kg/M ² trend		0.0760	+	BMI kg/M ² trend		0.0760
1	-	Processed meat KC/d trend	*	0.0000	-	Processed meat KC/d trend	*	0.00
2	+	Red meat KC/d trend	*	0.1068	+	Red meat KC/d trend	*	2.52
3	-	Fish KC/d trend	*	0.0064	-	Fish KC/d trend	*	0.15
4	+	Milk KC/d trend	*	0.0092	+	Milk KC/d trend	*	0.22
5	+	Poultry KC/d trend	*	0.2315	+	Poultry KC/d trend	*	5.46
6	+	Eggs KC/d trend	*	0.0801	+	Eggs KC/d trend	*	1.89
7	+	Alcohol KC/d trend	*	0.0029	+	Alcohol KC/d trend	*	0.07
8	+	Sugary beverages KC/d trend	*	0.0051	+	Sugary beverages KC/d trend	*	0.12
9	-	Corn KC/d trend	*	0.0006	-	Corn KC/d trend	*	0.02
10	+	Potatoes KC/d trend	*	0.0208	+	Potatoes KC/d trend	*	0.49
11	-	SFA KC/d trend	*	0.0003	-	SFA KC/d trend	*	0.01
12	+	PUFA KC/d trend	*	0.0429	+	PUFA KC/d trend	*	1.01
13	+	TFA trend	*	0.0005	+	TFA trend	*	0.01
14	+	Fruits KC/d trend	*	0.1651	+	Fruits KC/d trend	*	3.89
15	+	Vegetables KC/d trend	*	0.2511	+	Vegetables KC/d trend	*	5.92
16	+	Nuts and seeds KC/d trend	*	0.0047	+	Nuts and seeds KC/d trend	*	0.11
17	+	Whole grains KC/d trend	*	0.0021	+	Whole grains KC/d trend	*	0.05
18	-	Legumes KC/d trend	*	0.0374	-	Legumes KC/d trend	*	0.88
19	+	Rice KC/d trend	*	0.1389	+	Rice KC/d trend	*	3.28
20	+	Sweet potatoes KC/d trend	*	0.1311	+	Sweet potatoes KC/d trend	*	3.09
21	+	Physical activity METs trend	*	0.20825	+	Physical activity METs trend	*	4.91
22	-	Child underweight >2SD trend	*	0.03926	-	Child underweight >2SD trend	*	0.93
23		Sum		1.48519		Total percent weights		35.02
24		BMI trend formula R ²		0.35020				
25		BMI trend formula R ² /sum		0.23579				
26		H ₂₅ * 100 for percent weights		23.57947				

381
382
383

384 **Appendix 1. Bradford Hill causality criteria based** 385 **assessment methodology for BMI formula detailed**

386 The Bradford Hill causality criteria, enumerated by the English occupational physician and
387 epidemiologist Sir Austin Bradford Hill, are the gold standard assessment tools to test causality
388 of risk factors related to health outcomes.³ The relevant causality criteria included #1 strength, #2
389 experimentation, #3 consistency, #4 dose-response (biological gradient), #5 temporality, #6
390 analogy, #7 plausibility, (8) specificity, and #9 coherence.

391
392 A literature search revealed no published methodological precedents for statistically modeling
393 and validating the relationship between mean cohort BMIs of worldwide countries and
394 subnational regions/provinces/states and their corresponding dietary and other risk factors. In
395 considering many candidate methodologies involving univariate and multiple regression
396 analysis, we sought a methodology with good performance with as many Bradford Hill causality
397 criteria as possible. Of the candidate statistical modeling strategies, the methodology for deriving
398 the BMI multiple regression formula that will be detailed in Appendix 2 functioned best. So,
399 after deriving the BMI formula, we formulated the following Bradford Hill causality criteria
400 scoring methodology for this unprecedented purpose.

401 402 **Bradford Hill causal criteria testing methodology applied to current study**

403 The nine original Bradford Hill criteria were each scored as:

- 404 • “5” very strongly supporting causality,
- 405 • “4” strongly supporting causality,

- 406 • “3” moderately strongly supporting causality,
- 407 • “2” supporting causality,
- 408 • “1” weakly supporting causality, and
- 409 • “0” not supporting causality

410

411 The scoring for each Bradford Hill causal criterion was as follows:

- 412 1. Strength: The correlation coefficient, r , of the worldwide multiple regression derived
413 BMI formula with BMI (dependent variable) and BMI risk factors (independent
414 variables) assessed strength.

415 Scoring of strength:

416 5=BMI formula regressed with BMI $r \geq 0.50$ and $p < 0.0001$

417 4=BMI formula regressed with BMI $0.50 > r \geq 0.40$ and $p < 0.0001$

418 3=BMI formula regressed with BMI $0.40 > r \geq 0.30$ and $p < 0.0001$

419 2=BMI formula regressed with BMI $0.30 > r \geq 0.20$ and $p < 0.0001$

420 1=BMI formula regressed with BMI $0.20 > r \geq 0.10$ and $p < 0.0001$

421 0=BMI formula regressed with BMI $r < 0.10$ or $p \geq 0.0001$

422

- 423 2. Experiment: Dr. Hill thought that evidence drawn from experimentation, including in
424 epidemiologic studies, may lead to the strongest support for causal inference.³ We used
425 the bootstrap method to test Hill’s “experiment” criterion. Random number generation of
426 20 subgroups each with 100 cohorts (with replacements) derived 20 standardized BMI
427 formulas to compare with the standardized worldwide BMI formula.

428 Scoring of experiment:

- 429 5=In all 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20 of BMI
430 formulas regressed with RCT1-RCT20 of BMIs, respectively: $r \geq 0.50$ and $p < 0.0001$
431 4= In at least 15 out of 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20
432 of BMI formulas regressed with RCT1-RCT20 of BMIs, respectively: $r \geq 0.50$ and
433 $p < 0.0001$.
- 434 3= In at least 15 out of 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20
435 of BMI formulas regressed with RCT1-RCT20 of BMIs, respectively: $r \geq 0.40$ and
436 $p < 0.0001$.
- 437 2= In at least 10 out of 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20
438 of BMI formulas regressed with RCT1-RCT20 of BMIs, respectively: $r \geq 0.30$ and
439 $p < 0.0001$.
- 440 1= In at least 5 out of 20 bootstrap trials each with 100 random cohorts, RCT1-RCT20 of
441 BMI formulas regressed with RCT1-RCT20 of BMIs, respectively: $r \geq 0.30$ and
442 $p < 0.0001$.
- 443 0=None of the above.
- 444 3. Consistency: For the purposes of this study, consistency between BMI and BMI formula
445 output was determined by comparing the mean BMI and the mean BMI formula output in
446 each of the following 37 subgroups:
- 447 A. We divided the world's population by quartiles of socio-demographic index (SDI)—
448 see Supplementary Table 1 for definition of SDI.
- 449 B. A variable, “continents,” allowed for analyses of the cohorts from countries from
450 each of the six inhabited continents.

- 451 C. The four countries (UK, USA, Mexico, and Japan) with subnational data on BMI and
452 the risk factors were grouped together to compare the BMI formula output with the
453 overall mean BMI in those four countries.
- 454 D. Based on the total kcal/day of the 13 foods that increased BMI (six animal foods,
455 sugary beverages, alcohol, corn, potatoes, SFA, PUFA, and TFA), a combination
456 variable was constructed and the world's population divided into quartiles from the
457 highest to lowest total kcal/day of BMI increasing foods.
- 458 E. Similarly to D above, based on the total kcal/day of all seven foods that decreased
459 BMI (fruits, vegetables, nuts and seeds, whole grains, legumes, rice, and sweet
460 potatoes), we divided the world's population into quartiles from the highest to lowest
461 in BMI decreasing foods (kcal/day).
- 462 F. Based on physical activity (METs/week), we divided the world's population into
463 quartiles from the highest to the lowest.
- 464 G. We evaluated dose response by dividing the BMI formula output into quartiles from
465 the highest to lowest.
- 466 H. The four countries with subnational data were individually evaluated.
- 467 I. The first four of the 20 random number generated database subgroups were included
468 in the consistency analysis.
- 469 J. Male and female cohorts were individually assessed.
- 470 K. We assessed dose-response with the BMI formula outputs after the BMI formula was
471 harmonized with the mean and SD of worldwide mean BMI. BMI formula outputs
472 were divided into quartiles.
- 473

474 For each of the 37 subgroups, the absolute differences between the means of BMI and the
475 BMI formula output were totaled (e.g., continent Africa BMI formula output – mean BMI
476 for Africa, etc.).

477
478 Scoring of consistency for mean cohort BMI compared with BMI formula output for each
479 of the 37 subgroups:

480 5=The mean of the absolute differences between mean BMI and BMI formula output was
481 $\leq 0.30 \text{ kg/M}^2$.

482 4=The mean of the absolute differences between mean BMI and BMI formula output was
483 $\leq 0.40 \text{ kg/M}^2$.

484 3=The mean of the absolute differences between mean BMI and BMI formula output was
485 $\leq 0.50 \text{ kg/M}^2$.

486 2=The mean of the absolute differences between mean BMI and BMI formula output was
487 $\leq 0.60 \text{ kg/M}^2$.

488 1=The mean of the absolute differences between mean BMI and BMI formula output was
489 $\leq 1.0 \text{ kg/M}^2$.

490 0=The mean of the absolute differences between mean BMI and BMI formula output was
491 $> 1.0 \text{ kg/M}^2$.

492
493 4. Dose-response (biological gradient): Dr. Hill thought that a clear dose-response effect on
494 the incidence of disease with exposure to a single risk factor was the clearest evidence of
495 a causal relationship. In this analysis levels of a multivariable regression derived BMI

496 formula outputs in quartiles were related to mean BMIs in those quartiles (Table 3.
497 subgroups #34-#37).

498 As with consistency, scoring of dose-response was based on this mean absolute
499 difference of BMI and BMI formula output when the BMI formula output was divided
500 into quartiles:

501 5= The BMI formula output versus mean BMI absolute differences from each of the four
502 quartiles averages $\leq 0.30 \text{ kg/M}^2$.

503 4= The BMI formula output versus mean BMI absolute differences from each of the four
504 quartiles averages $\leq 0.40 \text{ kg/M}^2$.

505 3= The BMI formula output versus mean BMI absolute differences from each of the four
506 quartiles averages $\leq 0.50 \text{ kg/M}^2$.

507 2= The BMI formula output versus mean BMI absolute differences from each of the four
508 quartiles averages $\leq 0.60 \text{ kg/M}^2$.

509 1= The BMI formula output versus mean BMI absolute differences from each of the four
510 quartiles averages $\leq 1.0 \text{ BMI units}$.

511 0= The BMI formula output versus mean BMI absolute differences from each of the four
512 quartiles averages $> 1.0 \text{ kg/M}^2$.

513 5. Temporality: Dr. Hill said, "Temporality refers to the necessity that the cause precedes
514 the effect in time."³ Dr. Hill was an occupational physician before the current availability
515 in nutritional epidemiology of data on trends over 28 years of 20 components of
516 worldwide diets along with the global BMI trend. Consequently, now it is fair to test
517 temporality by deriving a standardized multiple regression formula with BMI trend,
518 measured by the slope of the least squared regression line (LSRL) over 1990-2017, as the

519 dependent variable. The independent variables consisted of the LSRL trends over 1990-
520 2017 of the same dietary components, physical activity, and childhood underweight as in
521 the original BMI formula—but not sex. The signs of the risk factors in the BMI trend
522 formula were determined as follows:

- 523 a. If the sign of the risk factor in Table 4 (BMI basic trend statistics) was “+” and
524 the r was “+”, then the sign in the BMI trend formula was +.
- 525 b. If the sign of the risk factor in Table 4 was “-“ and the r was “-,” then the sign in
526 the BMI trend formula was “+”. If the risk factor mean value was trending down
527 and the correlation coefficient of the risk factor with the BMI trend was negative,
528 this risk factor trend would be in alignment with an up-trending BMI.
- 529 c. If the sign of the risk factor in Table 4 was “-“ and the r was “+,” then the sign in
530 the BMI trend formula was “-”.
- 531 d. If the sign of the risk factor in Table 4 was “+“ and the r was “-,” then the sign in
532 the BMI trend formula was “-”.

533 Supplementary Table 6 shows the derivation of the BMI trend formula with standardized
534 BMI trend (dependent variable) versus standardized risk factor trends (independent
535 variables). As with the derivation of the BMI formula (Supplementary Table 5), the
536 coefficients of the risk factors in the formula were adjusted to equate to the trend percent
537 weights.

538 Scoring of temporality: The multiple regression derived BMI trend formula output versus
539 the BMI trend r (correlation coefficient) determined the score:

540 $5=r \geq 0.50$ and $p < 0.0001$.

541 $4=0.50 > r \geq 0.40$ and $p < 0.0001$.

542 3= $0.40 > r \geq 0.30$ and $p < 0.0001$.

543 2= $0.30 > r \geq 0.20$ and $p < 0.0001$.

544 1= $0.20 > r \geq 0.10$ and $p < 0.0001$.

545 0= $r < 0.10$ or $p \geq 0.0001$

546

547 6. Analogy: High BMI was among the four metabolic risk factors that were strongly associated
548 with cardiovascular diseases, cancers, and other non-communicable diseases. The other
549 major metabolic risk factors for non-communicable diseases were high fasting plasma
550 glucose (FPG), low density lipoprotein cholesterol (LDL-C), and high systolic blood pressure
551 (SBP).

552

553 We tested analogy by the strength of the correlations of BMI with FPG, BMI with LDL-C,
554 and BMI with SBP in univariate analysis.

555 Scoring of analogy:

556 5=BMI correlated with all three other metabolic risk factors showed $r \geq 0.500$, $p < 0.0001$
557 for all three.

558 4= BMI correlated with the three other metabolic risk factors showed $r \geq 0.500$, $p < 0.0001$
559 for two of the three.

560 3= BMI correlated with the three other metabolic risk factors showed $r \geq 0.400$, $p < 0.0001$
561 for two of the three.

562 2= BMI correlated with the three other metabolic risk factors showed $r \geq 0.300$, $p < 0.0001$
563 for two of the three.

- 564 1=BMI correlated with the three other metabolic risk factors showed $r \geq 0.200$, $p < 0.0001$
565 for two of the three.
566 0= None of the above.
- 567 7. Plausibility: To test plausibility that the BMI formula could accurately predict mean BMI
568 of cohorts, we looked to find if any of our findings were at variance with the
569 preponderance of studies published. We searched the medical literature particularly for
570 systematic reviews of the relationships of foods and other variables with BMI.
571 Scoring of plausibility:
572 5=None of the BMI formula risk factor percent weight coefficients was at variance with
573 the preponderance of the medical literature about risk factors for high BMI.
574 4= One of the BMI formula risk factor percent weight coefficients was at variance with
575 the preponderance of the medical literature about risk factors for high BMI.
576 3= Two of the BMI formula risk factor percent weight coefficients were at variance with
577 the preponderance of the medical literature about risk factors for high BMI.
578 2=Three of the BMI formula risk factor percent weight coefficients were at variance with
579 the preponderance of the medical literature about risk factors for high BMI.
580 1=Four of the BMI formula risk factor percent weight coefficients were at variance with
581 the preponderance of the medical literature about risk factors for high BMI.
582 0= Five or more of the BMI formula risk factor percent weight coefficients were at
583 variance with the preponderance of the medical literature about risk factors for high BMI.
- 584 8. Specificity: Either the BMI formula derived is unique or not. Scoring of specificity:
585 5=The BMI formula derived is unique and can model no other metabolic risk factor, other
586 risk factor, or health outcome.

587 0= The BMI formula derived is not unique and can model at least one other metabolic risk
588 factor, other risk factor, or health outcome.

589 9. Coherence: According to Dr. Hill, "...cause and effect interpretation of our data should not
590 seriously conflict with the generally known facts of the natural history and biology of the
591 disease."³ In this analysis of BMI associated with BMI formula estimates, coherence was the
592 numerical total score of the above eight relevant causality criteria each on a 0-5 scale. The
593 maximum score was 40.

594 Scoring of coherence:

595 5=Score on the first eight Bradford Hill causation criteria=35-40.

596 4=Score on the first eight Bradford Hill causation criteria=30-34.

597 3=Score on the first eight Bradford Hill causation criteria=25-29.

598 2=Score on the first eight Bradford Hill causation criteria=20-24.

599 1=Score on the first eight Bradford Hill causation criteria=15-24.

600 0=Score on the first eight Bradford Hill causation criteria<15.

601

602

603

604 **Appendix 2. The derivation of the BMI formula**

605 **Population attributable fraction (PAF) derivation methodology by WHO and IHME**

606 Because of wide variations in definitions of risk factors and health outcomes in reports of risk
607 factor-health outcome correlations, the World Health Organization in 2002 published a new
608 framework, termed comparative risk assessment (CRA), for quantifying deaths and burden of
609 disease caused by risk factors.¹⁵ Goals were to increase the rigor of population health estimates
610 and to improve the compatibility of the estimates of different groups of investigators studying
611 different populations. CRA can be divided into six key steps:¹⁶

- 612
- 613 1. Inclusion of risk–outcome pairs in the analysis;
- 614 2. estimation from literature reviews of relative risk as a function of exposure;
- 615 3. estimation from literature reviews of exposure levels and distributions;
- 616 4. determination from literature reviews of the counterfactual level of exposure, the
617 level of exposure with minimum risk called the theoretical minimum risk exposure
618 level (TMREL);
- 619 5. computation of population attributable fractions (PAFs) and attributable burden; and
- 620 6. estimation from literature reviews of mediation of different risk factors through other
621 risk factors

622

623 Originally, most PAFs were calculated for single risk factors related to single health outcomes
624 (e.g., cigarette smoking and lung cancer). This methodology has now evolved to encompass 550
625 risk factor-health outcome pairs in the latest risk factor article published by the IHME.¹⁶
626 Additionally, multiple risk factors with PAFs for the same health outcome have been reported.

627 Data inputs for CRAs have depended on systematic literature reviews rather than GBD data on
628 risk factors and health outcomes.

629
630 For the purposes of risk factor-health outcome pair analysis, high BMI has been considered a risk
631 factor for cardiovascular outcomes, cancer outcomes, etc., but not itself a health outcome for
632 modeling with CRA. The methodology in this paper is offered as an alternative to CRA to model
633 PFAs for the risk factors for BMI. The case will be made that PAFs can be derived directly from
634 IHME GBD data. For validation, the BMI formula derived were tested with the nine Bradford
635 Hill causality criteria (Appendix 2).

636
637 **Population attributable fraction (PAF) derivation methodology for this analysis**

638 Supplementary Table 5 shows the BMI multiple regression formula derivation with mean cohort
639 BMI (dependent variable) versus

- 640 1. a 20 dietary risk factor combination variable,
- 641 2. physical activity,
- 642 3. child underweight, and
- 643 4. sex (independent variables).

644 To maximize the functionality of the BMI formula when tested with the nine Bradford Hill
645 causality criteria, all 20 dietary variables underwent the following four empirically derived
646 adjustments:

- 647 1. multiplied times their mean kcal/day values (Column B),
- 648 2. multiplied times their R^2 values in univariate correlation with BMI (Column C),

- 649 3. SFA, PUFA, and TFA multiplied times 0.46, an adjustment for fatty acids extracted (i.e.,
650 oils and solid fats) versus coming from within dietary foods (Column D) In determining
651 the portion of SFA, PUFA, and TFA added in addition to these fatty acids in the animal
652 and plant foods, we used an adjustment factor (fatty acids * 0.46), adapted from the
653 website “Our World in Data.”¹⁷ This adjustment factor differentiated the fatty acids in
654 individual foods (54% of the total) from the added fatty acids (46% of the total), and
655 4. the parameter estimates from the BMI versus risk factors multiple regression analysis
656 (Column F).

657

658 To derive the PAFs of each of the risk factors in the BMI formula, we used the following steps:

- 659 1. Assign the signs of the risk factors in Column A according to the signs of the risk factor
660 correlations with BMI, including leaving positive signs for fruits, vegetables, and nuts
661 and seeds.
- 662 2. Calculate the adjusted dietary risk factor coefficients (Column F (F1-F23) = Column B
663 (B1-B23) * Column C (C1-C23 * Column D (D1=D23) * Column E (E1-E23)).
- 664 3. Copy Column F coefficients (F1-F23) to Column H (H1-H23) for a further adjustment.
- 665 4. Multiply the copied coefficients in Column H (H1-H23) times the R^2 value of the entire
666 BMI formula correlated with BMI (F26=H26=0.8585) and divide the result by the sum of
667 the 23 adjusted coefficients from step 3=1.4071 (F24) to get the BMI formula adjustment
668 factor (F26=0.6101).
- 669 5. Multiply (F26=0.6101) * 100 to give a multiplier (F27=61.01) to derive PAFs for the
670 coefficients in Column H (H1-H23).

671 6. Multiply (F27=61.01) times the coefficients in Column H to get the risk factor PAFs (H1-
672 H23) and the not multicollinearity adjusted BMI formula total percent weight
673 (H24=85.85%).

674 Note that the above BMI formula contains the multicollinearities of fruits, vegetables, and
675 nuts and seeds (i.e., fruits, vegetables, and nuts and seeds correlate positively with 10 BMI
676 increasing foods and correlate negatively with the four other plant foods). Consequently, we
677 need to adjust the BMI formula to account for these multicollinearities by the following:

- 678 1. Add Column I and Column J that initially have the same BMI risk factors and
679 coefficients as Column G and Column H.
- 680 2. Switch the signs on fruits, vegetables, and nuts and seeds from “+” in Column H to
681 “-“ in Column J to adjust for the multicollinearities.
- 682 3. Run the multicollinearities adjusted BMI formula in step 2 on SAS software.
- 683 4. Check the r of the resulting multicollinearity adjusted BMI formula correlated with
684 BMI (r=0.8998 (J25)).
- 685 5. Calculate the multicollinearity adjusted BMI formula R^2 by squaring the r
686 ($R^2=0.8096$ (J26)).
- 687 6. Multiply J26=0.8096 times 100 to derive an adjusted total PAF for the final BMI
688 formula (J28=80.96%).
- 689 7. Divide the final BMI formula total percent weight (J28=80.96%) by the sum of the
690 non-collinearity adjusted BMI formula Column H coefficients (H24=85.85) to derive
691 a BMI formula coefficient multiplier (J27=0.9432) to account for the
692 multicollinearities.

- 693 8. Multiply the Column J coefficients (J1-J23) by the multicollinearity adjusting BMI
694 formula coefficient multiplier (J27=0.9432) to derive the final BMI formula
695 coefficients (J1-J23) as PAFs, totaling the final BMI formula total percent weight
696 (J24=80.96%).
- 697 9. Test the validity of the BMI formula with the nine Bradford Hill causality criteria
698 (Appendix 1).

699

700

701

702

703

704 **Article Information**

705 **Corresponding Author:** David K. Cundiff, Independent researcher,

706 davidkcundiff@gmail.com

707• **Authors' contributions:** DKC acts as guarantor; conceived and designed the study, acquired
708 and analysed the data, interpreted the study findings, drafted the manuscript, critically reviewed
709 and edited the manuscript and tables, and approved the final version for publication.

710

711 CW designed software programs in R to format and population weight the data,
712 aided with the SAS statistical analysis, critically reviewed the manuscript, and
713 approved the final version for publication.

714

715• **Conflict of Interest Disclosures:** None reported. Both authors have completed the ICMJE
716 uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any
717 organisation for the submitted work; no financial relationships with any organisations that might
718 have an interest in the submitted work in the previous three years; no other relationships or
719 activities that could appear to have influenced the submitted work.

720

721• **Funding/Support:** This research received no specific grant from any funding agency in the
722 public, commercial or not-for-profit sectors. The Bill and Melinda Gates Foundation funded the
723 acquisition of the data for this analysis by the IHME. The data were provided to the authors as
724 volunteer collaborators with IHME.

725•

726

727 **Role of the Funder/Sponsor:** While IHME GBD faculty and staff by virtue of the Bill and
728 Melinda Gates Foundation grants provided the raw data for this analysis, they did not vet the
729 analysis or sponsor the manuscript.

730
731 **Additional Contributors:** Martin Sebera, from the Department of Kinesiology, Faculty of
732 Sports Studies Masaryk University, Czech Republic, critiqued statistical aspects of the
733 manuscript and provided useful input. Pavel Grasgruber, from Masaryk University, Czech
734 Republic, provided suggestions after reviewing the manuscript. We thank Scott Glenn and Brent
735 Bell from IHME who supplied us with the GBD risk factor exposure data for the risk factors and
736 for BMI data.

737
738 **Data sharing statement:** The raw, unformatted data used in this analysis is now out of date. The
739 2019 GBD data on all the variables in this analysis may be obtained from the IHME by volunteer
740 collaborating researchers. The formatted database, SAS codes and Excel spreadsheets on which
741 this analysis is based are posted on the Mendeley data repository:

742 <https://data.mendeley.com/v1/datasets/publish-confirmation/g6b39zxck4/6>

743
744 **STROBE checklist:** This report follows the STrengthening the Reporting of OBservational
745 studies in Epidemiology (STROBE) guidelines for reporting global health estimates.³⁴

746

747

748

749

750 References

- 751 1. Artificial Intelligence (AI). IBM Cloud Education. Accessed August 7, 2021,
752 <https://www.ibm.com/cloud/learn/what-is-artificial-intelligence>
- 753 2. Ioannidis JPA. Implausible results in human nutrition research. *BMJ : British Medical*
754 *Journal*. 2013;347doi:10.1136/bmj.f6698
- 755 3. Ioannidis JA. The challenge of reforming nutritional epidemiologic research. *JAMA*.
756 2018;320(10):969-970. doi:10.1001/jama.2018.11025
- 757 4. Hill AB. THE ENVIRONMENT AND DISEASE: ASSOCIATION OR CAUSATION?
758 *Proc R Soc Med*. 1965;58(5):295-300.
- 759 5. Global Burden of Disease Study 2017 (GBD 2017) Data Input Sources Tool. Institute of
760 Health Metrics and Evaluation. Updated 2019. Accessed August 10, 2019,
761 <http://ghdx.healthdata.org/gbd-2017/data-input-sources>
- 762 6. GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk
763 assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of
764 risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of
765 Disease Study 2017: Table of risk factor definitions. *The Lancet*. 2018;392(10159):1923-94.
- 766 7. PROTOCOL FOR THE GLOBAL BURDEN OF DISEASES, INJURIES, AND RISK
767 FACTORS STUDY (GBD) Version 3.0; Issue 26. Institute for Health Metrics and Evaluation.
768 Updated February 2018. Accessed August 2, 2019,
769 http://www.healthdata.org/sites/default/files/files/Projects/GBD/GBD_Protocol.pdf
- 770 8. Naghavi M, Abajobir AA, Abbafati C, et al. Global, regional, and national age-sex
771 specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global
772 Burden of Disease Study 2016. *The Lancet*. 2017;390 (10100):1151-1210.
- 773 9. Flaxman AD, Lee YY, Vos T, et al. *An Integrative Metaregression Framework for*
774 *Descriptive Epidemiology*. University of Washington Press; 2015.
775 <http://www.guilles.website/wp-content/uploads/2017/11/Metaregression-AFlaxman.pdf>
- 776 10. GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk
777 assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of
778 risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of
779 Disease Study 2017. . *The Lancet*. 2018;392(10159):1923-94.
- 780 11. Call for Collaborators. . Institute for Health Metrics and Evaluation at the University of
781 Washington. Updated 2019. Accessed April 25, 2019, [http://www.healthdata.org/gbd/call-for-](http://www.healthdata.org/gbd/call-for-collaborators)
782 [collaborators](http://www.healthdata.org/gbd/call-for-collaborators)
- 783 12. Omega 3 Fatty Acids: Fact Sheet for Health Professionals (The Office of Dietary
784 Supplements) (2018).
- 785 13. Nutritionix Track App. Syndigo LLC Updated 2019. Accessed April 25, 2019,
786 <https://www.nutritionix.com/>
- 787 14. POTATO PROCESSING AND USES. International Potato Center. Updated 2020.
788 Accessed May 28, 2020, <https://cipotato.org/potato/potato-processing-uses/>
- 789 15. Murray C, Lopez A, Rodgers A, et al. The World Health Report 2002: Reducing Risks,
790 Promoting Healthy Life. World Health Organization.
791 https://apps.who.int/iris/bitstream/handle/10665/42510/WHR_2002.pdf
- 792 16. Murray CJL, Aravkin AY, Zheng P, et al. Global burden of 87 risk factors in 204
793 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease
794 Study 2019. *The Lancet*. 2020;396(10258):1223-1249. doi:10.1016/S0140-6736(20)30752-2

- 795 17. Ritchie H, Roser M. Diet Compositions. *Our World in Data*. 2017;
- 796 18. Bloom JOJ. Bootstrap confidence intervals (Class 24, 18.05). MIT. Accessed July 25,
- 797 2021, [https://ocw.mit.edu/courses/mathematics/18-05-introduction-to-probability-and-statistics-](https://ocw.mit.edu/courses/mathematics/18-05-introduction-to-probability-and-statistics-spring-2014/readings/MIT18_05S14_Reading24.pdf)
- 798 [spring-2014/readings/MIT18_05S14_Reading24.pdf](https://ocw.mit.edu/courses/mathematics/18-05-introduction-to-probability-and-statistics-spring-2014/readings/MIT18_05S14_Reading24.pdf)
- 799 19. Ruotsalainen H, Kyngäs H, Tammelin T, Kääriäinen M. Systematic review of physical
- 800 activity and exercise interventions on body mass indices, subsequent physical activity and
- 801 psychological symptoms in overweight and obese adolescents. *Journal of Advanced Nursing*.
- 802 2015;71(11):2461-2477. doi:10.1111/jan.12696
- 803 20. Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review
- 804 and meta-analyses of randomised controlled trials and cohort studies. *BMJ* 2013;346:e7492.
- 805 doi:10.1136/bmj.e7492
- 806 21. Traversy G, Chaput J-P. Alcohol Consumption and Obesity: An Update. *Curr Obes Rep*.
- 807 2015;4(1):122-130. doi:10.1007/s13679-014-0129-4
- 808 22. Turner-McGrievy G, Mandes T, Crimarco A. A plant-based diet for overweight and
- 809 obesity prevention and treatment. *J Geriatr Cardiol*. 2017;14(5):369-374.
- 810 doi:10.11909/j.issn.1671-5411.2017.05.002
- 811 23. Darmon N, Drewnowski A. Contribution of food prices and diet cost to socioeconomic
- 812 disparities in diet quality and health: a systematic review and analysis. *Nutrition reviews*.
- 813 2015;73(10):643-660. doi:10.1093/nutrit/nuv027
- 814 24. United States Department of Agriculture FY 2019 BUDGET SUMMARY (US Office of
- 815 Budget and Program Analysis) (2019).
- 816 25. Why is Government Subsidizing Junk Food?--22.6 percent of SNAP dollars are spent on
- 817 unhealthy food. Foundation for Economic Education. Updated February 25, 2017. Accessed
- 818 August 11, 2019, <https://fee.org/articles/why-is-government-subsidizing-junk-food/>
- 819 26. Wakamo B. We Subsidize the Wrong Kind of Agriculture. Inequality.org. Updated JUNE
- 820 21, 2018. <https://inequality.org/great-divide/subsidize-wrong-kind-agriculture/>
- 821 27. Fruits and Veggies Now Chock Full of Marketing Power (2017).
- 822 28. Ebbeling CB, Feldman HA, Klein GL, et al. Effects of a low carbohydrate diet on energy
- 823 expenditure during weight loss maintenance: randomized trial. *BMJ*. Nov 14 2018;363:k4583.
- 824 29. FORESIGHT--Tackling Obesities: Future Choices –Follow Up Actions. Government
- 825 Office for Science (UK). Updated 2019. Accessed August 5, 2019,
- 826 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/295885/07-1526-obesity-action-plan.pdf)
- 827 [/295885/07-1526-obesity-action-plan.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/295885/07-1526-obesity-action-plan.pdf)
- 828 30. Mărginean CO, Mărginean C, Meliț LE. New Insights Regarding Genetic Aspects of
- 829 Childhood Obesity: A Minireview. *Front Pediatr*. 2018;6:271-271.
- 830 doi:10.3389/fped.2018.00271
- 831 31. Castaner O, Goday A, Park Y-M, et al. The Gut Microbiome Profile in Obesity: A
- 832 Systematic Review. *Int J Endocrinol*. 2018;2018:4095789-4095789. doi:10.1155/2018/4095789
- 833 32. Monteiro CA, Moubarac J-C, Levy RB, Canella DS, Louzada MLdC, Cannon G.
- 834 Household availability of ultra-processed foods and obesity in nineteen European countries.
- 835 *Public Health Nutrition*. 2017;21(1):18-26. doi:10.1017/S1368980017001379
- 836 33. Monteiro CA, Cannon G, Levy RB, et al. Ultra-processed foods: what they are and how
- 837 to identify them. *Public Health Nutrition*. 2019;22(5):936-941.
- 838 doi:10.1017/S1368980018003762
- 839 34. Stevens GA, Alkema L, Black PRE, et al. Guidelines for Accurate and Transparent
- 840 Health Estimates Reporting: the GATHER statement. *The Lancet*. 2016;388(10062):e19-e23.

841