

1 **Access to HIV-prevention in female sex workers in Ukraine between 2009 and 2017:**
2 **coverage, barriers and facilitators**

3 Natasha Blumer^{1#a,2}, Lisa M. Pfadenhauer^{1,2}, Jacob Burns^{1,2}

4
5 ¹ Institute for Medical Information Processing, Biometry and Epidemiology - IBE

6 Ludwig-Maximilians-Universität München, Munich, Germany

7 ² Pettenkofer School of Public Health Munich, Munich, Germany

8
9 ^{#a} Current address : Institute for Medical Information Processing, Biometry and Epidemiology

10 - IBE, Chair of Public Health and Health Services Research Pettenkofer School of Public

11 Health (PSPH), LMU Munich, Marchioninstrasse 17, 81377 Munich, Germany

12 **Abstract**

13 The provision of comprehensive prevention services is vital for reducing the high
14 burden of HIV amongst Ukrainian female sex workers (FSWs). To identify barriers and
15 facilitators that influence access to HIV prevention amongst this population between 2009
16 and 2017, we developed a literature-informed conceptual framework and conducted a
17 document analysis to identify the components of the Ukrainian prevention package (PP).
18 Using the Integrated Bio Behavioural Surveillance Surveys, we then conducted descriptive
19 analyses to explore PP coverage from 2009 to 2017 and the influence of factors, identified by
20 our conceptual framework. After increasing over four years, a drop in PP coverage was
21 observed from 2013 onwards. Being a client of a non-governmental organisation, street and
22 highway solicitation, non-condom use, and knowledge of HIV may influence access to HIV
23 prevention in the Ukrainian context. Future interventions should consider barriers and
24 facilitators to HIV prevention and the multiple structural levels on which they operate.

25 **Keywords:** HIV, health services accessibility, sex worker, Ukraine, cross sectional survey

26

27 Introduction

28 Although the global HIV incidence rate has decreased, Ukraine has developed one of
29 the most severe HIV epidemics in Europe and Central Asia [1]. As in most countries, the HIV
30 epidemic in Ukraine is concentrated amongst populations that are at heightened risk of
31 contraction, designated 'key populations'. These include people who inject drugs, men who
32 have sex with men, transgender people, incarcerated persons and sex workers (SWs) [2-4].

33 Globally, female sex workers (FSWs) bear a disproportionately high burden of the HIV
34 pandemic, with a global prevalence of over 10% [5]. FSWs are 13.5 times more likely to
35 contract HIV than of women of the same reproductive age [6]. In Ukraine, the annual
36 incidence rate lies at 0.56 per 1000 uninfected SWs and the prevalence at 5.2% within an
37 estimated population of 80.000 SWs [3]. An array of determinants increase the vulnerability
38 of a FSW to HIV, including gender-based violence (GBV) [7, 8], poverty [9, 10], multiple
39 sexual partners [11], criminalisation [12-15], stigma [16, 17] and substance abuse [18].
40 Recent literature has posited that determinants of HIV operate on different structural levels,
41 ranging from the macrostructural to the individual level [13, 14].

42 Although a multitude of prevention interventions have been designed to reduce the impact of
43 HIV amongst key populations, it has been established that no single prevention intervention
44 is sufficient to contain the spread of the disease and that a multi-pronged, multi-level
45 approach is necessary [19-21]. In 2014, the World Health Organisation (WHO) published
46 guidelines consolidating all existing recommendations relevant to the prevention of HIV
47 amongst key populations, including SWs. These guidelines outline the components of a
48 comprehensive, evidence-based package of interventions, which has been found to be
49 acceptable and effective for the global SW community. The package has two parts: essential
50 health care sector interventions and essential strategies for an enabling environment. Health
51 care interventions include components such as comprehensive condom and lubricant
52 provision and behavioural interventions (e.g. consultation with social workers), whilst
53 strategies addressing community empowerment and addressing violence fall beneath the

54 enabling environment section [4]. As one of the five prevention pillars, the United Nations
55 General Assembly prevention targets emphasised the importance of this comprehensive
56 prevention package (PP), by urging countries to ensure that 90% of people at risk of HIV
57 infection have access to the PP by 2020 [22]. The Alliance of Public Health (Alliance) has
58 been distributing HIV prevention amongst key populations in Ukraine since 2003,, which was
59 first coined the PP in 2007, has since been scaled up countrywide [23].

60 Despite these efforts, only 46% and 48% of Ukrainian SWs accessed the PP in 2015 and
61 2017 respectively [24, 25]. Recent literature has identified that FSWs encounter barriers on
62 multiple structural levels, which impede their access to HIV prevention services [21, 26]. In
63 order to improve the intervention and its implementation in Ukraine, further investigation is
64 necessary to determine the factors that hinder FSWs from accessing the PP.

65 The aim of this study was to therefore explore which factors influenced the access that
66 Ukrainian FSWs had to the PP between 2009 and 2017. More specifically, our four research
67 objectives were 1) to develop a conceptual framework illustrating the factors that influence
68 FSWs access to prevention and the structural levels they operate on 2) to identify which
69 components were included in the Ukrainian PP for SWs, 3) to measure the coverage of the
70 PP in Ukraine amongst the FSW population and finally, 4) to describe and explore the factors
71 associated with the access of Ukrainian FSW to the PP over this nine-year period.

72 **Method**

73 We undertook multiple steps to address our four research objectives, as outlined in
74 Fig 1. We began by developing a conceptual framework, informed by a literature review
75 (objective 1). We then conducted a document analysis, which served to describe the
76 Ukrainian PP and its components (objective 2). The questionnaires from five Ukrainian
77 Integrated Bio Behavioural Surveillance Surveys (IBBS) were then reviewed to identify item
78 that corresponded to both the factors identified in the conceptual framework and the
79 components identified in the document analysis. These then served as our independent

80 variables and outcome variables, respectively and were used to conduct exploratory
81 analyses. This included calculating descriptive statistics to measure the coverage of the PP
82 amongst the Ukrainian FSW population (objective 3) and a descriptive longitudinal
83 comparison between FSWs that did and did not access prevention, to explore which factors
84 influenced access to PP between 2009 and 2017 (objective 4). As no information about the
85 gender of Ukrainian SWs is available in the IBBS data prior to 2015 and that in 2017, 97.72%
86 of Ukrainian SWs were female (2.14% male and 0.14% transgender), we have referred to
87 SWs as FSWs when referring to IBBS data, unless stated otherwise.

88

89 **Fig 1. Flowchart of methodological approach.**

90

91 **Conceptual Framework**

92 In an iterative process, NB thematically grouped specific factors associated with access to
93 HIV prevention, which were identified in a literature review. The literature review method,
94 corresponding protocol, and results are available in the supporting information, S1.
95 Uncertainties about attributing factors to a certain thematic cluster were discussed with JB
96 and LMP on an ongoing basis. To structure these thematic clusters, we adopted the
97 framework structure developed by Shannon and colleagues, which visualises the
98 determinants of HIV for the global population of FSWs. This framework identifies six
99 structural levels that HIV determinants operate on, including the macrostructural, sex worker
100 community organisation, work environment, sexual networks and patterning, sex worker and
101 client individual level [13, 14]. As our literature review identified that factors associated with
102 the access to HIV prevention also operate on different structural levels, we considered the
103 structure of this framework appropriate and populated it with our thematic clusters
104 accordingly (Table in S2 Table).

105 **Document Analysis and Description of the Ukrainian PP**

106 Our second research objective aimed to identify components offered as part of the
107 Ukrainian PP between 2008 and 2017. To do so, we conducted a document analysis of all
108 annual general reports published by Alliance over this time period [24, 25, 27-34]. With
109 substantial financial support by of the Global Fund to Fight AIDS, Tuberculosis and Malaria
110 (TGF), Alliance has been responsible for the distribution of HIV prevention amongst SWs in
111 Ukraine since 2004, and each report provides a detailed list of components included in the
112 PP [35]. These components were extracted from the English version of each annual report.
113 After extraction, we grouped the components of similar nature into categories to gain a
114 clearer picture of the composition of the PP over the years. For a detailed list of the
115 categories and associated components, see table in S3 Table.

116 **Review of IBBS questionnaires and variable selection**

117 We reviewed the questionnaires of five Integrated Bio Behavioural Surveillance
118 Surveys (IBBS) to identify items that corresponded to both our outcome and independent
119 variables, which were drawn from the results of the conceptual framework and document
120 review, respectively. The IBBS are cross-sectional studies that were administered amongst
121 Ukrainian SWs in 15 cities in 2009, 25 cities in 2011, 27 cities in 2013 and 2015 and 32 cities
122 in 2017. The IBBS were developed, piloted and administrated by Alliance under the financial
123 support of the TGF in 2009, 2011 and 2013 and then from the The U.S. Centers for Disease
124 Control and Prevention (CDC) in 2015 and 2017. The Public Health Centre (PHC) of the
125 Ukrainian Ministry of Health are now the holders of the data. Ethical approval was granted by
126 the Institutional Review Board (IRB) at the Ukrainian Institute on Public Health Policy (Kiev,
127 Ukraine) and CDC. The surveys were administered as face-to-face interviews lasting
128 approximately thirty minutes. The interview was followed by a voluntary rapid HIV tests,
129 details of rapid tests can be found in the Alliance analytic reports [25, 28, 30, 32, 34].

130 Participants were considered eligible if they were fourteen or older, had received money,
131 goods or services in exchange for sex within the last six months and consented to participate
132 in the study. Three sampling strategies were employed for recruiting participants:
133 respondent-driven-sampling (RDS), time-location-sampling (TLS) and key-informant
134 sampling was employed in 2013, no follow up was conducted. Recruitment continued until
135 sample size was reached. Sample size was calculated for each study site using the specific
136 HIV prevalence rate as determined by the previous IBBS result. Further details about the
137 study design of each year can be found in the analytic reports of Alliance [25, 28, 30, 32, 34].
138 All five IBBS surveys were translated from Ukrainian into English by an associate professor
139 of Slavic languages at the Donetsk National Technical University.

140 **Independent Variables**

141 To identify our independent variables, namely, *factors associated with access to HIV*
142 *prevention*, we drew upon our conceptual framework (objective 1) and then screened each
143 IBBS questionnaire for a corresponding item. The single most fitting item was selected for
144 each independent variable.

145 We support the emphasis that Shannon and colleagues place upon factors that operate at
146 the macrostructural level when designing prevention policies and schemas. As all IBBS
147 questionnaires measure individual behaviour, we were unable to extract items operating on
148 the macrostructural level, and hence, resorted to designating the most fitting item at the
149 individual level to these macrostructural factors. The exact item used for each independent
150 variable can be found in the table in S5 table.

151 **Outcome Variable**

152 *Receipt of the Ukrainian PP* was our outcome of interest; hence we drew upon the
153 findings of our document analysis. As there is no universally accepted way to measure PP

154 receipt, we created two definitions, including “basic PP receipt” and “extended PP receipt”.
155 We initially used Alliance’s definition for basic PP receipt which consists of “the receipt of at
156 least one consultation from a social worker and the receipt of at least one male condom over
157 a six month period” [36]. However, due to limitations in the IBBS data, we altered this to “the
158 receipt of at least one male condom and any form of HIV testing over a 12-month period”.

159 For extended PP receipt, we adopted the Joint United Nations Programme on HIV/AIDS’
160 (UNAIDS) [37] definition. According to UNAIDS, PP receipt is defined as, “access to at least
161 two HIV prevention interventions within the last three months”. As the time frame referenced
162 in many IBBS question was twelve months, we adapted this definition by increasing the
163 three-month period to twelve months. Thus, we defined extended PP receipt as, “access to
164 at least two HIV prevention interventions within the last twelve months”. The number and
165 specific type of prevention services included in each year’s extended PP, alongside the
166 corresponding IBBS item, can be found in the table in S4 Table.

167

168 **Exploratory Analyses: PP Coverage and Descriptive**

169 **Longitudinal Group Comparison**

170 We conducted descriptive statistics to measure the coverage of the basic and
171 extended PP receipt between 2009 and 2017 (objective 3). We did so by calculating the
172 proportion of FSWs that received the PP per year according to the definition for basic PP
173 receipt and extended PP receipt. These results enabled us to undertake a descriptive
174 longitudinal group comparison to compare the FSWs who did and did not receive the PP,
175 with regard to the various independent variables (objective 4). Specifically, we calculated
176 proportions for binary variables and means for continuous variables, stratified by those who
177 received the basic PP and the extended PP and those who did not (table 3). All variables
178 were screened for missing data before undergoing descriptive analyses. For all but one

179 variable assessed, the proportion of missing data was low, ranging from 0% to 6.7%. For the
180 item regarding HIV status, a large proportion of the data, 54.3% was missing. We excluded
181 missing data from the analysis. All analyses were conducted using the statistical software R
182 (3.6.0) [38].

183 **Results**

184 **Conceptual Framework (objective 1)**

185 Many factors associated with access to HIV prevention were presented in multiple
186 studies, identified in our literature review. As these factors were largely aligned, we clustered
187 factors with similar themes to better structure our conceptual framework. For example, three
188 specific factors found to influence access to HIV prevention – knowledge of where to access
189 prevention [9], knowledge of HIV transmission routes [39-41] and the self-perception of being
190 at high risk of HIV infection [42] – were grouped into one broader thematic cluster,
191 knowledge of HIV. After discussing the conceptual framework with staff members from the
192 PHC, one thematic cluster, sex worker- family was excluded from the conceptual framework,
193 as the specific factors were applicable to a generalised epidemic (HIV prevalence of $\geq 1\%$
194 amongst the general population) and were not applicable in the Ukrainian context, where the
195 epidemic is classified as a concentrated or mixed epidemic [43]. Our conceptual framework
196 illustrates that most thematic clusters which influence access to HIV prevention operate at
197 the macrostructural level, as seen in Fig 2.

198

199 **Fig 2. Conceptual framework of factors and structural levels influencing access to HIV**
200 **prevention.**

201

202 Document Analysis and Description of the Ukrainian PP

203 (objective 2)

204 A total of 32 categories describing prevention components emerged from the
205 document analysis of the Alliance annual general reports from 2008 to 2017 (see Table1).
206 Many components included in the PPs could be grouped into broader categories. For
207 example, testing for chlamydia, assisted testing for syphilis and examination for sexually
208 transmitted diseases (STI) were coded as the overarching category, 'any form of STI testing'.
209 A list of specific components within each overarching prevention category can be found in S3
210 in the supporting information.

211 The number of components included in the Ukrainian PP increased from 2008 until 2013,
212 when the highest number of components (n=24) was included in both 2012 and 2013. A
213 steep drop was observed in 2014, with only seven components included. Between 2015 and
214 2017, eleven components were included in the PP.

215 The four prevention components 'any form of social worker counselling', 'male condoms',
216 'awareness raising materials' and 'HIV testing and counselling' were the most commonly
217 delivered components in the Ukrainian PP across all years. Lubricants, 'any form of STI
218 testing' and 'referrals to relevant specialists' were also consistently included in the prevention
219 package in all years, except for 2014. 'Overdose prevention' was only included once in 2012,
220 while 'online counselling', 'training about safe behaviour', 'case management', 'humanitarian
221 aid' and 'tea and meals' were only included twice. The level of coverage of each component
222 may vary based on region.

223

Table 1: Overview of components of the PP provided by Alliance from 2008 to 2017

224

Component	Year	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	Sum
Any form of counselling from a social worker												10
Male condoms												10
Awareness raising and educational material												10
Any form of HIV testing and counselling												10
Lubricants												9
Any form of STI testing												9
Referrals to relevant specialists												9
Some form of peer counselling												8
Any form of specialist medical or legal counselling												6
Any form of hepatitis B testing												6
Antiseptics												6
Pregnancy tests												6
Organisation of leisure activities												5
Childcare and services												5
Cosmetologist and hairdresser services												5
Any form of professional/ skilled work training												5
Participation in group work												4
Female condoms and training for usage												4
Some form of TB screening												4
Counteracting violence												4
Any form of hepatitis C testing												3
Referral to opioid substitution therapy (OST)												3
Syringe exchange and/or delivery												3
Distribution of general medication												3
Basic household services												3
Intimate hygiene items												3
Online counselling												2
Humanitarian aid												2
Tea and meals												2
Training about safe behaviour												2
Case Management												2
Overdose prevention												1
Total components per year		10	21	22	23	24	24	7	11	11	11	

225

226 Results of exploratory analysis of IBBS

227 26.79 % of FSWs included in the overall study sample were recruited in the West of
 228 Ukraine, 20.06% in the South, 22.74% in the East and 30.41% in Central Ukraine. The
 229 average age was 28.52 years, 27.58% of the sample was living with a partner and 23.63 %
 230 had obtained tertiary education (table 2). In 2015, 99.12% of participants were female and
 231 0.88% were male. In 2017, 97.72 % of participants were female, 2.14% male and 0.13%

232 transgender. No data about gender was available for 2009,2011 and 2013.

233 **Table 2: Demographic characteristics of participants in IBBS**

	Total N (%)	2009 N (%)	2011 N (%)	2013 N (%)	2015 N (%)	2017 N (%)
Participants	-	N = 2278	N = 5023	N = 4806	N = 4300	N = 5043
Mean age (range)	28.52 (14-62)	27.38 (14-55)	27.65 (14-53)	28.51 (14-62)	28.87 (15-61)	29.63 (15-61)
Gender						
Female	N/A	-	-	-	99.12	4928 (97.72)
Male	N/A	-	-	-	0.88	108 (2.14)
Transgender	N/A	-	-	-	-	7 (0.14)
Region of Ukraine						
Western	5747 (26.79)	703 (30.86)	1250 (24.89)	1000 (20.81)	1200 (27.91)	1594 (31.61)
Southern	4303 (20.06)	150 (6.58)	1103 (21.96)	1350 (28.09)	800(18.6)	900 (17.84)
Eastern	4877 (22.74)	462 (20.28)	1260 (25.08)	1356 (28.21)	900 (20.93)	899 (17.83)
Central	6523 (30.41)	963 (42.27)	1410 (28.07)	1100 (22.89)	1400 (32.56)	1650 (32.72)
Partnership status						
Living with a partner	5915 (27.58)	749 (32.88)	1290 (25.68)	1262 (26.26)	1332 (30.98)	1282 (25.42)
Not living with partner	15535 (72.42)	1529 (67.12)	3733 (74.32)	3544 (73.74)	2968 (69.02)	3761 (74.58)
Education						
< Tertiary education	16362 (76.37)	1805 (79.34)	3881 (77.26)	3856 (80.23)	3175 (74.06)	3645 (72.41)
≥Tertiary education (attended vocational training OR higher form of education)	5063 (23.63)	470 (20.66)	1142 (22.73)	950 (19.76)	1112 (25.94)	1389 (27.59)

234

235 **Prevention Package Coverage Analysis (objective 3)**

236 Fig 3 represents the coverage of both the basic and extended definitions of the PP
237 amongst Ukrainian FSW from 2009 to 2017. For both definitions of access to prevention,
238 coverage increased from 2009, peaked in 2013 and then decreased in 2015 and 2017. The
239 highest proportion of FSWs who had access to the PP was in 2013, where 51.46% and
240 67.42% accessed the basic and extended PP, respectively, whereas the lowest level of
241 coverage was seen in 2017 with only 26.71% received the basic and 32.10% the extended
242 PP, respectively. Between 48.4% and 54.09% of Ukrainian FSW received HIV testing
243 between 2009 and 2015, however, this drastically decreased in 2017 with only 27.92%
244 coverage. Condom coverage presented a similar trend, as between 52.85% and 65.36% of
245 FSW received condoms from 2009 to 2015, yet only 46.72% in 2017.

246

247 **Fig 3. Coverage of the basic and extended PP in FSW in the Ukraine between 2009 and**
248 **2017.**

249 **Descriptive Longitudinal Group Comparison (objective 4)**

250 When stratified by receipt and non-receipt of the basic and extended PP, our
251 descriptive longitudinal group comparison found only marginal differences and therefore, only
252 results for the basic PP are shown and discussed. Results for the extended PP are available
253 in the table in S6 S Table. Additionally, due to the volume of data, only results from the
254 descriptive longitudinal group comparison that (i) showed a clear difference between those
255 that did access the basic PP compared to those that did not or (ii) were frequently discussed
256 in the literature as being critical, are explicitly described in our results. For some thematic
257 clusters, no suitable variables were found in the IBBS datasets; for these clusters, we could
258 not conduct further analysis.

259 In relation to the factors that operate on a *macrostructural level*, our descriptive comparison
260 produced mixed results. Amongst the individual-level proxies that we used for representing
261 macrostructural relationships, the belief that HIV testing was not personally available, was
262 less common amongst PP recipients compared to PP non-recipients in all years. In all years
263 except 2009, of the FSWs who believed that HIV testing was not personally available, fear of
264 HIV status being exposed and no desire to know HIV test results, two individual-level proxies
265 addressing the level of HIV stigma amongst the population, were less common amongst the
266 PP recipients. In all years except 2011, belief that HIV prevention is too expensive, the
267 individual-level proxy referring to the national financial scheme for HIV prevention, was less
268 common amongst the PP recipients compared to the PP non-recipients. In 2009 and 2011,
269 dissatisfaction with HCF staff attitude and the opening hours of HCFs, two individual-level
270 proxies addressing health-related policy, were more common amongst PP recipients. In all
271 years, sex work as the sole form of income, the individual-level proxy used to address
272 support schemas for women (re)-entering the workforce, was consistently more common
273 amongst PP recipients than PP non-recipients.

274 Factors that operate on both the *community organisational level* and the *work environment*
275 *level* identified consistent differences across all years between PP recipients and PP non-
276 recipients. The proportion of FSWs, who were clients of an NGO was much larger in PP
277 recipients (83-97%) than in PP non-recipients (15-25%). Across all years, soliciting on the
278 street and on highways was more common amongst PP recipients, whereas soliciting in
279 affluent settings was less common. Of those who experienced GBV, PP recipients were
280 consistently more affected. Of these FSWs, GBV from clients or police officers was more
281 common amongst the PP recipients than the PP non-recipients.

282 Regarding factors that operate on the *interpersonal dynamic level*, having sex without a
283 condom was less common, while having a partner that knew of their FSW status was more
284 common amongst PP recipients compared to PP non-recipients in all years.

285 Analyses of the factors that operate on a *sex worker individual level* showed that sharing
286 needles was more common amongst PP recipients, except in 2011. PP recipients were in
287 2009, 2011 and 2013 on average one year – and in 2015 and 2017, two years older than
288 their counterparts. Both groups increased in age over time, from 28 to 32 and 27 to 29 years
289 amongst PP recipients and PP non-recipients, respectively. Knowledge of HIV transmission
290 routes and knowledge of where to access prevention was consistently higher amongst PP
291 recipients compared to non-recipients.

Table 3: Barriers and facilitators to accessing HIV prevention stratified by receipt or non-receipt of basic PP between 2009 and 2017

		2009 (%)	2011 (%)	2013 (%)	2015 (%)	2017 (%)
Macrostructural						
Socio-cultural						
<i>Level of HIV stigma amongst the population</i>						
Belief that HIV testing is personally not available	Received	0.44	0.32	0.20	0.34	24.05
	Not Received	9.28	8.37	5.19	3.78	25.49
Fear of HIV status being exposed ¹	Received ¹	53.85¹	9.09¹	-	0.05¹	1.19¹
	Not Received ¹	31.43¹	17.63¹	0.86¹	0.58¹	3.76¹
No desire to know HIV test result ¹	Received ¹	8.70 ¹	10.00 ¹	0.69 ¹	-	0.37 ¹
	Not Received ¹	8.23 ¹	10.63 ¹	1.03 ¹	0.18 ¹	2.76 ¹
Geographical						
<i>Location of HIV prevention service centres</i>						
Inconvenient location of prevention service centres ¹	Received ¹	-	-	0.04 ¹	-	0.59 ¹
	Not Received ¹	1.79 ¹	4.46 ¹	0.51 ¹	0.13 ¹	1.33 ¹
<i>Migration associated with FSW</i>						
History of travel for FSW purposes	Received	21.98	16.28	12.33	7.85	6.98
	Not Received	15.50	14.18	16.20	7.52	10.71
Health-Related Policy						
<i>Behaviour of health care staff</i>						
Dissatisfaction with staff attitude ¹	Received ¹	7.69¹	18.18¹	-	-	0.15¹
	Not Received ¹	3.57¹	7.59¹	0.30¹	0.18¹	0.19¹
<i>Functional hours of health care facility</i>						
Inappropriate opening hours ¹	Received ¹	7.69¹	18.18¹	0.08¹	-	1.19¹
	Not Received ¹	2.14¹	6.25¹	0.47¹	0.09¹	0.73¹
Economic						
<i>National financial scheme for HIV prevention</i>						
Belief that HIV prevention is too costly ¹	Received ¹	7.69¹	18.18¹	-	1.61¹	0.97¹
	Not Received ¹	18.57¹	15.62¹	1.11¹	2.80¹	2.57¹
<i>National support schemas for women in the workforce</i>						
FSW is the sole form of employment	Received	40.00	47.45	41.77	58.92	60.73
	Not Received	33.80	43.01	39.95	42.22	51.52
Community Organisation						
Sex Worker Collectivisation						
<i>Collective agency of FSW community</i>						
Client of an NGO	Received	82.86	94.14	95.51	97.03	94.43
	Not Received	14.69	20.49	24.82	25.18	15.21
Work Environment						
Physical						

		2009 (%)	2011 (%)	2013 (%)	2015 (%)	2017 (%)
<i>Soliciting setting</i>						
Street	Received	23.30	27.33	15.17	15.29	17.45
	Not Received	10.96	17.46	14.81	10.94	8.82
Highway	Received	35.05	22.69	28.43	26.27	19.75
	Not Received	20.98	17.64	19.07	16.18	8.33
Hotel	Received	4.95	4.29	3.81	2.92	4.53
	Not Received	6.80	3.49	3.14	2.58	6.30
Affluent settings (e.g. cafes, bars, modelling agencies)	Received	10.66	13.62	11.27	12.37	10.62
	Not Received	22.81	17.36	18.68	16.90	20.05
<i>Social</i>						
<i>Gender-based violence at workplace</i>						
History of gender violence whilst providing sexual services	Received	-	51.24	51.44	52.34	43.06
	Not Received	-	41.84	47.54	37.68	38.39
Experience of gender violence from client ²	Received ²	-	74.91 ²	45.09 ²	43.71 ²	34.00 ²
	Not Received ²	-	72.74 ²	42.69 ²	30.60 ²	29.95 ²
Experience of gender violence from a police officer ²	Received	-	32.04 ²	12.94 ²	7.55 ²	7.80 ²
	Not Received	-	24.96 ²	8.57 ²	5.16 ²	7.66 ²
<i>Interpersonal dynamic</i>						
<i>Sex worker- client</i>						
<i>Condom use</i>						
Sex without a condom with last client	Received	4.29	3.79	1.78	4.24	3.12
	Not Received	13.82	8.77	4.50	8.63	6.71
Pressure from client to have sex without a condom	Received	23.08	48.81	0.81	1.66	1.63
	Not Received	33.86	38.62	1.80	4.09	3.30
<i>Number of clients</i>						
Number of clients in last 24 hours ⁺	Received ⁺	2.26 ⁺	2.24 ⁺	2.11 ⁺	1.87 ⁺	2.42 ⁺
	Not Received ⁺	1.62 ⁺	1.61 ⁺	1.88 ⁺	1.61 ⁺	2.20 ⁺
<i>Sex worker – non-partner</i>						
<i>Partner's knowledge of occupational status</i>						
Partner knowledge of FSW status	Received	40.11	38.39	17.83	18.52	15.81
	Not Received	35.44	31.43	12.56	11.61	8.33
<i>Gender-based violence</i>						
Experience of gender violence from intimate partner ²	Received ²	-	10.39²	4.08²	2.49²	2.90²
	Not Received ²	-	13.71²	3.51²	1.69²	3.17²
<i>Partnership status</i>						
Living with a partner	Received	35.38	28.28	28.55	33.09	30.51
	Not Received	31.21	23.63	23.83	29.05	23.57
<i>Sex Worker Individual</i>						
<i>Behavioural</i>						

		2009 (%)	2011 (%)	2013 (%)	2015 (%)	2017 (%)
<i>Knowledge of HIV</i>						
Knowledge of HIV transmission routes (all questions correct)	Received	64.07	67.88	60.09	59.55	57.76
	Not Received	55.63	51.25	55.68	57.38	50.46
Knowledge of where to access HIV prevention	Received	98.46	99.59	99.51	99.17	98.74
	Not Received	81.80	84.78	90.44	90.57	86.42
<i>History of substance abuse</i>						
Drug Injecting						
History of sharing needles	Received	24.32	4.41	1.82	13.65	7.57
	Not Received	14.52	19.33	1.41	5.65	5.09
<i>Work life balance</i>						
Workdays per week ⁺	Received ⁺	4.12 ⁺	4.15 ⁺	4.37 ⁺	4.43 ⁺	4.27 ⁺
	Not Received ⁺	2.97 ⁺	3.22 ⁺	4.02 ⁺	3.77 ⁺	3.87 ⁺
<i>Duration in sex work</i>						
Age of FSW debut ⁺	Received ⁺	20.28 ⁺	21.24 ⁺	21.65 ⁺	21.77 ⁺	23.16 ⁺
	Not Received ⁺	20.59 ⁺	21.08 ⁺	21.41 ⁺	21.22 ⁺	22.11 ⁺
Age of sexual debut ⁺	Received ⁺	15.99 ⁺	15.87 ⁺	15.79 ⁺	-	16.40 ⁺
	Not Received ⁺	15.92 ⁺	15.99 ⁺	15.89 ⁺	-	16.17 ⁺
<i>Level of education</i>						
Higher education (tertiary education)	Received	19.49	22.10	17.91	24.56	22.35
	Not Received	21.43	23.24	21.73	27.20	29.51
<i>Biological</i>						
<i>HIV factors</i>						
HIV positive status	Received	13.75	8.57	3.80	3.17	2.08
	Not Received	8.82	8.03	1.54	3.11	2.44
<i>STI factors</i>						
Diagnosis of hepatitis C within the last 12 months	Received	5.93	6.18	8.98	7.16	6.38
	Not Received	4.46	4.13	4.59	4.54	3.81
Diagnosis of hepatitis B within the last 12 months	Received	2.97	1.94	2.39	2.24	2.30
	Not Received	1.54	1.07	1.41	1.51	2.41
<i>Age</i>						
Age ⁺	Received ⁺	27.73 ⁺	28.11 ⁺	28.90 ⁺	29.84 ⁺	31.58 ⁺
	Not Received ⁺	27.15 ⁺	27.28 ⁺	28.09 ⁺	27.98 ⁺	28.92 ⁺
<i>Reproductive status</i>						
Presence of dependents	Received	51.10	60.89	43.39	44.01	46.70
	Not Received	47.44	50.36	51.95	52.98	57.90
Number of dependents ⁺	Received ⁺	1.94 ⁺	1.83 ⁺	1.91 ⁺	1.03 ⁺	0.98 ⁺
	Not Received ⁺	1.82 ⁺	1.79 ⁺	1.75 ⁺	0.82 ⁺	0.59 ⁺
<i>Client individual</i>						

		2009 (%)	2011 (%)	2013 (%)	2015 (%)	2017 (%)
Biological						
Age						
Age (\leq 35 years)	Received	36.92	36.58	44.22	33.95	28.80
	Not Received	42.47	40.59	43.38	43.34	34.15
Bolded text are variables that are described in the results. ¹ Sub-question of "I believe that HIV testing is not available for me personally" ² Sub-question of "History of gender violence whilst providing sexual services" + Mean						

293 Discussion

294 In this study, we identified factors that acted as potential barriers and facilitators to the
295 access of HIV prevention amongst the Ukrainian FSW population between 2009 and 2017.
296 Our literature-informed conceptual framework identified various risk factor clusters, which
297 operate on six structural levels. Our exploratory analyses identified that these factors may
298 have partially been responsible for the changes in PP coverage, which peaked in 2013 and
299 then dropped in both 2015 and 2017. Our results suggest that sex work as the only form of
300 income, being a client of an NGO, street and highway solicitation, partner's knowledge of
301 occupation, knowledge of where to access prevention and knowledge of HIV transmission
302 routes may act as important facilitators to HIV prevention. Potential barriers to accessing the
303 PP were soliciting in affluent settings and unsafe sex.

304 The initial conceptual framework reflected the findings from empirical studies, which were
305 conducted in the Americas, the African, East Mediterranean, European, South-East Asia and
306 West Pacific regions between 2005 and 2018. To tailor this global conceptual framework to a
307 Ukrainian setting, we discussed our findings with staff members in the PHC of the Ministry of
308 Health of Ukraine and subsequently removed one determinant – sex worker family – from the
309 framework, as it was not applicable to the Ukrainian context. The resulting conceptual
310 framework, thus, reflects both current, global literature as well as local insights into barriers
311 and facilitators to accessing PP in Ukraine. This conceptual framework can be utilised to
312 inform future research on barriers and facilitators of accessing HIV prevention for FSWs and
313 contribute to advancing theories underlying this field of research.

314 Findings from our document analysis identified that the Ukrainian PP has changed over the
315 past nine years, which may be attributed to the fluctuation in international funding and the
316 armed conflict in the east of Ukraine. Between 2008 and 2013, Alliance's international
317 funding grew from US\$28.4 to US\$37.7 million, which coincided with an expansion in the PP

318 from 10 to 24 components in 2008 and 2013, respectively [27, 32]. Only seven components
319 were offered in 2014, the year in which Ukraine experienced a severe restructuring of the
320 political landscape [44], recession of the economy [45] and the onset of the continuing
321 armed-conflict in the east [46]. As one of the only NGOs permitted to work in the conflict
322 zones of the east, Alliance diverted a large proportion of international funding to the
323 emergency procurement of HIV and tuberculosis (TB) medicine in these eastern regions [32,
324 33]. This diversion of funds may be one factor to explain the sharp decrease in PP
325 components in regions of Ukraine from 2014 onwards, particularly, specialist medical or legal
326 counselling. Despite this cutback, the components that were included from 2015 to 2017 did
327 largely meet the WHO recommendations, which included condoms, HIV testing and
328 counselling, consultations with social workers as well as screening and prevention of co-
329 infections such as TB and Hepatitis B [47].

330 The coverage of both the basic and the extended PP also changed over this nine-year
331 period. Between 2009 and 2013, progress toward the “five pillars for achieving less than
332 500,000 new infections by 2020”, a sub-target of the 90-90-90 goals seemed promising as
333 the level of extended PP coverage amongst Ukrainian FSWs increased from 60% to 67%.
334 However, although Kyiv and other cities signed the Paris Declaration on Fast-Track Cities in
335 2014, a pact that committed cities to focusing on vulnerable and marginalized people,
336 including key populations, [22] extended PP coverage dropped to 32% in 2017.
337 Coincidentally, Alliance introduced a new policy in 2017 that deemed FSWs who knew of
338 their HIV positive status ineligible for testing [48]. This may explain the steep decline in both
339 HIV testing coverage and subsequently, the sharp drop in PP coverage in 2017.
340 Nonetheless, our analyses also identified a 18.05% decrease in condom coverage in 2017,
341 hence, future research is required to further investigate this trend. Alliance defines receipt of
342 basic PP as ‘the receipt of at least one male condom and at least one consultation with a
343 social worker’, however, due to limitations in the IBBS questionnaires, our definition included,

344 'receipt of male condoms and receipt of HIV testing and counselling'. This could explain the
345 discrepancy between the coverage we observed, and that reported by Alliance. Alliance
346 report that the basic PP coverage remained relatively stable over time (46.7% in 2013 and
347 48% in 2017), whereas we identified a clear decrease in basic PP coverage over this time
348 period (51.46% and 26.71% in 2013 and 2017, respectively). Furthermore, Alliance utilises
349 an internal database, Syrex for reporting [32] as opposed to the IBBS, which may also
350 explain this difference in coverage.

351 The fall in coverage between 2013 and 2017 may further be explained by the reduction in
352 international financial support and the redistribution of resources associated with emergency
353 programmes in the conflict zone, both of which operate on a *macrostructural level*.

354 Changes in the PP coverage may also be explained by specific factors associated with
355 access to prevention. Our findings from the *macrostructural level* suggest that relying upon
356 sex work as one's sole source of income consistently acted as a facilitator to the basic PP.
357 We identified that the proportion of FSWs relying upon sex work as their only form of income
358 increased in both groups between 2015 and 2017, which coincides with and could potentially
359 have arisen because of the outbreak of the armed conflict in the East of Ukraine. This spike
360 in reliance upon sex work aligns with previous studies that outline the catalytic effect that
361 armed conflict can have upon internal displacement and economic insecurity, which in turn,
362 often results in conditions that facilitate engagement in sex work as a source of income,
363 particularly amongst woman [49-51].

364 We observed a high level of stigma (54%) amongst PP recipients who feared their HIV status
365 being exposed in 2009, however we did not observe meaningful between-group differences.
366 Previous studies have deemed stigma to be one of the most salient barriers to HIV
367 prevention [17, 52, 53], fuelled by other factors on different structural levels such as
368 criminalisation [15, 54-56] on the *macrostructural level* and a lack of community
369 empowerment on the *community organisational level* [17, 57, 58].

370 Our results identified an encouraging trend underscoring the importance of facilitators that
371 operate on a *community organisation level*. In congruence with previous literature [17, 57-
372 59], our findings strongly imply that collective agency of the Ukrainian FSW community via
373 NGO membership largely facilitated access to the basic PP, as most PP recipients were
374 clients of an NGO. Knowledge of HIV transmission routes was higher amongst PP recipients
375 across all years, which may be attributable to consultations with social workers and the
376 distribution of HIV awareness-raising materials, two components that were consistently
377 included in the PP. Authors have also identified that social and community participation can
378 foster transparency and trust between FSWs and non-FSWs, which can increase their
379 access to prevention [39, 60]. Our results support this in all years, as disclosing one's FSW
380 status to a partner was more common amongst PP recipients, suggesting that this may have
381 acted as a facilitator.

382 Our results from the *interpersonal dynamic* and the *work environment level* not only elucidate
383 potential barriers and facilitators to PP in the Ukrainian context but highlight the interplay of
384 factors that operate on different structural levels. We found condom use to be consistently
385 more common amongst PP recipients than PP non-recipients, this may suggest that sex
386 without a condom could act as a barrier to the PP. An array of studies have identified that
387 unsafe sex is associated with GBV, a factor operating on both the *work environment* and the
388 *interpersonal dynamic level*. Authors have described that FSWs who have experienced GBV
389 in the work environment are less likely to instigate condom negotiation in fear of client
390 aggression [61-66] and therefore, may be less inclined to seek condoms, an important
391 component of the Ukrainian PP. We identified an alarmingly high prevalence of GBV
392 amongst both groups at the *work environment*, however, contrary to the literature, GBV at
393 the *work environment level* was more common amongst PP recipients than PP non-
394 recipients. Results measuring GBV at the *interpersonal level* were not meaningful. It is
395 possible that the criminalisation of sex work, a barrier that operates on the *macrostructural*

396 *level*, may have led to an increase in GBV [65, 67, 68] in the workplace, and in turn, an
397 increase in unsafe sex [69]. Although recent literature supports this hypothesis [15, 55, 70],
398 future research is required to further explore this within the Ukrainian context. Further
399 research is particularly warranted to investigate the higher prevalence of GBV amongst PP
400 recipients and this association with condom use.

401 Two additional factors on the *work environment level*, street- and highway-based soliciting
402 appeared to act as facilitators to the basic PP in all years. This contrasted with previous
403 literature, which has highlighted that street-based FSWs are less likely to access HIV
404 prevention due to an array of barriers, such as higher mobility, higher risk-taking behaviour
405 and higher exposure to GBV [9, 61, 71, 72]. The facilitating effect that street- and highway
406 soliciting had upon access to the PP in the Ukrainian context may be attributed to the
407 successful employment of mobile clinics, an intervention conducted by Alliance to ease
408 access to HIV prevention services, specifically targeting these vulnerable subgroups [34].
409 Affluent-based FSWs may be less likely to disclose their HIV status due to fear of being
410 associated with HIV and being banned from a venue, resulting in loss of income [21, 39, 73,
411 74]. Furthermore, affluent-based FSWs may be less interested in accessing the basic PP, as
412 they are able to afford these services. This warrants future qualitative research to investigate
413 the prevention needs of Ukrainian FSWs, relative to their soliciting setting.

414 Our findings from the *sex worker individual level* corroborate previous studies, which have
415 demonstrated that knowledge of where to access prevention acts as a key facilitator [9, 39,
416 56, 57, 59, 75]. However, contrary to our expectations, we found that more than 82% of
417 Ukrainian FSWs who did *not* access the basic PP *knew* where to access prevention. This
418 may suggest the need to further address macrostructural barriers in efforts to ameliorate
419 access to HIV prevention.

420 Although our study provides valuable insights into the factors associated with access HIV
421 prevention, future research is needed to deepen this knowledge. Our analyses were limited

422 by the quality of the data and further quantitative analyses, both in Ukrainian but also in other
423 specific and broader contexts, should focus on collecting high-quality and valid data to
424 address both specific associations, as well as more complex, multi-component associations.
425 Qualitative research could provide evidence on opinions, perceptions and motivations of
426 FSWs, which may help to gain a deeper insight into the source of barriers such as stigma,
427 GBV and the influence of the armed conflict upon access to HIV in a Ukrainian context.
428 Future qualitative studies exploring the specific needs of FSWs, namely, which PP
429 components are most necessary, may also elucidate important findings for future prevention
430 programmes.

431

432 **Strengths and Limitations**

433 The strength of this work lies in its theoretical foundation in current empirical literature
434 and the use of a conceptual framework to inform the analyses. Using a conceptual
435 framework is not only considered a means of conceptualizing and structuring research, it
436 also contributes to theory building around the topic of interest. Our conceptual framework
437 informed the selection of variables in the survey, hence contributing to the structure of the
438 exploratory analyses. Additionally, the use of multiple, large datasets from FSWs in almost all
439 regions of Ukraine allowed us to explore changes over time, another major strength of this
440 study.

441 Several limitations need to be taken into account when interpreting these results. Firstly, we
442 limited our literature searches to one database, PubMed. Our search strategy did not focus
443 on identifying conceptual papers that presented and discussed theories, models or
444 frameworks of barriers and facilitators but empirical papers assessing barriers and facilitators
445 in real-world settings. There were also limitations with regard to the assessed IBBS surveys.
446 Changes in the design of surveys made it difficult to compare particular factors across time,

447 particularly questions measuring barriers on the *macrostructural level*. Additionally, missing
448 data prevented us from measuring informative outcomes, such as the coverage of the basic
449 PP, as defined by Alliance. The nature of the questions themselves may have also
450 influenced results; both FSW and HIV are two highly sensitive issues and therefore, recall
451 and social desirability bias may have impacted the responses in the survey, particularly since
452 the survey were administered face-to-face. There were also limitations in our analyses. In
453 measuring PP coverage, changes in the specific components of the PP over time resulted in
454 our indicator for extended PP coverage being more volatile and less informative than the
455 basic PP. However, a similar trend was seen in both basic and extended PP coverage,
456 hence this did not substantially impact the measurement of coverage. Nonetheless, the level
457 of coverage of each PP component may vary based on region and therefore have had some
458 influence upon our results. Geographical sampling of cities included in the IBBS may have
459 also influenced the level of coverage. As in other studies that utilised IBBS data [74, 76-78],
460 we did not weight for sampling and therefore, our results are to be interpreted with this in
461 account. Only in 2017 and 2015 was a question on gender included in the IBBS and
462 although only a small proportion of SWs were male or transgender, limitations caused by the
463 misclassification need to be considered.

464 We conducted and reported descriptive statistical analyses, however, due to limitations in the
465 data, such as inconsistent response scales and a lack of internal consistency throughout all
466 five datasets alongside structural changes between 2013 and 2015. Hence, the analyses
467 should be considered exploratory rather than confirmatory and should be interpreted
468 cautiously. As the IBBS measured individual behaviours, we were limited to assigning
469 individual-level proxies analyse factors that influence access to the Ukrainian PP on the
470 macrostructural level. RDS and TLS have found to be effective sampling methods for
471 reaching marginalised populations. In accordance with others working with the IBBS data

472 [74, 79], we did not consider survey weights when calculating descriptive statistics, which
473 may have led to an over-representation of certain subpopulations.

474 **Conclusion**

475 To our knowledge, this is the first study to explore the factors associated with the
476 access of Ukrainian FSW to HIV prevention on a national scale across a time span of nine
477 years. Our study strengthens current knowledge that access to prevention is complex with
478 many factors found on multiple levels and that access to prevention has decreased since
479 2014. We also identified several factors such as relying solely upon sex work for income,
480 being a client of an NGO, knowledge of HIV, partner's knowledge of occupation, soliciting
481 setting and condom use that may be associated with access to HIV prevention within the
482 Ukrainian context. Future interventions targeting these factors may increase the coverage of
483 HIV prevention and thus, contribute to reducing the HIV burden within the FSW population.

484 **Acknowledgments**

485 The paper summarizes and analyses the data from studies funded by of the Global
486 Fund to Fight AIDS, Tuberculosis and Malaria (TGF) and implemented in Ukraine by the
487 International Charitable Foundation "Alliance for Public Health" (Alliance). The data for
488 analysis were provided to the authors by the data holder the Public Health Centre (PHC) of
489 the Ukrainian Ministry of Health. All opinions presented in this manuscript belong to the
490 authors alone and not represent position of the Global Fund to Fight AIDS, Tuberculosis and
491 Malaria, Alliance for Public Health or the Public Health Centre (PHC) of the Ukrainian
492 Ministry of Health We would like to thank Ihor Kuzin (PHC), Yana Sazonova (Alliance),
493 Roksolana Kulchynska (CDC) and Tetyana Salyuk (Alliance) for their comments on the
494 manuscript, , Oleksandr Korotych (PHC) and Oksana Kovtun (Alliance) for providing us with

495 the IBBS datasets and, Nadiya Yanhol (Alliance) for her insight into the PP. . Lastly and most
496 importantly, thank you to the SWs that partook in the IBBS studies.

497 **Author contributions**

498 Conceptualisation: NB, JB, LMF

499 Formal analysis: NB, JB

500 Methodology: JB, LMF, NB

501 Supervision: JB, LMF

502 Visualisation: NB

503 Writing- original draft: NB

504 Writing- review and editing: NB, JB, LMF

505 References

- 506 1. European Centre for Disease Prevention and Control/WHO Regional Office for
507 Europe. HIV/AIDS surveillance in Europe 2017 - 2016 data. Stockholm; 2017.
- 508 2. Tokar A, Sazonova I, Mishra S, et al. HIV testing behaviour and HIV prevalence
509 among female sex workers in Ukraine: findings from an Integrated Bio-Behavioural Survey,
510 2013–2014. *Sexually Transmitted Infections*. 2019;95(3):193-200.
- 511 3. ICF "Alliance For Public Health". Захворюваність на ВІЛ серед ключових груп.
512 2018.
- 513 4. World Health Organisation. Consolidated Guidelines on HIV prevention, diagnosis,
514 treatment and care for key populations. WHO Library Cataloguing-in-Publication data.
515 2014:p2-11.
- 516 5. Shannon K, Crago AL, Baral SD, et al. The global response and unmet actions for
517 HIV and sex workers. *Lancet*. 2018;392(10148):698-710.
- 518 6. Baral S, Beyrer C, Muessig K, et al. Burden of HIV among female sex workers in low-
519 income and middle-income countries: a systematic review and meta-analysis. *Lancet Infect*
520 *Dis*. 2012;12(7):538-49.
- 521 7. World Health Organisation. Chapter 2: Addressing violence against sex workers.
522 Implementing comprehensive HIV/ STI programmes with sex workers: practical approaches
523 from collaborative interventions. Geneva2013.
- 524 8. Prakash R, Bhattacharjee P, Blanchard A, et al. Effects of exposure to an intensive
525 HIV-prevention programme on behavioural changes among female sex workers in Nairobi,
526 Kenya. *Afr J AIDS Res*. 2018;17(2):99-108.
- 527 9. Huang W, Operario D, Dong Y, et al. HIV-related risk among female migrants working
528 in entertainment venues in China. *Prev Sci*. 2014;15(3):329-39.
- 529 10. Cange CW, LeBreton M, Saylor K, et al. Female sex workers' empowerment
530 strategies amid HIV-related socioeconomic vulnerabilities in Cameroon. *Cult Health Sex*.
531 2017;19(10):1053-65.
- 532 11. Beyrer C, Crago A-L, Bekker L-G, et al. An action agenda for HIV and sex workers.
533 *The Lancet*. 2015;385(9964):287-301.
- 534 12. Shannon K, Strathdee SA, Shoveller J, Rusch M, Kerr T, Tyndall MW. Structural and
535 environmental barriers to condom use negotiation with clients among female sex workers:
536 implications for HIV-prevention strategies and policy. *Am J Public Health*. 2009;99(4):659-65.
- 537 13. Shannon K, Strathdee SA, Goldenberg SM, et al. Global epidemiology of HIV among
538 female sex workers: influence of structural determinants. *The Lancet*. 2015;385(9962):55-71.
- 539 14. Shannon K, Goldenberg SM, Deering KN, Strathdee SA. HIV infection among female
540 sex workers in concentrated and high prevalence epidemics: why a structural determinants
541 framework is needed. *Curr Opin HIV AIDS*. 2014;9(2):174-82.
- 542 15. Maher L, Dixon T, Phlong P, Mooney-Somers J, Stein E, Page K. Conflicting Rights:
543 How the Prohibition of Human Trafficking and Sexual Exploitation Infringes the Right to
544 Health of Female Sex Workers in Phnom Penh, Cambodia. *Health Hum Rights*.
545 2015;17(1):E102-13.

- 546 16. Lazarus L, Deering KN, Nabess R, Gibson K, Tyndall MW, Shannon K. Occupational
547 stigma as a primary barrier to health care for street-based sex workers in Canada. *Cult*
548 *Health Sex.* 2012;14(2):139-50.
- 549 17. Biradavolu MR, Blankenship KM, Jena A, Dhungana N. Structural stigma, sex work
550 and HIV: contradictions and lessons learnt from a community-led structural intervention in
551 southern India. *J Epidemiol Community Health.* 2012;66 Suppl 2:ii95-9.
- 552 18. Strathdee SA, West BS, Reed E, Moazen B, Azim T, Dolan K. Substance Use and
553 HIV Among Female Sex Workers and Female Prisoners: Risk Environments and Implications
554 for Prevention, Treatment, and Policies. *J Acquir Immune Defic Syndr.* 2015;69 Suppl
555 2:S110-7.
- 556 19. Bekker L-G, Beyrer C, Quinn TC. Behavioral and biomedical combination strategies
557 for HIV prevention. *Cold Spring Harb Perspect Med.* 2012;2(8):a007435.
- 558 20. Krishnaratne S, Hensen B, Cordes J, Enstone J, Hargreaves JR. Interventions to
559 strengthen the HIV prevention cascade: a systematic review of reviews. *The lancet HIV.*
560 2016;3(7):e307-e17.
- 561 21. Bekker L-G, Johnson L, Cowan F, et al. Combination HIV prevention for female sex
562 workers: what is the evidence? *The Lancet.* 2015;385(9962):72-87.
- 563 22. UNAIDS. Prevention gap report. Geneva; 2016.
- 564 23. ICF "Alliance For Public Health". Annual Report 2006. Kyiv; 2006.
- 565 24. ICF "Alliance For Public Health". Annual Report 2016. Kyiv; 2016.
- 566 25. ICF "Alliance For Public Health". Annual Report 2015. Kyiv; 2015.
- 567 26. Das P, Horton R. Bringing sex workers to the centre of the HIV response. *The Lancet.*
568 2015;385(9962):3-4.
- 569 27. ICF "Alliance For Public Health". Annual Report 2008. Kyiv; 2008.
- 570 28. ICF "Alliance For Public Health". Annual Report 2009. Kyiv; 2009.
- 571 29. ICF "Alliance For Public Health". Annual Report 2010. Kyiv; 2010.
- 572 30. ICF "Alliance For Public Health". Annual Report 2011. Kyiv; 2011.
- 573 31. ICF "Alliance For Public Health". Annual Report 2012. Kyiv; 2012.
- 574 32. ICF "Alliance For Public Health". Annual Report 2013. Kyiv; 2013.
- 575 33. ICF "Alliance For Public Health". Annual Report 2014. Kyiv; 2014.
- 576 34. ICF "Alliance For Public Health". Annual Report 2017. Kyiv; 2017.
- 577 35. ICF "Alliance For Public Health". Annual Report 2004. Kyiv; 2004.
- 578 36. Yanhol N. Kyiv: Alliance of Public Health; 2019.
- 579 37. UNAIDS. Global AIDS Monitoring 2018: indicators for monitoring the 2016 United
580 Nations Political Declaration on HIV and AIDS. Geneva 2018.
- 581 38. Team RC. R: A language and environment for statistical computing. R Foundation for
582 Statistical Computing. Vienna, Austria 2017.
- 583 39. Chanda MM, Perez-Brumer AG, Ortblad KF, et al. Barriers and Facilitators to HIV
584 Testing Among Zambian Female Sex Workers in Three Transit Hubs. *AIDS Patient Care*
585 *STDS.* 2017;31(7):290-6.

- 586 40. Qiao S, Li X, Zhang C, et al. Psychological fears among low-paid female sex workers
587 in southwest China and their implications for HIV prevention. *PLoS One*.
588 2014;9(10):e1111012.
- 589 41. Johnston LG, Bonilla L, Caballero T, et al. Associations of HIV Testing, Sexual Risk
590 and Access to Prevention Among Female Sex Workers in the Dominican Republic. *AIDS*
591 *Behav*. 2017;21(8):2362-71.
- 592 42. Langa J, Sousa C, Sidat M, et al. HIV risk perception and behavior among sex
593 workers in three major urban centers of Mozambique. *PLoS One*. 2014;9(4):e94838.
- 594 43. World Health Organisation. Consolidated guidelines on the use of antiretroviral drugs
595 for treating and preventing HIV infection: recommendations for a public health approach.
596 2013.
- 597 44. Shveda Y, Park JH. Ukraine's revolution of dignity: The dynamics of Euromaidan.
598 *Journal of Eurasian Studies*. 2016;7(1):85-91.
- 599 45. The World Bank Group. Ukraine Country Profile 2019 [Available from:
600 <https://data.worldbank.org/country/ukraine>].
- 601 46. Holt E. Conflict in Ukraine and a ticking bomb of HIV. *The Lancet HIV*.
602 2018;5(6):e273-e4.
- 603 47. World Health Organisation. Consolidated guidelines on HIV prevention, diagnosis,
604 treatment and care for key populations, 2016 update. Geneva; 2017.
- 605 48. Salyuk T. Kyiv: Alliance of Public Health; 2020.
- 606 49. Ferguson A, Shannon K, Butler J, Goldenberg SM. A comprehensive review of
607 HIV/STI prevention and sexual and reproductive health services among sex Workers in
608 Conflict-Affected Settings: call for an evidence- and rights-based approach in the
609 humanitarian response. *Conflict and health*. 2017;11:25.
- 610 50. Cazabat C. A gender perspective on internal displacement. Geneva: Internal
611 Displacement Monitoring Centre; 2019.
- 612 51. Maclin B, Kelly J, Kabanga J, VanRooyen M. 'They have embraced a different
613 behaviour': transactional sex and family dynamics in eastern Congo's conflict. *Cult Health*
614 *Sex*. 2015;17(1):119-31.
- 615 52. Nnko S, Kuringe E, Nyato D, et al. Determinants of access to HIV testing and
616 counselling services among female sex workers in sub-Saharan Africa: a systematic review.
617 *BMC Public Health*. 2019;19(1):15.
- 618 53. Duby Z, Nkosi B, Scheibe A, Brown B, Bekker LG. 'Scared of going to the clinic':
619 Contextualising healthcare access for men who have sex with men, female sex workers and
620 people who use drugs in two South African cities. *South Afr J HIV Med*. 2018;19(1):701.
- 621 54. Tucker JD, Ren X. Sex worker incarceration in the People's Republic of China. *Sex*
622 *Transm Infect*. 2008;84(1):34-5; discussion 6.
- 623 55. Shahmanesh M, Wayal S, Andrew G, Patel V, Cowan FM, Hart G. HIV prevention
624 while the bulldozers roll: exploring the effect of the demolition of Goa's red-light area. *Soc Sci*
625 *Med*. 2009;69(4):604-12.
- 626 56. Scorgie F, Nakato D, Harper E, et al. 'We are despised in the hospitals': sex workers'
627 experiences of accessing health care in four African countries. *Cult Health Sex*.
628 2013;15(4):450-65.

- 629 57. Zeng H, Zhang L, Zhao Y, et al. HIV prevention among street-based sex workers
630 (SSWs) in Chongqing, China: interviews with SSWs, clients and healthcare providers. *Health*
631 *Soc Care Community*. 2016;24(6):e173-e80.
- 632 58. Lim RBT, Cheung ONY, Tham DKT, et al. Using qualitative and community-based
633 engagement approaches to gain access and to develop a culturally appropriate STI
634 prevention intervention for foreign female entertainment workers in Singapore. *Global Health*.
635 2018;14(1):36.
- 636 59. Wanyenze RK, Musinguzi G, Kiguli J, et al. "When they know that you are a sex
637 worker, you will be the last person to be treated": Perceptions and experiences of female sex
638 workers in accessing HIV services in Uganda. *BMC Int Health Hum Rights*. 2017;17(1):11.
- 639 60. Tucker JD, Peng H, Wang K, et al. Female sex worker social networks and STI/HIV
640 prevention in South China. *PLoS One*. 2011;6(9):e24816.
- 641 61. Panchanadeswaran S, Johnson SC, Sivaram S, et al. Intimate partner violence is as
642 important as client violence in increasing street-based female sex workers' vulnerability to
643 HIV in India. *Int J Drug Policy*. 2008;19(2):106-12.
- 644 62. Erausquin JT, Reed E, Blankenship KM. Police-related experiences and HIV risk
645 among female sex workers in Andhra Pradesh, India. *J Infect Dis*. 2011;204 Suppl 5:S1223-
646 8.
- 647 63. Lang DL, Salazar LF, DiClemente RJ, Markosyan K. Gender based violence as a risk
648 factor for HIV-associated risk behaviors among female sex workers in Armenia. *AIDS Behav*.
649 2013;17(2):551-8.
- 650 64. Pando MA, Coloccini RS, Reynaga E, et al. Violence as a barrier for HIV prevention
651 among female sex workers in Argentina. *PLoS One*. 2013;8(1):e54147.
- 652 65. Decker MR, Lyons C, Billong SC, et al. Gender-based violence against female sex
653 workers in Cameroon: prevalence and associations with sexual HIV risk and access to health
654 services and justice. *Sex Transm Infect*. 2016;92(8):599-604.
- 655 66. Prakash R, Manthri S, Tayyaba S, et al. Effect of Physical Violence on Sexually
656 Transmitted Infections and Treatment Seeking Behaviour among Female Sex Workers in
657 Thane District, Maharashtra, India. *PLoS One*. 2016;11(3):e0150347.
- 658 67. Duff P, Sou J, Chapman J, et al. Poor working conditions and work stress among
659 Canadian sex workers. *Occup Med (Lond)*. 2017;67(7):515-21.
- 660 68. Reeves A, Steele S, Stuckler D, McKee M, Amato-Gauci A, Semenza JC. National
661 sex work policy and HIV prevalence among sex workers: an ecological regression analysis of
662 27 European countries. *Lancet HIV*. 2017;4(3):e134-e40.
- 663 69. Criminal Code of Ukraine, (2018).
- 664 70. Gruskin S, Ferguson L. Government regulation of sex and sexuality: in their own
665 words. *Reprod Health Matters*. 2009;17(34):108-18.
- 666 71. Jayanna K, Washington RG, Moses S, et al. Assessment of attitudes and practices of
667 providers of services for individuals at high risk of HIV and sexually transmitted infections in
668 Karnataka, south India. *Sex Transm Infect*. 2010;86(2):131-5.
- 669 72. Yi H, Mantell JE, Wu R, Lu Z, Zeng J, Wan Y. A profile of HIV risk factors in the
670 context of sex work environments among migrant female sex workers in Beijing, China.
671 *Psychol Health Med*. 2010;15(2):172-87.

- 672 73. Munoz J, Adedimeji A, Alawode O. 'They bring AIDS to us and say we give it to them':
673 Socio-structural context of female sex workers' vulnerability to HIV infection in Ibadan,
674 Nigeria. SAHARA J. 2010;7(2):52-61.
- 675 74. Tokar A, Sazonova I, Mishra S, et al. HIV testing behaviour and HIV prevalence
676 among female sex workers in Ukraine: findings from an Integrated Bio-Behavioural Survey,
677 2013–2014. Sex Transm Infect. 2019;95(3):193-200.
- 678 75. Asadi-Aliabadi M, Abolghasemi J, Rimaz S, Majdzadeh R, Rostami-Maskopae F,
679 Merghati-Khoei E. Barriers to Health Service Utilization Among Iranian Female Sex Workers:
680 A Qualitative Study. Journal of Preventive Medicine and Public Health. 2018;51(2):64.
- 681 76. Makarenko I, Ompad DC, Sazonova Y, Saliuk T, DeHovitz J, Gensburg L. Trends in
682 Injection Risk Behaviors among People Who Inject Drugs and the Impact of Harm Reduction
683 Programs in Ukraine, 2007–2013. J Urban Health. 2017;94(1):104-14.
- 684 77. Smyrnov P, Williams LD, Korobchuk A, et al. Risk network approaches to locating
685 undiagnosed HIV cases in Odessa, Ukraine. J Int AIDS Soc. 2018;21(1):e25040.
- 686 78. Iakunchykova OP, Burlaka V. Correlates of HIV and Inconsistent Condom Use
687 Among Female Sex Workers in Ukraine. AIDS Behav. 2017;21(8):2306-15.
- 688 79. Vitek CR, Čakalo J-I, Kruglov YV, et al. Slowing of the HIV epidemic in Ukraine:
689 evidence from case reporting and key population surveys, 2005–2012. PLoS One.
690 2014;9(9):e103657.
- 691

692 **Supporting Information Captions**


693 **S1 Appendix. Literature review method, protocol and results S2 Table. Formation of**
694 **thematic clusters used to populate the conceptual framework**

695 **S3 Table. Categories and specific PP components as extracted from the Alliance**
696 **General Reports from 2008 to 2017**

697 **S4 Table. PP component and corresponding IBBS item for 2009, 2011, 2013, 2015 and**
698 **2017**

699 **S5 Table. Independent variables from IBBS and corresponding thematic clusters for**
700 **2009, 2011, 2013, 2015 and 2017**

701 **S6 Table. Barriers and facilitators to accessing HIV prevention stratified by receipt or**
702 **non-receipt of extended PP between 2009 and 2017**



Literature Review

Document Analysis

Conceptual Framework
(objective 1)

Description of Ukrainian
prevention package
(objective 2)

Review of IBBS
questionnaire

Variable Selection

Exploratory Analyses

PP Coverage Analysis
(objective 3)

Descriptive longitudinal
group comparison
(objective 4)

Figure 2: Conceptual framework of factors and structural levels influencing access to HIV prevention




Figure 3: Coverage of the basic and extended PP in FSW in Ukraine between 2009 and 2017

