

Title: Diagnostic accuracy at the first episode of psychosis in Uganda.

Authors: Angel Nanteza^{1*}, Emmanuel K. Mwesiga^{2,3,4}; Juliet Nakku¹; Noeline Nakasujja^{2,3}; Dickens Akena^{2,3}.

Author affiliation

1. Butabika National Referral Mental Hospital.
2. Department of Psychiatry, College of Health Sciences, Makerere University
3. Neuropsychiatric Genetics in African Populations (NeuroGAP) Psychosis project.
4. Department of Psychiatry and Mental Health, University of Cape Town.

Corresponding Author*: Miss Angel Nanteza, Butabika National Referral Mental Hospital;
Email: angelnanteza96@gmail.com

ABSTRACT

Background: Correct clinical diagnosis at the first episode of psychosis may be difficult due to many non-specific symptoms. We aimed to determine the factors associated with a correct diagnosis among patients with a first episode of psychosis in Uganda.

Methods: A cross sectional study design was performed at the Butabika National Referral Mental Hospital in Uganda. We included treatment naïve participants aged 18 to 60 years with a diagnosis of a psychotic disorder. Patients with organic disorders like HIV/AIDS, syphilis and substance use disorders were excluded. The MINI international neuropsychiatric inventory was administered to confirm the clinical diagnosis. Concordance was based on the percentage agreement and kappa statistic between the admission chart diagnosis and the MINI diagnosis.

Results: 178 participants with a first episode of psychosis were included into the final analysis. The agreement between the MINI diagnosis and clinician diagnosis was 0.385, ($P < 0.001$) with a concordance of 49.5%. After controlling for nationality and the household's source of income, duration of untreated psychosis, [p-value 0.028(95%CI: 0.07-0.89)], living with a primary family member, [p-value 0.038(95%CI:0.95-2.86)] and cadre of the clinician who made the initial diagnosis[Medical officer, [p-value 0.011(95%CI: 0.18-0.80)] were associated with a correct diagnosis.

Conclusion: We found low agreement between clinician diagnoses and MINI diagnoses at the first episode of psychosis. Improved training of staff while considering the duration of untreated psychosis and the living arrangements of the patient are required to improve diagnostic accuracy in this population.

KEY WORDS: first episode psychosis, diagnosis, diagnostic accuracy, low and middle income country, agreement.

BACKGROUND

Psychotic disorders account for 10% of global burden of disease (Rathod et al., 2017; Whiteford, Ferrari, Degenhardt, Feigin, & Vos, 2015). Approximately 80% of the burden of psychotic disorders is found in low and middle income countries (LMIC) like Uganda (Kigozi et al., 2016; Organization, 2017; Rathod et al., 2017). Psychotic disorders are associated with high productivity loss and mortality due to suicide (Laursen, Munk-Olsen, & Gasse, 2011; Lencz et al., 2014; Rost, Meng, & Xu, 2014). In order to reduce the burden associated with these disorders it is essential that they are identified, diagnosed and managed early in the course of the illness (Marshall & Rathbone, 2011; McGorry, 2015). It is preferable that the management is initiated early at the first episode of psychosis(FEP) (Anderson et al., 2018; Norman & Manchanda, 2016).

FEP, operationally defined as the time of initiation of antipsychotic medication; is an important time point in the course of a psychotic illness (Breitborde, Srihari, & Woods, 2009). Literature from high income countries indicate that correct management of psychotic disorders during FEP has been associated with improved outcomes like quality of life (Heeramun-Aubeeluck et al., 2015; Marshall & Rathbone). Differences in health seeking behaviour and health systems however imply that the time for initiation of antipsychotic medication varies widely (Mossaheb et al., 2013; Murru & Carpiniello, 2018; Oosthuizen, Emsley, Keyter, Niehaus, & Koen, 2005). Moreover, a number of patients present with varieties of clinical features that make it difficult to make a conclusive diagnosis and rate severity (Abbo, Ekblad, Waako, Okello, & Musisi, 2009; Mossaheb et al., 2013). Even with this challenge it is still imperative that an attempt is made to correctly classify patients according to some set diagnostic criteria. For psychotic disorders this might be difficult as many of the symptoms continuously change as the disease progresses (Dutta et al., 2007; Heckers et al., 2013; Tandon, 2013). The presence of many non-specific signs and symptoms are at FEP may therefore impact on clinicians diagnoses (Dutta et al., 2007).

The diagnostic accuracy for psychotic disorders at FEP in LMIC is not well described. In low resource settings, severe human resource constraints and greater need to task shift implies that few patients are reviewed by psychiatrists (Swartz, Kilian, Twesigye, Attah, & Chiliza, 2014). It is unclear if there is an association between clinician's level of training and correct diagnoses among patients with FEP. The association of patient illness factors like symptom severity, age of onset of illness and duration of untreated psychosis on correct diagnoses at FEP is also not well described. We therefore aimed to determine the diagnostic accuracy between diagnoses made by clinicians against a gold standard among patients with FEP at the country's national referral and teaching mental hospital. We also aimed to determine whether socio-demographic, clinician and illness factors were associated with the accuracy of the diagnosis.

METHODS

Study site: This was a cross sectional study conducted on the four acute admission wards at Butabika National Referral Mental Hospital in Uganda. Butabika National Referral and Teaching Mental Hospital is located about 12 kilometers south east of Kampala city in Nakawa sub-county. On average, a total of about 200 patients are attended to daily in the outpatient units of the hospital (Emmanuel Kiiza Mwesiga et al., 2019). The hospital has 4 general admission wards, 2 sick wards and 2 convalescent wards. There are specialized services for addiction psychiatry, social work, forensic psychiatry, child and adolescent mental health services and psychotherapy. Each general ward has about 150 in patients who are assigned to one of five different firms. These firms are supervised by clinical teams comprised of psychiatrists, residents in training, medical officers, clinical psychologists, psychiatric social workers, psychiatric nurses and psychiatric clinical officers. These teams review the patients in their firms on a regular basis.

Ethical approvals: Ethical approval was sought from and approved by the Uganda National Council for Science and Technology (UNCST) (#HS142ES), the School of Medicine Research and Ethics Committee (SOMREC), Makerere University (#2017-153) and institutional approval was received from Butabika hospital

Study participants: Participants for this study were selected from an ongoing study in which we aim to determine the factors associated with cognitive impairment among patients with a first episode of psychosis. This study enrolled consenting in-patients with a psychosis diagnosis aged 18 to 60 years and who were naïve to antipsychotic medication. Participants with HIV/AIDs, substance use and Syphilis were excluded.

Study tools: The MINI International Neuropsychiatric Inventory is a semi-structured interview guide for making DSM-5 diagnoses. It has been extensively used in our setting. Only Module D (mania), module J (alcohol use), module K (other substances) and module L (psychotic disorders) were administered during the study. Specifically, for module L an extended version of the MINI was used. This extended version has an additional 12 questions that helps a clinician to subtype a specific psychotic disorder. With the extended version, one can differentiate between schizophrenia and schizophreniform disorders, which may not be possible in the non-extended version of the MINI. The non-extended version is being used in the original study (Anne Stevenson et al., 2019).

In the primary study, participants had sociodemographic information collected using a standardized questionnaire. This questionnaire collected data on sociodemographic characteristics like age, gender, religion, source of income and marital status. It also collected data on onset of psychotic symptoms which was used to calculate the duration of untreated psychosis as well as alternative therapies used before admission.

Study procedure: All participants who were enrolled in the primary study and agreed to take part in this study had their psychosis diagnosis confirmed using the extended version of the MINI. The extended version of the MINI was administered by a psychiatric clinical officer (AN) who had been trained. A psychiatrist (EKM) supervised the assessment process. Later, chart abstraction was performed to determine the diagnosis made at admission as well the cadre (level of training) of staff who made the diagnosis.

Statistical methods: Data was analysed in June 2019 using StataCorp (2015). Statistical Software: Release 14. College Station, TX: StataCorp LP. Diagnoses made by the different clinician cadres in the different psychiatric teams were compared to diagnoses confirmed using the MINI. Our dependant variable was the participants whose admission diagnosis correlated with the MINI diagnosis. Patient sociodemographic, illness related (age of onset, duration of untreated psychosis and age at initial contact with mental health facility) and clinician-related factors like who made the diagnosis were our independent variables. Duration of untreated psychosis (DUP) can be defined as the time from the emergence of the first psychotic symptom to the start of antipsychotic treatment (Drancourt et al., 2013). DUP was calculated by subtracting the age at which the patient first recognised symptoms from their current age. At multi-variate analysis a level of significance of less than 0.05 was used to test for significance between different factors and a correct diagnosis on both the MINI and admission file while controlling for nationality and the household's source of income.

RESULTS

Our final dataset included 178 participants who were assessed using the MINI. Briefly the median age was 27 years (IOR 24-36) with 51% (92/178) of the participants being female. The commonest diagnosis at admission was Bipolar Affective disorder (28%). The proportions of different diagnoses in the chart and on the MINI are shown in table 1 below.

Table 1 Comparison between the diagnosis made at admission and the gold standard (MINI) diagnosis

Diagnosis	Dx at Admission Number (%)	MINI Dx Number (%)
Alcohol psychosis	0 (0)	14 (7.7)
Bipolar Affective Disorder	51 (28.0)	50 (27.5)
Brief Psychotic disorder	29 (15.9)	17 (9.3)
Major depressive disorder	0 (0)	1 (0.6)
Depression with psychotic features	15 (8.2)	28 (15.4)
Schizophrenia	42 (23.1)	57 (31.3)
Schizoaffective	2 (1.1)	4 (2.2)
Schizophreniform	2 (1.1)	11 (6.0)
Other	23 (12.7)	0 (0)
Unknown	18 (9.9)	0 (0)

Agreement: Cohen's Kappa = 0.385, $P < 0.001$ and concordance was 49.5%

Table 2 shows the bivariate associations between of concordance of the admission and MINI diagnoses with various sociodemographic characteristics. There were no associations between a correct diagnosis and age [prevalence ratio (PR) 0.99 (95% confidence interval (CI) 0.73-1.35; p value 0.984] or gender age [PR 1.08 (95% CI 0.79-1.46); p value 0.643].

Table 2: The factors associated with the concordance between clinical diagnosis at admission and MINI Diagnosis by trained RA.

Variable	Number (N)	Concordance [n(%)]	Prevalence ratio	95% CI	P-value
Age (categorised at median)					
Median = 27 (IQR 24 – 36)					
≤ 27	93	45 (48.4)	1		
> 27	85	41 (48.2)	0.99	0.73 – 1.35	0.984
Gender					
Male	86	40 (46.5)	1		
Female	92	46 (50.0)	1.08	0.79 – 1.46	0.643
Marital status					
Single	95	45 (47.4)	1		
Married	51	27 (52.9)	1.12	0.80 – 1.56	0.516
Divorced/Widowed	32	14 (43.8)	0.92	0.59 – 1.45	0.728
Current employment					
Student	15	8 (53.3)	1		
Formal employment	17	8 (47.1)	0.88	0.44 – 1.77	0.724
Non-formal employment	73	33 (45.2)	0.85	0.49 – 1.45	0.547
Unemployed	73	37 (50.7)	0.95	0.56 – 1.61	0.850
Nationality					
Uganda	167	78 (46.7)	1		
Other	10	7 (70.0)	1.50	0.97 – 2.32	0.070
Main source of income in household					
Self	63	26 (41.3)			
Father	29	16 (55.2)	1.34	0.86 – 2.08	0.198
Mother	21	12 (57.1)	1.38	0.86 – 2.23	0.179
Relative/Guardian	49	22 (44.9)	1.09	0.71 – 1.67	0.700
Non-relative/Organisation	13	8 (61.5)	1.49	0.88 – 2.51	0.134
Most import household source of earning					
Subsistence farming	51	25 (49.0)	1		
Commercial farming	3	2 (66.7)	1.36	0.58 – 3.18	0.478
Wage employment	65	31 (47.7)	0.97	0.67 – 1.42	0.887
Non-agricultural enterprises	16	10 (62.5)	1.23	0.79 – 2.05	0.314
Property income	19	7 (36.8)	0.75	0.39 – 1.45	0.392
Organisational support	2	2 (100.0)	2.04	1.54 – 2.70	<0.001
Others	19	7 (36.8)	0.75	0.39 – 1.45	0.392
Current living arrangements					
Renting	30	10 (33.3)	1		
Owns house	38	19 (50.0)	1.50	0.82 – 2.73	0.185
Living with primary family	71	39 (54.9)	1.65	0.95 – 2.86	0.075
Living with other family	29	13 (44.8)	1.34	0.70 – 2.57	0.371
Living with friends	3	1 (33.3)	1.00	0.19 – 5.38	1.000
No housing/Living in street	6	3 (50.0)	1.50	0.58 – 3.88	0.403
Others	1	1 (100.0)	3.00	1.81 – 4.98	<0.001
First time presenting with symptoms?					
No	54	29 (53.7)	1		
Yes	124	57 (46.0)	0.86	0.63 – 1.17	0.331

First time at Medical facility?					
No	22	12 (54.6)	1		
Yes	152	73 (48.0)	0.88	0.58 – 1.34	0.550
Given Antipsychotic medication before?					
No	151	73 (48.3)	1		
Yes	24	13 (54.2)	1.12	0.75 – 1.68	0.582
Age first sought help (categorised at median; Median = 26. IQR 22 – 33)					
≤ 26	89	43 (48.3)	1		
> 26	82	41 (50.0)	1.02	0.76 – 1.41	0.826
Previously used Traditional healer					
No	123	61 (49.6)	1		
Yes	53	24 (45.3)	0.91	0.65 – 1.29	0.607
Previously used a spiritual healer					
No	96	46 (47.9)	1		
Yes	80	39 (48.8)	1.02	0.75 – 1.38	0.912
Previously used a witch doctor					
No	143	68 (47.6)	1		
Yes	33	17 (51.5)	1.08	0.75 – 1.57	0.675
Who made initial diagnosis					
Psychiatrist	17	13 (76.5)	1.00		
Medical officer	14	6 (42.9)	0.56	0.9 – 1.09	0.086
Clinical officer	136	65 (47.8)	0.63	0.45 – 0.89	0.004
Nurse	11	2 (18.2)	0.24	0.07 – 0.89	0.028
DUP [Mean (SD)]	1.30 (3.71)		1.03	1.01 – 1.05	0.002

After controlling for nationality and household's most import source of income the variables that remain significantly associated with a correct diagnosis are shown in table 3 below.

Table 3: Multi-variable logistic regression between correct diagnosis and socio-demographic characteristics.

Variable	Prevalence ratio	95% CI	P-value
Main source of income in household			
Self	1.00		
Father	1.32	0.71 – 2.46	0.384
Mother	0.94	0.50 – 1.74	0.833
Relative/Guardian	0.96	0.60 – 1.55	0.882
Non-relative/Organisation	1.18	0.53 – 2.59	0.681
Most import household source of earning			
Subsistence farming	1.00		
Commercial farming	2.16	0.87 – 5.38	0.098
Wage employment	0.82	0.54 – 1.25	0.361
Non-agricultural enterprises	1.35	0.80 – 2.28	0.262
Property income	0.60	0.32 – 1.16	0.128
Organisational support	1.85	0.24 – 14.52	0.560
Others	0.83	0.45 – 1.51	0.534

Current living arrangements			
Renting	1.00		
Owns house	1.74	0.82 – 3.70	0.148
Living with primary family	2.21	1.04 – 4.69	0.038
Living with other family	2.02	0.87 – 4.70	0.103
Living with friends	1.48	0.36 – 6.03	0.585
No housing/Living in street	1.88	0.24 – 14.53	0.544
Others	1.56	0.16 – 14.94	0.698
DUP	1.05	1.02 – 1.08	0.003
Who made initial diagnosis			
Psychiatrist	1.00		
Medical officer	0.38	0.18 – 0.80	0.011
Clinical officer	0.51	0.34 – 0.75	0.001
Nurse	0.24	0.66 – 0.89	0.033
Nationality			
Uganda	1.00		
Other	1.25	0.68 – 2.28	0.201

DISCUSSION

Concordance of diagnoses at admission and on the MINI: About 50% of all diagnoses made at admission by health care workers agreed with the diagnoses confirmed using an extended version of the MINI. The low levels of agreement may be due to the instability of psychotic symptoms at FEP (Björkenstam, Björkenstam, Hjern, Reutfors, & Bodén, 2013; Kim et al., 2011). Bipolar disorder and Schizophrenia had the highest diagnostic accuracy which is in keeping with other studies from high income countries (Fusar-Poli et al., 2016; Hobbs, Stanbrook, & Chakraborty, 2017; Pope, Joobar, & Malla, 2013). Fourteen (14) participants (7.7%) met criteria for a substance induced psychotic disorder despite attempts to exclude all participants with substance use. Given the high prevalence of substance use among patients with FEP and its association with outcomes, it is essential that clinicians at the national referral hospital are advised to actively screen for substance use disorders in those with FEP (Mazzoncini et al., 2010; Myles, Myles, & Large, 2016)

Duration of untreated psychosis and concordance of diagnoses: A longer duration of untreated psychosis was associated with an accurate diagnosis. A diagnosis may be harder to make early in the course of psychotic disorders which are associated with many nonspecific symptoms before the onset of frank psychosis (Dutta et al., 2007; Heckers et al., 2013; Tandon, 2013). Unfortunately, longer durations of untreated psychosis have been associated with worse outcomes (Farooq, Large, Nielsens, & Waheed, 2009; Mossaheb et al., 2013; Oosthuizen et al., 2005; Penttila, Jaaskelainen, Hirvonen, Isohanni, & Miettunen, 2014). It is therefore imperative

that clinicians at the national referral hospital are taught to correctly diagnose psychotic disorders early in the course of the illness.

Living with primary family and diagnostic accuracy: Participants living with a primary family caregiver were two times more likely to have an accurate diagnosis as compared to those with other living arrangements. This is of public health importance in this setting as about half of participants were not living with a primary family member. Primary caregivers play a crucial role in the management of patients with psychotic disorders as they are an important resource of clearer history (Cairns, Reid, & Murray, 2015; McCann, Lubman, & Clark, 2011; Strand, Olin, & Tidefors, 2015). It is plausible that those who live with a primary caregiver were brought to the hospital and thus gave clearer medical history at admission. Also, family members are often not allowed to stay during the time of the admission due to space constraints and as such are not available to attend the ward round. In tools that assess the onset of psychotic symptoms like the Nottingham Onset Schedule of psychotic disorders, it is recommended that a family member who is conversant with the patient is interviewed when making an initial assessment (Singh et al., 2005). Future strategies need to be devised were a primary family member is available during the initial patient contact.

Level of training and a correct admission diagnosis: Psychiatrists were most accurate at diagnosing FEP followed by Psychiatric clinical officers and medical officers. However, most participants received a diagnosis from psychiatric clinical officers. This is due to the limited number of psychiatrists at the national referral and the country in general (Molodynski, Cusack, & Nixon, 2017). Effective and accurate diagnosis of FEP largely depends on the clinical skills possessed by the diagnosing clinician, with ground knowledge of all the possible underlying factors (Nasrallah, 2017). There should therefore be greater emphasis put in training the clinical officers to correctly diagnose a first episode of psychosis.

CONCLUSION: Early intervention for psychotic disorders is crucial to long term outcomes of psychotic disorders. This early intervention requires correct diagnosis to ensure initiation of appropriate management strategies. We have highlighted the poor agreement of the diagnoses of FEP at Uganda's national referral and teaching mental hospital. There is therefore need for additional skills training in the diagnosis and management of FEP. Specific emphasis must be placed on training lower cadre staff, emphasis on the duration of untreated psychosis in diagnostic accuracy as well as obtaining collateral history from primary family caregivers.

LIMITATIONS: As this was a cross sectional study design it is difficult to define causality. Longitudinal studies are required to determine the accuracy of diagnoses across the admission period. No standardized method was used to determine the duration of untreated psychosis and was only calculated by subtracting the age at which the patient remembers first developing symptoms from the current age. There are however ongoing studies to determine the true DUP of this cohort using a standardized instrument (E. K. Mwesiga et al., 2019).

ACKNOWLEDGEMENTS: We are indebted to Prof. David Sheehan who provided us with an extended version of the MINI to carry out this study.

DECLARATIONS: The authors declare no conflict of interests

FUNDING: The funding for this study was provided by the Neuropsychiatric Genetics in African Populations (NeuroGAP) Psychosis project (Anne Stevenson et al., 2019). The content is solely the responsibility of the authors and does not necessarily represent the official views of the project.

CONFLICT OF INTEREST: The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT: The fully anonymized data set is available on request from the corresponding author.

REFERENCES:

- Abbo, C., Ekblad, S., Waako, P., Okello, E., & Musisi, S. (2009). The prevalence and severity of mental illnesses handled by traditional healers in two districts in Uganda. *Afr Health Sci, 9 Suppl 1*, S16-22.
- Anderson, K. K., Norman, R., MacDougall, A., Edwards, J., Palaniyappan, L., Lau, C., & Kurdyak, P. (2018). Effectiveness of early psychosis intervention: comparison of service users and nonusers in population-based health administrative data. *American Journal of Psychiatry, 175*(5), 443-452.
- Anne Stevenson, Dickens Akena, Rocky E Stroud, Lukoye Atwoli, Megan M Campbell, Lori B Chibnik, . . . Karestan C Koenen. (2019). Neuropsychiatric Genetics of African Populations-Psychosis (NeuroGAPPsychosis): a case-control study protocol and GWAS in Ethiopia, Kenya, South Africa and Uganda. *BMJ Open*, e025469. doi:10.1136/bmjopen-2018-025469.
- Björkenstam, E., Björkenstam, C., Hjern, A., Reutfors, J., & Bodén, R. (2013). A five year diagnostic follow-up of 1840 patients after a first episode non-schizophrenia and non-affective psychosis. *Schizophrenia Research, 150*(1), 205-210.
- Breitborde, N. J. K., Srihari, V. H., & Woods, S. W. (2009). Review of the operational definition for first-episode psychosis. *Early intervention in psychiatry, 3*(4), 259-265. doi:10.1111/j.1751-7893.2009.00148.x
- Cairns, V. A., Reid, G. S., & Murray, C. (2015). Family members' experience of seeking help for first-episode psychosis on behalf of a loved one: a meta-synthesis of qualitative research. *Early Intervention in Psychiatry, 9*(3), 185-199.
- Drancourt, N., Etain, B., Lajnef, M., Henry, C., Raust, A., Cochet, B., . . . Zanouy, L. (2013). Duration of untreated bipolar disorder: missed opportunities on the long road to optimal treatment. *Acta Psychiatrica Scandinavica, 127*(2), 136-144.
- Dutta, R., Greene, T., Addington, J., McKenzie, K., Phillips, M., & Murray, R. M. (2007). Biological, Life Course, and Cross-Cultural Studies All point Toward the Value of Dimensional and Developmental Ratings in the Classification of Psychosis. *Schizophrenia Bulletin, 33*(4), 868-876. doi:10.1093/schbul/sbm059
- Farooq, S., Large, M., Nielssen, O., & Waheed, W. (2009). The relationship between the duration of untreated psychosis and outcome in low-and-middle income countries: a systematic review and meta analysis. *Schizophr Res, 109*(1-3), 15-23.
- Fusar-Poli, P., Cappucciati, M., Rutigliano, G., Heslin, M., Stahl, D., Brittenden, Z., . . . Carpenter, W. T. (2016). Diagnostic stability of ICD/DSM first episode psychosis diagnoses: meta-analysis. *Schizophrenia Bulletin, 42*(6), 1395-1406.
- Heckers, S., Barch, D. M., Bustillo, J., Gaebel, W., Gur, R., Malaspina, D., . . . Carpenter, W. (2013). Structure of the psychotic disorders classification in DSM-5. *Schizophrenia Research, 150*(1), 11-14. doi:<http://dx.doi.org/10.1016/j.schres.2013.04.039>
- Heeramun-Aubeeluck, A., Liu, N., Fischer, F., Huang, N., Chen, F., He, L., . . . Lu, Z. (2015). Effect of time and duration of untreated psychosis on cognitive and social functioning in Chinese patients with first-episode schizophrenia: A 1-year study. *Nord J Psychiatry, 69*(4), 254-261. doi:10.3109/08039488.2014.929738

- Hobbs, M., Stanbrook, R., & Chakraborty, N. (2017). Frequency of change of diagnoses in first-episode psychosis. *Progress in Neurology and Psychiatry, 21*(1), 6-8.
- Kigozi, F. N., Kizza, D., Nakku, J., Ssebunnya, J., Ndyabangi, S., Nakiganda, B., . . . Patel, V. (2016). Development of a district mental healthcare plan in Uganda. *The British journal of psychiatry, 208*(s56), s40-s46.
- Kim, J. S., Baek, J. H., Choi, J. S., Lee, D., Kwon, J. S., & Hong, K. S. (2011). Diagnostic stability of first-episode psychosis and predictors of diagnostic shift from non-affective psychosis to bipolar disorder: a retrospective evaluation after recurrence. *Psychiatry research, 188*(1), 29-33.
- Laursen, T. M., Munk-Olsen, T., & Gasse, C. (2011). Chronic somatic comorbidity and excess mortality due to natural causes in persons with schizophrenia or bipolar affective disorder. *PLoS one, 6*(9), e24597.
- Lencz, T., Knowles, E., Davies, G., Guha, S., Liewald, D. C., Starr, J. M., . . . Malhotra, A. K. (2014). Molecular Genetic Evidence for Genetic Overlap between General Cognitive Ability and Risk for Schizophrenia: A Report from the Cognitive Genomics Consortium (COGENT). *Mol Psychiatry, 19*(2), 168-174. doi:10.1038/mp.2013.166
- Marshall, M., & Rathbone, J. Early intervention for psychosis. (1469-493X (Electronic)). doi:D - nlm: ems57372
- D - NLM: PMC4163966 EDAT- 2011/06/17 06:00 MHDA- 2011/08/25 06:00 CRDT- 2011/06/17 06:00 AID - 10.1002/14651858.CD004718.pub3 [doi] PST - epublish
- Marshall, M., & Rathbone, J. (2011). Early intervention for psychosis. *Cochrane Database Syst Rev, 15*(6).
- Mazzoncini, R., Donoghue, K., Hart, J., Morgan, C., Doody, G. A., Dazzan, P., . . . Fearon, P. (2010). Illicit substance use and its correlates in first episode psychosis. *Acta Psychiatrica Scandinavica, 121*(5), 351-358.
- McCann, T. V., Lubman, D. I., & Clark, E. (2011). First-time primary caregivers' experience accessing first-episode psychosis services. *Early Intervention in Psychiatry, 5*(2), 156-162.
- McGorry, P. D. (2015). Early intervention in psychosis: obvious, effective, overdue. *J Nerv Ment Dis, 203*(5), 310-318. doi:10.1097/nmd.0000000000000284
- Molodynski, A., Cusack, C., & Nixon, J. (2017). Mental healthcare in Uganda: desperate challenges but real opportunities. *BJPsych international, 14*(4), 98-100.
- Mossaheb, N., Schloegelhofer, M., Kaufmann, R. M., Werneck-Rohrer, S., Zehetmayer, S., Malik, F., . . . Aschauer, H. (2013). Duration of untreated psychosis in a high-income versus a low- and middle-income region. *Aust N Z J Psychiatry, 47*(12), 1176-1182. doi:10.1177/0004867413504472
- Murru, A., & Carpiniello, B. (2018). Duration of untreated illness as a key to early intervention in schizophrenia: A review. *Neuroscience Letters, 669*, 59-67. doi:<https://doi.org/10.1016/j.neulet.2016.10.003>
- Mwesiga, E. K., Nakasujja, N., Nakku, J., Nanyonga, A., Gumikiriza, J. L., Bangirana, P., . . . Musisi, S. (2019). One year prevalence of psychotic disorders among first treatment contact patients at Butabika National Psychiatric Referral Hospital in Uganda. *bioRxiv, 669606*. doi:10.1101/669606
- Mwesiga, E. K., Nakasujja, N., Ongeru, L., Semeere, A., Loewy, R., & Meffert, S. (2019). A cross-sectional mixed methods protocol to describe correlates and explanations for a long duration of

- untreated psychosis among patients with first episode psychosis in Uganda. *BMJ Open*, 9(7), e028029. doi:10.1136/bmjopen-2018-028029
- Myles, H., Myles, N., & Large, M. (2016). Cannabis use in first episode psychosis: Meta-analysis of prevalence, and the time course of initiation and continued use. *Australian & New Zealand Journal of Psychiatry*, 50(3), 208-219.
- Nasrallah, H. A. (2017). For first-episode psychosis, psychiatrists should behave like cardiologists. *Current Psychiatry*, 16(8), 4-7.
- Norman, R. M., & Manchanda, R. (2016). Prevention and early intervention program for psychoses (PEPP). *Healthc Q*, 18, 37-41.
- Oosthuizen, P., Emsley, R. A., Keyter, N., Niehaus, D. J., & Koen, L. (2005). Duration of untreated psychosis and outcome in first-episode psychosis. Perspective from a developing country. *Acta Psychiatr Scand*, 111(3), 214-219.
- Organization, W. H. (2017). Depression and other common mental disorders: global health estimates.
- Penttila, M., Jaaskelainen, E., Hirvonen, N., Isohanni, M., & Miettunen, J. (2014). Duration of untreated psychosis as predictor of long-term outcome in schizophrenia: systematic review and meta-analysis. *Br J Psychiatry*, 205(2), 88-94.
- Pope, M. A., Joober, R., & Malla, A. K. (2013). Diagnostic stability of first-episode psychotic disorders and persistence of comorbid psychiatric disorders over 1 year. *The Canadian Journal of Psychiatry*, 58(10), 588-594.
- Rathod, S., Pinninti, N., Irfan, M., Gorczynski, P., Rathod, P., Gega, L., & Naeem, F. (2017). Mental health service provision in low-and middle-income countries. *Health services insights*, 10, 1178632917694350.
- Rost, K. M., Meng, H., & Xu, S. (2014). Work productivity loss from depression: evidence from an employer survey. *BMC health services research*, 14(1), 597.
- Singh, S. P., Cooper, J. E., Fisher, H. L., Tarrant, C. J., Lloyd, T., Banjo, J., . . . Jones, P. (2005). Determining the chronology and components of psychosis onset: The Nottingham Onset Schedule (NOS). *Schizophr Res*, 80(1), 117-130.
- Strand, J., Olin, E., & Tidefors, I. (2015). Mental health professionals' views of the parents of patients with psychotic disorders: a participant observation study. *Health & social care in the community*, 23(2), 141-149.
- Swartz, L., Kilian, S., Twesigye, J., Attah, D., & Chiliza, B. (2014). Language, culture, and task shifting—an emerging challenge for global mental health. *Global health action*, 7(1), 23433.
- Tandon, R. (2013). Definition of psychotic disorders in the DSM-5 too radical, too conservative, or just right! *Schizophrenia Research*, 150(1), 1-2. doi:<http://dx.doi.org/10.1016/j.schres.2013.08.002>
- Whiteford, H. A., Ferrari, A. J., Degenhardt, L., Feigin, V., & Vos, T. (2015). The global burden of mental, neurological and substance use disorders: an analysis from the Global Burden of Disease Study 2010. *PLoS one*, 10(2), e0116820.

