

Cohort Profile Update: GendAge and Berlin Aging Study II (BASE-II)

Ilja Demuth^{1,2,*}, Verena L. Banszerus¹, Johanna Drewelies³, Sandra Düzel⁴, Ute Seeland⁵, Dominik Spira¹, Esther Tse⁵, Julian Braun^{6,7}, Lars Bertram^{8,9}, Andreas Thiel^{6,7}, Ulman Lindenberger^{4,10}, Vera Regitz-Zagrosek^{5,11,12}, Denis Gerstorff³

Additional BASE-II/GendAge investigators (=collaborators) are listed on page 20 of this manuscript

¹Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health; Department of Endocrinology and Metabolism; 10117 Berlin, Germany.

²BCRT - Berlin Institute of Health Center for Regenerative Therapies, Charité - Universitätsmedizin Berlin, Berlin, Germany.

³Department of Psychology, Humboldt-Universität zu Berlin

⁴Center for Lifespan Psychology, Max Planck Institute for Human Development, Berlin, Germany

⁵Berlin Institute for Gender in Medicine, Charité – Universitätsmedizin Berlin

⁶Si-M / “Der Simulierte Mensch” a science framework of Technische Universität Berlin and Charité - Universitätsmedizin Berlin, Berlin, Germany

⁷Regenerative Immunology and Aging, BIH Center for Regenerative Therapies, Charité Universitätsmedizin Berlin, Berlin, Germany

⁸Lübeck Interdisciplinary Platform for Genome Analytics, Institutes of Neurogenetics and Cardiogenetics, University of Lübeck, Lübeck, Germany;

⁹Center for Lifespan Changes in Brain and Cognition (LCBC), Dept of Psychology, University of Oslo, Oslo, Norway;

¹⁰Max Planck UCL Centre for Computational Psychiatry and Ageing Research, Berlin, Germany, and London, United Kingdom

¹¹DZHK (German Centre for Cardiovascular Research), partner site Berlin

¹²Department of Cardiology, University Hospital Zürich, University of Zürich, Switzerland

*Corresponding author:

Ilja Demuth (Ph.D.)
Charité - Universitätsmedizin Berlin
Augustenburger Platz 1
13353 Berlin
Email: ilja.demuth@charite.de
Phone: ++49 30 450 569 143

Abstract

The study “Sex- and gender-sensitive prevention of cardiovascular and metabolic disease in older adults in Germany”, the GendAge study, focusses on major risk factors for cardiovascular and metabolic diseases and on the development of major outcomes from intermediate phenotypes in the context of biological sex and gender differences. It is based on a re-investigation of participants of the Berlin Aging Study II (BASE-II). The Berlin Aging Study II (BASE-II) is aiming at identifying factors that distinguish healthy from unhealthy ageing and completed baseline assessments in 2,200 adult volunteers (1,600 participants aged 60-80 years and 600 participants aged 20-35 years) in 2014. The BASE-II follow-up assessments of 1,100 men and women aged 65-94 years in 2018-2020 were part of GendAge. In addition to re-assessing most baseline measures (geriatrics, internal medicine, immunology and psychology) we implemented a comprehensive gender questionnaire covering socio-cultural gender characteristics and added high-quality echocardiography.

The original BASE-II cohort

The Berlin Aging Study II (BASE-II) was launched as a multidisciplinary study aiming at identifying factors that distinguish ‘healthy’ from ‘unhealthy’ ageing. Baseline recruitment of 2,200 adult volunteers from the Berlin metropolitan area and baseline assessments were completed in 2014¹. The ascertainment protocol included the collection of data from different domains for each of the 1,600 participants aged 60-80 years and 600 participants aged 20-35 years, namely geriatrics and internal medicine, immunology, genetics, psychology, sociology, and economics^{1,2}.

BASE-II baseline data were used in a multitude of analysis projects focusing on key questions revolving around age and aging. Research topics of the ongoing study include, but are not limited to, cognitive aging³⁻⁵, cardiovascular and metabolic health⁶⁻⁸, sarcopenia and frailty^{9,10}, psychosocial factors of aging^{11,12}, genetic risk factors of aging and disease¹³⁻¹⁵, the impact of characteristics of the neighborhood people are living in¹⁶, as well as indicators of biological age^{17,18} and immune biomarkers¹⁹. For an overview of the BASE-II research foci and publications, refer to²⁰ and the BASE-II website (<https://www.base2.mpg.de/en/project-information/publications>).

What is the reason for the new data collection?

The study “Sex- and gender-sensitive prevention of cardiovascular and metabolic disease in older adults in Germany”, the GendAge study, focusses on major risk factors for cardiovascular and metabolic diseases and on the development of major outcomes from intermediate phenotypes in the context of biological sex and gender differences. Major outcomes include but are not limited to myocardial infarction (MI), heart failure (HF) and T2D, as well as mortality and quality of life. Major aim is the first systematic collection of follow-up data in BASE-II participants and the analysis of sex- and gender-related differences. Gender was obtained by a novel comprehensive gender questionnaire covering a range of socio-cultural gender characteristics as a central instrument. This questionnaire contains an

adapted version of the gender questionnaire developed by Pelletier and colleagues²¹ and additionally included the variables used to calculate a gender score retrospectively by making use of gender-related variables assessed at baseline²². This gender questionnaire will be central to develop a second gender score (GS-II).²³

What will be the new areas of research?

There is new knowledge showing that sex differences play a role in all major diseases, their prevention and treatment²⁴. Other studies showed that gender as the sociocultural dimension of disease affects disease and treatment outcomes and also well-being^{21,25}. The new areas of research cover the systemic inclusion of sex-specific analysis and the inclusion of gender. Aging interacts with sex and gender differences in health, but it is not clear, which mechanisms are most important.

GendAge aims to better understand, which mechanisms affect cardio-metabolic morbidity, mortality, and quality of life among older adults in a sex- and gender-sensitive manner.

While on different occasions follow-up data were ascertained for questionnaire and cognitive data^{5,26-30}, as being part of the GendAge study this cohort profile update describes, the first comprehensive follow-up assessments in BASE-II that also includes a re-assessment of central variables in the areas of internal medicine and geriatrics. At the same time, the other BASE-II research foci established over the past 10 years as described above (and in²⁰) will be continued and strengthened by the transition of BASE-II into a longitudinal study.

Who is in the cohort?

The original BASE-II sample consisted of 2,200 participants from the greater metropolitan area of Berlin, Germany (baseline assessment between 2009 and 2014). The follow-up assessments within the GendAge study took place between 22 June 2018 and 10 March

2020 at the Charité Universitaetsmedizin Berlin. During the recruitment of the follow-up cohort, we approached all participants of the remaining pool of 1,428 subjects out of the originally 1,671 subjects who completed the baseline medical assessments at an age of 60 years and older. Potential follow-up participants were contacted via telephone and an invitation letter containing the comprehensive participant's information sheet and the letter of consent was sent at least five days before the scheduled first of two assessment days to all subjects who agreed to participate at least five days before the scheduled first of two assessment days. Additionally, we included 17 subjects of the older age group who were not medically assessed at baseline, but were part of the BASE-II sample with data collected from at least one of the BASE-II partner sites. As presented in the flow chart (Figure 1), this resulted in a total of 1,100 participants of the older BASE-II group investigated in the follow-up. These older participants had a mean age of 75.60 years (SD \pm 3.77, range 64.91 - 94.07 years), with up to 10.37 years of follow-up (N=1,083, mean follow-up at 7.35 years, (SD \pm 1.46), range 3.91 - 10.37). At baseline, the BASE-II included a group of 600 younger subjects aged 20-35 years serving as a reference population ¹, of which 500 completed baseline medical assessments. Between 7 February 2020 and 13 March 2020, we performed follow-up assessments in a total of 64 participants of this younger group until these assessments were suspended because of the SARS-Cov2 pandemic. These younger participants had a mean age of 36.81 years (SD \pm 3.46, range 29.32 - 44.05 years), with up to 10.7 years of follow-up (minimum 6.08 years, mean follow-up at 8.16 years, SD \pm 1.58).

At follow-up, almost all of the older participants were retired (97.3%) as compared with 86% at the time of baseline assessment. At baseline, BASE-II participants were characterized by higher education and better self-reported health status than the general population of Berlin and Germany ¹. At follow-up, this selection seems to have increased, with 68.8% of the participants reported to have a high school degree (51% at baseline) and about 61% rated their health as *very good* or *good* (40% at baseline). The rate of divorce had been above average at baseline with 29% and had dropped to 21.7% at follow-up, which is still

significantly above the German and Berlin average (i.e. 12.0% and 17.4%, respectively)³¹, while the proportion of widowed participants increased from 5% at baseline to 10.5% in the follow-up dataset of older BASE-II participants. As shown in Table 1, differences between men and women are evident with respect to the sociodemographic status and psychosocial functioning in the follow-up cohort: Men reported significantly higher school degrees and higher satisfaction with life in general than women. Interestingly, self-rated health did not differ between men and women, which matches to the overall morbidity estimated by an adapted version of the Charlson morbidity index^{17,32}, which also did not differ between men and woman ($p=0.981$, Table 1). This morbidity index, however, increased between baseline and follow-up ($p<0.001$, Wilcoxon signed-rank test and data not shown). Differences between men and women exist in the follow-up dataset with respect to the prevalence of some, but not all cardiovascular risk factors and diseases (Table 1). Men for example had a higher BMI and a higher proportion of men reported to have T2D and myocardial infarction than women. No significant differences between men and women were evident in the reporting of hypertension, peripheral artery disease, and stroke. With the aim of achieving a particularly high quality standard in the assessment of participant's medical history at baseline and follow-up, including past and current diseases, the information given by the participants was recorded from study physicians as part of a structured one-to-one interview, allowing to consider its plausibility. This, however, does not cover the gap between reported (anamnestic) diseases and the diseases diagnosed in the course of the study. This is exemplified by T2D, which was reported by 15.6% and 8.8% of men and women, respectively. In contrast, this disease was diagnosed in 20.7% of men and 13.3% of women based on the ESC criteria 2019³³, indicating that a substantial proportion of almost 30% was unaware of the disease. The T2D prevalence increased from 12.7% at baseline to 16.7% at follow-up in the total cohort.

What has been measured?

With the aim of investigating human ageing processes in BASE-II under consideration of different disciplines and longitudinally, the baseline investigation aimed at the most comprehensive data collection possible. At follow-up, most of these data in the field of geriatrics, internal medicine and psychology were again part of the study protocol (for a select overview, see Table 2). As the follow-up assessment being part of the GendAge study, we have implemented a comprehensive gender questionnaire covering a range of socio-cultural *gender* characteristics as a central instrument. This questionnaire contains an adapted version of the gender questionnaire developed by Pelletier and colleagues²¹ and additionally included the variables used to calculate a gender score retrospectively by making use of gender-related variables assessed at baseline²². This gender questionnaire will be central to develop a second gender score (GS-II) and to reach the GendAge goals.

With a focus on cardio-metabolic diseases in GendAge, we extended the broad range of data assessed in this area at baseline by echocardiography. Data on right and left ventricular and atrial morphology and systolic and diastolic function and vascular stiffness were obtained by a trained investigator (US) on a General Electric Vivid T8 R3 System and analyzed by a trained analyst (ET). 10% of echo analysis were controlled by an independent supervisor. The intraclass correlation coefficient (ICC) between analyst and independent supervisor was 0.87 (LV IVSd: 0.86, LVEDV Biplane: 0.95, LVEF Biplane: 0.87, E/A: 0.93). LVEF was higher in women than in men, in agreement with most recent publications, and probably partially due to higher rate of MI in men³⁴.

Peripheral blood mononuclear cells were prepared from 903 participants at follow-up and frequencies as well as absolute counts of recent thymic emigrants (RTEs), T_{EMRA} effector T cell subsets (T_{EMRA}) and cytotoxic $CD4^+$ T cells were directly assessed. While RTEs are known to decrease with ageing³⁵, alterations in T_{EMRA} and specialised cytotoxic $CD4^+$ T cell compartments can be indicative of age-related perturbations of systemic T cell immunity³⁶.

Similar to baseline, we determined numerous routine laboratory parameters from blood and urine (Table 2), and also stored blood plasma/serum and urine samples for future analyses.

Genomic DNA was already extracted from EDTA-blood and buccal swab samples from GendAge participants, which will be used e.g. for the profiling of genome-wide DNA methylation signatures and new genome-wide single nucleotide polymorphism genotyping experiments (Table 2). In-between the two assessment days at the Charité, participants were asked to fill out a comprehensive psychosocial take-home questionnaire and return this at their second Charité visit. Moreover, another wave of cognitive assessments (see Table 2) carried out by the Max Planck Institute for Human Development (MPIB) has been tightly linked to the GendAge assessment of BASE-II participants. The cognitive session (= third study visit) was followed 7 days after the second medical examination and lasted about 4.5 to 5 hours. Subjects were tested in groups of 4–6 individuals. The cognitive battery included 17 measures of learning and memory performance, attention/processing speed, working memory, executive functioning, and perceptual speed (see Table 2). Within the week between study visit 2 (Charité) and 3 (MPIB) accelerometers (ActiGraph wGT3X-BT) have been used to track participants' physical activity and sleep in a subset of our participants (N=750).

After the cognitive session, participants were invited to take part in a one-to-one interview on a different day. This additional individual assessment took up to 60 minutes and serves as a cohort comparison between the BASE and BASE-II study populations. This additional data collection will also contribute to the BASE-II cognitive waves, allowing us to further investigate individual differences in aging trajectories (for an overview, refer to²⁰).

Furthermore, and as part of a collaboration with the Lifebrain study, a consortium of European studies funded by the EU Horizon 2020 Framework Programme³⁷, we collected blood samples using dried blood cards, in order to determine laboratory parameters with identical methods used for all Lifebrain participating sites. Lifebrain aims at identifying determinants of healthy lifespan development by integrating and harmonizing data and results from 11 large and predominantly longitudinal European samples from 7 countries.

This has yielded a database of fine-grained measures focusing on brain and cognition from more than 7,000 individual participants.

The GendAge study was approved by the Ethics Committee of the Charité–Universitätsmedizin Berlin (approval number EA2/144/16) and all participants gave written informed consent. GendAge is registered in the German Clinical Trials Register (Study-ID: DRKS00016157). The cognitive battery was approved by the Ethics Committee of the Max-Planck-Institute and the genomics experiments were approved by the Ethics Committees of the Charité (approval number EA2/144/16) and the University of Lübeck (approval numbers AZ19-390A and 19-391A).

What has been found?

GendAge was initiated in 2017 with a focus on sex and gender differences in the ongoing analyses of baseline BASE-II data in the area of cardio-metabolic risk factors and disease, some of which we would like to highlight here. The metabolic syndrome (MetS) combines the five cardiovascular risk factors of high waist circumference, low high density lipoprotein cholesterol, high fasting blood glucose, high triglycerides, and high systolic/diastolic blood pressure to diagnose the MetS based on cutoff values for each of the risk factors³⁸. While this approach is certainly useful in the clinical setting, it does not consider the full range of information of the continuously measured risk factors. We therefore applied confirmatory factor analysis to establish a latent metabolic load factor (MetL), representing a single continuous measure based on the five MetS risk factors. The MetL was on average significantly higher in men compared to women, and the factor was associated with more morbidity and poorer kidney function in both men and women, and with lower performance on a measure of cognitive function in men⁷. Strong evidence suggests that higher levels of Lipoprotein (a) [Lp(a)] are an independent risk factor for coronary heart disease³⁹. Interestingly, at the same time higher Lp(a) is associated with a reduced type 2 diabetes (T2D) risk, a finding which has been reported from numerous studies, including BASE-II

^{6,40,41}. We recently analyzed the relationship between Lp(a) and cognitive performance assessed with the *Consortium to Establish a Registry for Alzheimer's Disease (CERAD)-Plus* test battery. The results suggested that lower Lp(a) levels associate with better cognitive performance, at least in men ⁸. The analysis of arterial stiffness data from BASE-II assessed at baseline revealed sex differences in measures of arterial wave reflection, with an on average higher augmentation index in women than in men, a measure of endothelial function of the small and medium arteries. The results also suggested that women with exogenous intake of sex hormones for oral contraception and suppressed endogenous estradiol levels had higher levels of the augmentation index than non-users ⁴². T2D is an age-associated disease which is reflected by the 4.8% increase in the proportion of affected BASE-II participants within the 7.35 (SD \pm 1.46) years between baseline ⁶ and follow-up assessments. This is even more dramatic since almost one third of the participants was not aware of the disease (Table 1). In addition being a risk factor for cardiovascular disease and cognitive decline, a recent analysis of BASE-II data suggested that the capacity to regulate the glucose metabolism is associated with well-being among older men ⁴³.

The immunological screening has so far revealed significantly higher frequencies of RTEs in women as compared to men, indicating a higher thymic T cell production even at the advanced ages of the GendAge participants. In men, more CD45RA⁺ re-expressing TEMRAs were detected than in women (Table 1). These cells are associated with chronic viral infections (e.g. CMV) and can serve as a signature of immune-senescence ⁴⁴. We found no significant difference in the frequencies of cytotoxic CD4⁺ T cells. Together, these preliminary findings confirm the better immune status of aged women as compared to men. A detailed analysis of the datasets will identify additional correlates of sex and gender, aging, and the immune system.

In GendAge, we have developed a retrospective gender score taking BASE-II baseline data reflecting sociocultural aspects (e.g., level of education, marital status, and chronic stress) into account. This gender score (GS-I) was associated with a number of clinical and psychosocial variables and performed better in predicting differences in a subset of variables

compared to biological sex²². As mentioned above, we have implemented a comprehensive gender questionnaire as part of the follow-up assessments described here, including not only the variables required to re-calculate the GS-I based on the follow-up data, but also the items to develop an adapted version of the gender questionnaire proposed by Pelletier and colleagues²¹. This gender questionnaire will be central to develop a second gender score (GS-II) and to reach the GendAge goals.

What are the main strengths and weaknesses?

The BASE-II follow-up assessments covered most of the medical, psycho-social and cognitive domains and variables assessed at baseline, and thereby taking the BASE-II characteristic of an exceptionally broad and in-depth data collection to a next, longitudinal level. In addition, and in the context of the GendAge focus on cardio-metabolic disease, we extended the assessments in this area e.g. by including high-quality echocardiography resulting in a unique data collection. This strength with respect to comprehensive and longitudinal data offers the potential to answer a number of questions that are of crucial relevance for the health of old women and men. Thus, GendAge will make important contributions for improvements in understanding the health and well-being of older adults in both genders. BASE-II was initiated as a multidisciplinary study with expertise in a broad range of fields relevant for aging research (e.g. internal medicine and geriatrics, biology, psychology, genetics, immunology, socio-economics, and now in GendAge further extended by socio-cultural aspects of gender). The past ten years of BASE-II research have shown that multidisciplinary collaboration is not only a statement of intent, but a fruit-bearing working posture and a clear strength of BASE-II.

Sampling bias is a challenge which cohort studies have to deal with, and this is especially an issue in the follow-up of older study populations such as the older group of BASE-II participants. To address this, we have made a considerable effort (e.g. offering a taxi service

for participants not able to travel independently) to include as many participants in the follow-up as possible. Additionally, and similar to baseline, we are able to systematically quantify the sampling bias and even account for it when it comes to the question of generalizability of study results to a population as a whole (e.g., Berlin or Germany), due to the evaluation of selectivity and representativeness via the German Socio-Economic Panel Study (SOEP)¹. Despite these possibilities we cannot rule out the possibility of a selection bias completely, which certainly is a weakness of this study, a weakness that applies to all cohort studies relying on voluntary participants who have been non-randomly recruited. With our direct comparability to the national representative SOEP study, we are in a position though to quantify the amount of selectivity and, if need, take measures to correct and adjust our results.

Can I get hold of the data? Where can I find out more?

More details on GendAge can be found at <https://gendage.charite.de/en/> and information on the BASE-II as a whole is available at <https://www.base2.mpg.de/en>. BASE-II has a tradition of sharing data and biobank samples in joint collaborative projects which will be continued with respect to BASE-II data assessed in GendAge. Interested groups are invited to contact the study coordinating PI Ilja Demuth at ilja.demuth@charite.de for the data-sharing application form. Each application will be reviewed by the GendAge PIs (currently ID, VRZ and DG) and the decision communicated to the applicants usually within 4 weeks of submission.

Funding statement

The GendAge study research project (Co-PIs are Ilja Demuth, Vera Regitz-Zagrosek, and Denis Gerstorff) is supported by the German Federal Ministry of Education and Research (Bundesministerium für Bildung und Forschung, BMBF) grant numbers 01GL1716A (ID and VRZ) and 01GL1716B (DG). Genomics assessments are funded by the Cure Alzheimer's Fund (as part of the "CIRCUITS-AD" consortium project) and the European Research Council's "Horizon2020" funding scheme (as part of the "Lifebrain" consortium project; both to LB). Additional contributions (e.g., equipment, logistics, personnel) are made from each of the participating sites.

Acknowledgements

We are deeply indebted to all individuals of the BASE-II cohort who have agreed to participate in the follow-up assessments. We would further like to thank all staff members and students involved in the medical examinations of participants and documentation, especially Nora Berger, Janina Dombrowski, Mergim Domuzeti, Elisa Dreiig, Ilona Enarovic, Anthony Ganswindt, Thomas Grenkowitz, Anna Hunold, Ilias Katsianas, Yoo-Ri Kim, Jrn Kiselev, Paula Krull, Christine Kytmanow, Elisa Lemke, Julia Mtzkow, Charlotte Mentzel, Nadja Mielke, Narantuya Mishig, Angela Motz, Eduard Nitschke, Aabi Okute, Danai Pantelakis, Sophie Poser, Johanne Spieker and Taleka Vollmar (all Charit Universitaetsmedizin Berlin, Biology of Aging group); and Luisa Lth, Mariebelle Kaus, Isabel Ganter, Marlene Rosendahl, Jasmin Boneberger, Antonia Sprenger, Hania El-Kersh (all Humboldt University, Department of Psychology), and Kirsten Becker for the excellent daily organization of the running study within and between the research units, the Telefonstudio for their daily contact with the participants, Martin Becker for the continuous quality checks of our sample and Berndt Wischnewski for updating the database and cognitive battery (all Max Planck Institute for Human Development).

We kindly acknowledge the excellent cooperation with the Central Biomaterial Bank, the joint core facility of the Charité-Universitätsmedizin Berlin and the Berlin Institute for Health (BIH). Furthermore, we appreciate the great support by the BIH-REDCap team and especially from Andreas Hetey.

All co-workers listed above have confirmed their agreement to be acknowledged in this paper.

Conflict of interest: None declared.

Tables

Table 1: Selection of BASE-II follow-up characteristics as assessed of the GendAge study – older group of participants

	Total number of observations	Women¹ (N=573, 52.1%)	Men¹ (N=527, 47.9%)	p-value²
Age (years)	1,100	75.7 (± 3.5)	75.5 (± 4.0)	0.276
Highest school degree Elementary school Intermediate school High school	1,095	35 (6.1%) 183 (32.0%) 354 (61.9%)	18 (3.4%) 104 (19.9%) 401 (76.7%)	8.4×10 ⁻⁷
Family status Married Not married, in partnership Single Divorced Widowed Other	1,098	218 (38.0%) 12 (2.1%) 60 (10.5%) 187 (32.6%) 89 (15.5%) 7 (1.2%)	386 (73.5%) 19 (3.6%) 33 (6.3%) 51 (9.7%) 26 (5.0%) 10 (1.9%)	3.0×10 ⁻³⁴
Employment status Retired	1,055	540 (97.6)	486 (96.8)	0.689
Self-rated health Very good Good Fair Poor or very poor	1,096	56 (9.8%) 284 (49.7%) 166 (29.0%) 66 (11.5%)	65 (12.4%) 262 (50.0%) 143 (27.3%) 54 (10.3%)	0.499
Satisfaction with life in general	1,097	7.9 (±1.6)	8.1 (±1.4)	0.034
Digit Symbol Substitution Test³	1,095	41.82(±8.28)	39.65(±8.95)	3.4×10 ⁻⁵
Depression (ever diagnosed)	1,095	122 (21.3%)	63 (12.0%)	5×10 ⁻⁵
BMI	1,098	26.6 (±4.7)	27.4 (±3.7)	0.005
Diabetes mellitus type II (self-reported)	1,097	49 (8.6%)	82 (15.6%)	3.7×10 ⁻⁴
Diabetes mellitus type II (diagnosed/ ESC criteria 2019)	1,097	76 (13.3%)	109 (20.7%)	0.001
Metabolic Syndrome (diagnosed, AHA/IDF/NHLBI criteria 2009)	1,074	252 (45.5%)	327 (62.9%)	1.2×10 ⁻⁸
Hypertension	1,097	296 (51.7%)	311 (59.2%)	0.013
Myocardial infarction	1,097	11 (1.9%)	24 (4.6%)	0.015
Stroke	1,096	13 (2.3%)	20 (3.8%)	0.158
Peripheral artery disease	1,094	8 (1.4%)	15 (2.9%)	0.138
Morbidity index	955	1.0 (IQR 2.0)	1.0 (IQR 2.0)	0.594
Pulse wave velocity (m/s)	932	11.21 (±0.92)	11.04 (±0.91)	0.006
Left ventricular ejection fraction (%)	774	64.16 (±6.42)	62.90 (±5.75)	0.004

Frailty (Fried) Not frail Pre-frail Frail	1,087	260 (45.4%) 280 (48.9%) 28 (4.9%)	251 (47.6%) 248 (47.1%) 20 (3.8%)	0.542
Maximal hand grip strength (kg)	1,098	20.5 (±4.4)	35.1 (±6.8)	3.1×10 ⁻²³⁶
Recent thymic emigrants (naïve CD4⁺ T cells)	395 ³	64.69 (± 16.34)	51.03 (± 13.69)	6.3×10 ⁻¹⁶
T_{EMRA} (effector memory T cells re-expressing CD45RA)	395 ⁴	32.82 (± 19.96)	34.94 (± 21.29)	0.309
Cytotoxic SLAMF7+ CD4⁺ T cells	181 ⁵	6.02 (± 5.98)	6.05 (± 6.60)	0.974

¹Data are presented as N (%), mean ±standard deviation or median (interquartile range, IQR). ²Differences between women and men were assessed using the parametric t test, the nonparametric Mann-Whitney U test or the χ^2 where appropriate. ³assessed at study visit 1 ⁴903 expected to be available after completion of the analyses. ⁵700 expected to be available after completion of the analyses.

Table 2: BASE-II follow-up assessments during the three study visits in the GendAge study

Type of assessment/ domain	Example assessments/tests
Physical examination and medical history	Medical history structured by organ systems, medication, body weight, height, lifestyle (including smoking status, alcohol consumption, physical activity)
Physical status and functional tests	Tinetti Mobility Test, Timed up & Go Test, Barthel Index (ADL), Lawton Instrumental Activities of Daily Living Scale (IADL), hand grip strength, anthropometric parameters, pulse wave velocity/arterial stiffness (Mobil-o-Graph), echocardiography, electrocardiography (ECG), spirometry, motion monitoring (Actigraph), dual-energy X-ray absorptiometry (DXA) ¹ ,
Psychological screening tests	Mini Mental State Examination (MMSE), Digit Symbol Substitution Test (DSST) ² , Center for Epidemiologic Studies Depression Scale (CESD)
Questionnaires	EPIC (food-frequency questionnaire), Gender Questionnaire, Pittsburgh Sleep Quality Index, Rapid Assessment of Physical Activity, SARC-F , SF-36
Laboratory values ³	Blood, serum or plasma: 25-hydroxyvitamin D, apolipoprotein A1, apolipoprotein B, basophiles, calcium, cortisol, creatinine, creatinine kinase, C-reactive protein, cystatin C, dehydroepiandrosterone, eosinophils, erythrocytes, ferritin, folic acid, gamma-glutamyltransferase, glucose 1, glucose 2 ⁴ , glutamate, oxalacetate transaminase, glutamate-pyruvate transaminase, HbA1c, HDL-cholesterol, hematocrit, hemoglobin, homocysteine, international normalized ratio, iron, LDL-

	<p>cholesterol, leukocytes, lipoprotein (a), lymphocytes, magnesium, MCH, MCHC, MCV, monocytes, neutrophils, oestradiol, osteocalcin, partial thromboplastin time, RDW, sex hormone-binding globulin, testosterone, thrombocytes, thyroid-stimulating hormone, thyroxine, total-cholesterol, triglycerides, triiodothyronine, urea, uric acid, vitamin B12, zinc.</p> <p>Urine: albumine, creatinine, desoxypyridinoline, teststrip: bilirubin, blood (erythrocytes), glucose, ketones, leukocytes, nitrite, pH value, protein, specific weight, urobilinogen</p> <p>Dried blood cards: arsenic, brain derived neurotropic-factor, cadmium, chromium, fatty acids (C12:0, C14:0, C15:0, C16:0, C16:1n7, C17:0, C18:0, C18:1,t6-11, C18:1,c9, C18:1,c11, C18:2,n-6, C20:0, C18:3,n-6, C18:3,n-3, C20:1,n-9, C20:2,n-6, C22:0/C20:3,n-6, C20:4,n-6, C20:5,n-3, C24:0, C22:5,n-3, C22:6,n-3, unknown), HbA1c, hsCRP, lead, mercury, nickel, total-cholesterol</p>
Genomics	Genome-wide single nucleotide polymorphism genotyping using the “Global Screening Array” (Illumina, Inc.); genome-wide DNA methylation profiling using the “Infinium MethylationEPIC” array (Illumina, Inc.)
Psychosocial questionnaire	Well-being, positive affect and negative affect, emotion regulation, stress, personality, control beliefs, domain-specific control, time perception, embitterment, loneliness, solitude, social activities, network structure, sexuality, risk behavior, etc.

Biobanking	Blood plasma & serum, urine, DNA extracted from EDTA-blood and buccal swaps
Cognitive tests	<i>Episodic memory</i> (Picture-Word-Task, Face-Profession-Task, Object Location Task, Scene-Encoding, Verbal learning and memory test), <i>Working memory</i> (Letter Updating, Spatial Updating, Number-N-Back), <i>Executive functioning / processing speed</i> (Multi-Source-Interference Task, Digit Symbol Substitutions Test ² , Fluid intelligence (Letter series, Number series, Practical Problems), <i>Subjective Health Horizon Questionnaire</i> (SHH-Q)
Immunological assessment	Cryopreservation of whole blood (SmartTube system) or isolated PBMCs, and serum samples. Direct ex vivo staining of recent thymic emigrants (RTE, CD31+ CD45RA+ CD4+ T cells), TEMRA (CD45RA+ CD8+ T cells), Tregs (CD25bright CD127- CD4+ T cells), cytotoxic CD4+ T cells, amongst others using four different panels: 1) ImmunoCount Panel (CD45, CD3, CD56, CD19, CD16, CD14, CD123, CD1c); 2) RTE panel (CD3, CD4, CD8, CD45RA, CCR7, CD31, CD95, CD11a); 3) TREG panel (CD3, CD4, CD8, CD25, CD127); 4) Effector T cell panel (CD3, CD4, CD8, CD45RA, CCR7, SLAM-F7, IL-6R, CD57, PD-1). Panels were measured on MacsQuant 10 (Miltenyi), MacsQuant 16 (Miltenyi) or LSR II (BD)

¹only available for older group; ²assessed at study visit 1 and 3, ³blood samples were drawn after a fasting period of at least 8 hours (if not otherwise indicated), ⁴post-load (75g glucose, 2h), not assessed in participants with known diabetes

Additional BASE-II/GendAge investigators:

Nikolaus Buchmannⁱ, Peter Eibichⁱⁱ, Friederike Kendelⁱⁱⁱ, Maximilian König^{iv}, Christina M. Lill^v,
^{vi}, Maïke Mangold^{vii}, Ahmad Tauseef Naumanⁱⁱⁱ, Kristina Norman^{viii, ix}, Graham Pawelec^{x, xi},
Elisabeth Steinhagen-Thiessen^{iv}, Sarah Toepfer^{iv}, Valentin Max Vetter^{iv}, Gert G. Wagner^{xii},
Ursula Wilkenschoffⁱ, Kilian Wistuba-Hamprecht^{xiii}

ⁱCharité; Department of Cardiology, Charité - University Medicine Berlin (Campus Benjamin Franklin), Berlin, Germany; ⁱⁱMax Planck Institute for Demographic Research, Rostock, Germany; ⁱⁱⁱBerlin Institute for Gender in Medicine, Charité – Universitätsmedizin Berlin; ^{iv}Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health; Department of Endocrinology and Metabolism, Berlin, Germany; ^vSection for Translational Surgical Oncology and Biobanking, Department of Surgery, University of Lübeck and University Medical Center Schleswig-Holstein, Campus Lübeck, 23552 Lübeck, Germany; ^{vi}Ageing Epidemiology Research Unit, School of Public Health, Imperial College, London SW71, UK; ^{vii}Regenerative Immunology and Aging, BIH Center for Regenerative Therapies, Charité Universitätsmedizin Berlin, Berlin, Germany; ^{viii}German Institute of Human Nutrition, Department of Nutrition and Gerontology, Potsdam-Rehbruecke (DIfE), Germany; ^{ix}Charité - Universitätsmedizin Berlin, Forschungsgruppe Geriatrie am EGZB, Berlin, Berlin, Germany; ^xDepartment of Immunology, University of Tübingen, Tübingen, Germany; ^{xi}Health Sciences North Research Institute, Sudbury, ON, Canada; ^{xii}German Socio-Economic Panel Study (SOEP); ^{xiii}Division of Dermatocology Department of Dermatology, University of Tübingen, Tübingen, Germany.

References

- 1 Bertram, L. *et al.* Cohort profile: The Berlin Aging Study II (BASE-II). *Int J Epidemiol* **43**, 703-712, doi:10.1093/ije/dyt018 (2014).
- 2 Gerstorf, D. *et al.* Editorial. *Gerontology* **62**, 311-315, doi:10.1159/000441495 (2016).
- 3 Hensel, C. *et al.* Influence of nutritional tyrosine on cognition and functional connectivity in healthy old humans. *Neuroimage* **193**, 139-145, doi:10.1016/j.neuroimage.2019.03.005 (2019).
- 4 Bender, A. R. *et al.* Hippocampal Subfields and Limbic White Matter Jointly Predict Learning Rate in Older Adults. *Cereb Cortex* **30**, 2465-2477, doi:10.1093/cercor/bhz252 (2020).
- 5 Dahl, M. J. *et al.* Rostral locus coeruleus integrity is associated with better memory performance in older adults. *Nat Hum Behav* **3**, 1203-1214, doi:10.1038/s41562-019-0715-2 (2019).
- 6 Buchmann, N. *et al.* Association between lipoprotein(a) level and type 2 diabetes: no evidence for a causal role of lipoprotein(a) and insulin. *Acta Diabetol* **54**, 1031-1038, doi:10.1007/s00592-017-1036-4 (2017).
- 7 Duzel, S. *et al.* Validation of a single factor representing the indicators of metabolic syndrome as a continuous measure of metabolic load and its association with health and cognitive function. *PLoS One* **13**, e0208231, doi:10.1371/journal.pone.0208231 (2018).
- 8 Rohr, F. *et al.* Relationship between Lipoprotein (a) and cognitive function - Results from the Berlin Aging Study II. *Sci Rep* **10**, 10636, doi:10.1038/s41598-020-66783-3 (2020).
- 9 Spira, D. *et al.* Sex-specific differences in the association of vitamin D with low lean mass and frailty: Results from the Berlin Aging Study II. *Nutrition* **62**, 1-6, doi:10.1016/j.nut.2018.11.020 (2019).
- 10 Spira, D. *et al.* Association of Low Lean Mass With Frailty and Physical Performance: A Comparison Between Two Operational Definitions of Sarcopenia-Data From the Berlin Aging Study II (BASE-II). *J Gerontol A Biol Sci Med Sci* **70**, 779-784, doi:10.1093/gerona/glu246 (2015).
- 11 Gerstorf, D. *et al.* Secular changes in late-life cognition and well-being: Towards a long bright future with a short brisk ending? *Psychol Aging* **30**, 301-310, doi:10.1037/pag0000016 (2015).
- 12 Hurler, G. *et al.* Cohort Differences in Psychosocial Function over 20 Years: Current Older Adults Feel Less Lonely and Less Dependent on External Circumstances. *Gerontology* **62**, 354-361, doi:10.1159/000438991 (2016).
- 13 Lu, Y. *et al.* New loci for body fat percentage reveal link between adiposity and cardiometabolic disease risk. *Nat Commun* **7**, 10495, doi:10.1038/ncomms10495 (2016).
- 14 Nikpay, M. *et al.* A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet* **47**, 1121-1130, doi:10.1038/ng.3396 (2015).
- 15 Schmidt, A. F. *et al.* PCSK9 genetic variants and risk of type 2 diabetes: a mendelian randomisation study. *Lancet Diabetes Endocrinol* **5**, 97-105, doi:10.1016/S2213-8587(16)30396-5 (2017).
- 16 Eibich, P., Krekel, C., Demuth, I. & Wagner, G. G. Associations between Neighborhood Characteristics, Well-Being and Health Vary over the Life Course. *Gerontology* **62**, 362-370, doi:10.1159/000438700 (2016).
- 17 Meyer, A. *et al.* Leukocyte telomere length is related to appendicular lean mass: cross-sectional data from the Berlin Aging Study II (BASE-II). *Am J Clin Nutr* **103**, 178-183, doi:10.3945/ajcn.115.116806 (2016).
- 18 Vetter, V. M., Spira, D., Banszerus, V. L. & Demuth, I. Epigenetic Clock and Leukocyte Telomere Length are Associated with Vitamin D Status, but not with Functional Assessments and Frailty in the Berlin Aging Study II. *J Gerontol A Biol Sci Med Sci*, doi:10.1093/gerona/glaa101 (2020).

- 19 Wistuba-Hamprecht, K., Haehnel, K., Janssen, N., Demuth, I. & Pawelec, G. Peripheral blood T-cell signatures from high-resolution immune phenotyping of gammadelta and alphabeta T-cells in younger and older subjects in the Berlin Aging Study II. *Immun Ageing* **12**, 25, doi:10.1186/s12979-015-0052-x (2015).
- 20 Demuth, I. *et al.* Berlin Aging Study II (BASE-II). *Encyclopedia of Gerontology and Population Aging*, 1-8, doi:10.1007/978-3-319-69892-2_27-1 (2019).
- 21 Pelletier, R., Ditto, B. & Pilote, L. A composite measure of gender and its association with risk factors in patients with premature acute coronary syndrome. *Psychosom Med* **77**, 517-526, doi:10.1097/PSY.000000000000186 (2015).
- 22 Nauman, A. T. *et al.* Gender score development in a retrospective approach in the Berlin Aging Study II. *medrxiv*, doi:<https://doi.org/10.1101/2019.12.16.19015032> (2019).
- 23 Moran, A. E. *et al.* Temporal trends in ischemic heart disease mortality in 21 world regions, 1980 to 2010: the Global Burden of Disease 2010 study. *Circulation* **129**, 1483-1492, doi:10.1161/CIRCULATIONAHA.113.004042 (2014).
- 24 Mauvais-Jarvis, F. *et al.* Sex and gender: modifiers of health, disease, and medicine. *Lancet* **396**, 565-582, doi:10.1016/S0140-6736(20)31561-0 (2020).
- 25 Lacasse, A. *et al.* Conducting gender-based analysis of existing databases when self-reported gender data are unavailable: the GENDER Index in a working population. *Can J Public Health* **111**, 155-168, doi:10.17269/s41997-019-00277-2 (2020).
- 26 Gerstorff, D. *et al.* Cohort differences in adult-life trajectories of internal and external control beliefs: A tale of more and better maintained internal control and fewer external constraints. *Psychol Aging* **34**, 1090-1108, doi:10.1037/pag0000389 (2019).
- 27 Potter, S. *et al.* Trajectories of multiple subjective well-being facets across old age: The role of health and personality. *Psychol Aging*, doi:10.1037/pag0000459 (2020).
- 28 Graham, E. K. *et al.* Trajectories of Big Five Personality Traits: A Coordinated Analysis of 16 Longitudinal Samples. *European Journal of Personality* **34**, 301-321, doi:10.1002/per.2259 (2020).
- 29 Mantantzis, K. *et al.* Dehydration predicts longitudinal decline in cognitive functioning and well-being among older adults. *Psychol Aging* **35**, 517-528, doi:10.1037/pag0000471 (2020).
- 30 Mueller, S. *et al.* Personality development in old age relates to physical health and cognitive performance: Evidence from the Berlin Aging Study II. *Journal of Research in Personality* **65**, 94-108, doi:<https://doi.org/10.1016/j.jrp.2016.08.007> (2016).
- 31 Goebel, J. *et al.* The German Socio-Economic Panel (SOEP). *Jahrbücher für Nationalökonomie und Statistik* **239**, 345, doi:<https://doi.org/10.1515/jbnst-2018-0022> (2019).
- 32 Charlson, M. E., Pompei, P., Ales, K. L. & MacKenzie, C. R. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* **40**, 373-383, doi:10.1016/0021-9681(87)90171-8 (1987).
- 33 Cosentino, F. *et al.* 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J* **41**, 255-323, doi:10.1093/eurheartj/ehz486 (2020).
- 34 Asch, F. M. *et al.* Similarities and Differences in Left Ventricular Size and Function among Races and Nationalities: Results of the World Alliance Societies of Echocardiography Normal Values Study. *Journal of the American Society of Echocardiography* **32**, 1396-1406.e1392, doi:<https://doi.org/10.1016/j.echo.2019.08.012> (2019).
- 35 Kohler, S. & Thiel, A. Life after the thymus: CD31+ and CD31- human naive CD4+ T-cell subsets. *Blood* **113**, 769-774, doi:10.1182/blood-2008-02-139154 (2009).
- 36 Weyand, C. M., Brandes, J. C., Schmidt, D., Fulbright, J. W. & Goronzy, J. J. Functional properties of CD4+ CD28- T cells in the aging immune system. *Mech Ageing Dev* **102**, 131-147, doi:10.1016/s0047-6374(97)00161-9 (1998).
- 37 Walhovd, K. B. *et al.* Healthy minds from 0-100 years: Optimising the use of European brain imaging cohorts ("Lifebrain"). *Eur Psychiatry* **47**, 76-87, doi:10.1016/j.eurpsy.2017.10.005 (2018).

- 38 Alberti, K. G. *et al.* Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* **120**, 1640-1645, doi:10.1161/CIRCULATIONAHA.109.192644 (2009).
- 39 Kronenberg, F. Prediction of cardiovascular risk by Lp(a) concentrations or genetic variants within the LPA gene region. *Clin Res Cardiol Suppl* **14**, 5-12, doi:10.1007/s11789-019-00093-5 (2019).
- 40 Kamstrup, P. R. & Nordestgaard, B. G. Lipoprotein(a) concentrations, isoform size, and risk of type 2 diabetes: a Mendelian randomisation study. *Lancet Diabetes Endocrinol* **1**, 220-227, doi:10.1016/S2213-8587(13)70064-0 (2013).
- 41 Ye, Z. *et al.* The association between circulating lipoprotein(a) and type 2 diabetes: is it causal? *Diabetes* **63**, 332-342, doi:10.2337/db13-1144 (2014).
- 42 Seeland, U., Demuth, I., Regitz-Zagrosek, V., Steinhagen-Thiessen, E. & König, M. Sex differences in arterial wave reflection and the role of exogenous and endogenous sex hormones: results of the Berlin Aging Study II. *J Hypertens* **38**, 1040-1046, doi:10.1097/HJH.0000000000002386 (2020).
- 43 Mantantzis, K. *et al.* Poor glucose regulation is associated with declines in well-being among older men, but not women. *Psychol Aging* **35**, 204-211, doi:10.1037/pag0000404 (2020).
- 44 Di Benedetto, S. *et al.* Impact of age, sex and CMV-infection on peripheral T cell phenotypes: results from the Berlin BASE-II Study. *Biogerontology* **16**, 631-643, doi:10.1007/s10522-015-9563-2 (2015).

Figure 1:

Flow chart explaining the final BASE-II sample with follow-up assessments completed in GendAge

