1	TITLE PAGE							
2								
3	<u>Title</u> :							
4	Beneficial and harmful outcomes of tocilizumab in severe COVID-19:							
5	a systematic review and meta-analysis							
6								
7	Running title:							
8	Tocilizumab in COVID-19: a meta-analysis							
9								
10	Author list:							
11	Manuel Rubio-Rivas ^{a,*} , Jose María Mora-Luján ^a , Abelardo Montero ^a , Narcís A. Homs ^a ,							
12	Jordi Rello ^{b,c,d} , Xavier Corbella ^{a,d}							
13								
14	Affiliations:							
15								
16	^a Department of Internal Medicine, Bellvitge University Hospital, Bellvitge Biomedical							
17	Research Institute-IDIBELL, University of Barcelona, Barcelona, Spain.							
18	^b Centro de Investigación Biomédica en Red (CIBERES), Instituto de Salud Carlos III,							
19	Madrid, Spain							
20	CRIPS, Vall d'Hebrón Institute of Research, Barcelona, Spain.							
21	^d School of Medicine, Universitat Internacional de Catalunya, Barcelona, Spain.							
22								
23								
24	*Corresponding author: Dr Manuel Rubio-Rivas, Internal Medicine Department,							
25	Bellvitge University Hospital, 08907 L'Hospitalet de Llobregat, Barcelona, Spain.							
26								
27	<i>E-mail addresses</i> : <u>mrubio@bellvitgehospital.cat</u> (M. Rubio-Rivas),							
28	jmora@bellvitgehospital.cat (J.M. Mora-Luján), <u>asaez@bellvitgehospital.cat</u> (A.							
29	Montero), <u>nhoms@bellvitgehospital.cat</u> (N.A. Homs), <u>irello@crips.es</u> (J. Rello),							
30	xcorbella@bellvitgehospital.cat (X. Corbella)							
31								

J. Rello and X. Corbella contributed equally as last authors 32

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

33 ABSTRACT

34

Objectives: Pending for randomized control trials, the use of tocilizumab (TCZ) in COVID-35 19 remains controversial. We performed a systematic review and meta-analysis to 36 investigate the effect on clinical outcomes of TCZ to treat severe COVID-19. 37

38

Methods: From 1 January to 21 August 2020, we searched PubMed (via MEDLINE), 39 Scopus, and medRxiv repository databases for observational studies in any language 40 41 reporting efficacy and safety of TCZ use in hospitalized adults with COVID-19. Independent and dually data extraction and quality assessment were performed. 42

43

Results: Of 57 eligible studies, 27 controlled and 30 not. The overall included patients 44 were 8,128: 4,021 treated with TCZ, in addition to standard of care (SOC), and 4,107 only 45 receiving SOC. The pooled mortality was lower in the TCZ-group, with a relative risk (RR) 46 47 of 0.73 (95%CI 0.57-0.93; p=0.010). TCZ-treated patients were transferred to the intensive care unit (ICU) in a higher proportion, but ICU mortality was lower than in the 48 control group. Conversely, a higher proportion of TCZ-treated patients developed 49 secondary infections after TCZ use. 50

51

52 **Conclusions:** TCZ seems beneficial in preventing in-hospital mortality in severe, non-53 critically ill COVID-19 patients. However, patients receiving TCZ appear to be at higher 54 risk for secondary infections, especially those admitted to ICU.

- 55
- 56
- 57

Keywords: Coronavirus, SARS-CoV-2, COVID-19, Tocilizumab, Meta-analysis 58

59

60 Abstract Word count: 200

Text Word count: 3469 61

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

62 **INTRODUCTION**

63

Since early 2020, when the SARS-CoV-2 pandemic hit the world, a variety of treatments 64 have been suggested for COVID-19 [1-5]. However, to date, only remdesivir [4] and 65 dexamethasone [5] have demonstrated evidence-based efficacy on randomized, 66 67 controlled clinical trials (RCTs). This double strategy combining antiviral and immunomodulatory therapy is in accordance with the two pathological mechanisms 68 that appear to coexist in the disease; the first triggered by the virus itself and the second 69 70 by the cytokine storm and systemic dysregulated host-immune hyperinflammatory 71 response [6].

72

While the pandemic continues to spread globally, a worrying 15% of patients continue to transit into the most severe stage of the disease, requiring hospitalization or intensive care unit (ICU) admission. This advanced clinical-stage presents as severe pulmonary injury and multi-organ failure, causing fatality in nearly half of cases, resembling complications from CAR T cell therapy [7].

78

79 Among other pro-inflammatory cytokines, IL-6 plays a part in innate immunity, but excessive production by the host facing SARS-CoV-2 is detrimental [8-10]. Accordingly, 80 81 the use of immunomodulatory agents such as tocilizumab (TCZ), a monoclonal antibody 82 to the recombinant human IL-6 receptor, was initially reported as successful among 21 83 patients in China [11]. Since then, an emerging number of observational studies from 84 America and Europe have been published or registered assessing the effect of TCZ in 85 severe COVID-19 [12-67]. In most of them, the authors report an association between earlier use of TCZ and reduced mortality; however, interpretation of these results is 86 87 limited because several of them did not describe a comparison group.

88

Conversely, given preliminary results from the industry-sponsored Phase 3 COVACTA trial (ClinicalTrials.gov Identifier NCT04320615), the COVID-19 Treatment Guidelines Panel by the National Institute of Health, has taken a position against the use of TCZ [68]. This RCT is the first global, randomized, double-blind, placebo-controlled phase III trial investigating TCZ in hospitalized patients with severe COVID-19, but it failed to demonstrate improvement in clinical status as the primary endpoint, or several key secondary outcomes such as 4-week mortality [69].

96

In the current emergency, while waiting for additional data from RCTs, the fact that most
institutions and physicians worldwide are still tackling COVID-19 based only on realworld reported data, prompted us the present review. Our aim was to summarize the
updated results from available observational studies on the effect of TCZ on clinical
outcomes in hospitalized patients with COVID-19.

102 METHODS

103

104 This systematic review and meta-analysis followed the guidance of the Preferred 105 Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) statement [70]. The protocol was published in the National Institute for Health Research international 106 107 register of systematic reviews (PROSPERO); registration number CRD42020204934. A clinical question under the PICO framework format (Population-Intervention-108 Comparison-Outcome) was created (Table 1). 109

110

111 Data Sources and Searches

112

The search strategy was developed by three investigators (M.R-R., J.R., X.C.), which was 113 114 revised and approved by the other investigators (J.M.M-L., A.M, N.A.H.). We searched 115 the following databases from 1 January to 21 August 2020: MEDLINE database through 116 the PubMed search engine, Scopus, and the medRxiv repository, using the terms "COVID-19" [MesH]) AND "Tocilizumab" [MesH]. 117

Study Eligibility Criteria 119

120

118

121 Full-text observational studies in any language reporting beneficial or harmful outcomes 122 from the use of TCZ in adults hospitalized with COVID-19 were included. Two 123 investigators (M.R-R., J.M.M-L.) independently screened each record title and abstract 124 for potential inclusion. Restriction of publication type was manually applied: secondary analyses of previously reported trials, protocols, abstracts-only and experimental 125 studies were excluded. Potentially relevant articles were retrieved for full-text review. 126 127 Two investigators (M.R-R., A.M.) read the full text of the abstracts selected. 128 Discrepancies were resolved through discussion or by a third investigator (X.C.).

129

130 Publications were included if they met all the following criteria: 1) the study reported data on adults ≥18 years-old with COVID-19, diagnosed by polymerase chain reaction 131 (PCR), admitted hospital-wide or in ICUs; 2) the study design was an observational 132 133 investigation providing real-world original data on TCZ use in COVID-19, either 134 intravenous or subcutaneous; and 3) the study data collection finished after 1 January 135 2020.

136

Those studies reported being "case-control studies", in which subjects from the control 137 138 group also presented COVID-19, just as those from the TCZ group, were also included in the present review. Studies focusing on a sole subgroup of patients (e.g. renal transplant 139 recipients) were excluded. A careful revision was also performed of patients' origins 140 141 included in studies from the same country/hospital to avoid overlapping data, and only 142 the latest and largest study was selected. The search was completed by the bibliography 143 review of every paper selected for full-text examination.

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

144 Data Extraction and Quality Assessment

145

146 Two investigators (M.R-R., J.M.M-L.) independently abstracted the following details: study characteristics, including setting; intervention or exposure characteristics, 147 148 including medication dose and duration; patient characteristics, including the severity 149 of disease; and outcomes, including mortality, admission to ICU, adverse events such as 150 secondary infections, and length of hospital stay. Discrepancies were resolved by discussion in consultation with a third investigator (X.C.). Quality assessment was 151 152 performed by two investigators (M.R-R., N.A.H) using the Newcastle-Ottawa Scale (NOS) for observational studies [71]. In case of disagreement, a third author (J.R.) 153 154 independently determined the quality assessments.

155

156 Data Analysis

157

Categorical variables were described as absolute numbers and percentages. When the number of events and the sample size was small (and followed a Poisson distribution), confidence intervals were estimated using Wilson's method [72-74]. We carried out a meta-analysis of the pooled mortality ratio by including all comprised studies. Those studies with a control group were also meta-analyzed to assess the relative risk (RR) of mortality in TCZ-treated patients vs. those non-TCZ treated (RR of 1).

164

165 The inverse variance-weighted method was initially performed using a fixed-effects model. Heterogeneity between studies was assessed using the Q statistic. The 166 percentage of variability between studies by the Higgins I² parameter and between-167 study variability was measured by the Tau² parameter and, when confirmed ($p \le 0.05$), 168 the analysis was completed by using the random-effects model [75]. Studies with 0 169 events in the only arm (uncontrolled studies) or both arms (controlled studies) were not 170 included in the meta-analysis. Forest plots were depicted accordingly. Publication bias 171 172 was assessed using the Egger method [76]. Statistical analysis was performed by IBM 173 SPSS Statistics for Windows, Version 26.0. Armonk, NY: IBM Corp.

- 174
- 175

176 **RESULTS**

177

A total of 781 articles were identified in our search. Of these, 81 qualified for full-text
review following title and abstract screening, of which 57 [11-67] were included in the
analysis. The PRISMA flow diagram is detailed in Figure 1.

181

182 Study characteristics

183

The majority of included studies were carried out in different hospitals in high-income
countries in America and Europe, such as the US [13, 18, 22, 28, 31, 34, 35, 48, 49, 5360, 66], Italy [14, 15, 17, 19, 20, 41, 44-47, 50, 51, 63, 65], Spain [21, 26, 27, 32, 33, 39,
40, 42, 43], and France [25, 29, 30, 61, 62, 64]. A lesser number were conducted in China
[11, 12]. The distribution of the studies worldwide is shown in Figure 2.

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

189 Search results

190

191 Of the total of 57 studies included, 12 were prospective and 45, retrospective. In 30 of 192 these cohort investigations, a control group was not described [10-38] and 27 added a 193 comparison group [40-67]. The overall results provided data from 8,128 hospitalized 194 patients with COVID-19: 4,021 TCZ-treated, in addition to standard of care (SOC) (including 711 patients admitted to ICU), and 4,107 only receiving SOC (including 694 195 patients admitted to ICU). SOC was basically antiviral therapy (remdesivir, 196 197 lopinavir/ritonavir), antimalarials (hydroxychloroquine), or azythromicin. Characteristics of patients in the TCZ-group vs. the control group are detailed in Tables 2-7. Of the TCZ 198 199 group, 2,645 (65.8%) were men with a mean age of 61.8 (SD 6.1) and median age 62.6 200 [range 59-65], according to the data provided. TCZ was given as a single dose in 201 2,030/2,952 patients (68.8%), and in 922 (31.2%), as two or more doses.

202

Concomitantly to TCZ use, additional treatment with steroids was given in 1,560/3,073
patients in the TCZ-group (50.8%) vs. 592/2,733 (21.7%) in the control group (p<0.001).
Comparing both groups, remdesivir was used in 37/3,511 (1.05%) vs. 23/3,945 (0.58%)
(p=0.023) patients. Finally, the administration of TCZ has prescribed a median of 10 days
[range 9-11] after the onset of COVID-19 symptoms, in those studies in which this data
was provided. The median follow-up of the overall cohort was 10.3 days [range 12-19].

209

210 Mortality

211

After applying the random-effects model, hospital-wide (including ICUs) pooled 212 mortality of patients with COVID-19 treated with TCZ was 19.2% (95%CI 16.4-22.5) 213 214 (I²=83.6% Q=305.3 tau²=0.23 p<0.001). Egger's method A=-3.354 p<0.001 (Figure 2). In the control group, overall mortality was 27.4% (95%CI 21.1-35.6%) (I²=95.9% Q=569.1 215 $tau^2=0.38 p<0.001$). These differences between the TCZ-group and the control group 216 217 achieved statistical significance (p<0.001). The RR of mortality in the TCZ-group was 0.73 (95%CI 0.57-0.93; p=0.010) (I²=77.7% Q=107.4 tau²=0.24 p<0.001). Egger's method A=-218 0.712 p=0.380 (Figure 3). The number needed to treat (NNT) to avoid one death was 20. 219 220

221 When the analysis was restricted to high-quality controlled observational studies, 222 excluding those non-controlled, those with <20 patients, or those showing NOS <7, the 223 pooled mortality in the TCZ-group was 17.7% (95%Cl 13.9-22.6) (I^2 =85% Q=168 224 tau²=0.295 p<0.001). Egger's method A=-3.727 p<0.001. The RR of death in the TCZ-225 group vs. the control group was 0.58 (95%Cl 0.40-0.85) (I^2 =79.7% Q=108.3 tau²=0.60 p=0.004). Egger's method A=-1.511 p=0.153.

227

228 **Overall Mortality in hospital wards.** The pooled mortality of COVID-19 patients 229 receiving TCZ in conventional wards was 17% (95%CI 13.9-20.8) (I^2 =58.5% Q=50.6 230 tau2=0.108 p<0.001). Egger's method A=-2.082 p<0.001. In contrast, mortality in the 231 control group after being admitted for COVID-19 in conventional wards was 23.8% 232 (95%CI 14.8-38.4) (I^2 =90.3% Q=51.8 tau²=0.314 p<0.001). Egger's method A=5.433 p=0.044. These differences did not achieve statistical significance (p=0.261). The RR of

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

234 mortality in hospital wards in the TCZ-group vs. the control group was 0.64 (95%CI 0.26-1.54; p=0.314) (I²=80.8% Q=26 tau²=0.91 p=0.314). Egger's method A=-3.576 p=0.005. 235 When analysis was restricted to high-quality controlled studies, the pooled mortality in 236 hospital wards in the TCZ-group was 13.3% (95%Cl 8.7-20.4) (I²=76.2% Q=25 tau²=0.235 237 p<0.001). Egger's method A=-2.653 p=0.074. 238 239 Overall Mortality in ICUs. The pooled mortality of COVID-19 patients receiving TCZ once 240 241 admitted to ICU was 31.7% (95%CI 24.7-40.8) (I²=82.6% Q=97.6 tau²=0.185 p<0.001). 242 Egger's method A=-2.995 p<0.001. When compared, mortality of ICU control group patients was 39.1% (95%Cl 29.1-52-4) (l²=91.6% Q=59.4 tau²=0.101 p<0.001). Egger's 243 244 method A=-1.766 p=0.290. These differences achieved statistical significance (p<0.001). 245 The RR for mortality in those ICU patients receiving TCZ vs. SOC was 0.78 (95%CI 0.44-1.35; p=0.369) (I²=65.4% Q=14.4 tau²=0.29 p=0.639). Egger's method A=0.693 p=0.727. 246 The NNT in ICU to avoid one death was 9. 247 248 When the analysis was restricted to high-quality controlled studies, the pooled mortality 249 in the ICU in the TCZ-group was 29.3% (95%CI 19.4-44.2) (I²=89.1% Q=55 tau²=0.235 250 p<0.001). Egger's method A=-4.564 p=0.009. 251 252 253 Mortality in the TCZ-group receiving additional corticosteroids. Concomitantly to TCZ 254 use, additional treatment with steroids was given in 1,560/3,073 patients in the TCZ-255 group (50.8%) vs. 592/2,733 (21.7%) in the control group (p<0.001). Since outcomes 256 were not reported separately in most of the included studies, differences in mortality 257 could only be compared between those studies in which steroids were not prescribed in 258 addition of TCZ use and those in which 100% of TCZ-treated patients received 259 concomitant therapy with steroids. 260 The pooled mortality of TCZ patients without corticosteroids was 10.8% (95%CI 6.3-18.5) 261 262 (I²=91% Q=136.3 tau²=0.820 p<0.001). Egger's method A=-4.438 p=0.006. Alternatively, 263 the pooled mortality of TCZ patients with corticosteroids was 16.3% (95%CI 12.8-20.8) 264 (I²=35.9% Q=6.2 tau²=0.046 p<0.001). Egger's method A=-0.342 p=0.895. 265 266 Mortality in the TCZ-group receiving early vs. late TCZ administration. Hospital-wide

Mortality in the TCZ-group receiving early vs. late TCZ administration. Hospital-wide (including ICUs) pooled mortality in TCZ-treated patients in whom TCZ was early administered (<10 days from symptoms onset) was 15.9% (95%Cl 10.9-23) (l^2 =74.9% Q=31.9 tau²=0.222 p<0.001). Egger's method A=-2.306 p=0.239. Hospital-wide (including ICUs) pooled mortality in patients with COVID-19 illness in whom TCZ was administered later (\geq 10 days) was 23.3% (95%Cl 17.9-30.3) (l^2 =75.5% Q=36.8 tau²=0.115 p<0.001). Egger's method A=-2.983 p=0.021. Differences between groups did not achieve statistical significance (p=0.252).

274

275 When the analysis was restricted to high-quality controlled studies, the pooled mortality 276 after early TCZ was 11% (95%Cl 8.4-14.3) (I^2 =42.4% Q=5.2 tau²=0.057 p<0.001). Egger's 277 method A=-2.824 p=0.372. Alternatively, the pooled mortality after late TCZ was 22.6% 278 (95%Cl 16.4-31.1) (I^2 =3.4% Q=2.1 tau²=0.004 p<0.001). Egger's method A=-2.756 279 p=0.034.

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

280 **Risk of ICU admission**

281

In patients with COVID-19 initially admitted to hospital wards, the pooled ICU admission 282 rate after TCZ administration was 17.1% (95%CI 11.5-25.5) (I²=90% Q=149.5 tau²=0.51 283 p<0.001). Egger's method A=-3.272 p=0.015. Conversely, the risk for ICU admission in 284 285 the control group, initially admitted outside the ICUs, was 9.5% (95%Cl 2.9-31.2) (l²=96.2% Q=78.8 tau²=1.37 p<0.001). Egger's method A=-1.315 p=0.886. These 286 differences achieved statistical significance (p<0.001). Accordingly, in the subset of 287 288 patients initially admitted outside the ICUs, those receiving TCZ showed a RR of ICU admission of 1.49 (95%Cl 0.30-7.34; p=0.621) (I²=93% Q=43 tau²=2.35 p=0.62). Egger's 289 290 method A=-4.026 p=0.435.

291

292 When the analysis was restricted to high-quality controlled studies, the pooled risk of 293 ICU admission after receiving TCZ at the ward was 15.6% (95%CI 11.8-20.9) (I^2 =22.4% 294 Q=3 tau²=0.022 p<0.001). Egger's method A=-1.839 p=0.113.

295 296 **Safety**

290 **3a**

The pooled rate of reported secondary viral, bacterial or opportunistic fungal infections in those patients with COVID-19 treated with TCZ was 18.9% (95%CI 14.5-24.8) (I^2 =88.1% Q=218.8 tau²=0.391 p<0.001). Egger's method A=-3.852 p=0.001. The RR of secondary infections in TCZ-treated vs. the control group was 1.47 (95%CI 0.99-2.19; p=0.058) (I^2 =66.1% Q=38.3 tau²=0.34 p=0.058). Egger's method A=0.039 p=0.977

303

When the analysis was restricted to high-quality controlled studies, the pooled percentage of secondary infections was 20.7% (95%Cl 14.6-29.2) (I^2 =88.2% Q=118.5 tau²=0.351 p<0.001). Egger's method A=-4.128 p=0.003.

308 Length of hospital stay

309

307

Among survivors, the length of hospital stay in the TCZ-group was a median of 15.3 days [range 12.4-19.4] vs. 14 days [range 9-20] in the control group. These differences were not statistically significant (p=0.953).

- 313
- 314

315 **DISCUSSION**

316

Pending published evidence from RCTs, this SRMA focused on available real-world
 observational studies, revealing a beneficial effect of TCZ use in preventing mortality in
 hospitalized adults with COVID-19. However, the present results also showed a higher
 relative risk for ICU admission and the occurrence of secondary infections in such COVID 19 patients receiving TCZ.

322

To date, two existing SRMA have summarized current evidence on the beneficial and harmful effects of TCZ in COVID-19. The first by Lan SH Zhang et al., included 7 studies, with no conclusive evidence that TCZ would provide any additional benefit to patients

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

with severe COVID-19 [77]. The second, registered in the medRxiv repository by
 Boregowda et al., included 16 studies, concluding that the addition of TCZ to SOC might
 reduce mortality in COVID-19 patients requiring hospitalization [78].

329

330 The present SRMA updated and expanded the revision to 57 observational studies. As 331 expected, most included studies emerged recently and were performed in hospitals from high-income European countries and the US. The finding that most of the included 332 patients were men in their 6-8th decades of life was consistent with what has already 333 334 been described in the general population requiring hospital admission due to COVID-19 [79]. The vast majority of included studies used a single dose of 400-800 mg intravenous 335 336 or 162-324 mg subcutaneous, and a few of them allowed a second or even a third dose 337 in case of worsening. Interestingly, the present SRMA showed that TCZ was mainly 338 prescribed as a second step to treat those patients at risk of transition to a more severe 339 condition, after showing poor response to antiviral agents. Accordingly, in a notable 340 proportion of included studies, TCZ was indicated in combination with corticosteroids. In this respect, in view of the RECOVERY trial [5] which reported significant benefit from 341 the use of steroids in severe COVID-19, it is difficult to distinguish in depth the 342 contribution of TCZ on the outcomes in such included patients receiving TCZ and 343 steroids. On the other hand, in addition to TCZ, a very small proportion (<1%) of included 344 345 patients from more recent studies also received remdesivir, as a proven antiviral against 346 SARS-Cov-2.

347

In addition to concomitant drugs, other relevant factors should be taken into account when considering the impact of TCZ use on clinical outcomes in COVID-19. In this respect, in addition to the age and underlying comorbidities of included patients, one of the most important factors to be considered at the clinical level is the severity of the clinical-stage when indicating TCZ to treat the COVID-19 illness [6,79].

353

354 Unfortunately, the severity of patients could not be accurately inferred from the clinical, 355 laboratory, or radiological parameters documented in the included studies of the 356 present SRMA. Therefore, mortality was compared between those patients admitted to 357 hospital wards and ICUs, in those studies in which the hospital site from where TCZ was 358 administered was specified. Logically, patients in whom TCZ was indicated during ICU 359 admission showed higher mortality in comparison to those treated in hospital wards. 360 However, since the COVID-19 pandemic induced an unprecedented influx of patients into the ICUs, the particular emergency and resource availability of each hospital 361 362 involved in the included studies most likely conditioned either the criteria when 363 transferring patients from hospital wards to ICUs, or the ethical decisions related to the 364 withdrawal of life support decisions.

365

366 Outside of RCTs, observational data from the included studies only reflected the clinical 367 practice of physicians when indicating TCZ use. Consequently, the present results show 368 that TCZ was mostly prescribed in those more seriously-ill patients presenting at a more 369 advanced stage of COVID-19, with severe lung injury and systemic hyperinflammatory 370 multi-organ failure. In this regard, it would be unfair to infer that the higher the use of 371 TCZ and the average of concomitant corticosteroids, the higher the risk of ICU admission

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

or death. Therefore, further RCTs on TCZ use in COVID-19 should clarify the interaction
of confounding variables, it being crucial to know which patients are the best candidates
to eventually receive TCZ as an immunomodulatory agent, as well as the beneficial and
harmful effects of its use in the absence of or in combination with steroids.

376

377 Moreover, in the particular case of the ICU setting, most of the included studies showed insufficient data to appropriately assess the effects when TCZ was prescribed in 378 critically-ill patients. However, since a majority of such seriously-ill patients with COVID-379 380 19 admitted to ICUs are submitted to mechanical ventilation and other multiple invasive 381 procedures, and receive concomitant wide-spectrum antibiotics or steroid treatment, it 382 is of great concern the well-known risk of TCZ favoring the occurrence of life-threatening 383 secondary bacterial, viral or fungal opportunistic infections, as it has also been 384 documented in the present SRMA in up to one-fifth of cases in the TCZ-group [53,80,81]. 385

386 As mentioned, the first and major limitation of this SRMA is the lack of data from RCTs. 387 In their absence, the present revision was based on observational studies; therefore, conclusions should be considered with caution. Moreover, a second limitation is the fact 388 389 that most of the included studies were retrospective in nature. A third limitation is the heterogeneity regarding the study population (I² index) and the potential risk of 390 391 detected bias. Fourth, variations in criteria for prescribing TCZ may not be ruled out in 392 the included studies, although most of them indicated TCZ use to treat those patients 393 with severe COVID-19 with the systemic hyperinflammatory state. Fifth, important 394 factors influencing the effect of TCZ on clinical outcomes such as the baseline 395 characteristics of the patients included, the average time from symptoms onset to TCZ administration, the clinical severity of the disease at the time of TCZ administration, the 396 397 doses and the form of administration used, the hospital site from where TCZ was 398 indicated, or the use of concomitant drug regimens could not be evaluated in-depth, since they were not uniformly provided by the included studies. Sixth, in the vast 399 400 majority of included studies, there is a lack of subgroup analyses according to age, sex 401 or underlying conditions, concomitant treatments, the requirement of mechanical 402 ventilation or ICU admission, and comparisons between ventilated and non-ventilated 403 patients. Finally, there is a wide range in the median time of follow-up after TCZ 404 administration, which hinders assessment of consistent improvement, late-onset 405 adverse events, and real in-hospital mortality in those patients with prolonged 406 evolution. Thus, some patients considered "survivors" in some included studies may 407 have ended up dying.

408

409 In conclusion, pending evidence from RCTs, this systematic review provides updated and 410 extended data from observational studies on the use of TCZ in COVID-19. The present 411 results showed TCZ to be beneficial in reducing overall in-hospital mortality in adults 412 with COVID-19, with an NNT to save one life of 20. These findings were more apparent 413 in those non-critically-ill COVID-19 patients admitted to hospital wards, receiving TCZ at 414 the early stage of hyperinflammatory response syndrome. By contrast, the TCZ-group 415 was at higher risk for secondary infections, especially in those patients admitted to ICU. 416 Notwithstanding these results, conclusions should be considered as weak evidence since they are based on observational studies, most of them retrospective. However, these 417

findings may help physicians and researchers to optimize strategies towards precision medicine when designing further RCTs focused on the use of TCZ in COVID-19.

ACKNOWLEDGMENTS

We dedicate this work to the memory of those patients worldwide who have not survived COVID-19. We thank CERCA Programme/Generalitat de Catalunya for institutional support.

FUNDING SOURCE

No funding or sponsorship was received for this study or publication of this article.

DECLARATION OF INTERESTS

Jordi Rello has received consultancy honoraria from Roche. The rest of the authors declare no competing interests.

AUTHORS' CONTRIBUTION

- Study conception and design: M.R-R., X.C.
- Acquisition, analyses and interpretation of data: M.R-R., J.M.M-L, A.M., N.A.H., J.R., X.C.
- Manuscript draft and critical revision: M.R-R., J.R., X.C.

REFERENCES

[1] Azoulay E, de Waele J, Ferrer R, Staudinger T, Borkowska M, Povoa P et al. International variation in the management of severe COVID-19 patients. Crit Care 2020;24(1):486. <u>https://doi.org/10.1186/s13054-020-03194-w. Accessed 12 August</u> 2020.

 [2] Gautret P, Cagier JC, Parola P, Hoang VT, Meddeb L, Mailhe M et al.
 Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an openlabel non-randomized clinical trial. Int J Antimicrob Agents 2020:105949.
 <u>https://doi.org/10.1016/j.ijantimicag.2020.105949</u>.
 Accessed 25 March 2020.

[3] Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al. A Trial of Lopinavir-Ritonavir in adults hospitalized with severe Covid-19. N Engl J Med 2020; 382:1787-99.

[4] Grein J, Ohmagari N, Shin D, Diaz G, Asperges E, Castagna A, et al. Compassionate Use of Remdesivir for Patients with Severe Covid-19. N Engl J Med. 2020;382:2327-36.

[5] Horby P, Lim WS, Emberson JR, Mafham M, Bell J, Linsell L, et al. Dexamethasone in hospitalized patients with Covid-19. Preliminary report. N Engl J Med 2020. <u>https://doi.org/10.1056/NEJMoa2021436</u>. Accessed 25 August 2020

[6] Siddiqi HK, Mehra MR. COVID-19 illness in native and immunosuppressed states: A clinical-therapeutic staging proposal. J Heart Lung Transplant 2020;39(5):405-7.

[7] Azoulay E, Darmon M, Valade S. Acute life-threatening toxicity from CAR T-cell therapy. Intensive Care Med 2020. <u>https://doi.org/10.1007/s00134-020-06193-1</u>. Accessed 25 August 2020

[8] Jamilloux Y, Henry T, Belot A, Viel S, Fauter M, El Jammalet T, al. Should we stimulate or suppress immune responses in COVID-19? Cytokine and anti-cytokine interventions. Autoimmun Rev 2020;19:102567. <u>https://doi.org/10.1016/j.autrev.2020.102567</u>. Accessed 25 August 2020

[9] Aziz M, Fatima R, Assaly R. Elevated Interleukin-6 and Severe COVID-19: A Meta-Analysis. J Med Virol. 2020. <u>https://doi.org/10.1002/jmv.25948</u>. Accessed 21 August 2020.

[10] Zhang C, Wu Z, Li JW, Zhao H, Wang GQ. The cytokine release syndrome (CRS) of severe COVID-19 and Interleukin-6 receptor (IL-6R) antagonist Tocilizumab may be the key to reduce the mortality. Int J Antimicrob Agents 2020;55:105954.

[11] Xu X, Han M, Li T, Sun W, Wang D, Fu B, et al. Effective treatment of severe COVID-19 patients with tocilizumab. Proc Natl Acad Sci USA 2020;117:10970-5.
[12] Luo P, Liu Y, Qiu L, Liu X, Liu D, Li J. Tocilizumab treatment in COVID-19: a single center experience. J Med Virol 2020;92:814-8.

[13] Antony SJ, Davis MA, Davis MG, Almaghlouth NK, Guevara R, Omar F, et al. Early use of tocilizumab in the prevention of adult respiratory failure in SARS - CoV - 2 infections and the utilization of interleukin - 6 levels in the management. IDCases. 2020; 21: e00888. https://doi.org/10.1016/j.idcr.2020.e00888. Accessed 25 August 2020

[14] Quartuccio L, Sonaglia A, Pecori D, Peghin M, Fabris M, Tascini C, et al. Higher levels of IL-6 early after tocilizumab distinguish survivors from non-survivors in COVID-19 pneumonia: a possible indication for deeper targeting IL-6. J Med Virol. 2020:10.1002/jmv.26149. https://doi.org/10.1002/jmv.26149. Accessed 25 August 2020

[15] Toniati P, Piva S, Cattalini M, Garrafa E, Regola F, Castelli F, et al. Tocilizumab for the treatment of severe COVID-19 pneumonia with hyperinflammatory syndrome and acute respiratory failure: a single center study of 100 patients in Brescia, Italy. Autoimmun Rev 2020; 19:102568. Accessed 25 August 2020

[16] Alattar R, Ibrahim TBH, Shaar SH, Abdalla S, Shukri K, Daghfalt JN, et al. Tocilizumab for the treatment of severe COVID-19. J Med Virol 2020: 10.1002/jmv.25964. https://doi.org/10.1002/jmv.25964. Accessed 25 August 2020

[17] Sciascia S, Aprà F, Baffa A, Baldovino S, Boaro D, Boero R, et al. Pilot prospective open, single-arm multicentre study on off-label use of tocilizumab in patients with severe COVID-19. Clin Exp Rheumatol 2020;38:529-32.

[18] Morrison AR, Johnson JM, Griebe KM, Jones MC, Stine JJ, Hencken LN, et al. Clinical characteristics and predictors of survival in adults with coronavirus disease 2019 receiving tocilizumab. J Autoimmun 2020:102512. https://doi.org/10.1016/j.jaut.2020.102512. Accessed 25 August 2020

[19] Morena V, Milazzo L, Oreni L, Bestetti G, Fossali T, Bassoli C, et al. Off-label use of tocilizumab for the treatment of SARS-CoV-2 pneumonia in Milan, Italy. Eur J Intern Med 2020;76:36-42.

[20] Marfella R, Paolisso P, Sardu C, Bergamaschib L, D'Angelob EC, Barbieri M, et al. Negative impact of hyperglycaemia on tocilizumab therapy in Covid-19 patients. Diabetes Metab 2020: S1262-3636(20)30082-3. https://doi.org/10.1016/j.diabet.2020.05.005. Accessed 25 August 2020

[21] Sanchez-Montalvà A, Selares-Nadal J, Espinosa-Pereiro J, Fernandez-Hidalgo N, Perez-Hoyos S, Salvador F, et al. Early outcomes of tocilizumab in adults hospitalized with severe COVID19. An initial report from the Vall d'Hebron COVID19 prospective cohort study. medRxiv preprint at

https://www.medrxiv.org/content/10.1101/2020.05.07.20094599v2. Accessed 31 July 2020

[22] Jordan SC, Zakowski P, Tran HP, Smith EA, Gaultier C, Marks G, et al. Compassionate Use of Tocilizumab for Treatment of SARS-CoV-2 Pneumonia. Clin Infect Dis 2020: ciaa812. https://doi.org/10.1093/cid/ciaa812. Accessed 25 August 2020

[23] Nasir N, Syed Faisal M, Kiren H, Khanum I, Jamil B. Treatment of ARDS and hyperinflammation in COVID-19 with IL-6 antagonist Tocilizumab: a tertiary care experience from Pakistan. medRxiv preprint at

https://www.medrxiv.org/content/10.1101/2020.06.23.20134072v1 Accessed 25 August 2020

[24] Borku Uysal B, Ikitimur H, Yavuzer S, Ikitimur B, Uysal H, Islamoglu MS, et al. Tocilizumab challenge: A series of cytokine storm therapy experiences in hospitalized COVID-19 pneumonia patients. J Med Virol 2020 Jun 2:10.1002/jmv.26111. https://doi.org/10.1002/jmv.26111. Accessed 25 August 2020

[25] Issa N, Dumery M, Guisset O, Mourissoux G, Bonnet F, Camou F. Feasibility of Tocilizumab in ICU patients with COVID-19. J Med Virol 2020:10.1002/jmv.26110. https://doi.org/10.1002/jmv.26110. Accessed 25 August 2020

[26] Campins L, Boixeda R, Perez-Cordon L, Aranega R, Lopera C, Force L. Early tocilizumab treatment could improve survival among COVID-19 patients. Clin Exp Rheumatol 2020;38:578. Accessed 25 August 2020

[27] Sanz Herrero F, Puchades Gimeno F, Ortega García P, Ferrer Gómez C, Ocete Mochón MD, García Deltoro M. Methylprednisolone added to tocilizumab reduces mortality in SARS-CoV-2 pneumonia: An observational study. J Intern Med 2020; 10.1111/joim.13145. https://doi.org/10.1111/joim.13145. Accessed 25 August 2020

[28] Knorr JP, Colomy V, Mauriello CM, Ha S. Tocilizumab in patients with severe COVID-19: a single-center observational analysis. J Med Virol. 2020 Jun 17:10.1002/jmv.26191. https://doi.org/10.1002/jmv.26191. Accessed 25 August 2020

[29] Lohse A, Klopfenstein T, Balblanc JC, Royer PY, Bossert M, Gendrin V, et al. Predictive factors of mortality in patients treated with tocilizumab for acute respiratory distress syndrome related to coronavirus disease 2019 (COVID-19). Microbes Infect 2020: S1286-4579(20)30123-4. https://doi.org/10.1016/j.micinf.2020.06.005. Accessed 25 August 2020

[30] Conrozier T, Lohse A, Balblanc JC, Dussert P, Royer P, Bossert M, et al. Biomarker variation in patients successfully treated with tocilizumab for severe coronavirus disease 2019 (COVID-19): results of a multidisciplinary collaboration. Clin Exp Rheumatol 2020; 38:742-7.

[31] Petrak R, Skorodin N, Van Hise N, Fliegelman R, Pinsky J, Didwania V, et al. Tocilizumab as a Therapeutic Agent for Critically III Patients Infected with SARS-CoV-2. medRxiv preprint at

https://www.medrxiv.org/content/10.1101/2020.06.05.20122622v1 Accessed 25 August 2020

[32] Górgolas M, Cabello A, Prieto Perez L, Villar Alvarez F, Alvarez Alvarez B, Rodriguez Nieto MJ, et al. Compassionate Use of Tocilizumab in Severe SARS-CoV2 Pneumonia. When late administration is too late. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.06.13.20130088v1 Accessed 25 August 2020

[33] Jiménez-Brítez G, Ruiz P, Soler X. Tocilizumab plus glucocorticoids in severe and critically COVID-19 patients. A single center experience. Med Clin 2020:S0025-7753(20)30441-3. https://doi.org/10.1016/j.medcli.2020.07.001. Accessed 25 August 2020

[34] Rimland CA, Morgan CE, Bell GJ, Kim MK, Hedrick T, Marx A, et al. Clinical characteristics and early outcomes in patients with COVID-19 treated with tocilizumab at a United States academic center. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.05.13.20100404v2 Accessed 25 August 2020

[35] Sinha P, Mostaghim A, Bielick CG, McLaughlin A, Hamer DH, Wetzler LM, et al. Early administration of Interleukin-6 inhibitors for patients with severe Covid-19 disease is associated with decreased intubation, reduced mortality, and increased discharge. Int J Infect Dis 2020:S1201-9712(20)30568-3. https://doi.org/10.1016/j.ijid.2020.07.023. Accessed 25 August 2020

[36] Patel A, Shah K, Dharsandiya M, Patel K, Patel T, Patel M, et al. Safety and efficacy of tocilizumab in the treatment of severe acute respiratory syndrome coronavirus-2 pneumonia: A retrospective cohort study. Indian J Med Microbiol 2020; 38:117-23.

[37] Hashimoto S, Kitajima H, Arai T, Tamura Y, Nagai T, Morishita H, et al. A retrospective study evaluating efficacy and safety of compassionate use of tocilizumab in 13 patients with severe-to-critically ill COVID-19: analysis of well-responding cases and rapidly-worsening cases after tocilizumab administration. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.06.24.20134288v1 Accessed 25 August 2020

[38] Formina DS, Lysenko MA, Beloglazova IP, Mutinova ZY, Poteshkina NG, Samsonova IV, et al. Temporal clinical and laboratory response to interleukin-6 receptor blockade with Tocilizumab in 89 hospitalized patients with COVID-19 pneumonia. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.06.12.20122374v1 Accessed 25 August 2020

[39] Rubio-Rivas M, Ronda M, Padulles A, Riera-Mestre A, Garcia-Forero C, Iriarte A, et al. Beneficial Effect of Corticosteroids in Preventing Mortality in Patients Receiving Tocilizumab to Treat Severe COVID-19 Illness. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.08.31.20182428v1 Accessed 25 August 2020

[40] Rodríguez-Baño J, Pachón J, Carratalà J, Ryan P, Jarrín I, Yllescas M, et al. Treatment with tocilizumab or corticosteroids for COVID-19 patients with hyperinflammatory state: a multicentre cohort study (SAM-COVID-19). Clin MIcrobiol Infect 2020 Aug 26:S1198-743X(20)30492-4. https://doi.org/10.1016/j.cmi.2020.08.010. Accessed 25 August 2020

[41] Quartuccio L, Sonaglia A, Mcgonagle D, Fabris M, Peghin M, Pecori D, et al. Profiling COVID-19 pneumonia progressing into the cytokine storm syndrome: Results from a single Italian Centre study on tocilizumab versus standard of care. J Clin Virol 2020;129:104444. https://doi.org/10.1016/j.jcv.2020.104444. Accessed 25 August 2020

[42] Martínez-Sanz J, Muriel A, Ron R, Herrera S, Ron R, Perez-Molina JA, et al. Effects of Tocilizumab on Mortality in Hospitalized Patients with COVID-19: A Multicenter Cohort Study. medRxiv preprint at

https://www.medrxiv.org/content/10.1101/2020.06.08.20125245v1 Accessed 31 August 2020

[43] Moreno-García E., Rico V, Albiach L, Aguero D, Ambrosioni J, Bodro M, et al. Tocilizumab is associated with reduced risk of ICU admission and mortality in 1 patients with SARS-CoV-2 infection. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.06.05.20113738v1 Accessed 31 August 2020

[44] Guaraldi G, Meschiari M, Cozzi-lepri A, Milic J, Tonelli R, Menozzi M, et al. Tocilizumab in patients with severe COVID-19 : a retrospective cohort study. The Lancet Rheumatology 2020;9913(20):1-11.

[45] Campochiaro C, Della-Torre E, Cavalli G, De Luca G, Ripa M, Boffini N, et al. Efficacy and safety of tocilizumab in severe COVID-19 patients: a single-center retrospective cohort study. Eur J Intern Med 2020;76:43-9.

[46] Mikulska M, Nicolini LA, Signori A, Di Biagio A, Sepulcri C, Russo C, et al. Tocilizumab and steroid treatment in patients with COVID-19 pneumonia. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.06.22.20133413v1. Accessed 31 August 2020

[47] Colaneri M, Bogliolo L, Valsecchi P, Sacchi P, Zuccaro V, Brandolino F, et al.; COVID IRCCS San Matteo Pavia Task Force. Tocilizumab for treatment of severe COVID-19 patients: preliminary results from SMAtteo COvid19 REgistry (SMACORE). Microorganisms 2020; 8: 695.

[48] Price CC, Altice FL, Shyr Y, Koff A, Pischel L, Goshua G, et al. Tocilizumab Treatment for Cytokine Release Syndrome in Hospitalized COVID-19 Patients: Survival and Clinical Outcomes. Chest 2020:S0012-3692(20)31670-6.

[49] Maeda T, Obata R, Rizk D, Kuno T. The Association of Interleukin-6 value, Interleukin inhibitors and Outcomes of Patients with COVID-19 in New York City. J Med Virol 2020; https://doi.org/10.1002/jmv.26365 Accessed 31 August 2020

[50] Capra R, De Rossi N, Mattioli F, Romanelli G, Scarpazza C, Sormani MP, et al. Impact of low dose tocilizumab on mortality rate in patients with COVID-19 related pneumonia. Eur J Intern Med 2020;76:31-5.

[51] Rossotti R, Travi G, Ughi N, Corradin M, Baiguera C, Fumagalli R, et al. Safety and efficacy of anti-il6-receptor tocilizumab use in severe and critical patients affected by coronavirus disease 2019: a comparative analysis. J Infect. 2020:S0163-4453(20)30467-9. https://doi.org/10.1016/j.jinf.2020.07.008. Accessed 31 August 2020

[52] Eimer J, Vesterbacka J, Svensson AK, Stojanovic B, Wagrell C, Sönnerborg A, et al. Tocilizumab shortens time on mechanical ventilation and length of hospital stay in patients with severe COVID-19: a retrospective cohort study. J Int Med 2020; https://doi.org/10.1111/joim.13162 Accessed 31 August 2020

[53] Kimmig LM, Wu D, Gold M, Pettit NN, Pitrak D, Mueller J, et al. IL6 inhibition in critically ill COVID-19 patients is associated with increased secondaryInfections. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.05.15.20103531v2 Accessed 31 August 2020

[54] Somers EC, Eschenauer GA, Troost JP, Golob JL, Gandhi TN, Wang Lu, et al. Tocilizumab for treatment of mechanically ventilated patients with COVID-19. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.05.29.20117358v1 Accessed 25 August 2020

[55] Tsai A, Diawara O, Nahass RG, Brunetti L. Impact of tocilizumab administration on mortality in severe COVID-19. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.07.30.20114959v1 Accessed 31 August 2020

[56] Kewan T, Covut F, Al-Jaghbeer MJ, Rose L, Gopalakrishna KV, Akbik B. Tocilizumab for treatment of patients with severe COVID19: A retrospective cohort study. EClinicalMedicine 24 (2020) 100418

[57] Patel K, Gooley TA, Bailey N, Bailey M, Hegerova L, Batchelder A, et al. Use of the IL-6R Antagonist Tocilizumab in Hospitalized COVID-19 Patients. J Inter Med 03 Aug 2020, https://doi.org/10.1111/joim.13163 Accessed 31 August 2020

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

[58] Wadud N, Ahmed N, Shergil M, Khan M, Krishna MG, Gilani A, et al. Improved survival outcome in SARs-CoV-2 (COVID-19) Acute Respiratory Distress Syndrome patients with Tocilizumab administration. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.05.13.20100081v1 Accessed 31 August 2020

[59] Ramaswamy M, Mannam P, Comer R, Sinclair E, McQuaid DB, Schmidt ML, et al. Off-Label Real World Experience Using Tocilizumab for Patients Hospitalized with COVID-19 Disease in a Regional Community Health System: A Case-Control Study. medRxiv preprint at

https://www.medrxiv.org/content/10.1101/2020.05.14.20099234v1 Accessed 31 August 2020

[60] Rojas-Marte GR, Khalid M, Mukhtar O, Hashmi AT, Waheed MA, Ehrlich S, et al. Outcomes in Patients with Severe COVID-19 Disease Treated with Tocilizumab - A Case-Controlled Study. QJM. 2020: hcaa206. https://doi.org/10.1093/qjmed/hcaa206 Accessed 31 August 2020

[61] Roumier M, Paule R, Groh M, Vallee A, Ackermann F. Interleukin-6 blockade for severe COVID-19. medRxiv preprint at

https://www.medrxiv.org/content/10.1101/2020.04.20.20061861v1 Accessed 31 August 2020

[62] Rossi B, Nguyen LS, Zimmermann P, Boucenna F, Baucher L, Dubret L, et al. Effect of tocilizumab in hospitalized patients with severe pneumonia COVID-19: a cohort study. medRxiv preprint at

https://www.medrxiv.org/content/10.1101/2020.06.06.20122341v1 Accessed 31 August 2020

[63] Potere N, Di Nisio M, Cibelli D, Scurti R, Frattari A, Porreca E, et al. Interleukin-6 receptor blockade with subcutaneous tocilizumab in severe COVID-19 pneumonia and hyperinflammation: a case-control study. Ann Rheum Dis. 2020:218243. https://doi.org/10.1136/annrheumdis-2020-218243. Accessed 31 August 2020

[64] Klopfenstein T, Zayet S, Lohse A, Balblanc JC, Badie J, Royer PY, et al. Tocilizumab 300 therapy reduced intensive care unit admissions and/or mortality in COVID-19 patients. Med Mal Infect 2020:S0399-077X(20)30129-3.

[65] Canziani LM, Trovati S, Brunetta E, De Santis M, Bombardieri E, Guidelli G, et al. Interleukin-6 receptor blocking with intravenous tocilizumab in COVID-19 severe acute respiratory distress syndrome: A retrospective case-control survival analysis of 128 patients. J Autoimmun 2020 Jul 8;102511. https://doi.org/10.1016/j.jaut.2020.102511 Accessed 31 August 2020

[66] Ip A, Berry DA, Hansen E, Goy AH, Pecora AL, Sinclaire BA, et al. Hydroxychloroquine and Tocilizumab Therapy in COVID-19 Patients – An Observational Study. medRxiv preprint at

https://www.medrxiv.org/content/10.1101/2020.05.21.20109207v1 Accessed 31 August 2020

[67] Carvalho V, Turon R, Gonçalves B, Ceotto V, Kurtz P, Righy C. Effects of Tocilizumab in Critically III Patients With COVID-19: A Quasi-Experimental Study. medRxiv preprint at https://www.medrxiv.org/content/10.1101/2020.07.13.20149328v1 Accessed 31 August 2020

[68] COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. National Institutes of Health. Available at https://www.covid19treatmentguidelines.nih.gov/. Accessed 25 August 2020.

[69] Roche. Roche provides an update on the Phase III COVACTA trial of Actemra/RoActemra in hospitalized patients with severe COVID-19 associated pneumonia. 2020. Available at: https://www.roche.com/investors/updates/inv-update-2020-07-29.htm. Accessed 10 August 2020.

[70] Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. BMJ 2009 Jul 21;339:b2700. https://doi.org/10.1136/bmj.b2700 Accessed 30 July 2020

[71] Wells G, Shea B, O'Connell D. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analysis. Available at http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp. Accessed 29 August 2020

[72] Wilson EB. Probable inference, the law of succession, and statistical inference. J Am Stat Assoc 1927;22:209-12.

[73] Garwood F. Fiducial Limits for the Poisson Distribution. Biometrika 1936,28:437-42.

[74] Kahn HA, Sempos CT. Statistical Q14 methods in epidemiology. Oxford University Press; 1989.

[75] Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in metaanalyses. BMJ 2003;327:557-60.

[76] Begg CB, Berlin JA. Publication bias and dissemination of clinical research. J Natl Cancer Inst 1989;81:107-15.

[77] Lan SH, Lai CC, Huang HT, Chang SP, Lu LC, Hsuehft PR. Tocilizumab for severe COVID-19: a systematic review and metaanalysis. International Journal of Antimicrobial Agents, https://doi.org/10.1016/j.ijantimicag.2020.106103 Accessed 30 July 2020

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

[78] Boregowda U, Perisetti A, Nanjappa A, Gajendran M, Goyal H. Addition of Tocilizumab to the standard of care reduces mortality in severe COVID-19: A systematic review and meta-analysis. medrxiv preprint at

https://www.medrxiv.org/content/10.1101/2020.07.10.20150680v3 Accessed 30 July 2020

[79] Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet 2020;395(10229):1054-62. https://doi.org/10.1016/S0140-6736(20)30566-3. Accessed 30 July 2020

[80] Antinori S, Bonazzetti C, Gubertini G, Capetti A, Pagani C, Morena V, et al. Tocilizumab for cytokine storm syndrome in COVID-19 pneumonia: an increased risk for candidemia? Autoimmun Rev. 2020;19:102564. https://doi.org/10.1016/j.autrev.2020.102564. Accessed 30 July 2020

[81] Han W, Quan B, Guo Y, Zhang J, Lu Y, Feng G, et al. The course of clinical diagnosis and treatment of a case infected with coronavirus disease 2019. J Med Virol. 2020;92:461-3. https://doi.org/10.1002/jmv.25711 Accessed 31 August 2020

Table 1. PICO framework format

Patient Problem or	Hospitalized adults with severe COVID-19						
Population	-Observational studies						
Intervention or	Tocilizumab (TCZ) administration ("TCZ-group")						
Exposure							
Comparison or	Standard of care ("Control group")						
Control							
Outcome Measures	 All-cause in-hospital mortality Overall mortality restricted to high-quality observational studies Mortality at hospitalization ward vs. ICU Mortality in the TCZ-group receiving steroids Mortality in the TCZ-group receiving early vs. late administration Risk of ICU admission Safety: Rate of secondary infections Length of hospital stay 						

Table 2. Non-controlled studies. General data.

Author	Country	Type of study	NOS	N TCZ/Control	Study period	TCZ criteria
Xu et al .[11]	CHI	Retrospective cohort	6	21/0	FEB 5-FEB 14	≥30bpm or SpO2≤93% or PaO2/FiO2≤300mmHg or MV or ICU
Luo et al.[12]	CHI	Retrospective cohort	6	15/0	JAN 27-MAR 5	High IL-6 and/or CRP
Antony et al.[13]	US	Prospective cohort	6	80/0	FEB 1-MAR 31	O2 >3L+ PSI score ≤130
Quartuccio et al.[14]	ITA	Prospective cohort	6	24/0	FEB 29*-NA	High CRP and IL-6
Toniati et al.[15]	ITA	Prospective cohort	7	100/0	MAR 9-MAR 20	BCRSS≥3
Alattar et al.[16]	QAT	Retrospective cohort	7	25/0	MAR*	≥30bpm or SpO2≤93% or PaO2/FiO2≤300mmHg +ICU+ high CRP
Sciascia et al.[17]	ITA	Prospective cohort	6	63/0	MAR*	SpO2<93%+PaO2/FiO2<300mmHg +
						3 out of: CRPx10; ferritin>1,000ng/ml; D-dimerx10; LDHx2
Morrison et al.[18]	US	Retrospective cohort	7	81/0	MAR 1-APR 3	Persistent fever+ PaO2/FiO2<200mmHg+
						1 out of:ferritin>1,000µg/l,D-dimer>5mg/ml+LDH≥500U/l, IL6x5
Morena et al.[19]	ITA	Prospective cohort	7	51/0	MAR 10-MAR 23	≥30bpm+SpO2<93%+PaO2/FiO2<250mmHg+IL-6>40pg/mI
Marfella et al.[20]	ITA	Retrospective cohort	6	78/0	MAR 1-APR 10*	Moderate-to-severe respiratory illness
Sánchez-Montalvá et al. [21]	SPA	Retrospective cohort	8	82/0	MAR 13 - MAR 18	PaO2/FiO2<300mmHg+ IL6>40pg/ml or D-dimer>1,500ng/ml
Jordan et al.[22]	US	Prospective cohort	7	27/0	MAR 13-APR 1	SpO2<90% on 4I O2 + 2 out of: IL-6>10 pg/ml; CRP>35mg/l; Ferritin>600ng/ml; D-
						dimer>1mcg/l; Neutrophil-Lymphocyte Ratio>4; LDH>200U/l
Nasir et al.[23]	РАК	Retrospective cohort	6	30/0	FEB 26-MAY 15	PaO2/FiO2<300 +CRP≥100mg/l or ferritin≥900ng/ml
Borku et al.[24]	TUR	Prospective cohort	7	12/0	MAR 18-APR 8	increased acute phase reactants, persistent fever, decreased SpO2 and deepened
						respiratory distress progress
Issa et al.[25]	FR	Retrospective cohort	6	10/0	MAR 15-APR 30	Persistent fever or PaO2/FiO2<300 + 2 out of: fibrinogen>8g/l, ferritin>1,000ng/ml, D-
						dimer>3,000ng/ml, CRP>150mg/l
Campins et al.[26]	SPA	Prospective cohort	6	58/0	MAR 21-APR 27	ΝΑ
Sanz et al.[27]	SPA	Prospective cohort	7	72/0	MAR 15-APR 8*	1 out of: PaO2/FiO2<300, SpO2<92, tachypnea or high ferritin levels
Knorr et al.[28]	US	Retrospective cohort	7	66/0	MAR 25-APR 24	D-dimer > 1.5 mg/l
Lohse et al. [29]	FR	Retrospective cohort	7	34/0	APR 1-MAY 11	\geq 5l/min O2 + \geq 2 out of: high ferritin/CRP/D-dimer/LDH and lymphopenia
Conrozier et al.[30]	FR	Retrospective cohort	7	40/0	APR 1-MAY 11	\geq 5l/min O2 + \geq 2 out of: high ferritin/CRP/D-dimer/LDH and lymphopenia
Petrak et al.[31]	US	Retrospective cohort	7	145/0	MAR 13-APR 16	NA
Górgolas et al.[32]	SPA	Retrospective cohort	7	186/0	MAR 8-APR 19	SpO2<93% on FiO2 >24%
Jiménez et al.[33]	SPA	Retrospective cohort	6	25/0	MAR 26-APR 17	≥30bpm+SpO2≤93%+PaO2/FiO2≤300mmHg+cytokine storm
Rimland et al.[34]	US	Retrospective cohort	6	11/0	MAR 21-APR 25	NA
Sinha et al.[35]	US	Prospective cohort	7	255/0	MAR 17-APR 30	FiO2>45% + CRP>100mg/l, LDH>450 U/l or ferritin>700 ng/ml
Patel A et al.[36]	IND	Retrospective cohort	6	20/0	MAR 21-MAY 20	SpO2<94% or CRPx10 or D-dimer>2,500ng/ml
Hashimoto et al. [37]	JAP	Retrospective cohort	6	13/0	NA-MAY 31	CRP>5mg/dl or ferritin>1000ng/ml
Formina et al. [38]	RUS	Retrospective cohort	6	89/0	NA	ΝΑ
Rubio-Rivas et al. [39]	SPA	Retrospective cohort	7	186/0	MAR 17-APR 7	PaO2/FiO2<300mmHg or SpO2/FiO2<315 + 2 out of: ferritin>1,000 ng/ml,
						CRP>100 mg/l, IL-6>70 ng/l, D-dimer>1,000 mcg/l, LDH>400 U/l

*Estimate. ALT: alanine aminotransferase. BCRSS: Brescia COVID-19 Respiratory Severity Scale. CRP: C-reactive protein. HFNC: high flow nasal cannula. ICU: intensive care unit. IL-6:interleukin 6. LDH: lactate deshydrogenase. MV: mechanical ventilation. NA: not applicable/available. NOS: Newcastle-Ottawa scale. PSI: pneumonia severity index. TCZ: tocilizumab

Author	Country	Type of study	NOS	N TCZ/Control	Study period	TCZ criteria
Rodríguez-Baño et al. [40]	SPA	Retrospective cohort	8	88/344	FEB 2-MAR 31	≥38ºC + increase in oxygen support to maintain SpO2>92%+ 1 out of:
				151/344		ferritin>2,000ng/ml, D-dimer>1,500mcg/ml, IL-6>50pg/ml
Quartuccio et al.[41]	ITA	Retrospective cohort	8	42/69	FEB 29-APR 6	High CRP and IL-6
Martínez-Sanz et al.[42]	SPA	Retrospective cohort	7	260/969	JAN 31-APR 23	NA
Moreno-García et al.[43]	SPA	Retrospective cohort	8	77/94	FEB 19-APR 16	CRP≥8mg/dl, ferritin≥800ng/ml or lymphocyte count<800cells/mm ³
Guaraldi et al.[44]	ITA	Retrospective cohort	8	179/365	FEB 21-APR 30	≥30bpm+SpO2<93%+PaO2/FiO2<300mmHg
Campochiaro et al.[45]	ITA	Retrospective cohort	8	32/33	MAR 13-MAR 19	SpO2≤92% + PaO2/FiO2 ≤300 mmHg +
						LDH>220 U/I + CRP≥100mg/I or ferritin ≥900 ng/mI
Mikulska et al.[46]	ITA	Retrospective cohort	7	85/66	MAR 11-MAR 24	≥30bpm+SpO2≤93%+PaO2/FiO2≤300mmHg+cytokine storm
Colaneri et al.[47]	ITA	Retrospective cohort	8	21/91	MAR 14-MAR 27	CRP>5 mg/dl+Procalcitonin<0.5 ng/mL+PaO2/FiO2<300mmHg+ALT<500 U/L.
Price et al.[48]	US	Retrospective cohort	8	153/86	MAR 10-MAR 31	O2 ≥ 3 l to maintain SpO2>93% or MV
Maeda et al.[49]	US	Retrospective cohort	6	23/201	MAR 13-MAR 31	NA
Capra et al.[50]	ITA	Retrospective cohort	8	62/23	MAR 13-APR 2	1 out of:≥30bpm+SpO2≤93%+PaO2/FiO2≤300mmHg
Rossotti et al.[51]	ITA	Retrospective cohort	8	74/148	MAR 13-APR 3	≥30bpm+SpO2≤93%+PaO2/FiO2≤300mmHg or ICU +
						CRP>1 mg/dl or IL-6>40 pg/ml or D-dimer >1.5 mcg/ml or ferritin >500 ng/ml
Eimer et al.[52]	SWE	Retrospective cohort	8	29/58	MAR 11-APR 15	SpO2<94 on 5l O2 + 1 out of: CRP>100mg/L, LDH>8 μkat/L, IL-6>40 ng/L, D-
						Dimer>2mg/L, troponin T>15 ng/L,ferritin>500 µg/L
Kimmig et al.[53]	US	Retrospective cohort	7	48/63	MAR 1-APR 27	Progressive clinical deterioration + inflammation markers
Somers et al.[54]	US	Prospective cohort	7	78/76	MAR 9-APR 20	MV
Tsai et al.[55]	US	Retrospective cohort	7	66/66	MAR 1-MAY 5	SpO2≤94%+ferritin>300 mcg/ml
Kewan et al.[56]	US	Retrospective cohort	7	28/23	MAR 13-APR 19	CRP≥3g/dl or ferritin>400ng/ml
Patel K et al.[57]	US	Retrospective cohort	7	42/41	MAR 16-APR 17	NA
Wadud et al.[58]	US	Retrospective cohort	6	44/50	MAR 15-APR 20	NA
Ramaswamy et al.[59]	US	Retrospective cohort	7	21/65	MAR 16-APR 22	SpO2≤88%+CRP≥7mg/dl
Rojas-Marte et al.[60]	US	Retrospective cohort	8	96/97	MAR 8-APR 25	O2 mask/HFNC up to 10l to maintain SpO2≥95% or MV
Roumier et al.[61]	FR	Retrospective cohort	8	30/29	MAR 21-APR 2	>6l O2 + high CRP
Rossi et al.[62]	FR	Retrospective cohort	7	106/140	MAR 23-NA	SpO2≤96%+on 6l O2
Potere et al.[63]	ITA	Retrospective cohort	7	40/40	MAR 28-APR 21	Pneumonia+CRP≥20mg/dl+SpO2<90%
Klopfenstein et al.[64]	FR	Retrospective cohort	8	20/25	APR 1-APR 13	\geq 5l/min O2 + \geq 2 out of: high ferritin/CRP/D-dimer/LDH and lymphopenia
Canziani et al.[65]	ITA	Retrospective cohort	7	64/64	FEB 23-MAY 9	Respiratory worsening + High CRP, ferritin, CK, ALT, D-dimer or lymphopenia
lp et al.[66]	US	Retrospective cohort	7	134/413	APR 22-MAY 5	ΝΑ
Carvalho et al.[67]	BRA	Prospective cohort	7	29/24	MAR 21-MAY 31	ICU + fever + High CRP≥5mg/dl

Table 3. Controlled studies. General data.

ALT: alanine aminotransferase. BCRSS: Brescia COVID-19 Respiratory Severity Scale. CRP: C-reactive protein. HFNC: high flow nasal cannula. ICU: intensive care unit. IL-6:interleukin 6. LDH: lactate deshydrogenase. MV: mechanical ventilation. NA: not applicable/available. NOS: Newcastle-Ottawa scale. PSI: pneumonia severity index. TCZ: tocilizumab

 Table 4. Non-controlled studies. Drugs and mortality.

Author	Age Mean (SD or range) or median [IQR]	Gender, males n (%)	Days from onset to TCZ, Mean (SD) or median [IQR]	TCZ doses	TCZ Infusion 1/ ≥2	Steroids n (%)	Remdesivir n (%)	Follow-up, days Mean (SD) or median [IQR]	Mortality TCZ n (%)
Xu et al.[11]	56.8 (16.5)	18 (85.7)	NA	4-8 mg/kg/d iv	18/3	NA	0	15.1 (5.8)	0
Luo et al.[12]	73 [62-80]	12 (80)	NA	80-600 mg iv	10/5	8 (53.3)	0	7 (NA)	3 (20)
Antony et al.[13]	63 [51-72]	45 (56.9)	NA	4 mg/kg/d q12h x 72h	0/80	80 (100)	0	NA	7 (8.8)
Quartuccio et al.[14]	66 (NA)*	19 (79.2)	NA	8 mg/kg/d iv	23/1	0	0	30 (NA)	6 (25)
Toniati et al.[15]	62 [57-71]	88 (88)	12 [9-14]	8 mg/kg/d iv	0/100	100 (100)	NA	10 (NA)	20 (20)
Alattar et al.[16]	58 [50-63]	23(92)	6 [6-8]	400-600 mg iv	16/9	0	0	14 (NA)	3 (12)
Sciascia et al.[17]	62.6 (12.5)	56 (88)	NA	8 mg/kg/d iv or 325 mg sc	11/52	0	NA	14 (NA)	7 (11)
Morrison et al.[18]	64 [58-71]	56 (69.1)	10 [8-13]	8 mg/kg/d iv	62/19	57 (70.4)	2 (2.5)	28 (NA)	35 (43.2)
Morena et al.[19]	60 [50-70]	40 (78.4)	12 [10-16]	400 mg-8 mg/kg/d iv	0/51	0	0	34 [32-37]	14 (27)
Marfella et al.[20]	66 (NA)*	NA	NA	8 mg/kg/d iv	66/12	0	0	20 (NA)*	39 (50)
Sánchez-Montalvá et al.[21]	59.1 (19.8)	52 (63.5)	9 [6-11]	400-600 mg iv	0/82	NA	0	7 (NA)	22 (26.8)
Jordan et al.[22]	63 [51-75]	24 (85)	NA	400 mg iv	27/0	0	7 (26)	12 (NA)*	2 (7.4)
Nasir et al.[23]	62.5 (13.5)	25 (83)	NA	600 mg iv median (range 320-680 mg)	30/0	21 (70)	0	12 (6.7)	7 (23)
Borku et al.[24]	65.8 (11.3)	6 (50)	NA	400 mg iv	0/12	0	0	14 (NA)*	0
Issa et al.[25]	66 [NA]	10 (100)	10 [NA]	8 mg/kg/d iv	10/0	3 (30)	0	11 [NA]	1 (10)
Campins et al.[26]	60.6 (NA)	42 (72.4)	NA	NA	NA	57 (98.3)	NA	NA	8 (13.8)
Sanz et al.[27]	67.5 [61-76.7]	45 (62.5)	9.2 [6-11]	400 mg iv	72/0	56 (77.8)	0	30 (NA)	21(29.2)
Knorr et al.[28]	61 [54.5-67]	41(62.1)	NA	8 mg/kg/d iv	57/9	44 (66.7)	0	14 (NA)	6 (9.1)
Lohse et al.[29]	75.3 [52-93]	24 (70.6)	NA	8 mg/kg/d iv	0/34	16 (47.1)	0	45 (NA)*	10 (29.4)
Conrozier et al.[30]	75 [65-89]*	NA	NA	8 mg/kg/d iv	0/40	NA	0	6 (NA)	10 (25)
Petrak et al.[31]	58.1 (NA)	93 (64.6)	NA	4-8 mg/kg/d iv	123/22	87 (60)	0	15.3 (NA)	41 (28.3)
Górgolas et al.[32]	65 (11.4)	126 (67.7)	11 [8-13]	400-600 mg iv	169/17	178 (95.7)	0	10 [7-15]	36 (19.4)
Jiménez et al.[33]	62 (9)	14 (56)	NA	NA	21/4	25 (100)	NA	25 [15-38]	5 (20)
Rimland et al.[34]	59 [48-65]	9 (82)	9 [7-14]	NA	10/1	3 (27)	0	18 [15-25]	3 (27)
Sinha et al.[35]	59 [47-70]	161 (63.1)	NA	400-8 mg/kg/d iv	255/0	0	0	12.9 (NA)	28 (10.9)
Patel A et al.[36]	54 [NA]	17 (85)	9.5 [8-10]	8 mg/kg/d iv	NA	13 (65)	0	12 (NA)	4 (20)
Hashimoto et al.[37]	63 (NA)	10 (76.9)	NA	400 mg iv	13/0	1 (7.7)	0	19 [14-25]	1 (7.7)
Formina et al.[38]	NA	53 (60)	10 [7-11]	400 mg iv	89/0	NA	0	NA	11 (12.4)
Rubio-Rivas et al. [39]	64.3 (13)	129 (69.4)	12 (4.3)	400-600 mg iv	172/14	155 (83.3)	4 (2.2)	60 (NA)	62 (33.3)

*Estimate. CS: corticosteroids. NA: not applicable/available. TCZ: tocilizumab

Author	Age TCZ patients Mean (SD or range) or median [IQR]	Males TCZ n (%)	Days from onset to TCZ infusion, Mean (SD) or median [IQR]	TCZ doses	TCZ Infusion 1/ ≥2	CS TCZ vs. the control group n (%)	Remdesivir TCZ vs. the control group n (%)	Follow-up, days Mean (SD) or median [IQR]	Mortality TCZ vs. the control group n (%)
Rodríguez-Baño et al. [40]	66 [56-72]	64 (72.7)	10 [8-13]	NA	NA	0 vs. 0	0	21 [16-21]	2 (2.3) vs. 41 (11.9)
Ouartuccio et al.[41]	65 [58-74] 62.4 (11.8)	109 (71.9) 33 (78.6)	11 [8-13] 8.4 (3.7)	8 mg/kg/d iv	42/0	88 (100) vs. 0 16 (38.1) vs. 0	0 3 (7.1)/0	20 [11-21] NA	19 (12.6) vs. 41 (11.9) 4 (9.5) vs. 0
Martínez-Sanz et al.[42]	65 [55-76]	191 (73)	NA	NA	NA	NA	0	NA	61 (23) vs. 120 (12)
Moreno-García et al.[43]	40 (52)	53 (68.8)	NA	400-600 mg iv	NA	39(50.6) vs.26(27.7)	0	11.2 (6.2)	8 (10.3) vs. 17 (18)
Guaraldi et al.[44]	64 [54-72]	127 (71)	7 [4-10]	8 mg/kg/d iv or 162 mg sc	0/179	0 vs. 0	0	12 [6-17]	13 (7) vs. 73 (20)
Campochiaro et al.[45]	64 [53-75]	29 (91)	11 [8-14]	400 mg iv	23/9	0 vs. 0	0	28 (NA)	5 (16) vs. 11 (33)
Mikulska et al.[46]	NA	NA	NA	400 mg-8 mg/kg/d iv or 162 mg sc	85/0	56 (65.9) vs. 0	0	30 (NA)	9 (10.6) vs. 23 (34.8)
Colaneri et al.[47]	62.3 [18-68]	19 (90.5)	NA	8 mg/kg/d iv	21/0	21 (100) vs. 91(100)	0	7 (NA)	5 (23.8) vs. 19 (20.9)
Price et al.[48]	65 [NA]	88 (58)	7 [4.5-10]	8 mg/kg/d iv	NA	47 (31) vs. 1 (1.2)	0	12 [8-22]	23 (15) vs. 10 (11.6)
Maeda et al.[49]	NA	NA	NA	NA	NA	NA	0	NA	8 (33.3) vs. 33 (16.4)
Capra et al.[50]	63 [54-73]	45 (73)	NA	400 mg iv	62/0	0 vs. 0	0	9 [5-19]	2 (3.2) vs. 11 (47.8)
Rossotti et al.[51]	59 [51-71]	61 (82.4)	NA	8 mg/kg/d iv	NA	0 vs. 0	7(9.5)/12(8.1)	7 (NA)	8 (11.8) vs. NA
Eimer et al.[52]	58 [49-63]	28 (96.6)	11 [8-14]	8 mg/kg/d iv	NA	8(27.6) vs. 23 (39.7)	0	30 (NA)	5 (17.2) vs. 19 (32.8)
Kimmig et al.[53]	64.4 (14.2)	34 (70.8)	NA	400-800 mg iv	45/3	NA	0	NA	19 (39.6) vs. 11 (17.4)
Somers et al.[54]	55 (14.9)	53 (68)	NA	8 mg/kg/d iv	78/0	23 (29) vs. 15 (19.7)	2 (3)/2 (3)	28 (NA)	14 (18) vs. 27 (36)
Tsai et al. [55]	62.4 (13.5)	46 (69.7)	NA	400-800 mg iv	62/4	NA	0	NA	18 (27.3) vs. 18 (27.3)
Kewan et al.[56]	62 [53-71]	20 (71)	NA	4-8 mg/kg/d iv	28/0	20 (71) vs. 11 (47.8)	0	11 [6-22]	3 (11) vs. 2 (9)
Patel K et al.[57]	68 [NA]	21 (50)	NA	1800 mg median [IQR 910] iv	NA	12 (29) vs. 14 (34.1)	NA	19 [14-25]	11 (26.2) vs. 11 (26.8)
Wadud et al.[58]	55.5 [NA]	37 (84.1)	NA	NA	NA	NA	0	17.9 (NA)	17 (38.4) vs. 31 (62)
Ramaswamy et al.[59]	63.2 (15.6)	13 (61.9)	NA	400-800 mg iv	21/0	9 (42.9) vs. 9 (13.8)	0	NA	3 (14.3) vs. 8 (12.3)
Rojas-Marte et al.[60]	58.8 (13.6)	74 (77.1)	NA	NA	96/0	41(42.7) vs. 32 (33)	12(12.5)/9(9.3)	NA	43 (44.8) vs. 55 (56.7)
Roumier et al.[61]	58.8 (12.4)	24 (80)	NA	8 mg/kg/d iv	NA	0 vs. 2 (6.7)	0	8 [6-9.8]	3 (10) vs. 9 (31)
Rossi et al.[62]	64.3 (13)	70 (66)	8.3 (4.2)	400 mg iv	106/0	43(40.6) vs 47(33.6)	0	28 (NA)	36 (34) vs. 80 (57.1)
Potere et al.[63]	56 [50-73]	26 (65)	NA	324 mg sc	0/40	26(65) vs. 23(57.5)	0	35 (NA)	2 (5) vs. 11 (27.5)
Klopfenstein et al. [64]	76.8 (11)	NA	13 [4-21]	NA	NA	NA	0	13 (7)	5 (25) vs. 12 (48)
Canziani et al.[65]	63 (10)	47 (73)	13 (5)	8 mg/kg/d iv	3/61	31 (48) vs. 26 (40.6)	0	30 (NA)	17 (27) vs. 24 (38)
Ip et al.[66]	62 [53-70]	99 (74)	NA	400 mg-8mg/kg/d iv	104/30	89(66) vs. 263(63.7)	0	30 (NA)	62 (46) vs. 280 (67.8)
Carvalho et al.[67]	55 [44-65]	18 (62)	NA	400 mg iv	0/29	24 (83) vs. 9 (37.5)	0	NA	5 (17) vs. 4 (17)

Table 5. Controlled studies. Drugs and mortality.

CS: corticosteroids. NA: not applicable/available. TCZ: tocilizumab

 Table 6. Non-controlled studies. Length of hospital stay, Mortality ICU vs. ward and secondary infections.

Author	Length of stay Days, mean (SD) or median [IQR]	ICU admission TCZ n/N (%)	TCZ prescribed at ICU mortality n/N (%)	TCZ prescribed at the ward mortality n/N (%)	Secondary Infections n/N (%)
Xu et al .[11]	15.1 (5.8)	2/20 (10)	NA	0/20 (0)	Overall 0/20 (0)
Luo et al.[12]	NA	NA	3/7 (42.9)	0/8 (0)	NA
Antony et al.[13]	NA (range 5-10)	9/80 (11.3)	NA	7/80 (8.7)	Overall 12/80 (15) Bacterial 12/80 (15)
Quartuccio et al.[14]	NA	NA	NA	NA	NA
Toniati et al.[15]	NA	3/57 (5.3)	10/43 (24)	10/57 (17.5)	Overall 2/100 (2) Bacterial 2/100 (2)
Alattar et al.[16]	NA	NA	3/25 (12)	ΝΑ	Overall 8/25 (32) Fungal 8/25 (32)
Sciascia et al.[17]	NA	NA	1/5 (20)	6/58 (10.3)	NA
Morrison et al.[18]	18 [13-29]	NA	32/70 (45.7)	3/11 (27.3)	Overall 20/81 (24.7) Bacterial 17/81 (21) Fungal 3/81 (3.7)
Morena et al.[19]	NA	NA	5/6 (83)	9/45 (20)	Overall 14/51 (27) Bacterial 14/51 (27)
Marfella et al.[20]	NA	NA	NA	NA	NA
Sánchez-Montalvá et al. [21]	NA	14/82 (17.1)	NA	22/82 (26.8)	Overall 0/82 (0)
Jordan et al.[22]	NA	1/27 (3.7)	2/21 (9.5)	0/6 (0)	Overall 2/27 (7.4) Bacterial 2/27 (7.4)
Nasir et al.[23]	12 (6.7)	0/20 (0)	NA	NA	Overall 16/30 (54) Bacterial 9/30 (30) Fungal 7/30 (24)
Borku et al.[24]	NA	2/12 (16.7)	NA	0/12 (0)	NA
Issa et al.[25]	NA	0	1/10 (10)	NA	Overall 2/10 (20) Bacterial 1/10 (10) Fungal 1/10 (10)
Campins et al.[26]	NA	19/58 (32.4)	NA	8/58 (13.8)	NA
Sanz et al.[27]	16.4 [11-20]	43/72 (59.7)	NA	21/72 (29.2)	NA
Knorr et al.[28]	16.4 [8.8-23]	16/48 (33.3)	1/18 (5.6)	3/48 (6.25)	NA
Lohse et al.[29]	NA	NA	NA	10/34 (29.4)	NA
Conrozier et al.[30]	NA	NA	NA	NA	NA
Petrak et al.[31]	15.3 (NA)	NA	35/81 (43.2)	6/64 (9.4)	Overall 0/145 (0)
Górgolas et al.[32]	NA	19/186 (10.2)	NA	36/186 (19.4)	Overall 13/186 (6.3) Bacterial 6/186 (3.2)
Jiménez et al.[33]	25 [15-38]	NA	NA	NA	Overall 8/25 (32)
Rimland et al.[34]	NA	5/7 (71.4)	NA	NA	Overall 2/11 (18.2)
Sinha et al.[35]	12.9 (NA)	NA	NA	ΝΑ	Overall 34/255 (13.3) Bacterial 34/255 (13.3)
Patel A et al.[36]	NA	2/20 (10)	NA	4/20 (20)	Overall 1/20 (5)
Hashimoto et al [27]	17 (range 12-27)	4/11 (36 1)	0/2 (0)	1/11 (9 1)	
Formina at al [38]		4/ 11 (30.4) NA	10/17 (58 8)	1/72 (1 /)	NA
Rubio-Rivas et al. [39]	NA	NA	NA	NA	NA

ICU: intensive care unit. NA: Not available/applicable. TCZ: tocilizumab

Table 7.	Controlled studies.	. Length of hospital stay	y, Mortality ICU vs. ward and secondary infections	s

Author	Length of stay TC7 vs. Control days		TC7 prescribed at ICU	TC7 prescribed at the ward	Secondary Infections
Aution	mean (SD) or median [IQR]	TCZ vs. Control, n/N (%)	mortality TCZ vs. Control, n/N (%)	mortality TCZ vs. Control, n/N (%)	TCZ vs. control group n/N (%)
Rodriguez-Baño et al.[40]	NA	NA	NA	NA	Overall 11/88 (12.5) vs. 36/339 (10.3)
					Bacterial 11/88 (12.5) vs. 36/339 (10.3)
					Overall 18/150 (12) vs. 36/339 (10.3)
					Bacterial 18/150 (12) vs. 36/339 (10.3)
Quartuccio et al.[41]	NA	0/15 (0) vs. 0/69 (0)	3/27 (11.1) vs. 0/0 (0)	1/15 (6.7) vs. 0/69 (0)	Overall 18/27 (16.2) vs. 0 (0) Bacterial 18/27 (16.2) vs. 0 (0)
Martínez-Sanz et al.[42]	19 [NA] vs. 10 [NA]*	50/260 (19) vs. 32/969 (3)	NA	61/260 (23) vs. 120/969 (12)	NA
Moreno-García et al.[43]	NA	8/77 (10.3) vs. 26/94 (27.6)	NA	8/77 (10.3) vs. 17/94 (18)	NA
Guaraldi et al.[44]	NA	NA	NA	NA	Overall 24/132 (14.4) vs. 14/222 (6) Bacterial 12/132 (9.1) vs.11/222 (4.6) Fungal 7/132 (5.3) vs. 3/222 (1.4)
Campochiaro et al.[45]	13.5[10-16.7] vs. 14 [12-15.5]	4/32 (13) vs. 2/33 (6)	NA	5/32 (16) vs. 11/33 (33)	Overall 5/32 (16.1) vs. 4/33 (12) Bacterial 4/32 (13) vs. 4/33 (12) Fungal 1/32 (3.1) vs. 0/33 (0)
Mikulska et al.[46]	NA	NA	NA	NA	NA
Colaneri et al.[47]	2[6] vs. 14 [4]	3/21 (14.3) vs. 12/91 (13.2)	NA	5/21 (23.8) vs. 19/91 (20.9)	Overall 0/21 (0) vs. 0/91 (0)
Price et al.[48]	12[8-22] vs. NA	NA	NA	NA	Overall 4/153 (2.6) vs.NA Bacterial 4/153 (2.6) vs.NA
Maeda et al.[49]	NA	NA	ΝΑ	ΝΑ	Overall 3/23 (13) vs. 2/201 (1.1) Fungal 3/23 (13) vs. 2/201 (1.1)
Capra et al.[50]	12.5 [4-18] vs. 8 [7-15]	NA	NA	NA	Overall 0/62 (0) vs. NA
Rossotti et al.[51]	NA	NA	NA	NA	Overall 24/74 (32.4) vs. NA
Eimer et al.[52]	20.5 [16.5-30] vs. 30[21.5-30]	NA	5/29 (17.2) vs. 19/58 (32.8)	NA	Overall 5/29 (17.2) vs. 14/58 (24.1) Bacterial 5/29 (17.2) vs. 14/58 (24.1)
Kimmig et al.[53]	NA	NA	19/48 (39.6) vs. 11/63 (17.4)	NA	Overall 27/48 (56.3) vs. 18/63 (28.6) Bacterial 24/48 (50) vs. 18/63 (28.6)
Somers et al.[54]	20.4 [13.8-35.8] vs. 22.9[16.3-28.5]	NA	14/78 (18) vs. 27/76 (36)	NA	Overall 41/78 (52.5) vs. 22/76 (28.9) Bacterial 38/78 (48.7) vs. 21/76 (27.6) Fungal 3/78 (3.8) vs. 1/76 (1.3)
Tsai et al.[55]	NA	NA	NA	NA	NA
Kewan et al.[56]	11[6-22.3] vs. 7 [5-13.5]	NA	NA	NA	Overall 5/28 (18) vs. 5/51 (22)
Patel K et al.[57]	NA	NA	NA	NA	NA
Wadud et al.[58]	NA	NA	NA	NA	NA
Ramaswamy et al.[59]	NA	NA	NA	NA	NA
Rojas-Marte et al.[60]	14.5 (8.8) vs. 16.5 (10.8)	NA	41/61 (67.2) vs. 45/60 (75)*	2/33(6.1) vs. 9/34 (26.5)*	Overall 16/61 (16.7) vs. 26/60 (26.8) Bacterial 12/61 (12.5) vs. 23/60 (23.7) Fungal 4/61 (4.2) vs.3/60 (3.1)
Roumier et al.[61]	NA	NA	NA	NA	NA
Rossi et al.[62]	NA	NA	NA	NA	NA
Potere et al.[63]	NA	NA	NA	NA	Overall 1/40 (2.5) vs. 3/40 (7.5) Bacterial 1/40 (2.5) vs. 3/40 (7.5)
Klopfenstein et al. [64]	13 [4-32] vs. 17 [5-41]	0/20 (0) vs. 11/25 (44)	NA	5/20 (25) vs. 12/25 (48)	NA
Canziani et al.[65]	NA	NA	NA	NA	Overall 20/64 (31) vs. 25/64 (39)
lp et al.[66]	NA	NA	62/134 (46) vs. 231/413 (56)	NA	Overall 30/134 (22) vs. 69/413 (17) Bacterial 30/134 (22) vs. 69/413 (17)
Carvalho et al.[67]	NA	NA	5/29 (17) vs. 4/24 (17)	NA	Overall 16/29 (55) vs. 4/24 (16) Bacterial 10/29 (34) vs. 3/24 (12) Fungal 6/29 (21) vs. 1/24 (4)

*Estimate. ICU: intensive care unit. NA: Not available/applicable. TCZ: tocilizumab.



Figure 2. Worldwide Distribution of the Included Studies.

