

**Brief communication: A meta-analysis of randomized trials of
hydroxychloroquine for the prevention of COVID-19**

Xabier García-Albéniz^{1,2}, Julia del Amo³, Rosa Polo³,

José Miguel Morales-Asencio⁴, Miguel A Hernán^{2,5}

¹ RTI Health Solutions. Barcelona, Spain

² Department of Epidemiology. Harvard T.H. Chan School of Public Health. Boston, MA, USA

³ National Plan Against AIDS, Ministry of Health. Madrid, Spain

⁴ Department of Nursing and Podiatry, Universidad de Málaga, Málaga, Spain.

⁵ Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA, USA; Harvard-MIT Division of Health Sciences and Technology, Boston, MA, USA

Correspondence:

Xabier Garcia-Albeniz. RTI Health Solutions. Av. Diagonal, 605, 9-1. 08028. Barcelona, Spain. xgarcia@rti.org

Abstract

Background: There is disagreement about whether hydroxychloroquine (HCQ) is effective as prophylaxis for COVID-19. We conducted a meta-analysis of randomized trials that study the effectiveness of HCQ to prevent COVID-19.

Methods and Findings: A search of PubMed, Embase, medRxiv, and clinicaltrials.gov found three completed randomized trials: one pre-exposure prophylaxis trial and two post-exposure prophylaxis trials. We obtained or calculated the risk ratio of COVID-19 diagnosis for assignment to HCQ versus no HCQ (either placebo or usual care) for each trial, and then pooled the risk ratio estimates. The risk ratio estimated for each of the individual trials were 0.74 (95% CI 0.50-1.10), 0.83 (95% CI: 0.58-1.18), and 0.69 (95% CI: 0.35-1.37). The pooled risk ratio estimate was 0.78 (95% CI: 0.61-0.99). All three trials found a similar rate of adverse effects in the HCQ and no HCQ groups.

Discussion: The available evidence indicates that HCQ reduces the risk of COVID-19 by about 20%. Yet the findings from the randomized trials were widely interpreted as evidence of lack of effectiveness of HCQ, simply because they were not “statistically significant” when taking them individually. Completion of the ongoing prophylaxis trials is needed to generate more precise estimates of the effectiveness of HCQ to prevent COVID-19.

Background

Hydroxychloroquine (HCQ) is not an effective treatment for established COVID-19 (1-4). However, there is disagreement about whether HCQ is effective as prophylaxis for COVID-19. We conducted a meta-analysis of randomized trials that study the effectiveness of HCQ to prevent COVID-19 either before known exposure to an infected individual (pre-exposure prophylaxis) or after known exposure to an infected individual (postexposure prophylaxis).

Methods and Findings

A search of PubMed, Embase, medRxiv, and clinicaltrials.gov found 9 ongoing randomized trials (5-13) and three that have been completed: one pre-exposure prophylaxis trial (14) and two post-exposure prophylaxis trials (15, 16). We obtained or calculated the risk ratio of COVID-19 diagnosis for assignment to HCQ versus no HCQ (either placebo or usual care) for each of the completed trials. When the trial included both PCR-positive and PCR-negative individuals at baseline, we restricted our attention to the PCR-negative individuals. We then pooled the risk ratio estimates using both fixed effect and random effects meta-analytic approaches.

The Figure shows the risk ratio estimates from each of the three trials trial and the pooled estimates.

The first trial was a double-blinded, placebo-controlled trial of pre-exposure prophylaxis (available as a preprint) that included healthcare workers with ongoing exposure to patients with COVID-19. There were 58 cases (11 confirmed by PCR) among 989 (5.9%) participants assigned to HCQ (two 400 mg doses on day 1, followed by 400 mg either once or twice weekly for 12 weeks) and 39 cases (6 confirmed by PCR) among 494 (7.9%) participants

assigned to placebo. The risk difference can be calculated as -2.0% (95% CI -4.8 to 0.8) and the risk ratio can be calculated as 0.74 (95% CI 0.50-1.10).

The second trial was a double-blind, placebo-controlled trial of post-exposure prophylaxis (15) included asymptomatic contacts of confirmed COVID-19 cases. The risk of COVID-19 diagnosis (largely based on compatible symptoms) was 11.8% in 414 contacts assigned to HCQ (800 mg. on day 1, 600 mg. daily for four days), and 14.3% in 407 contacts assigned to placebo. The risk difference was -2.4% (95% CI: -7.0 to 2.2) and the risk ratio can be calculated as 0.83 (95% CI: 0.58-1.18).

The third trial was an open-label, cluster-randomized trial of post-exposure prophylaxis (16) (available as a preprint) that included asymptomatic contacts of confirmed COVID-19 cases. The COVID-19 risk was 3.0% in 958 contacts assigned to HCQ (800 mg. on day 1, 400 mg. daily for six days), and 4.3% in 1042 contacts assigned to usual care. The risk ratio (adjusted for age, gender, region and time of exposure) was 0.69 (95% CI: 0.35-1.37); the corresponding risk difference cannot be directly calculated from the data provided.

The pooled risk ratio estimated from the three trials was 0.78 (95% CI: 0.61-0.99). All three trials found a similar rate of adverse effects in the HCQ and no HCQ groups.

Discussion

When considered together, the available randomized trials of HCQ for the prevention of COVID-19 indicate that HCQ reduces the risk of COVID-19 by about 20%. The actual effect of HCQ may be underestimated because none of these trials adjusted for losses to follow-up or other deviations from protocol. Also, effectiveness may be underestimated in the post-exposure prophylaxis trials because the time from exposure to initiation of prophylaxis was relatively long: in one trial, about a third of participants were enrolled 4 days after exposure

(none was enrolled later); in the other, participants were enrolled up to 7 days after exposure.


For comparison, post-exposure prophylaxis for HIV is recommended in the first 6-72h after the exposure (17).

Pending final publication, these trials suggests a 20% reduction of COVID-19 risk attributed to HCQ, with no increased risk of serious side effects. Yet the findings from the trials were widely (and incorrectly) interpreted as evidence of lack of effectiveness of HCQ, simply because they were not “statistically significant” when taking them individually. Completion of the ongoing prophylaxis trials is needed to generate more precise estimates of the effectiveness of HCQ to prevent COVID-19.

References

1. Ramsay LE, Yeo WW, Jackson PR. Dietary reduction of serum cholesterol concentration: time to think again. *British Medical Journal*. 1991;303:953-7.
2. Boyer PJ, Dillon M, Navaie M, Deveikis A, Keller M, O'Rourke S, et al. Factors predictive of maternal-fetal transmission of HIV-1. Preliminary analysis of zidovudine given pregnancy and/or delivery. *Journal of the American Medical Association*. 1994;271(24):1925-30.
3. Rogers MF, Ou C-Y, Rayfield M, Thomas PA, Schoenbaun EE, Abrams E, et al. Use of the polymerase chain reaction for early detection of the proviral sequences of human immunodeficiency virus in infants born to seropositive mothers. *New England Journal of Medicine*. 1989;320:1649-54.
4. Greenberg BL, Semba RD, Vink PE, Farley JJ, Sivapalasingam M, Steketee RW, et al. Vitamin A deficiency and maternal-infant transmission of HIV in two metropolitan areas in the United States. *AIDS*. 1997;11:325-32.
5. Grau-Pujol B, Camprubí D, Marti-Soler H, Fernández-Pardos M, Guinovart C, Muñoz J. Pre-exposure prophylaxis with hydroxychloroquine for high-risk healthcare workers during the COVID-19 pandemic: A structured summary of a study protocol for a multicentre, double-blind randomized controlled trial. *Trials*. 2020;21(1):688.
6. University SZABM. Efficacy of Various Doses of Hydroxychloroquine in Pre-Exposure Prophylaxis for COVID 19. <https://ClinicalTrials.gov/show/NCT04359537>; 2020.
7. Foundation G. HEalth Care Worker pROphylaxis Against COVID-19: The HERO Trial. <https://ClinicalTrials.gov/show/NCT04352946>; 2020.
8. Landes M, University Health Network T. Does Hydroxychloroquine Before & During Patient Exposure Protect Healthcare Workers From Coronavirus? : <https://ClinicalTrials.gov/show/NCT04374942>; 2020.
9. Sida PNse, S.L. ESPII. Randomized Clinical Trial for the Prevention of SARS-CoV-2 Infection (COVID-19) in Healthcare Personnel. <https://ClinicalTrials.gov/show/NCT04334928>; 2020.
10. Ceara NDPEDMDUFD. COVID-19 Prophylaxis With Hydroxychloroquine Associated With Zinc For High-Risk Healthcare Workers. <https://ClinicalTrials.gov/show/NCT04384458>; 2020.
11. Oxford Uo. Chloroquine/ Hydroxychloroquine Prevention of Coronavirus Disease (COVID-19) in the Healthcare Setting. <https://ClinicalTrials.gov/show/NCT04303507>; 2020.
12. Education PloM, Research. Efficacy of Hydroxychloroquine (HCQ) as Post Exposure Prophylaxis (PEP) for Prevention of COVID-19. <https://ClinicalTrials.gov/show/NCT04408456>; 2020.
13. System HFH. Will Hydroxychloroquine Impede or Prevent COVID-19. <https://ClinicalTrials.gov/show/NCT04341441>; 2020.
14. Rajasingham R, Bangdiwala AS, Nicol MR, Skipper CP, Pastick KA, Axelrod ML, et al. Hydroxychloroquine as pre-exposure prophylaxis for COVID-19 in healthcare workers: a randomized trial. *medRxiv*. 2020:2020.09.18.20197327.
15. Boulware DR, Pullen MF, Bangdiwala AS, Pastick KA, Lofgren SM, Okafor EC, et al. A Randomized Trial of Hydroxychloroquine as Postexposure Prophylaxis for Covid-19. *New England Journal of Medicine*. 2020;383(6):517-25.
16. Mitja O, Ubals M, Corbacho M, Alemany A, Suner C, Tebe C, et al. A Cluster-Randomized Trial of Hydroxychloroquine as Prevention of Covid-19 Transmission and Disease. *medRxiv*. 2020:2020.07.20.20157651.
17. Sempos CT, Cleeman JI, Carroll MD, Johnson CL, Bachorik PS, Gordon DJ, et al. Prevalence of high blood cholesterol among US adults. *Journal of the American Medical Association*. 1993;269(23):3009-14.

Figure: Risk ratio estimates of COVID-19 risk for hydroxychloroquine vs. no hydroxychloroquine in randomized trials of pre- and post-exposure prophylaxis, pooled and by study.



TE: treatment effect; seTE: standard error of the treatment effect; RR: risk ratio; CI: confidence interval