Vulnerability to rumors during the COVID-19 pandemic:

Results of a national survey

Victoria Jane En Long¹, MPH, Medical student Wei Shien Koh¹, BSocSc, Medical student Young Ern Saw², BA

*Jean CJ Liu^{1,2}, PhD, Assistant Professor

¹Duke-NUS Medical School, Singapore

²Division of Social Sciences, Yale-NUS College, Singapore

Victoria Jane En Long: victoria.long@u.duke.nus.edu

Wei Shien Koh: weishienkoh@u.duke.nus.edu

Young Ern Saw: sawyoungern@u.yale-nus.edu.sg

Jean CJ Liu: jeanliu@yale-nus.edu.sg

Affiliation addresses: **Duke-NUS Medical School** 8 College Rd, Singapore 169857 Yale-NUS College 28 College Ave West

Running head: COVID-19 rumors

*Please address correspondence to:

Jean CJ Liu Yale-NUS College 28 College Ave West #01-501

Singapore 138533

Email: jeanliu@yale-nus.edu.sq

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

- 1 **Funding**
- 2 This research was funded by a grant awarded to JCJL from the JY Pillay
- 3 Global Asia Programme [grant number: IG20-SG002].
- 6 Highlights

4

5

- 7 Prior studies linked exposure to COVID-19 rumors with poor mental health.
- 8 In a community sample, most participants reported having heard rumors.
- 9 Few participants shared or believed rumors.
- 10 Sharing sometimes occurred in the absence of belief.
- 11 More educated individuals believed and shared fewer rumors.

Abstract

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

media; infodemic; fake news

Amidst the COVID-19 pandemic, many rumors have emerged. Given prior research linking rumor exposure to mental well-being, we conducted a nation-wide survey to document the base rate of rumor exposure and factors associated with rumor vulnerability. Between March to July 2020, 1237 participants were surveyed on 5 widely-disseminated COVID-19 rumors (that drinking water frequently could be preventive, that eating garlic could be preventive, that the outbreak arose because of bat soup consumption, that the virus was created in an American lab, and that the virus was created in a Chinese lab). For each rumor, participants reported whether they had heard, shared or believed each rumor. Although most participants had been exposed to COVID-19 rumors, few shared or believed these. Sharing behaviors sometimes occurred in the absence of belief; however, education emerged as a protective factor for both sharing and belief. Together, our results suggest that campaigns targeting skills associated with higher education (e.g. epistemology) may prove more effective than counter-rumor messages. KEYWORDS: Infectious diseases; public health; COVID-19; misinformation; social

All rights reserved. No reuse allowed without permission.

COVID-19 rumors

Vulnerability to rumors during the COVID-19 pandemic:

Results of a national survey

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

1. Introduction The global outbreak of coronavirus disease 2019 (COVID-19) has come with increased psychological burden. In several meta-analyses, depression and anxiety symptoms have been found to be elevated amongst healthcare workers and the general population since the start of the pandemic (da Silva and Neto, 2021; Pappa et al., 2020; Salari et al., 2020). Others have reported a higher incidence of stressrelated symptoms or post-traumatic stress disorder (Cooke et al., 2020; Torales et al., 2020). These findings highlight the urgent need to understand factors these predicting anxiety and mood outcomes, allowing vulnerable individuals to be identified and interventions to be developed. In terms of predictors, exposure to COVID-19 rumors has emerged as a risk factor for poor mental health (Gao et al., 2020; Liu and Tong, 2020; Vardanjani et al., 2020; Xiong et al., 2020). This negative mental health impact has occurred against the backdrop of an "infodemic" - a surge of COVID-19 misinformation created and shared primarily via social media (World Health Organization, 2020). In particular, the fast-changing nature of the pandemic means that accurate information has not always been accessible (Cuan-Baltazar et al., 2020; Hou et al., 2020), resulting in many uncertainties. This has given rise to a large number of rumors (Depoux et al., 2020; Larson, 2018; Vosoughi et al., 2018). To date, several publications have used publicly available data to analyze and document the spread of rumors. For example, during the early stage of the pandemic (December 2019 to April 2020), search engine keywords reflected popular myths (Rovetta and Bhagavathula, 2020; Singh et al., 2020), with a large number of searches pertaining to alternative medicines that had been speculated to prevent COVID-19 (e.g., garlic, Chinese medicinal herbs, or the malaria medication chloroquine; Hou et al., 2020). On the social media platform Twitter, conspiratorial

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

theories were posted regarding disease origins – suggesting, for example, that the virus had been developed as a bioweapon or had resulted from the introduction of 5G mobile networks (Abd-Alrazaq et al., 2020; Ahmed et al., 2020). In turn, the spread of COVID-19 rumors has led to deleterious consequences. In Iran for example, a myth that alcohol consumption could prevent or treat COVID-19 resulted in over 700 deaths related to methanol poisoning, with deaths attributed to methanol poisoning exceeding those attributed to COVID-19 in some provinces (Aghababaeian et al., 2020). Returning to mental health outcomes, the extent to which an individual has been exposed to, has shared, or believed in COVID-19 rumors has also been found to predict anxiety symptoms (Liu and Tong, 2020). While demographic predictors of pandemic-related mental health are difficult to address (e.g., age, gender, pre-existing medical conditions; (Torales et al., 2020), a person's exposure to rumors may constitute a modifiable risk factor (Abdoli, 2020; Ebrahim et al., 2020; Holmes et al., 2020). Correspondingly, efforts to develop interventions would benefit from an understanding of rumor vulnerability: (i) the base rates by which individuals are exposed to, believe in, or share rumors; and (ii) factors predicting these rumor-related experiences (Chua and Banerjee, 2018). At present, little is known about individual vulnerability to COVID-19 rumors. While a handful of studies have surveyed individuals on their social media usage and enquired about rumor dissemination via these platforms (Banakar et al., 2020), we are not aware of any study that has identified persons most likely to encounter, believe in, or to share COVID-19 rumors. To address this gap in the literature, we thus conducted a nationwide survey examining rumor vulnerability during the COVID-19 pandemic. 2. Methods 2.1 Study design and population Our study was conducted across five months in Singapore (7 March to 27 July 2020), a city-state in Asia that experienced a high number of COVID-19 cases in

the early stage of the pandemic. During this time, we recruited 1237 participants who met the following eligibility criteria: (1) aged ≥21 years old, and (2) had lived in Singapore for ≥2 years. All participants were recruited via social media advertisements within community groups (e.g., groups for residential estates, universities, and workplaces), or through paid Facebook advertisements targeting Singapore-based users.

Upon study enrolment, participants provided informed consent and completed a 20-minute online survey via Qualtrics. As part of a larger study, participants reported their: (i) demographics, (ii) responses to the pandemic; (iii) sources of COVID-19 news; and (as we report in this paper) (iv) familiarity with rumors (Liu and Tong, 2020; Long and Liu, 2020; Saw et al., 2020). The study protocol was approved by the Yale-NUS College Ethics Review Committee (#2020-CERC-001) and was pre-registered on ClinicalTrials.gov (NCT04305574).

2.2 Outcome variables

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

As the primary outcome variables, we assessed participants' familiarity with five rumors that had been widely spread during the COVID-19 pandemic: that (1) drinking water frequently will help prevent infection (COVID-19 prevention); (2) eating garlic can help prevent infection (COVID-19 prevention); (3) the outbreak arose from people eating bat soup (COVID-19 origins); (4) the virus was created in a US lab to affect China's economy (COVID-19 origins); and (5) the virus was created in a Chinese lab as a bioweapon (COVID-19 origins). These rumors were presented in the survey as claims, rather than rumors to avoid influencing participants' response. Rumors were selected for their widespread distribution both internationally and within the local context (Taylor, 2020; World Health Organisation, 2020).

For each rumor, participants indicated whether they: (1) had heard the claim before (yes/no); (2) thought the claim was true (yes/no); or (3) had shared the claim on social media (e.g., Facebook, WhatsApp) (yes/no). We assigned a score of 1 for "yes" responses, and summed across the rumors to create three scores: the total

perpetuity.
All rights reserved. No reuse allowed without permission.

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

excluded from statistical analyses.

COVID-19 rumors 7

number of claims heard, the total number of claims believed, and the total number of claims shared. Finally, participants also indicated which of 13 possible sources they had encountered the rumors (e.g., Facebook, WhatsApp, online forums, television). 2.3 Predictor variables As predictor variables, participants reported the following demographic details: age, gender, ethnicity, religion, country of birth, marital status, education, house type (a proxy of socio-economic status), and household size. Using the survey timestamp, we also recorded two situation-related variables: the total number of local cases reported to date, and whether the country had been in a lock-down when participants completed the survey. 2.4 Statistical analysis Using counts (%), we first summarized the baseline rates of rumor familiarity and rumor sources. As further exploratory analyses, we conducted Fisher's exact test exploring the relationship between believing and sharing each rumor. We then ran linear regression models to predict the following outcome measures: the total number of claims heard [Model 1], the total number of claims believed [Model 2], and the total number of claims shared [Model 3]. Each model involved the full set of predictor variables (described in 2.3), with the number of COVID-19 cases log-transformed for linearity. For each model, we applied Bonferroni correction to control the type 1 familywise error rate at 0.05 (Bonferroni-adjusted alpha level of 0.05/22 predictors = 0.002). All statistical analyses were conducted using R (Version 4.0). 3. Results 3.1 Response rate Out of 1751 individuals who accessed the survey link, 1446 (82.6%) provided informed consent and participated in the survey. However, 209 (14.5%) participants did not complete the primary outcome measures (on COVID-19 rumors) and were

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

COVID-19 rumors 8

The final sample of 1237 participants is comparable to the resident population with regards to: the proportion of participants born in Singapore, ethnicity, and household size (≤10% difference) and age. However, our sample had more participants who were female (63.9% vs. 51.1%), single (41.9% vs 18.8%), and university graduates (70.7% vs. 32.4%); and fewer participants who lived in 1-3 room public housing flats (8.2% vs. 23.7%) or who had Buddhist beliefs (14.6% vs 33.2%) (Table 1). [Table 1 about here.] 3.2 Base rates of familiarity with COVID-19 rumors Out of 5 widely-disseminated rumors, the average participant had heard of 3.34 (SD = 1.33) rumors. The most commonly-heard rumor – reported by 8 in 10 participants (84.6%) – was that the outbreak had arisen from individuals eating bat soup. Despite high exposure to COVID-19 rumors, however, participants only believed an average of 0.27 claims (SD = 0.59) and shared 0.18 (SD = 0.63). The most commonly-believed rumor was that drinking water could prevent infection (11.4%), whereas the most commonly-shared rumor was again that the disease had arisen from bat soup consumption (7.1%) (Figure 1). [Figures 1 and 2 about here.] In other words, most participants who had heard each of the 5 rumors neither believed in nor shared the claims. Using Fisher's exact test, we conducted exploratory analyses to examine how belief and sharing behaviors were related. First, for the claim about the United States manufacturing the coronavirus to affect China's economy, none who shared this rumor believed that it was true (p-value of 1 for Fisher's exact test). In the case of the other 4 rumors, however, there was a significant association between belief and sharing (p < 0.001 for the rumors on drinking water and bat soup; p = 0.001 for the rumor on garlic, and p = 0.02 for the rumor on China creating the virus). Nonetheless, even with these 4 rumors, not all

who propagated the rumors believed that they were true (Figure 2).

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

Finally, Figure 3 depicts how participants had encountered COVID-19 rumors. As has been previously reported (Ippolito et al., 2020; Pew Research Center, 2018), social media platforms emerged as the leading sources, with 1 in 2 individuals reporting exposure through Facebook (55.5%) or WhatsApp (53.6%). [Figure 3 about here.] 3.3 Predicting rumor hearing, sharing, and believing Model 1 examined if any demographic or situational factors predicted the number of rumors heard. As shown in Table 2, participants reported hearing more rumors when confirmed local cases were few (early in the pandemic) (b = -0.621, t(1190) = -3.588, p < 0.001) or as lockdown restrictions were lifted (b = 1.129, t(1190)= 4.289, p < 0.001). Additionally, there was a trend for education to predict rumor exposure, with those with higher education hearing more rumors, (b = 0.077, t(1190)= 2.625, p = 0.009). (However, this association was not observed with Bonferroni correction.) In Models 2 and 3, the same set of predictors were used to predict the number of rumors shared and believed, respectively. For both these models, those who were more educated shared or believed fewer rumors (Model 2: b = -0.046, t(1190) = -3.289, p = 0.001; Model 3: b = -0.046, t(1190) = -3.488, p = 0.001). [Table 2 about here.] 4. Discussion During times of crisis, rumors have the potential to transmit misinformation and induce anxiety (Jin et al., 2014; Tran et al., 2020). Against the backdrop of the COVID-19 pandemic, we thus documented how individuals in the community were vulnerable to receive, believe in, or share specific COVID-19 rumors. First, we observed that rumor exposure was endemic. Nearly all participants had heard at least one rumor and were familiar with an average of 3 out of 5 popular claims assessed. Additionally, most rumor transmission occurred via social media

channels (e.g., Facebook and WhatsApp), as others have noted (Islam et al., 2020; Wang et al., 2019).

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

Extending previous research, we further described how the base rate of believing or sharing rumors was far lower than the rate of exposure (with an average of <1 rumor believed or shared). Notably, belief and sharing behaviors did not always co-occur. In the extreme case of one rumor in particular (that the COVID-19 crisis had been manufactured by the United States), not a single participant who reported forwarding the rumor actually believed in it. Although belief and sharing were linked for the other rumors we assessed, there continued to be – in each case – individuals who shared rumors without believing their veracity.

Our finding that COVID-19 rumors were disseminated even when disbelieved highlights the sheer difficulty of managing the so-called 'infodemic'. Although similar findings had been reported outside the COVID-19 context (Chua and Banerjee, 2017), the World Health Organization and individual governments continue to issue as the prevailing strategy - fact-checking statements to debunk rumors (Wong et al., 2020; World Health Organisation, 2020; Zarocostas, 2020). Our results bring to question the utility of such statements, since individuals continue to share claims despite perceiving them to be untrue.

Based on our findings, an alternative strategy might be to target individual vulnerabilities instead of rumor content. Given that rumor exposure changed with pandemic severity (e.g., the number of cases) and most individuals had encountered COVID-19 rumors, the ensuing question is why only certain individuals fell prey to either belief in or the sharing of these rumors. In our analyses, we found that educational level was a consistent predictor of vulnerability: although higher education predicted that an individual would hear more rumors, higher education was nonetheless protective, associated with fewer rumors shared or believed in. Consequently, it may be profitable to increase public awareness of knowledge and skillsets associated with higher education – for example, epistemology or scientific

thinking (Chong et al., 2020; Chua and Banerjee, 2017). We note, however, that the correlational nature of our dataset precludes causal inferences, and further research will need to examine the efficacy of such strategies in curbing pandemic-related rumors.

4.1 Limitations

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

In describing these findings, we highlight several limitations of our research methods. First, we relied on participants' self-reports regarding rumor exposure and behaviors. Although this strategy provided individual-level information (e.g., beliefs, demographics) not available in studies of actual rumor posts (e.g., when Twitter posts are mined), the survey method is vulnerable to recollection and reporting biases. Moving forward, future studies may opt to integrate digital documentation of rumor posts alongside self-reported measures.

As a second limitation, we only sampled rumors that were not time-sensitive. Given the limitations of the survey methodology, we could not track rumors that arose from fast-changing events on the ground – for example, rumors about the first COVID-19-related death in Singapore, or rumors about the availability of face masks (Asokan, 2020; Ministry of Communications and Information, 2020). It thus remains to be seen whether our findings can generalize to these forms of rumors.

4.2 Strengths

These study limitations need to be viewed alongside the putative strengths of our research methodology. To the best of our knowledge, our study represents the first attempt to identify individual vulnerabilities in the spread of COVID-19 rumors. The research involved a large sample size (1237 participants), captured pandemicrelated dynamics over a long duration (5 months), and examined specific rumors that had been widely disseminated.

4.3 Conclusions

In conclusion, our study revealed that educational level was a protective factor amidst an onslaught of COVID-19 rumors. At a time when information

regulation is crucial to resilience and well-being (Abdoli, 2020; Garfin et al., 2020), our findings provide a basis to manage the spread of rumors. In other words, it is not apparent veracity that makes a rumor 'have it'. Instead, COVID-19 rumors are shared even when disbelieved, but may be stemmed through higher education.

253

254

255

256

257 References 258 Abd-Alrazaq, A., Alhuwail, D., Househ, M., Hamdi, M., Shah, Z., 2020. Top Concerns 259 of Tweeters During the COVID-19 Pandemic: Infoveillance Study. J. Med. 260 Internet Res. 22, e19016. https://doi.org/10.2196/19016 261 Abdoli, A., 2020. Gossip, Rumors, and the COVID-19 Crisis. Disaster Med. Public 262 Health Prep. 1–2. https://doi.org/10.1017/dmp.2020.272 263 Aghababaeian, H., Hamdanieh, L., Ostadtaghizadeh, A., 2020. Alcohol intake in an 264 attempt to fight COVID-19: A medical myth in Iran. Alcohol 88, 29–32. 265 https://doi.org/10.1016/j.alcohol.2020.07.006 266 Ahmed, W., Vidal-Alaball, J., Downing, J., Sequí, F.L., 2020. COVID-19 and the 5G 267 Conspiracy Theory: Social Network Analysis of Twitter Data. J. Med. Internet 268 Res. 22, e19458. https://doi.org/10.2196/19458 269 Asokan, A., 2020. MOH debunks rumour about COVID-19 death in Singapore - CNA. 270 Channel News Asia. URL 271 https://www.channelnewsasia.com/news/singapore/covid19-coronavirus-272 death-singapore-rumour-moh-12481200 (accessed 9.30.20). 273 Banakar, M., Sadati, A.K., Zarei, L., Shahabi, S., Heydari, S.T., Lankarani, K.B., 2020. 274 Public sphere attitudes toward rumor sources on COVID-19 epidemics: 275 Evidence from community perceptions in Iran (preprint). In Review. 276 https://doi.org/10.21203/rs.3.rs-30297/v1 277 Chong, Y.Y., Cheng, H.Y., Chan, H.Y.L., Chien, W.T., Wong, S.Y.S., 2020. COVID-278 19 pandemic, infodemic and the role of eHealth literacy. Int. J. Nurs. Stud. 279 108, 103644. https://doi.org/10.1016/j.ijnurstu.2020.103644 280 Chua, A.Y.K., Banerjee, S., 2018. Intentions to trust and share online health rumors: 281 An experiment with medical professionals. Comput. Hum. Behav. 87, 1–9. 282 https://doi.org/10.1016/j.chb.2018.05.021

283 Chua, A.Y.K., Banerjee, S., 2017. To share or not to share: The role of epistemic 284 belief in online health rumors. Int. J. Med. Inf. 108, 36-41. 285 https://doi.org/10.1016/j.ijmedinf.2017.08.010 286 Cooke, J.E., Eirich, R., Racine, N., Madigan, S., 2020. Prevalence of posttraumatic 287 and general psychological stress during COVID-19: A rapid review and meta-288 analysis. Psychiatry Res. 292, 113347. 289 https://doi.org/10.1016/j.psychres.2020.113347 290 Cuan-Baltazar, J.Y., Muñoz-Perez, M.J., Robledo-Vega, C., Pérez-Zepeda, M.F., 291 Soto-Vega, E., 2020. Misinformation of COVID-19 on the internet: 292 Infodemiology study. JMIR Public Health Surveill. 6, e18444. 293 https://doi.org/10.2196/18444 294 da Silva, F.C.T., Neto, M.L.R., 2021, Psychological effects caused by the COVID-19 295 pandemic in health professionals: A systematic review with meta-analysis. 296 Prog. Neuropsychopharmacol. Biol. Psychiatry 104, 110062. 297 https://doi.org/10.1016/j.pnpbp.2020.110062 298 Depoux, A., Martin, S., Karafillakis, E., Preet, R., Wilder-Smith, A., Larson, H., 2020. 299 The pandemic of social media panic travels faster than the COVID-19 300 outbreak. J. Travel Med. 27. https://doi.org/10.1093/jtm/taaa031 301 Ebrahim, S.H., Ahmed, Q.A., Gozzer, E., Schlagenhauf, P., Memish, Z.A., 2020. 302 Covid-19 and community mitigation strategies in a pandemic. BMJ m1066. 303 https://doi.org/10.1136/bmj.m1066 304 Gao, J., Zheng, P., Jia, Y., Chen, H., Mao, Y., Chen, S., Wang, Y., Fu, H., Dai, J., 305 2020. Mental health problems and social media exposure during COVID-19 306 outbreak. PLoS ONE 15. https://doi.org/10.1371/journal.pone.0231924 307 Garfin, D.R., Silver, R.C., Holman, E.A., 2020. The novel coronavirus (COVID-2019) 308 outbreak: Amplification of public health consequences by media exposure. 309 Health Psychol. Off. J. Div. Health Psychol. Am. Psychol. Assoc. 39, 355–357. 310 https://doi.org/10.1037/hea0000875

311 Holmes, E.A., O'Connor, R.C., Perry, V.H., Tracey, I., Wessely, S., Arseneault, L., 312 Ballard, C., Christensen, H., Cohen Silver, R., Everall, I., Ford, T., John, A., 313 Kabir, T., King, K., Madan, I., Michie, S., Przybylski, A.K., Shafran, R., 314 Sweeney, A., Worthman, C.M., Yardley, L., Cowan, K., Cope, C., Hotopf, M., 315 Bullmore, E., 2020. Multidisciplinary research priorities for the COVID-19 316 pandemic: a call for action for mental health science. Lancet Psychiatry 7, 317 547–560. https://doi.org/10.1016/S2215-0366(20)30168-1 318 Hou, Z., Du, F., Zhou, X., Jiang, H., Martin, S., Larson, H., Lin, L., 2020. Cross-319 country comparison of public awareness, rumors, and behavioral responses 320 to the COVID-19 epidemic: Infodemiology study. J. Med. Internet Res. 22, 321 e21143. https://doi.org/10.2196/21143 322 Ippolito, G., Hui, D.S., Ntoumi, F., Maeurer, M., Zumla, A., 2020, Toning down the 323 2019-nCoV media hype—and restoring hope. Lancet Respir. Med. 8, 230-324 231. https://doi.org/10.1016/S2213-2600(20)30070-9 325 Islam, M.S., Sarkar, T., Khan, S.H., Mostofa Kamal, A.-H., Hasan, S.M.M., Kabir, A., 326 Yeasmin, D., Islam, M.A., Amin Chowdhury, K.I., Anwar, K.S., Chughtai, A.A., 327 Seale, H., 2020. COVID-19-Related Infodemic and Its Impact on Public 328 Health: A Global Social Media Analysis. Am. J. Trop. Med. Hyg. 329 https://doi.org/10.4269/ajtmh.20-0812 330 Jin, F., Wang, W., Zhao, L., Dougherty, E., Cao, Y., Lu, C.-T., Ramakrishnan, N., 331 2014. Misinformation propagation in the age of twitter. Computer 47, 90-94. 332 https://doi.org/10.1109/MC.2014.361 333 Larson, H.J., 2018. The biggest pandemic risk? Viral misinformation. Nature 562, 334 309-309. https://doi.org/10.1038/d41586-018-07034-4 335 Liu, J.C., Tong, E.M., 2020. COVID-19 news exposure as a modifiable risk factor of 336 psychological symptoms: Can an official WhatsApp channel help? (Preprint). 337 J. Med. Internet Res. 22, e22142. https://doi.org/10.2196/preprints.22142

338 Long, V.J., Liu, J.C., 2020. Behavioural changes during the COVID-19 pandemic: 339 Results of a national survey in Singapore. medRxiv 2020.08.06.20169870. 340 https://doi.org/10.1101/2020.08.06.20169870 341 Ministry of Communications and Information, 2020. Corrections and clarifications 342 regarding falsehoods published by States Times Review on availability of face 343 masks. gov.sq. URL https://www.gov.sq/article/factually-clarifications-on-344 falsehoods-posted-by-str-on-availability-of-face-masks (accessed 9.30.20). 345 Pappa, S., Ntella, V., Giannakas, T., Giannakoulis, V.G., Papoutsi, E., Katsaounou, 346 P., 2020, Prevalence of depression, anxiety, and insomnia among healthcare 347 workers during the COVID-19 pandemic: A systematic review and meta-348 analysis. Brain. Behav. Immun. 88, 901-907. 349 https://doi.org/10.1016/j.bbi.2020.05.026 350 Pew Research Center, 2018. News use across social media platforms 2018. Pew 351 Research Center. 352 Rovetta, A., Bhagavathula, A.S., 2020. COVID-19-related web search behaviors and 353 infodemic attitudes in Italy: Infodemiological study. JMIR Public Health 354 Surveill. 6, e19374. https://doi.org/10.2196/19374 355 Salari, N., Hosseinian-Far, A., Jalali, R., Vaisi-Raygani, A., Rasoulpoor, Shna, 356 Mohammadi, M., Rasoulpoor, Shabnam, Khaledi-Paveh, B., 2020. 357 Prevalence of stress, anxiety, depression among the general population 358 during the COVID-19 pandemic: a systematic review and meta-analysis. Glob. 359 Health 16, 57. https://doi.org/10.1186/s12992-020-00589-w 360 Saw, Y.E., Tan, E.Y.-Q., Liu, J.S., Liu, J.C., 2020. Towards a digital solution: 361 Predicting public take-up of Singapore's contact tracing application during the 362 COVID-19 crisis. medRxiv 2020.08.26.20182386. 363 https://doi.org/10.1101/2020.08.26.20182386

Singh, L., Bansal, S., Bode, L., Budak, C., Chi, G., Kawintiranon, K., Padden, C., 364 365 Vanarsdall, R., Vraga, E., Wang, Y., 2020. A first look at COVID-19 366 information and misinformation sharing on Twitter. ArXiv200313907 Cs. 367 Taylor, J., 2020. Bat soup, dodgy cures and "diseasology": the spread of coronavirus 368 misinformation. The Guardian. URL 369 https://www.theguardian.com/world/2020/jan/31/bat-soup-dodgy-cures-and-370 diseasology-the-spread-of-coronavirus-bunkum 371 Torales, J., O'Higgins, M., Castaldelli-Maia, J.M., Ventriglio, A., 2020. The outbreak 372 of COVID-19 coronavirus and its impact on global mental health. Int. J. Soc. 373 Psychiatry 66, 317–320. https://doi.org/10.1177/0020764020915212 374 Tran, T., Valecha, R., Rad, P., Rao, H.R., 2020. An Investigation of Misinformation 375 Harms Related to Social Media During Humanitarian Crises, in: Sahav, S.K., 376 Goel, N., Patil, V., Jadliwala, M. (Eds.), Secure Knowledge Management In 377 Artificial Intelligence Era, Communications in Computer and Information 378 Science. Springer, Singapore, pp. 167-181. https://doi.org/10.1007/978-981-379 15-3817-9 10 380 Vardanjani, H.M., Heydari, S.T., Dowran, B., Pasalar, M., 2020. A cross-sectional 381 study of Persian medicine and the COVID-19 pandemic in Iran: Rumors and 382 recommendations. Integr. Med. Res., Integrative Medicine for COVID-19: 383 Researches and Evidence 9, 100482. 384 https://doi.org/10.1016/j.imr.2020.100482 385 Vosoughi, S., Roy, D., Aral, S., 2018. The spread of true and false news online. 386 Science 359, 1146-1151. https://doi.org/10.1126/science.aap9559 387 Wang, Y., McKee, M., Torbica, A., Stuckler, D., 2019. Systematic Literature Review 388 on the Spread of Health-related Misinformation on Social Media. Soc. Sci. 389 Med. 240, 112552. https://doi.org/10.1016/j.socscimed.2019.112552

| 390 | Wong, J.E.L., Leo, Y.S., Tan, C.C., 2020. COVID-19 in Singapore—Current |
|-----|---|
| 391 | Experience: Critical Global Issues That Require Attention and Action. JAMA |
| 392 | 323, 1243. https://doi.org/10.1001/jama.2020.2467 |
| 393 | World Health Organisation, 2020. COVID-19 mythbusters. World Health Organ. URL |
| 394 | https://www.who.int/westernpacific/emergencies/covid- |
| 395 | 19/information/mythbusters |
| 396 | World Health Organization, 2020. Novel coronavirus (2019-nCoV): Situation report - |
| 397 | 13. World Health Organization, Geneva, Switzerland. |
| 398 | Xiong, J., Lipsitz, O., Nasri, F., Lui, L.M.W., Gill, H., Phan, L., Chen-Li, D., Iacobucci, |
| 399 | M., Ho, R., Majeed, A., McIntyre, R.S., 2020. Impact of COVID-19 pandemic |
| 400 | on mental health in the general population: A systematic review. J. Affect. |
| 401 | Disord. 277, 55-64. https://doi.org/10.1016/j.jad.2020.08.001 |
| 402 | Zarocostas, J., 2020. How to fight an infodemic. The Lancet 395, 676. |
| 403 | https://doi.org/10.1016/S0140-6736(20)30461-X |
| | |

Table 1. Baseline demographics of participants

| Characteristic | N | (%) | | | |
|--------------------------------------|------|--------|--|--|--|
| Age (Mean = 39.3, SD = 12.7) | | | | | |
| Gender | | | | | |
| Female | 791 | (63.9) | | | |
| Male | 445 | (36.0) | | | |
| Did not answer | 1 | (0.1) | | | |
| Ethnicity | | | | | |
| Chinese | 1047 | (84.6) | | | |
| Indian | 55 | (4.4) | | | |
| Malay | 71 | (5.7) | | | |
| Others | 63 | (5.1) | | | |
| Did not answer | 1 | (0.1) | | | |
| Religion | | | | | |
| Buddhism | 181 | (14.6) | | | |
| Taoism/ Chinese traditional beliefs | 51 | (4.1) | | | |
| Islam | 71 | (5.7) | | | |
| Hinduism | 41 | (3.3) | | | |
| Roman Catholicism | 125 | (10.1) | | | |
| Christianity (Protestant) | 405 | (32.7) | | | |
| No religion | 331 | (26.7) | | | |
| Others | 30 | (2.4) | | | |
| Did not answer | 2 | (0.2) | | | |
| Married status | | | | | |
| Single | 518 | (41.9) | | | |
| Married | 667 | (53.9) | | | |
| Widowed/ separated/ divorced | 47 | (3.8) | | | |
| Did not answer | 5 | (0.4) | | | |
| Educational level | | | | | |
| Primary school | 4 | (0.3) | | | |
| Secondary school | 65 | (5.3) | | | |
| Junior college | 91 | (7.4) | | | |
| Vocational training | 22 | (1.8) | | | |
| Polytechnic/ diploma | 154 | (12.4) | | | |
| University (undergraduate) | 623 | (50.3) | | | |
| University (postgraduate) | 252 | (20.4) | | | |
| Did not answer | 26 | (2.1) | | | |
| House type | | | | | |
| HDB flat: 1-2 rooms | 12 | (1.0) | | | |
| HDB flat: 3 rooms | 89 | (7.2) | | | |
| HDB flat: 4 rooms | 305 | (24.6) | | | |
| HDB flat: 5 rooms or executive flats | 358 | (28.9) | | | |
| Condominium or private apartments | 321 | (25.9) | | | |
| Landed property | 131 | (10.6) | | | |
| Did not answer | 21 | (1.7) | | | |
| | | ` ' | | | |

| Househ | old size 1 | 55 | (4.4) |
|---------|----------------|------|--------|
| | 2 | 178 | (14.4) |
| | 3 | 279 | (22.6) |
| | 4 | 359 | (29.0) |
| | 5+ | 364 | (29.4) |
| | Did not answer | 2 | (0.2) |
| Country | of birth | | |
| - | Singapore | 1004 | (81.1) |
| | Other | 232 | (18.8) |
| | Did not answer | 1 | (0.1) |

Table 2. Predicting the number of rumors, heard, shared, and believed during the COVID-19 outbreak

| | Outcome Measure ^a | | | | | |
|--|---|---|--|---|---|---|
| | Model 1: Number | | | Number | | Number |
| | of rumo | rs heard | of rumors shared | | of rumors believed | |
| Age | 0.003 | (0.004) | 0.005 | (0.002) | 0.002 | (0.002) |
| Gender (base = Female) Male | -0.084 | (0.080) | -0.025 | (0.038) | 0.017 | (0.036) |
| Ethnicity (base = Chinese) Indian Malay Others | -0.456 -0.537 -0.234 | (0.353) (0.279) (0.206) | -0.060 0.349 0.022 | (0.169) (0.133) (0.098) | 0.152 0.184 -0.059 | (0.159) (0.125) (0.092) |
| Religion (base = No religion) Christianity (Protestant) Buddhism Roman Catholicism Taoism/ Chinese traditional beliefs Islam | 0.093 0.159 0.115 0.171 0.330 | (0.101) (0.125) (0.144) (0.207) (0.325) | -0.018 0.115 0.036 0.018 0.006 | (0.048) (0.060) (0.069) (0.099) (0.155) | 0.012 -0.025 0.135 0.078 -0.104 | (0.045) (0.056) (0.065) (0.093) (0.146) |
| Hinduism Others | 0.215 0.560 | (0.351) (0.281) | -0.259 0.047 | (0.167) (0.134) | 0.122 0.072 | (0.157) (0.126) |
| Marital status (base = Single) Married Widowed/ separated/ divorced | -0.041 0.002 | (0.093) (0.217) | -0.021 0.037 | (0.045) (0.104) | -0.036 -0.170 | (0.042) (0.098) |
| Education level | 0.077 | (0.029) | -0.046 | (0.014)* | -0.046 | (0.013)* |
| House type | 0.092 | (0.036) | -0.030 | (0.017) | -0.042 | (0.016) |
| Household size | -0.018 | (0.036) | 0.011 | (0.017) | 0.013 | (0.016) |
| Country of birth (base = Singapore) Other | 0.271 | (0.108) | 0.071 | (0.051) | 0.072 | (0.048) |
| Lockdown (base = Lockdown period) Before lockdown After lockdown | -0.437 1.129 | (0.206) (0.263)* | -0.017 0.011 | (0.098) (0.126) | 0.075 -0.151 | (0.092) (0.118) |
| Number of local COVID-19 cases (log transformed) | -0.621 | (0.173)* | -0.052 | (0.083) | 0.079 | (0.078) |
| π | 0.474 | | 0.435 | | 0.388 | |

^aData reported as beta estimates (standard error)

^{*}Indicates significance at p < .002 (following Bonferroni corrections).

Figures and captions




Figure 1: Proportion of participants hearing, sharing, and believe each COVID-19 rumor (that the virus originated from the consumption of bat soup, from an American lab, or from a Chinese lab; or that the virus can be cured by eating garlic or drinking water). Vertical lines represent the 95% confidence interval.




Figure 2: For each rumor, the vertical bar depicts the quantity of participants who shared each rumor, represented as a percentage of participants who believed or disbelieved each rumor. Vertical lines represent the 95% confidence interval.




Figure 3: Sources of where participants heard COVID-19 rumors. Horizontal lines represent 95% confidence interval.