

1 Title: Organophosphate and organohalogen flame-retardant exposure and thyroid  
2 hormone disruption in a cohort of female firefighters and office workers from San  
3 Francisco.

4  
5 Jessica Trowbridge<sup>1,2</sup>, Roy Gerona<sup>3</sup>, Michael McMaster<sup>4,5</sup>, Catherine Ona<sup>5</sup>, Cassidy  
6 Clarity<sup>1,2</sup>, Vincent Bessonneau<sup>6</sup>, Ruthann Rudel<sup>6</sup>, Heather Buren<sup>7</sup>, Rachel Morello-  
7 Frosch<sup>1,2</sup>

8  
9 <sup>1</sup>Department of Environmental Science, Policy and Management, University of California,  
10 Berkeley, Berkeley, CA

11 <sup>2</sup>School of Public Health, University of California, Berkeley, Berkeley, CA

12 <sup>3</sup>Clinical Toxicology and Environmental Biomonitoring Lab, Department of Obstetrics,  
13 Gynecology and Reproductive Sciences, University of California, San Francisco, San  
14 Francisco, CA

15 <sup>4</sup>Department of Cell and Tissue Biology, University of California, San Francisco, San  
16 Francisco, CA

17 <sup>5</sup>Center for Reproductive Sciences, Department of Obstetrics, Gynecology and  
18 Reproductive Sciences, University of California, San Francisco, San Francisco, CA

19 <sup>6</sup>Silent Spring Institute, Newton, MA

20 <sup>7</sup>United Fire Service Women, San Francisco, CA

21

---

22 \*Corresponding author:

23 Rachel Morello-Frosch

24 [rmf@berkeley.edu](mailto:rmf@berkeley.edu)

25 510-643-6358

26 Department of Environmental Science, Policy and Management

27 130 Mulford, Hall #3144, University of California, Berkeley

28 Berkeley CA, 94720

29

30

31 Acknowledgements: The authors thank all of the WFBC study participants for their  
32 contribution to the study. This work is supported by the California Breast Cancer  
33 Research Program #19BB-2900 and #23BB-1700 & 1701 & 1702 (JT, RG, CC, MM,  
34 CA, RAR, HB, VB, RMF), the National Institute of Environmental Health Sciences  
35 R01ES027051 (RMF), the National Institute for Occupational Safety and Health,  
36 Targeted Research Training Program T42 OH008429 (JT), the San Francisco  
37 Firefighter Cancer Prevention Foundation (HB) and the International Association of  
38 Firefighters-Local 798. We thank Anthony Stefani, Emily O'Rourke, Nancy Carmona,  
39 Karen Kerr, Julie Mau, Natasha Parks, Lisa Holdcroft, San Francisco Fire Chief Jeanine  
40 Nicholson, former San Francisco Fire Chief Joanne Hayes-White, Sharyle Patton,  
41 Connie Engel and Nancy Buermeyer for their contributions to the study.  
42 RAR and VB, are employed at the Silent Spring Institute, a scientific research  
43 organization dedicated to studying environmental factors in women's health. The  
44 Institute is a 501(c)3 public charity funded by federal grants and contracts, foundation  
45 grants, and private donations, including from breast cancer organizations. HB is former  
46 president and member of United Fire Service Women, a 501(c)3 public charity  
47 dedicated to supporting the welfare of women in the San Francisco Fire Department.  
48 The authors declare they have no actual or potential competing financial interests.

49 **Abstract**

50 **Background:** Occupational exposures to flame retardants (FR), which are suspected  
51 endocrine disrupting compounds, may be of particular concern for firefighters as they  
52 are commonly found in consumer products and have been detected in fire station dust  
53 and firefighter gear.

54 **Objectives:** The Women Workers Biomonitoring Collaborative is a community based  
55 participatory research study that sought to measure environmental chemicals relevant  
56 to firefighting and evaluate their effects on thyroid hormone levels.

57 **Methods:** We measured 10 FR or their metabolites in urine of female firefighters and  
58 office workers from San Francisco: bis(1,3-dichloro-2-propyl) phosphate (BDCPP),  
59 bis(2-chloroethyl) phosphate (BCEP), dibutyl phosphate (DBuP), dibenzyl phosphate  
60 (DBzP), di-p-cresyl phosphate (DpCP), di-o-cresyl phosphate (DoCP), 2,3,4,5-  
61 tetrabromobenzoic acid (TBBA), tetrabromobisphenol-A (TBBPA), 5-OH-BDE 47, and 5-  
62 OH-BDE 100. We assessed potential predictors of exposure levels and the association  
63 between FR exposures and thyroxine ( $T_4$ ) and thyroid stimulating hormone (TSH).

64 **Results:** BDCPP, BCEP, and DBuP were the most commonly detected FRs, among all  
65 study participants, with intermediate BMI and college educated women having the  
66 highest levels, and Black women having higher BDCPP levels than White women.  
67 Firefighters had higher detection frequencies (DF) and exposure levels compared to  
68 office workers; median BDCPP levels were five times higher in firefighters than in office  
69 workers. Among firefighters, occupational activities were not significantly associated  
70 with FR levels, although position (i.e. officer and firefighter versus driver), being on-duty  
71 (versus off-duty) and assigned to the airport suggested a positive association with FR

72 levels. Among firefighters, a doubling of BDCPP was associated with a 2.88% decrease  
73 (95%CI -5.28,-0.42) in T<sub>4</sub>. We did not observe significant associations between FR and  
74 T<sub>4</sub> among office workers.

75 **Discussion:** Firefighters had significantly higher exposures to FR compared to office  
76 workers, and we observed a negative association between BDCPP and thyroxine in  
77 firefighters. Future research should elucidate occupational sources of FR exposure and  
78 opportunities for exposure reduction.

79

## 80 **Introduction**

81 Studies show that firefighters have elevated risk for many cancers. For instance,  
82 a meta-analysis of 32 studies identified increased rates of lymphoma, and testicular and  
83 prostate cancer among male firefighters (LeMasters et al. 2006). A study of over 19,000  
84 male firefighters in a pooled cohort from San Francisco, Chicago and Philadelphia found  
85 that the occupation of firefighting was associated with elevated lung cancer incidence  
86 and mortality, and leukemia mortality (Daniels et al. 2015). Breast cancer incidence was  
87 non-significantly elevated among men and among the 991 women included in the study  
88 and in both groups the increases were largest among those at younger ages (<65 years  
89 old for men and ages 50-55 for women (Daniels et al. 2014)). In a published update  
90 adding 7 years of follow up to the pooled cohort, Pinkerton et al. (2020) found that  
91 women firefighters had elevated mortality rates for bladder cancer and non-statistically  
92 elevated mortality from Non-Hodgkin's lymphoma, multiple myeloma, lung cancer and  
93 breast cancer (Pinkerton et al. 2020). A study of Florida firefighters that included 5,000  
94 women found elevated incidence of cervical and thyroid cancer and Hodgkin's disease

95 compared to the general Florida population (Ma et al. 2006) and a more recent study of  
96 Florida firefighters using that state's cancer registry found that women firefighters had  
97 elevated risk of melanoma, thyroid, and brain cancer (Lee et al. 2020).

98 In addition to limited studies of cancer risk among women firefighters, few studies  
99 have assessed their exposures to occupational hazards, including environmental  
100 chemicals, although occupational exposures have been well documented among male  
101 firefighters. Common exposures measured in firefighters and fire stations include  
102 polyaromatic hydrocarbons (PAH), formaldehyde, benzene, dioxins, diesel, per- and  
103 poly-fluoroalkyl substances (PFAS), and flame retardants including organohalogen  
104 flame retardants, like polybrominated diphenyl ethers (PBDE), and organophosphate  
105 flame retardants (OPFR) (Caux et al. 2002; Dobraca et al. 2015; Fent et al. 2014;  
106 Grashow et al. 2020; Jin et al. 2011; Laitinen et al. 2014; Park et al. 2015; Shaw et al.  
107 2013; Shen et al. 2015, 2018; Trowbridge et al. 2020). Many of these chemicals have  
108 demonstrated their potential for breast tumor development in animal and human studies  
109 (Rodgers et al. 2018; Rudel et al. 2011, 2014) and their capacity as endocrine  
110 disrupting compounds (EDC) (Gore et al. 2015; Rudel et al. 2011).

111 Flame retardants are of particular interest for firefighters because they have been  
112 found in firefighting gear (Alexander and Baxter 2016) and in fire station dust (Shen et  
113 al. 2015, 2018). Additionally, firefighter biomonitoring and studies of fire station dust  
114 samples have found elevated levels of brominated flame retardants compared to homes  
115 and offices (Park et al. 2015; Shen et al. 2015). One study found that dust collected  
116 from 26 fire stations across the US had elevated levels of tris(1,3-dichloro-isopropyl)-

117 phosphate (TDCPP), an OPFR flame retardant, at levels higher than those measured in  
118 homes and offices (Shen et al. 2018).

119 California's furniture flammability standard known as TB117 was in effect  
120 between 1977 and 2013 and required furniture filling such as foam to withstand an open  
121 flame test without igniting. In order to meet the standard, manufacturers typically added  
122 chemical flame retardants to filling materials such as flexible polyurethane foam (PUF).  
123 Additionally, furniture manufacturers often complied with the California standard for their  
124 products sold nationwide, impacting everyone in the U.S. (Castorina et al. 2017;  
125 Dodson et al. 2014; Stapleton et al. 2012). Because flame retardants are added to  
126 products post-production and are not bound to fabric and foams, they can leach out of  
127 materials and contaminate air and dust (Dodson et al. 2012; Stapleton et al. 2012).  
128 California's revised furniture flammability standard TB117-2013 removed the open flame  
129 test in favor of a smolder test designed to mimic a lit cigarette on furniture fabric, and  
130 manufacturers have been able to meet this standard without adding chemical flame  
131 retardants in PUF (Bureau of Electronic and Appliance 2018). At the same time,  
132 concerns about toxicity and bioaccumulation of PBDE flame retardants in the early  
133 2000s led to an increase in the use of other flame retardants such as OPFRs in both  
134 furniture and fabrics (van der Veen and de Boer 2012a). Although use of flame  
135 retardants in furniture has declined, many are persistent and bioaccumulative, and both  
136 legacy and replacement FR remain in durable consumer products, such as furniture,  
137 and exposures remain a concern (Dodson et al. 2012; Zota et al. 2013). OPFRs are  
138 commonly used as flame retardants in fabric, electronics, PUF, as plasticizers, and  
139 engine lubricants (Covaci et al. 2011; Van den Eede et al. 2012; van der Veen and de

140 Boer 2012a). Due to OPFR use in fabrics and that PBDEs have been found in firefighter  
141 gear (Alexander and Baxter 2016), OPFRs may conceivably be used in firefighter  
142 clothing and protective gear. Organophosphate and brominated flame retardants may  
143 be associated with cancer (Lerro et al. 2015; Rudel et al. 2014) and have also been  
144 identified as endocrine disruptors that are associated with altered thyroid hormone (TH)  
145 levels in both *in vitro* and *in vivo* studies (**Table 1**) (Dishaw et al. 2014; Farhat et al.  
146 2013; Hill et al. 2018; Meeker and Stapleton 2010; Wang et al. 2013b).

**Table 1.** Select flame retardant compounds, their metabolites, potential sources and uses and evidence of toxicity

Parent compound	Metabolite	Abbr.	Potential sources	Prop 65 <sup>a</sup> list	Evidence of TH effects <sup>b</sup>
tris(2-chloroethyl) phosphate (TCEP)	bis(2-chloroethyl) phosphate	BCEP	PUF, plastics, polyester resin and textiles (Van den Eede et al. 2012; van der Veen and de Boer 2012b)		
tris(1-chloro-2-propyl) phosphate (TCIPP)	bis(1-chloro-2-propyl) phosphate	BCPP	Polyurethane foams (Van den Eede et al. 2012)		•
tris(1,3-dichloro-2-propyl) phosphate (TDCIPP)	bis(1,3-dichloro-2-propyl) phosphate	BDCPP	Plastics polyurethane foam and textiles (Van den Eede et al. 2012; van der Veen and de Boer 2012b)	•	•
tri-n-butyl phosphate	di-n-butyl phosphate	DBuP	Lubricants and greases, Paints and coatings (Schindler et al. 2013)		•
2-ethylhexyl-2,3,4,5-tetrabromobenzoate	2,3,4,5-tetrabromobenzoic acid	TBBA	Component of Firemaster® 550 (Hoffman et al. 2014) Plastics, water barriers, kitchen hoods and electronics (Covaci et al. 2011)		•
tetrabromobisphenol-A	NA	TBBPA	Reactive in circuit boards, additive in polymers, among most widely used FR (Birnbaum and Staskal 2004)	•	•
tri-o-cresyl phosphate	Di-o-cresyl phosphate	DoCP	Plasticizer and flame retardant (van der Veen and de Boer 2012b)		•
tri-p-cresyl phosphate	Di-p_cresyl phosphate	DpCP	Plasticizer and flame retardant (van der Veen and de Boer 2012a)		
tri-benzyl phosphate	Dibenzyl phosphate	DBzP	---		

<sup>a</sup>prop 65 <https://oehha.ca.gov/proposition-65/about-proposition-65>; <sup>b</sup>(Boas et al. 2012; Calsolaro et al. 2017; Dishaw et al. 2014)



147 **Thyroid hormone disruption.**

148           Studies suggest that TH levels, including thyroid stimulating hormone (TSH) and  
149 thyroxine (T<sub>4</sub>), are affected by exposure to environmental chemicals including OPFRs  
150 (Diamanti-Kandarakis et al. 2009; Dishaw et al. 2014; Gore et al. 2015; Rudel et al.  
151 2014). TH production is controlled via a negative feedback loop: as TH levels decline in  
152 the body, the pituitary gland secretes TSH, which induces an increase in production of  
153 T<sub>4</sub> by the thyroid gland. TSH secretion is suppressed when T<sub>4</sub> levels reach a “set point”  
154 that varies by individual (Zoeller et al. 2007).

155           Studies of zebrafish and chicken embryos show that OPFRs can disrupt thyroid  
156 hormone homeostasis and decrease total T<sub>4</sub> concentrations (Farhat et al. 2013; Wang  
157 et al. 2013a). Human studies suggest impacts of exposure to OPFRs on thyroid  
158 hormone function, although research remains limited and the direction of the  
159 association is inconclusive. Meeker and Stapleton found decreased T<sub>4</sub> levels with  
160 higher TDCPP concentrations in house dust among men (Meeker and Stapleton 2010).  
161 A study of e-waste recycling workers found that a two-fold increase in exposure to tert-  
162 butyl diphenyl phosphate was associated with lower total T<sub>4</sub> levels in men (Gravel et al.  
163 2020); and Preston et al. (2017) found that high levels of diphenyl phosphate, a  
164 metabolite of triphenyl phosphate, were associated with an increase in total T<sub>4</sub>  
165 concentration in women (Preston et al. 2017). Thyroid hormones are important for fetal  
166 neurodevelopment and for TH-mediated gene expression. Additionally, TH may be  
167 relevant to downstream adverse health impacts such as cardiovascular disease and  
168 cancer (Gore et al. 2015; Krashin et al. 2019; Rodondi et al. 2006).

169 While biomonitoring studies show that firefighters have higher body burdens of  
170 flame retardant chemicals than the general United States population (Alexander and  
171 Baxter 2016; Brown et al. 2014; Park et al. 2015), very little research has investigated  
172 the extent and health implications of exposure among women firefighters due to the  
173 limited number of women in most fire departments. Assessing the potential health risks  
174 of flame-retardant exposures among women firefighters, including for outcomes such as  
175 breast cancer poses methodological challenges due to the low numbers of women in  
176 the fire service and the long latency period from exposure to onset of disease.  
177 Assessing thyroid hormone disruption associated with flame retardant exposures  
178 enables identification of early biological perturbations of potential relevance to thyroid  
179 dysfunction or disease (Ward et al. 2010), and long-term adverse health outcomes such  
180 as cancer (Krashin et al. 2019).

181 San Francisco has one of the largest forces of women firefighters among large  
182 urban fire departments in the U.S.—approximately 15% of firefighters in SFFD are  
183 women. As firefighting and other first responder professions continue to diversify and  
184 increase the number of women in their ranks, it is important to understand occupational  
185 exposures and potential health implications for women firefighters. The Women  
186 Workers Biomonitoring Collaborative (WWBC) is a community-based participatory  
187 research study that aims to characterize occupational exposure to potential breast  
188 carcinogens among women workers. This study sought to characterize exposures to  
189 OPFRs and replacement organohalogen flame retardants (henceforth referred to  
190 together as flame retardants (FR)) among women firefighters and office workers in San  
191 Francisco and to assess exposure to FR and their effect on total T<sub>4</sub> and TSH.

192 **Methods:**

193 Study design and participant recruitment protocols have been described  
194 elsewhere (Grashow et al. 2020; Trowbridge et al. 2020). Briefly, participant  
195 recruitment, interviews and sample collections took place between 2014 and 2015.  
196 Participants were employees of the City and County of San Francisco or the San  
197 Francisco Fire Department (SFFD). Firefighter collaborators and researchers actively  
198 recruited study participants through the Fire Department, as well as firefighter advocacy  
199 organizations including the San Francisco Firefighters Cancer Prevention Foundation,  
200 United Fire Service Women, and the International Association of Firefighters Union  
201 Local 798. Office workers, who are non-first responder employees of the City and  
202 County of San Francisco, were recruited by emails to employees city-wide, tabling, and  
203 presentations by research staff and firefighter collaborators. Study participants were  
204 eligible to participate if they were 18 years or older and non-smokers. Firefighters were  
205 required to have worked in SFFD for a minimum of 5 years and be on “active duty” (i.e.  
206 currently assigned to a fire station at the time of recruitment). Participants were  
207 consented into the study following protocols approved by the Institutional Review Board  
208 of the University of California, Berkeley (# 2013-07-5512).

209 **Data collection**

210 All participants completed an hour-long in-person exposure assessment interview  
211 with research staff. This interview collected demographic and basic health information  
212 including body mass index (BMI) and the use of hormone replacement medications. We  
213 also asked about possible sources of FR exposure including consumer product use,  
214 diet, and occupational activities. A subset of participants (N = 66) gave researchers

215 permission to access their departmental firefighting history records, from which we  
216 abstracted the number of fires fought in the 7 days and month prior to the sample  
217 collection date.

218 We collected most biospecimen samples, including blood and urine, between  
219 8AM and 11AM. A trained phlebotomist collected blood in EDTA-treated lavender top  
220 tubes. Urine specimens were collected by participants in 60mL polypropylene  
221 biospecimen cups. Prior to sample collection, we gave participants a biospecimen cup  
222 and instructions for collecting a morning void sample (first urine sample after a night's  
223 sleep) which they brought with them to their sample collection appointment. Research  
224 staff then asked for a second, spot urine, sample at the time of the blood collection.  
225 Biospecimens were put on ice and transferred to the lab at the University of California,  
226 San Francisco where research staff processed samples within 3 hours of their  
227 collection. Blood collection tubes were spun at 3000 rpm for 10 minutes and plasma  
228 was aliquoted into 1.1 mL cryovial tubes. Urine samples were aliquoted into 3.5 mL  
229 cryovial tubes. All samples were stored at  $-80^{\circ}\text{C}$  until analysis.

### 230 **Laboratory analysis:**

#### 231 *FR analysis*

232 We sought to quantify metabolites in urine of six OPFR chemicals: bis(1,3-  
233 dichloro-2-propyl) phosphate (BDCPP), bis(2-chloroethyl) phosphate (BCEP), dibutyl  
234 phosphate (DbuP), dibenzyl phosphate (DBzP), di-p-cresyl phosphate (DpCP), di-o-  
235 cresyl phosphate (DoCP), and 4 brominated flame retardants: 2,3,4,5-  
236 tetrabromobenzoic acid (TBBA), tetrabromobisphenol a (TBBPA), 5-OH-BDE 47, and 5-  
237 OH-BDE 100.

238 Quantification of the 10 analytes was performed using liquid-chromatography-  
239 tandem mass spectrometry (LC-MS/MS) on an Agilent LC 1260 (Agilent Technologies,  
240 Sta. Clara, CA)- AB Sciex 5500 system (Sciex, Redwood City, CA). Freshly thawed  
241 urine specimens (1 mL) were deconjugated prior to LC-MS/MS analysis by addition of  
242 450 U *H. pomatia* glucuronidase (Sigma-Aldrich, St Louis, MO) and incubated at 37 °C  
243 for two hours with constant shaking. Deconjugated urine samples were then prepared  
244 for LC-MS/MS analysis by solid phase extraction (SPE) using Waters Oasis WAX  
245 cartridges (10 mg, 30 µm, 1 cc). An Agilent ZORBAX Eclipse XDB-C8 column (2.1x100  
246 mm, 3.5µm) maintained at 50°C was used in reversed-phase chromatography. The  
247 analytes were separated by gradient elution using water with 20 mM ammonium acetate  
248 as mobile phase A (MPA) and acetonitrile as mobile phase B (MPB). The gradient used  
249 for analyte separation consisted of 5% MPB at 0–0.5 min, gradient to 75% MPB from  
250 0.5 to 7.5 min, gradient to 100% MPB from 7.5-9 min, 100% MPB at 9-11 min, and 5%  
251 MPB at 11.1–15 min. The analytes were ionized in the negative mode using  
252 electrospray ionization (ESI) and mass scanning was performed by multiple reaction  
253 monitoring. Each analyte was monitored using two transitions and retention time.  
254 Quantitation of each analyte was performed by isotope dilution method with their  
255 deuterated or C-13 isotopologues as internal standards.

256 Each batch of samples was injected in duplicate. Procedural quality control  
257 materials and procedural blanks were run along with the calibration curve at the start,  
258 middle, and end of each run. Two QC materials were used at low and high  
259 concentrations. To accept the results of a batch run, QC materials measurements must  
260 be within 20% of their target values and the precision of their measurements have  $\leq 20\%$

261 CV (coefficient of variation). Analyte identification from total ion chromatograms was  
262 evaluated using AB Sciex Analyst v2.1 software while quantification of each analyte was  
263 processed using AB Sciex MultiQuant v2.02 software. Analysts were blinded to  
264 firefighter and office worker status of the urine samples during the analysis.

265

#### 266 *T<sub>4</sub> and TSH measurement*

267 The TSH and T<sub>4</sub> levels were measured in blood plasma using ELISA (Antibodies-  
268 online, cat. No. ABIN2773773) following manufacturer's protocol (Antibodies-online  
269 2020). Room temperature calibrators, controls and samples (25 µl for T<sub>4</sub> and 50µl for  
270 TSH) were loaded onto streptavidin coated wells followed by the addition of biotinylated  
271 antibody for T<sub>4</sub> or TSH. Standard curves were constructed in duplicate with calibrators  
272 supplied in the kit. The reaction was incubated at room temperature for 1 hr. and then  
273 washed three times. Substrate solution was added and stopped after 15 minutes.  
274 Absorbance at 450 nm was read immediately using a microplate reader. The T<sub>4</sub> and  
275 TSH concentrations of each sample, run in duplicate, were obtained from the standard  
276 curve. According to the manufacturer's instructions, samples below the LOD were rerun  
277 with a 30 min development time.

#### 278 **Statistical analysis:**

279 The goal of this analysis was twofold: First to characterize exposure to FR  
280 chemicals and identify potential predictors of exposure among firefighters and office  
281 workers. Second, to assess the relationship between FR exposures and T<sub>4</sub> and TSH.

282 We tested the distributions for FRs, T<sub>4</sub> and TSH visually inspecting the  
283 distributions and with the Shapiro-Wilks test. Because of the evidence for skewed  
284 distributions, we used nonparametric approaches to test differences between groups  
285 (permutation, Wilcoxon) and we natural log-transformed values which improved  
286 normality for use in linear models. All regression models were adjusted for log-  
287 transformed creatinine to account for urine dilution when quantifying the FR (Barr et al.  
288 2005). We used two different linear regression models based on whether FR levels  
289 were being evaluated as the outcome or exposure and to account for FR levels below  
290 the limit of detection (LOD). In bivariate analyses looking at the relationship between  
291 covariates (e.g. food consumption or position in the fire department) and FR levels as  
292 the outcome, we used the maximum likelihood estimation (MLE) model from the NADA  
293 package in R, which accounts for levels below the LOD without the need for  
294 substitution, when the chemical measurement is the outcome (Helsel 2005). To  
295 assessed the relationship between FR levels (exposure) and thyroid hormone  
296 (outcome), we used ordinary least squares (OLS) regression models and  
297 operationalized levels below the LOD in the following ways: We included all LC-MS/MS  
298 reported values (even if those values reported were below the LOD) and substituted  
299  $LOD/\sqrt{2}$  for any remaining non-detect values.

300 First, we treated FR as the outcome to assess differences between firefighters  
301 and office workers and evaluate predictors of FR levels. We calculated summary  
302 statistics for FRs including the geometric mean (GM), Geometric Standard Deviation  
303 (GSD), and distribution percentiles for each group. We plotted FR levels from WFBC  
304 firefighters and office workers along with FR levels from women ages 18-65 from the

305 National Health and Nutrition Examination Survey (NHANES) (2013-2014) to see how  
306 WFBC participants compared with a nationally representative sample of the U.S.  
307 population. In the full cohort we assessed the impact of variables collected from the  
308 exposure assessment interview on FR levels as the outcome using MLE regression. We  
309 limited analyses to FRs with a detection frequency (DF) of 70% or higher in at least one  
310 group (firefighters or office workers) and applied separate MLE regression models on  
311 each FR chemical (continuous outcome) controlling for occupation and log(creatinine).  
312 We analyzed the relationship of FR concentrations and the following variables: age,  
313 race/ethnicity, body mass index (BMI), and educational attainment and we assessed the  
314 association of eating certain foods and packaged foods based on prior literature  
315 suggesting an association with FR exposures (Kim et al. 2020).

316 We then limited the analysis to firefighters to explore specific occupational  
317 activities that might be associated with FR levels such as the participant's assigned  
318 position in the fire department (i.e. firefighter, officer, or driver), the frequency of using a  
319 self-contained breathing apparatus (SCBA) during fire suppression, salvage and  
320 overhaul, the frequency of showering or washing up after a fire event, and the number  
321 of fires fought in the week and month prior to the sample collection. We applied MLE  
322 regression and exponentiated the beta coefficients and 95% CI to find the proportional  
323 change in geometric mean for each unit increase or category change versus referent.

324 Next, we assessed the impact of FR (exposure) on TH (outcome) using OLS  
325 regression and substitution for FR values below the LOD. We ran separate models for  
326 each FR chemical and each TH outcome, TSH and T<sub>4</sub>. We considered variables to  
327 adjust for in our models if they demonstrated a statistically significant association (p-



328 value < 0.05) with the exposure (at least one FR) and the outcome (TSH or T<sub>4</sub>) in our  
329 data or if previous literature identified an association. Although age was not associated  
330 with FR in our data, it is associated with TH levels in other studies (Hollowell et al. 2002)  
331 and therefore it was included as a covariate in regression models. We did not control for  
332 body-mass index (BMI) in regression models because literature suggests that both FRs  
333 and thyroid hormone disruption can lead to increased BMI (Boyle et al. 2019; Knudsen  
334 et al. 2005), implying that it may be a collider for which adjustment could induce a  
335 spurious association. Final models were adjusted for age and log(creatinine). Due to  
336 large differences in DF between firefighters and office workers, we stratified the analysis  
337 by occupation. We used the continuous exposure when the DF ≥70% and we  
338 categorized FR values when DF < 70%. FR was categorized into the following groups:  
339 <LOD and ≥LOD (for FR with DF of 25% to 50%) and <LOD, LOD to median, and  
340 >median (for FR with DF between 50% up to 70%). Compounds with DF below 25%  
341 were excluded from the multivariate analysis.

342 Results from the OLS regression models with continuous exposure and outcome  
343 were converted to the percent change in the outcome for a twofold increase in FR  
344 exposure with the formula:  $(2^\beta - 1) * 100$ . From OLS regression models with  
345 categorical FR exposure and continuous outcomes we calculated the percent change of  
346 the outcome for each category compared to the referent (< LOD) with the formula:  
347  $(e^\beta - 1) * 100$ .

348 Analyses were conducted using R version 3.6.1 and R-studio version 1.2.1335  
349 (R Core Team 2015; RStudio Team 2016).


## 350 **Results**

351 We had chemical measurements for 170 participants and TH measurements for  
352 168 participants. This difference in sample size was due to limited biospecimen samples  
353 available to measure TH. We retained the full cohort when characterizing FR levels  
354 among firefighters and office workers and assessing the relationship between FR and  
355 covariates. When we assessed the relationship between FR levels and TH, we  
356 excluded the participants without TH measurements and three participants who  
357 reported taking TH replacement medications. The final number analyzed for thyroid  
358 hormone disruption was N = 165 (84 firefighters and 81 office workers).

### 359 *Flame retardant levels and predictors of exposure*

360 BDCPP had the highest DF and was found in all firefighters (100%) and almost  
361 all office workers (90%). BCEP and DBuP were detected in most firefighters (DF>70%)  
362 while among office workers the DF was below 50%. TBBPA was detected in 45% of  
363 firefighters and 42% percent of office workers while DpCP was detected in 40% of  
364 firefighters and 17% of office workers. TBBA, DoCP and DBzP was found in fewer than  
365 30% of firefighters and office workers. OH-BDE and OH-BDE47 were not detected  
366 above the LOD in any of our participants' urine samples. We plotted the distribution  
367 (median, interquartile range (IQR) and 95<sup>th</sup>) of each FR detected in our cohort (**Figure**  
368 **1**; GM (GSD) and percentiles listed in **Table S1**). Firefighters had both higher detection  
369 frequencies and higher average levels of FR chemicals compared to office workers with  
370 the largest median differences observed for DBuP, BDCPP and BCEP compared to  
371 office workers. Indeed, BDCPP levels in firefighters were five times higher than the  
372 levels measured in office workers. When compared to NHANES, firefighters had higher

373 levels of DBuP, BDCPP, and BCEP, while office workers had levels similar levels of  
374 BDCPP, but lower levels of DBuP and BCEP than NHANES women (**Figure S1**).



375 Figure 1: Distribution (median, interquartile range, and 95th percentile) of flame-  
376 retardant metabolite levels ( $\mu\text{g/g-creatinine}$ ) in urine from 86 firefighters and 84  
377 office workers of the WFBC (2014-15). We substituted values below the limit of detection  
378 (LOD) with  $\text{LOD}/\sqrt{}$ . Significance stars represent the p-value from permutation test of  
379 the difference in average chemical level between firefighter and office workers: \*\*\*  
380  $<0.001$ ; \*\*  $<0.05$ ; \*  $<0.1$ .  
381

382 We used MLE regression models to assess the relationship between  
383 demographic variables and FR levels, controlling for occupation and log(creatinine)  
384 (**Table 2**) These models focused on BDCPP, BCEP and DBuP since these compounds  
385 had sufficient detection frequency (DF>70% in at least one group) to use in regression  
386 models. We found that having a BMI of 25.0 to 29.0 as compared to 18.5 to 24.9 was  
387 associated with increased levels of BDCPP, BCEP, and DBuP ( $\exp(\beta)$  (95%CI) = 1.71  
388 (1.02, 2.84), 2.70 (1.06, 6.84) and 2.45 (1.28, 4.86) respectively). Likewise, higher  
389 education was associated with FR levels; those who completed a bachelor's degree or  
390 higher education had higher BDCPP ( $\exp(\beta)$  (95%CI) = 1.65 (1.00, 2.73); BCEP, 3.63  
391 (1.49, 8.84); and DBuP, 2.24 (1.17, 4.31)) compared to participants who had completed  
392 some college or less. Race and ethnicity were not associated with most OPFRs except  
393 for BDCPP, where Black participants had a 2.52 (95%CI 1.10, 5.75) times higher  
394 geometric mean compared to White participants.

395

**Table 2.** Description of covariates among firefighters and office workers and the adjusted<sup>a</sup> proportional change in geometric mean (95% CI) of urinary chemical concentration (ng/mL) (DF>70%) for each unit increase, or each category increase compared to the referent, from maximum likelihood estimation (MLE) models

Variable	Mean ( $\pm$ SD) or N (%)		MLE exponentiated $\beta$ (95%CI)		
	office workers (N = 84)	firefighters (N= 86)	BDCPP	BCEP	DBuP
Age (years)	48.3 ( $\pm$ 10.5)	47.5 ( $\pm$ 4.6)	1.00 (0.97,1.03)	0.99 (0.94,1.04)	1.00 (0.96,1.04)
Time lived in CA (years)	35.5 ( $\pm$ 14.5)	40.0 ( $\pm$ 10.1)	0.99 (0.97,1.01)	1.00 (0.97,1.04)	1.01 (0.99,1.04)
U.S. Born	62 (73.8%)	77 (89.5%)	0.80 (0.44,1.46)	0.69 (0.24,2.02)	0.90 (0.40,2.02)
<b>BMI<sup>b,c</sup></b>					
18.5 – 24.9	43 (51.2%)	33 (38.4%)	reference	reference	reference
25.0 - 29.0	23 (27.4%)	35 (40.7%)	<b>1.71 (1.02,2.84)</b>	<b>2.70 (1.06,6.84)</b>	<b>2.45 (1.28,4.68)</b>
>30	16 (19.0%)	13 (15.1%)	0.84 (0.44,1.58)	0.75 (0.23,2.50)	1.55 (0.68,3.54)
<b>Race/ethnicity</b>					
White	37 (44.0%)	40 (46.5%)	reference	reference	reference
Black	5 (6.0%)	9 (10.5%)	<b>2.52 (1.10,5.75)</b>	0.76 (0.17,3.46)	0.73 (0.25,2.17)
Latina	13 (15.5%)	19 (22.1%)	0.56 (0.31,1.03)	0.89 (0.30,2.67)	0.82 (0.37,1.79)
Asian	19 (22.6%)	11 (12.8%)	1.02 (0.55,1.88)	1.60 (0.52,4.93)	1.15 (0.50,2.63)
Other/multi	10 (11.9%)	7 (8.1%)	0.66 (0.31,1.42)	0.43 (0.10,1.89)	0.29 (0.09,0.89)
<b>Education</b>					
Some college or less	15 (17.9%)	48 (55.8%)	reference	reference	reference
Bachelors or greater	69 (82.1%)	38 (44.2%)	<b>1.65 (1.00,2.73)</b>	<b>3.63 (1.49,8.84)</b>	<b>2.24 (1.17,4.31)</b>

<sup>a</sup> Models adjusted for occupation (firefighter or office worker) and log(creatinine)

<sup>b</sup> CDC guidelines for BMI classification: normal weight 18.5-24.9; overweight 25.0-29.9, obese >30; BMI units: kg/m<sup>2</sup>

<sup>c</sup> seven participants declined to answer height weight questions

397 When we limited the analysis to firefighters, several variables were associated  
 398 with increased FR levels, however none of these relationships were statistically  
 399 significant (p-value  $\leq$ 0.05) (**Table 3**). We observed that FR levels were higher  
 400 firefighters who were on duty at the time of the sample collection (BDCPP exp( $\beta$ )  
 401 (95%CI) = 1.97 (0.91, 4.23) and BCEP (2.76 (0.91, 8.36)). We found slight associations  
 402 by firefighters' assigned role; officers and firefighters showed slightly elevated levels of  
 403 BDCPP and BCEP compared to drivers. SCBA use in general was associated with  
 404 lower FR levels with the exception of SCBA use during exterior fire suppression which  
 405 was associated with higher mean BCEP compared to those who responded that they

406 sometimes or less frequently did so. Firefighters assigned to one of the San Francisco  
 407 Airport fire stations had elevated levels of BDCPP, BCEP, and DBuP. Fighting a fire  
 408 within the 24 hours and 7 days prior to the sample collection was also slightly  
 409 associated with BDCPP and BCEP levels. Use of firefighting foam, number of hours  
 410 spent in vehicles per week for both home and work and the number of years worked  
 411 with SFFD was not associated with FR levels.

**Table 3.** Adjusted<sup>a</sup> proportional change in geometric mean of urine flame retardant concentration (ng/mL, DF >70%) by unit increase, or category change from the referent, of firefighter occupational activities and characteristics from individual MLE regression models

Variable	number responded	BDCPP	BCEP	DBuP
Assigned to airport (yes)	14	1.51 (0.66,3.44)	1.73 (0.50,5.99)	1.73 (0.69,4.33)
Years worked with SFFD	86	0.93 (0.87,1.00)	0.89 (0.80,1.00)	0.92 (0.85,0.99)
Used firefighting foam past year (yes)	25	0.99 (0.50,1.94)	0.88 (0.32,2.45)	0.80 (0.38,1.71)
On-duty at sample collection	21	1.97 (0.91,4.23)	2.76 (0.91,8.36)	0.98 (0.41,2.33)
Hours spent in vehicle per week	86	1.01 (0.98,1.04)	1.04 (1.00,1.09)	1.01 (0.98,1.05)
Fire in last 24hrs (yes)	15	1.31 (0.58,2.94)	1.52 (0.45,5.17)	0.98 (0.40,2.43)
Fire in last 7 days (yes) <sup>b</sup>	18	1.32 (0.61,2.89)	1.86 (0.59,5.86)	0.75 (0.30,1.90)
Fire in last 1 month (yes) <sup>b</sup>	44	1.24 (0.59,2.57)	1.21 (0.40,3.66)	0.95 (0.39,2.30)
<b>Position</b>				
Driver	21	reference	reference	reference
Firefighter	40	1.40 (0.65,3.01)	1.32 (0.41,4.23)	1.19 (0.50,2.83)
Officer	25	1.62 (0.71,3.73)	1.88 (0.53,6.70)	1.40 (0.55,3.58)
<b>SCBA use with:<sup>c</sup></b>				
Interior fire suppression (always vs often or less)	60	0.75 (0.39,1.47)	1.26 (0.45,3.48)	0.95 (0.45,2.02)
Exterior fire suppression (often/always vs sometimes or less)	32	0.90 (0.48,1.70)	2.52 (0.92,6.88)	1.01 (0.49,2.08)
Salvage & overhaul (often/always vs sometimes or less)	26	0.51 (0.27,0.99)	0.78 (0.28,2.16)	0.50 (0.24,1.05)

<sup>a</sup> Adjusted for log(creatinine)

<sup>b</sup> n = 66 firefighters who consented to giving researchers access to their SFFD fire history records; <sup>c</sup> responses collected as 'never, rarely, sometimes, often, always' and were combined due to low frequency in individual response categories

#### 412 *FR exposure and thyroid hormone levels*

413 T<sub>4</sub> and TSH were slightly negatively correlated with each other (Spearman  
 414 correlation coefficient: -0.13, p-value = 0.1). We found that most participants were within

415 the reference range (i.e. the range of levels that are considered normal)(Chiovato et al.  
416 2019; Hollowell et al. 2002) for both TSH and T<sub>4</sub> (**Table 4**). Only 6% (N = 11) of  
417 participants had TSH levels outside (below or above) of the reference range and sixteen  
418 participants (9%) had T<sub>4</sub> levels outside the reference range, however the majority of  
419 those outside the reference range were firefighters (N = 14).

**Table 4.** Geometric mean, geometric standard deviation and select percentiles for thyroid hormones TSH and T<sub>4</sub> in WFBC study participants N = 165

Hormone	Reference range <sup>a</sup>	GM (GSD)	Min	Max	Percentiles			
					25%	50%	75%	95%
TSH	0.4 to 4.0 mIU/L	1.26 (2.06)	0.03	11.4	0.88	1.36	1.96	3.36
T <sub>4</sub>	4.6 to 12 ug/dL	6.41 (1.30)	3.69	20.58	5.43	6.36	7.26	9.29

Abbreviations: GM = Geometric mean, GSD = geometric Standard deviation.

<sup>a</sup>Chiovato et al. 2019; Hollowell et al. 2002

420 TSH levels were not significantly different between firefighters and office workers  
421 (permutation test p-value = 0.6) however T<sub>4</sub> levels did vary slightly by occupation  
422 (permutation test p-value = 0.06). We assessed the relationship of potential  
423 confounders with TSH and T<sub>4</sub> as a consideration for inclusion into OLS regression  
424 models. Since neither race/ethnicity nor education were associated with TSH or T<sub>4</sub> in  
425 our data (data not shown), we did not include them in final models. Although age was  
426 not associated with TH in our study population, it was included as a potential  
427 confounder based on prior literature indicating it may be an important factor for  
428 determining TH levels (Hollowell et al. 2002).

429 We applied OLS regression models to assess the impact of FR levels on T<sub>4</sub> and  
430 TSH levels controlling for log(creatinine) and age. FR levels were not associated with  
431 TSH in our models (**Table S2**); however, we did see a relationship between several FR  
432 and T<sub>4</sub> levels. BDCPP, which we defined as continuous (DF>70%) in OLS regression

433 models was negatively associated with T<sub>4</sub> levels (**Table 5**). In the full cohort, a two-fold  
434 increase in BDCPP levels was associated with a decrease of 1.95% (95% CI 3.57, 0.29)  
435 in T<sub>4</sub>, and in models limited to firefighters, a two-fold increase in BDCCP was associated  
436 with a T<sub>4</sub> decrease of 2.88% (5.278, 0.417), controlling for age and log(creatinine). The  
437 percent change among office workers was smaller and not significant.

**Table 5.**  $\beta$ (95% confidence interval (CI)) and percent change (95% CI) in T<sub>4</sub> levels for each doubling of BDCPP in adjusted<sup>a</sup> OLS regression models

Model	$\beta$ (95%CI)	percent change (95%CI)
Full cohort	-0.028 (-0.053,-0.004)	-1.95 (-3.57,-0.29)
Firefighters only	-0.042 (-0.078,-0.006)	-2.88 (-5.28,-0.42)
Office workers only	0.003 (-0.036,0.043)	0.23 (-2.49,3.026)

<sup>a</sup>Models adjusted for age and log(creatinine); values below LOD replaced with LOD/sqrt(2)

438 Due to large differences in DF for DBuP, BCEP, TBBPA and DPCP between  
439 firefighters and office workers, we ran separate OLS regression models by occupation  
440 for each FR as a predictor and T<sub>4</sub> as a continuous outcome. For office workers, we  
441 categorized DBuP, BCEP, TBBPA and DPCP into <LOD,  $\geq$ LOD. Among firefighters we  
442 categorized TBBPA and DPCP as <LOD,  $\geq$ LOD and categorized DBuP and BCEP as  
443 <LOD, LOD to 50<sup>th</sup> %, >50<sup>th</sup> %. Because the DF for DBuP and BCEP was >70% for  
444 firefighters we also ran OLS regression models with the continuous DBuP and BCEP.  
445 Among firefighters, DBuP was slightly associated with a decrease in T<sub>4</sub> levels among  
446 those at the LOD or above compared to below the LOD. DBuP was not significantly  
447 associated with T<sub>4</sub> among office workers nor were BCEP, TBBPA and DPCP associated  
448 with T<sub>4</sub> levels in either firefighters or office workers (**Table 6**).

449



450

**Table 6.** Percent change (95% CI) in T4 for each category increase in FR compared to <LOD for firefighters and office workers from adjusted OLS regression models<sup>a</sup>.

Firefighters		Office Workers	
<b>DBuP</b>			
<LOD	reference	<LOD	reference
LOD-50th%	-6.07 (-19.45,9.53)	>LOD	5.57 (-6.21,18.83)
>50th%	-12.68 (-24.94,1.59)		
<i>Continuous</i>	-2.01 (-4.63,0.68)		
<b>BCEP</b>			
<LOD	reference	<LOD	reference
LOD-50th %	8.05 (-7.13,21.61)	>LOD	-1.16 (-11.59,10.49)
>50th%	5.23 (-1.79,0.61)		
<i>Continuous</i>	0.623 (-1.58,2.88)		
<b>TBBPA</b>			
<LOD	reference	<LOD	reference
>LOD	4.30 (-6.29,16.08)	>LOD	6.28 (-4.71,18.54)
<b>DPCP</b>			
<LOD	reference	NA (DF <25%)	
>LOD	-9.21 (-19.29,2.14)		

<sup>a</sup>Models adjusted age and log(creatinine); If DF < 50% OPFR values were categorized into <= LOD or >LOD; those OPFRs with a DF of at least 50% were categorized as <=LOD, >LOD to 50th, and >50th; DBuP and BCEP DF >70 %

451

## 452 Discussion

453 To our knowledge this is the first study to measure exposures to replacement  
 454 FR and effects in thyroid function among women firefighters compared to office workers.  
 455 In addition, this paper confirms prior reports that firefighters have higher levels of  
 456 OPFRs compared to the general population and contributes further evidence that  
 457 OPFRs are an endocrine disrupting compound.

458 *Flame retardant levels and predictors of exposure*

459 FR were detected in all firefighters tested and most office workers. Women  
460 firefighters had higher detection frequencies and higher average levels of flame  
461 retardants compared to office workers. BDCPP, BCEP, and DBuP were detected more  
462 frequently and at higher concentrations among firefighters compared to office workers.  
463 BDCPP is a metabolite of TDCPP, a FR that is commonly detected in household dust  
464 and in furniture foam (Hammel et al. 2017), which may help explain why office workers  
465 also had a high detection frequency of this compound. In fact, studies of adult women  
466 (age 18+) and children (ages 6-12) from the 2013-2014 National Health and Nutrition  
467 Examination Survey (NHANES), a nationally representative sample of the US  
468 population, had similar levels to the office workers of our study (Ospina et al. 2018).

469 Few demographic variables were associated with FR levels. Age, for example  
470 was not associated with FR in our study; Conversely, the positive relationship between  
471 OFPR exposures and BMI is consistent with other studies (Boyle et al. 2019). While  
472 Black participants had higher levels of BDCPP compared to White participants when  
473 controlling for age and occupation, this relationship may in fact reflect other unknown  
474 exposures and be a function of the limited numbers of Black participants in both the  
475 firefighters and office worker groups.

476 California's flammability standards and the changes implemented in 2013 may  
477 have impacted the levels of chemicals that we observed in our study population.  
478 Concerns about the bioaccumulative properties of brominated flame retardants and the  
479 updated regulation TB177-2013, led to a shift in the types of FR used in furniture.  
480 Indeed, studies have observed a decrease in PBDEs in people and the environment,  
481 while levels of OPFRs and other replacement flame retardants are increasing in dust

482 and human biomonitoring studies (Dodson et al. 2012; Ospina et al. 2018; Stapleton et  
483 al. 2008; Zota et al. 2013). The half-lives of these replacement chemicals are not  
484 known, though they are generally considered to be shorter than that of PBDEs and  
485 potentially on the order of hours or days (Carignan et al. 2013; Nomeir et al. 1981).  
486 However, furniture in homes and offices can be a reservoir for both legacy and  
487 replacement FR; furniture is not frequently replaced and the chemicals applied to them  
488 are relatively stable enabling FR to persist in homes and offices for decades (Zota et al.  
489 2013). The elevated levels of BDCPP and other flame retardants in furniture, homes,  
490 and offices, may also be an important exposure source among firefighters as well as the  
491 office workers in our study. Firefighter turnout gear and protective equipment is  
492 plausibly another source of FR for firefighters, since they have been shown to be  
493 treated with flame retardant chemicals. Similarly, FR-containing furniture may add to the  
494 toxic burden of fighting fires and translate to higher exposures among firefighters when  
495 they respond to fires. Studies have shown that fighting fires can contaminate firefighter  
496 gear, trucks, engines and equipment, bringing chemical exposures indoors (Banks et al.  
497 2020; Mayer et al. 2019; Shen et al. 2018).

498 Fire station dust is potentially an important source of FR exposure among  
499 firefighters. The increased levels of OPFR found in the women firefighters compared to  
500 office workers is consistent with studies measuring elevated levels of OPFRs in fire  
501 station dust in the U.S. and abroad. Shen *et al.* (2018) found higher levels of TDCPP,  
502 tri-n-butyl phosphate (TNBP), and tris(2-chloroisopropyl)phosphate (TCPP), in dust  
503 collected in 2015 from fire stations across the U.S., compared to levels measured in  
504 homes and other occupational settings (Shen et al. 2018). Similarly, a study of

505 Australian fire station dust found higher median OPFR levels in fire stations than from  
506 dust samples collected in homes and offices (Banks et al. 2020). Firefighters who were  
507 on duty at the time of the sample collection had higher levels of two FR chemicals  
508 supporting evidence that being at work and potentially in the fire station increases their  
509 exposures. We expect that firefighter gear may also be a source of exposure; however,  
510 we do not have information on the age, type, manufacturer or chemical composition of  
511 turnout gear that firefighters use in San Francisco. A 2016 study of firefighter gear found  
512 that some new hoods and gloves had detectable levels of brominated flame retardants  
513 (Alexander and Baxter 2016) but to our knowledge studies have not analyzed gear for  
514 OPFRs. Nevertheless, firefighter gear may still contain FR compounds that can  
515 additionally contribute to exposures from dust accumulated on the gear from fires and  
516 calls (Alexander and Baxter 2016). A recent study of firefighting gear found that fighting  
517 fires may contaminate the hoods firefighters wear under their protective equipment with  
518 OPFRs (Mayer et al. 2019). Additionally, laundering may have a limited impact on  
519 reducing exposures and, in some cases, may cross-contaminate gear (Mayer et al.  
520 2019).

#### 521 *FR exposure and thyroid hormone levels*

522 The toxicity of most replacement chemical flame retardants has not been fully  
523 characterized. One of the few FRs that has been assessed for toxicity and adverse  
524 health effects is TDCPP which, due to its potential for carcinogenicity has been included  
525 in California's Prop 65 list (Office of Environmental Health Hazard Assessment 2015). In  
526 fact TDCPP, which has been used as a FR replacement in PUF and fabrics, is  
527 structurally similar to tris(2,3-dibromopropyl) phosphate (TDBPP or brominated "Tris")

528 which was banned in the 1970's from use in children's pajamas because of its  
529 mutagenic and carcinogenic properties (Dodson et al. 2012).

530 Our study demonstrated that firefighters in San Francisco have higher levels of  
531 FR than office workers and contributes to the limited literature of thyroid hormone  
532 disruption in women in relation to FR exposure. This association of FR with decreased  
533 T<sub>4</sub> was primarily observed among firefighters, and particularly for BDCPP, which had the  
534 highest DF of the FR we measured. The broader range and much higher levels of  
535 exposure among the firefighters in our cohort may have allowed us to see an effect of  
536 FR exposure on T<sub>4</sub> that we did not observe among office workers.

537 *In vitro* studies provide evidence for the mechanistic plausibility of TH  
538 disruption from OPFR exposures. However, the mechanism of effect is not yet fully  
539 understood. Hill et al. (2018) tested the competition of binding of T<sub>4</sub> and TTR (one of  
540 many transport proteins for T<sub>4</sub>). They found that TDCPP and other organophosphate tri-  
541 and di-esters increased the binding affinity of T<sub>4</sub> with TTR. They hypothesized that  
542 organophosphate compounds may bind to the surface of TTR, creating a  
543 conformational change allowing more T<sub>4</sub> binding and increasing the delivery of T<sub>4</sub> to  
544 target cells resulting in lower circulating levels of T<sub>4</sub> and disrupting T<sub>4</sub> homeostasis (Hill  
545 et al. 2018). Our study provides evidence that exposure to TDCPP and, to a lesser  
546 extent, DBuP exposure can affect TH levels. While our study does not have a clinical  
547 outcome, and the changes we observed were relatively small, even small disruptions to  
548 TH can have multiple adverse downstream health effects (Boas et al. 2012) even within  
549 normal ranges (Taylor et al. 2013). Thyroid dysfunction affects up to 5% of the  
550 population and is more likely to affect women than men (Hollowell et al. 2002). Over the

551 past two decades the incidence of thyroid cancer has also increased and cannot be  
552 fully explained by improved testing and diagnosis (Ward et al. 2010). Identifying  
553 environmental chemical exposures that may be associated with biological changes,  
554 such as thyroid disruption could be relevant to adverse health outcomes such as  
555 cardiovascular disease (Rodondi et al. 2006) and thyroid disease (Ward et al. 2010),  
556 brain development of offspring during gestation, and long-term adverse health  
557 outcomes such as cancer (Krashin et al. 2019).

558 We were unable to identify specific sources of FRs or why firefighters had  
559 higher levels of exposure than office workers. While durable consumer goods such as  
560 couches and mattresses may contribute to levels in the general population and fire  
561 station dust may be an important source in firefighters, this does not fully explain where  
562 the FR exposures are coming from and why the levels are so much higher in fire  
563 fighters. Future studies need to elucidate potential sources of exposure to legacy and  
564 replacement FR which would facilitate the development and promotion of effective  
565 exposure prevention strategies.

#### 566 **Limitations:**

567 This was a cross sectional study of FR exposure and effects on thyroid  
568 function, which precludes making inferences regarding causality. Nevertheless, our  
569 findings confirm an association between TH levels and TDCPP, one that has also been  
570 identified in studies conducted *in vitro*, *in vivo* and in limited human studies (Farhat et al.  
571 2013; Meeker and Stapleton 2010; Wang et al. 2013a).

572 While we did not detect PBDE metabolites in our study population, analyzing these  
573 metabolites in urine is insensitive, and therefore these results were somewhat expected.

574 OH-BDE metabolites are more commonly measured in serum and these chemicals  
575 have been found in participants of other studies conducted during the same time as  
576 WFBC study (Park et al. 2015; Parry et al. 2018). Therefore, although we did not find  
577 OH-BDE metabolites in any of our samples, this may be due to the type of biological  
578 matrix we measured rather than an absence of exposure. Another potential limitation is  
579 that we used a mix of morning void and spot urine samples when measuring FR levels,  
580 potentially increasing the variability of the urine concentration and of chemicals  
581 measured in our samples. However, morning void or spot samples were selected at  
582 random for analysis, therefore the variability would be non-differential between  
583 firefighter and office workers and not likely to affect the differences we observed in their  
584 chemical levels nor the FR's relationship with potential covariates. In addition, we  
585 measured and adjusted for creatinine to account for urine dilution and reduce some of  
586 this variability (Barr et al. 2005).

587 Thyroid disease and taking thyroid hormone replacement medications may  
588 alter the TH levels. We asked participants "do you take any hormones other than birth  
589 control, such as premarin? if so what?" Some participants disclosed taking TH  
590 medications or having thyroid problems, however, the question may not have accurately  
591 captured all the participants with thyroid dysfunction or those who take TH  
592 replacements. Therefore, we may not have excluded everyone with artificial or  
593 abnormal TH levels due to illness or medication rather than chemical exposure. This  
594 could over- or under-estimate thyroid hormone levels depending on the thyroid problem,  
595 and we would not be able to predict in what direction this could affect our results.  
596 Finally, we may also have had limited statistical power, due to our modest sample size,

597 to assess the relationship between FR levels and predictors of exposure. This may have  
598 precluded our ability to see the full effect of FR exposure on TH disruption, especially at  
599 the lower exposure levels we observed in office workers.

600

## 601 **Conclusion**

602 This is the first study to measure flame retardant exposure in a cohort of  
603 women firefighters. Most participants had detectable levels of at least one  
604 organophosphate flame retardant, and both detection frequencies and levels were much  
605 higher among women firefighters than in women office workers; Median levels of  
606 BDCPP were five times higher in firefighters than for office workers. Additionally,  
607 exposure to BDCPP, and to a smaller extent DBuP, was associated with decreased  
608 levels of thyroxine particularly among firefighters—a twofold increase in BDCPP was  
609 associated with a 2.88% decrease in  $T_4$ . The observed thyroid hormone disruption may  
610 indicate potential biological perturbations resulting from occupational exposure to these  
611 flame retardants. Further research is needed to understand sources in occupational  
612 settings and fully characterize potential health impacts of these replacement flame  
613 retardants.



## References

- Alexander BM, Baxter CS. 2016. Flame-retardant contamination of firefighter personal protective clothing - A potential health risk for firefighters. *J Occup Environ Hyg* 13:D148-155; doi:10.1080/15459624.2016.1183016.
- Antibodies-online. 2020. T3, T4, TSH ELISA Kit - Product details.
- Banks APW, Engelsman M, He C, Wang X, Mueller JF. 2020. The occurrence of PAHs and flame-retardants in air and dust from Australian fire stations. *J Occup Environ Hyg* 17:73–84; doi:10.1080/15459624.2019.1699246.
- Barr DB, Wilder LC, Caudill SP, Gonzalez AJ, Needham LL, Pirkle JL. 2005. Urinary Creatinine Concentrations in the U.S. Population: Implications for Urinary Biologic Monitoring Measurements. *Environ Health Perspect* 113:192–200; doi:10.1289/ehp.7337.
- Birnbaum LS, Staskal DF. 2004. Brominated flame retardants: cause for concern? *Environ Health Perspect* 112:9–17; doi:10.1289/ehp.6559.
- Boas M, Feldt-Rasmussen U, Main KM. 2012. Thyroid effects of endocrine disrupting chemicals. *Mol Cell Endocrinol* 355:240–248; doi:10.1016/j.mce.2011.09.005.
- Boyle M, Buckley JP, Quiros-Alcala L. 2019. Associations between urinary organophosphate ester metabolites and measures of adiposity among U.S. children and adults: NHANES 2013-2014. *Environ Int* 127:754–763; doi:10.1016/j.envint.2019.03.055.
- Brown FR, Whitehead TP, Park J-S, Metayer C, Petreas MX. 2014. Levels of non-polybrominated diphenyl ether brominated flame retardants in residential house dust samples and fire station dust samples in California. *Environmental Research* 135:9–14; doi:10.1016/j.envres.2014.08.022.
- Bureau of Electronic and Appliance. 2018. Technical Bulletin 117-2013. Available: [http://www.bearhfti.ca.gov/about\\_us/tb117\\_2013.pdf](http://www.bearhfti.ca.gov/about_us/tb117_2013.pdf) [accessed 3 March 2018].
- Calsolaro V, Pasqualetti G, Niccolai F, Caraccio N, Monzani F. 2017. Thyroid Disrupting Chemicals. *Int J Mol Sci* 18; doi:10.3390/ijms18122583.
- Carignan CC, McClean MD, Cooper EM, Watkins DJ, Fraser AJ, Heiger-Bernays W, Stapleton HM, Webster TF. 2013. Predictors of Tris(1,3-dichloro-2-propyl) phosphate Metabolite in the Urine of Office Workers. *Environ Int* 55:56–61; doi:10.1016/j.envint.2013.02.004.
- Castorina R, Bradman A, Stapleton HM, Butt C, Avery D, Harley KG, Gunier RB, Holland N, Eskenazi B. 2017. Current-use flame retardants: Maternal exposure and neurodevelopment in children of the CHAMACOS cohort. *Chemosphere* 189:574–580; doi:10.1016/j.chemosphere.2017.09.037.

- Caux C, O'Brien C, Viau C. 2002. Determination of firefighter exposure to polycyclic aromatic hydrocarbons and benzene during fire fighting using measurement of biological indicators. *Appl Occup Environ Hyg* 17:379–386; doi:10.1080/10473220252864987.
- Chiovato L, Magri F, Carlé A. 2019. Hypothyroidism in Context: Where We've Been and Where We're Going. *Adv Ther* 36:47–58; doi:10.1007/s12325-019-01080-8.
- Covaci A, Harrad S, Abdallah MA-E, Ali N, Law RJ, Herzke D, de Wit CA. 2011. Novel brominated flame retardants: A review of their analysis, environmental fate and behaviour. *Environment International* 37:532–556; doi:10.1016/j.envint.2010.11.007.
- Daniels RD, Bertke S, Dahm MM, Yiin JH, Kubale TL, Hales TR, Baris D, Zahm SH, Beaumont JJ, Waters KM, Pinkerton LE. 2015. Exposure-response relationships for select cancer and non-cancer health outcomes in a cohort of U.S. firefighters from San Francisco, Chicago and Philadelphia (1950-2009). *Occup Environ Med* 72:699–706; doi:10.1136/oemed-2014-102671.
- Daniels RD, Kubale TL, Yiin JH, Dahm MM, Hales TR, Baris D, Zahm SH, Beaumont JJ, Waters KM, Pinkerton LE. 2014. Mortality and cancer incidence in a pooled cohort of US firefighters from San Francisco, Chicago and Philadelphia (1950-2009). *Occup Environ Med* 71:388–397; doi:10.1136/oemed-2013-101662.
- Diamanti-Kandarakis E, Bourguignon J-P, Giudice LC, Hauser R, Prins GS, Soto AM, Zoeller RT, Gore AC. 2009. Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocr Rev* 30:293–342; doi:10.1210/er.2009-0002.
- Dishaw LV, Macaulay LJ, Roberts SC, Stapleton HM. 2014. Exposures, mechanisms, and impacts of endocrine-active flame retardants. *Curr Opin Pharmacol* 19:125–133; doi:10.1016/j.coph.2014.09.018.
- Dobraca D, Israel L, McNeel S, Voss R, Wang M, Gajek R, Park J-S, Harwani S, Barley F, She J, Das R. 2015. Biomonitoring in California firefighters: metals and perfluorinated chemicals. *J Occup Environ Med* 57:88–97; doi:10.1097/JOM.0000000000000307.
- Dodson RE, Perovich LJ, Covaci A, Van den Eede N, Ionas AC, Dirtu AC, Brody JG, Rudel RA. 2012. After the PBDE phase-out: a broad suite of flame retardants in repeat house dust samples from California. *Environ Sci Technol* 46:13056–13066; doi:10.1021/es303879n.
- Dodson RE, Van den Eede N, Covaci A, Perovich LJ, Brody JG, Rudel RA. 2014. Urinary biomonitoring of phosphate flame retardants: levels in California adults and recommendations for future studies. *Environ Sci Technol* 48:13625–13633; doi:10.1021/es503445c.

- Farhat A, Crump D, Chiu S, Williams KL, Letcher RJ, Gauthier LT, Kennedy SW. 2013. In Ovo effects of two organophosphate flame retardants--TCPP and TDCPP--on pipping success, development, mRNA expression, and thyroid hormone levels in chicken embryos. *Toxicol Sci* 134:92–102; doi:10.1093/toxsci/kft100.
- Fent KW, Eisenberg J, Snawder J, Sammons D, Pleil JD, Stiegel MA, Mueller C, Horn GP, Dalton J. 2014. Systemic exposure to PAHs and benzene in firefighters suppressing controlled structure fires. *Ann Occup Hyg* 58:830–845; doi:10.1093/annhyg/meu036.
- Gore AC, Chappell VA, Fenton SE, Flaws JA, Nadal A, Prins GS, Toppari J, Zoeller RT. 2015. EDC-2: The Endocrine Society's Second Scientific Statement on Endocrine-Disrupting Chemicals. *Endocr Rev* 36:E1–E150; doi:10.1210/er.2015-1010.
- Grashow R, Bessonneau V, Gerona RR, Wang A, Trowbridge J, Lin T, Buren H, Rudel RA, Morello-Frosch R. 2020. Integrating exposure knowledge and serum suspect screening as a new approach to biomonitoring: An application in firefighters and office workers. *Environ Sci Technol*; doi:10.1021/acs.est.9b04579.
- Gravel S, Lavoue J, Bakhiyi B, Lavoie J, Roberge B, Patry L, Bouchard MF, Verner M-A, Zayed J, Labreche F. 2020. Multi-exposures to suspected endocrine disruptors in electronic waste recycling workers: Associations with thyroid and reproductive hormones. *Int J Hyg Environ Health* 225:113445; doi:10.1016/j.ijheh.2019.113445.
- Hammel SC, Hoffman K, Lorenzo AM, Chen A, Phillips AL, Butt CM, Sosa JA, Webster TF, Stapleton HM. 2017. Associations Between Flame Retardant Applications in Furniture Foam, House Dust Levels, and Residents' Serum Levels. *Environ Int* 107:181–189; doi:10.1016/j.envint.2017.07.015.
- Hill KL, Hamers T, Kamstra JH, Willmore WG, Letcher RJ. 2018. Organophosphate triesters and selected metabolites enhance binding of thyroxine to human transthyretin in vitro. *Toxicol Lett* 285:87–93; doi:10.1016/j.toxlet.2017.12.030.
- Hoffman K, Fang M, Horman B, Patisaul HB, Garantziotis S, Birnbaum LS, Stapleton HM. 2014. Urinary tetrabromobenzoic acid (TBBA) as a biomarker of exposure to the flame retardant mixture Firemaster® 550. *Environ Health Perspect* 122:963–969; doi:10.1289/ehp.1308028.
- Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, Braverman LE. 2002. Serum TSH, T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J Clin Endocrinol Metab* 87:489–499; doi:10.1210/jcem.87.2.8182.

- Jin C, Sun Y, Islam A, Qian Y, Ducatman A. 2011. Perfluoroalkyl acids including perfluorooctane sulfonate and perfluorohexane sulfonate in firefighters. *J Occup Environ Med* 53:324–328; doi:10.1097/JOM.0b013e31820d1314.
- Kim H, Rebholz CM, Wong E, Buckley JP. 2020. Urinary organophosphate ester concentrations in relation to ultra-processed food consumption in the general US population. *Environmental Research* 182:109070; doi:10.1016/j.envres.2019.109070.
- Knudsen N, Laurberg P, Rasmussen LB, Bülow I, Perrild H, Ovesen L, Jørgensen T. 2005. Small Differences in Thyroid Function May Be Important for Body Mass Index and the Occurrence of Obesity in the Population. *J Clin Endocrinol Metab* 90:4019–4024; doi:10.1210/jc.2004-2225.
- Krashin E, Piekietko-Witkowska A, Ellis M, Ashur-Fabian O. 2019. Thyroid Hormones and Cancer: A Comprehensive Review of Preclinical and Clinical Studies. *Front Endocrinol (Lausanne)* 10; doi:10.3389/fendo.2019.00059.
- Laitinen JA, Koponen J, Koikkalainen J, Kiviranta H. 2014. Firefighters' exposure to perfluoroalkyl acids and 2-butoxyethanol present in firefighting foams. *Toxicol Lett* 231:227–232; doi:10.1016/j.toxlet.2014.09.007.
- Lee DJ, Koru-Sengul T, Hernandez MN, Caban-Martinez AJ, McClure LA, Mackinnon JA, Kobetz EN. 2020. Cancer risk among career male and female Florida firefighters: Evidence from the Florida Firefighter Cancer Registry (1981-2014). *Am J Ind Med* 63:285–299; doi:10.1002/ajim.23086.
- LeMasters GK, Genaidy AM, Succop P, Deddens J, Sobeih T, Barriera-Viruet H, Dunning K, Lockey J. 2006. Cancer risk among firefighters: a review and meta-analysis of 32 studies. *J Occup Environ Med* 48:1189–1202; doi:10.1097/01.jom.0000246229.68697.90.
- Lerro CC, Koutros S, Andreotti G, Friesen MC, Alavanja MC, Blair A, Hoppin JA, Sandler DP, Lubin JH, Ma X, Zhang Y, Beane Freeman LE. 2015. Organophosphate insecticide use and cancer incidence among spouses of pesticide applicators in the Agricultural Health Study. *Occup Environ Med* 72:736–744; doi:10.1136/oemed-2014-102798.
- Ma F, Fleming LE, Lee DJ, Trapido E, Gerace TA. 2006. Cancer incidence in Florida professional firefighters, 1981 to 1999. *J Occup Environ Med* 48:883–888; doi:10.1097/01.jom.0000235862.12518.04.
- Mayer AC, Fent KW, Bertke S, Horn GP, Smith DL, Kerber S, La Guardia MJ. 2019. Firefighter hood contamination: Efficiency of laundering to remove PAHs and FRs. *J Occup Environ Hyg* 16:129–140; doi:10.1080/15459624.2018.1540877.

- Meeker JD, Stapleton HM. 2010. House dust concentrations of organophosphate flame retardants in relation to hormone levels and semen quality parameters. *Environ Health Perspect* 118:318–323; doi:10.1289/ehp.0901332.
- Nomeir AA, Kato S, Matthews HB. 1981. The metabolism and disposition of tris(1,3-dichloro-2-propyl) phosphate (Fyrol FR-2) in the rat. *Toxicol Appl Pharmacol* 57:401–413; doi:10.1016/0041-008x(81)90238-6.
- Office of Environmental Health Hazard Assessment. 2015. Tris(1,3-dichloro-2-propyl) Phosphate (TDCPP). OEHHA. Available: <https://oehha.ca.gov/proposition-65/chemicals/tris13-dichloro-2-propyl-phosphate-tdcpp> [accessed 25 April 2020].
- Ospina M, Jayatilaka NK, Wong L-Y, Restrepo P, Calafat AM. 2018. Exposure to organophosphate flame retardant chemicals in the U.S. general population: Data from the 2013-2014 National Health and Nutrition Examination Survey. *Environ Int* 110:32–41; doi:10.1016/j.envint.2017.10.001.
- Park J-S, Voss RW, McNeel S, Wu N, Guo T, Wang Y, Israel L, Das R, Petreas M. 2015. High Exposure of California Firefighters to Polybrominated Diphenyl Ethers. *Environmental Science & Technology* 49:2948–2958; doi:10.1021/es5055918.
- Parry E, Zota AR, Park J-S, Woodruff TJ. 2018. Polybrominated diphenyl ethers (PBDEs) and hydroxylated PBDE metabolites (OH-PBDEs): A six-year temporal trend in Northern California pregnant women. *Chemosphere* 195:777–783; doi:10.1016/j.chemosphere.2017.12.065.
- Pinkerton L, Bertke SJ, Yiin J, Dahm M, Kubale T, Hales T, Purdue M, Beaumont JJ, Daniels R. 2020. Mortality in a cohort of US firefighters from San Francisco, Chicago and Philadelphia: an update. *Occup Environ Med* 77:84–93; doi:10.1136/oemed-2019-105962.
- Preston EV, McClean MD, Claus Henn B, Stapleton HM, Braverman LE, Pearce EN, Makey CM, Webster TF. 2017. Associations between urinary diphenyl phosphate and thyroid function. *Environ Int* 101:158–164; doi:10.1016/j.envint.2017.01.020.
- R Core Team. 2015. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing:Vienna, Austria.
- Rodgers KM, Udesky JO, Rudel RA, Brody JG. 2018. Environmental chemicals and breast cancer: An updated review of epidemiological literature informed by biological mechanisms. *Environ Res* 160:152–182; doi:10.1016/j.envres.2017.08.045.
- Rodondi N, Aujesky D, Vittinghoff E, Cornuz J, Bauer DC. 2006. Subclinical hypothyroidism and the risk of coronary heart disease: a meta-analysis. *Am J Med* 119:541–551; doi:10.1016/j.amjmed.2005.09.028.

- RStudio Team. 2016. *RStudio: Integrated Development for R*. RStudio Inc.:Boston, MA.
- Rudel RA, Ackerman JM, Attfield KR, Brody JG. 2014. New exposure biomarkers as tools for breast cancer epidemiology, biomonitoring, and prevention: a systematic approach based on animal evidence. *Environ Health Perspect* 122:881–895; doi:10.1289/ehp.1307455.
- Rudel RA, Fenton SE, Ackerman JM, Euling SY, Makris SL. 2011. Environmental exposures and mammary gland development: state of the science, public health implications, and research recommendations. *Environ Health Perspect* 119:1053–1061; doi:10.1289/ehp.1002864.
- Schindler BK, Weiss T, Schütze A, Koslitz S, Broding HC, Bünger J, Brüning T. 2013. Occupational exposure of air crews to tricresyl phosphate isomers and organophosphate flame retardants after fume events. *Arch Toxicol* 87:645–648; doi:10.1007/s00204-012-0978-0.
- Shaw SD, Berger ML, Harris JH, Yun SH, Wu Q, Liao C, Blum A, Stefani A, Kannan K. 2013. Persistent organic pollutants including polychlorinated and polybrominated dibenzo-p-dioxins and dibenzofurans in firefighters from Northern California. *Chemosphere* 91:1386–1394; doi:10.1016/j.chemosphere.2012.12.070.
- Shen B, Whitehead TP, Gill R, Dhaliwal J, Brown FR, Petreas M, Patton S, Hammond SK. 2018. Organophosphate flame retardants in dust collected from United States fire stations. *Environment International* 112:41–48; doi:10.1016/j.envint.2017.12.009.
- Shen B, Whitehead TP, McNeel S, Brown FR, Dhaliwal J, Das R, Israel L, Park J-S, Petreas M. 2015. High levels of polybrominated diphenyl ethers in vacuum cleaner dust from California fire stations. *Environ Sci Technol* 49:4988–4994; doi:10.1021/es505463g.
- Stapleton HM, Allen JG, Kelly SM, Konstantinov A, Klosterhaus S, Watkins D, McClean MD, Webster TF. 2008. Alternate and new brominated flame retardants detected in U.S. house dust. *Environ Sci Technol* 42:6910–6916; doi:10.1021/es801070p.
- Stapleton HM, Sharma S, Getzinger G, Ferguson PL, Gabriel M, Webster TF, Blum A. 2012. Novel and high volume use flame retardants in US couches reflective of the 2005 PentaBDE phase out. *Environ Sci Technol* 46:13432–13439; doi:10.1021/es303471d.
- Taylor PN, Razvi S, Pearce SH, Dayan CM. 2013. Clinical review: A review of the clinical consequences of variation in thyroid function within the reference range. *J Clin Endocrinol Metab* 98:3562–3571; doi:10.1210/jc.2013-1315.
- Trowbridge J, Gerona RR, Lin T, Rudel RA, Bessonneau V, Buren H, Morello-Frosch R. 2020. Exposure to Perfluoroalkyl Substances in a Cohort of Women Firefighters

- and Office Workers in San Francisco. *Environ Sci Technol* 54:3363–3374; doi:10.1021/acs.est.9b05490.
- Van den Eede N, Dirtu AC, Ali N, Neels H, Covaci A. 2012. Multi-residue method for the determination of brominated and organophosphate flame retardants in indoor dust. *Talanta* 89:292–300; doi:10.1016/j.talanta.2011.12.031.
- van der Veen I, de Boer J. 2012a. Phosphorus flame retardants: properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* 88:1119–1153; doi:10.1016/j.chemosphere.2012.03.067.
- van der Veen I, de Boer J. 2012b. Phosphorus flame retardants: Properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* 88:1119–1153; doi:10.1016/j.chemosphere.2012.03.067.
- Wang Q, Liang K, Liu J, Yang L, Guo Y, Liu C, Zhou B. 2013a. Exposure of zebrafish embryos/larvae to TDCPP alters concentrations of thyroid hormones and transcriptions of genes involved in the hypothalamic-pituitary-thyroid axis. *Aquat Toxicol* 126:207–213; doi:10.1016/j.aquatox.2012.11.009.
- Wang Y, Starling AP, Haug LS, Eggesbo M, Becher G, Thomsen C, Travlos G, King D, Hoppin JA, Rogan WJ, Longnecker MP. 2013b. Association between Perfluoroalkyl substances and thyroid stimulating hormone among pregnant women: a cross-sectional study. *Environ Health* 12:76; doi:10.1186/1476-069X-12-76.
- Ward EM, Jemal A, Chen A. 2010. Increasing incidence of thyroid cancer: is diagnostic scrutiny the sole explanation? *Future Oncology* 6:185–188; doi:10.2217/fon.09.161.
- Zoeller RT, Tan SW, Tyl RW. 2007. General background on the hypothalamic-pituitary-thyroid (HPT) axis. *Crit Rev Toxicol* 37:11–53; doi:10.1080/10408440601123446.
- Zota AR, Linderholm L, Park J-S, Petreas M, Guo T, Privalsky ML, Zoeller RT, Woodruff TJ. 2013. Temporal Comparison of PBDEs, OH-PBDEs, PCBs, and OH-PCBs in the Serum of Second Trimester Pregnant Women Recruited from San Francisco General Hospital, California. *Environ Sci Technol* 47:11776–11784; doi:10.1021/es402204y.