

Evaluation of different stool extraction methods for metabolomics measurements in human fecal samples

Authors: Vanessa Erben^{1,2}, Gernot Poschet³, Petra Schrotz-King¹, Hermann Brenner^{1,4,5}

Author Affiliations:

¹ Division of Preventive Oncology, German Cancer Research Center (DKFZ) and National Center for Tumor Diseases (NCT), Heidelberg, Germany;

² Medical Faculty Heidelberg, Heidelberg University, Heidelberg, Germany;

³ Center for Organismal Studies (COS), Heidelberg University, Heidelberg, Germany;

⁴ Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany;

⁵ German Cancer Consortium (DKTK), Heidelberg, Germany.

Authors' last names: Erben, Poschet, Schrotz-King, Brenner

Corresponding author: Dr. Petra Schrotz-King, NCT, Im Neuenheimer Feld 460, 69120 Heidelberg;
+49 (0)6221 5635970, petra.schrotz-king@nct-heidelberg.de

Short running head: Stool processing methods for metabolomics

Keywords: Stool, feces, metabolomics, processing

Word count: 2939; **Abstract word count:** 258; **Tables:** 4; **Supplementary material:** 2.

Abbreviations: CRC, colorectal cancer; MTBE, methyl tert-butyl ether.

1 **Abstract**

2 **Background:** Measurement of metabolomics in human stool samples is of great interest for a broad
3 range of applications in biomedical research including early detection of colorectal neoplasms.
4 However, due to the complexity of metabolites there is no consensus on how to process samples for
5 stool metabolomics measurements to obtain a broad coverage of hydrophilic and hydrophobic
6 substances.

7 **Methods:** We used frozen stool samples (50mg) from healthy study participants. Stool samples were
8 processed after thawing using 8 different processing protocols and different solvents. Metabolites were
9 measured afterwards using the MxP® Quant 500 kit (Biocrates). The best performing protocol was
10 subsequently applied to compare stool samples of participants with different dietary habits.

11 **Results:** In this study, we were able to determine up to 340 metabolites of various chemical classes
12 extracted from stool samples of healthy study participants with 8 different protocols. Polar metabolites
13 such as amino acids could be measured with each method while other metabolite classes, particular
14 lipid species, are more dependent on the solvent or combination of solvents used. Only a small number
15 of triglycerides or acylcarnitines were detected in human feces. Extraction efficiency was higher for
16 protocols using isopropanol or those using ethanol or methanol and MTBE including an evaporation
17 and concentration step than for other protocols. We detected significant fecal metabolite differences
18 between vegetarians, semi-vegetarians and non-vegetarians.

19 **Conclusion:** For the evaluation of metabolites in fecal samples we found protocols using solvents like
20 isopropanol and those using ethanol or methanol and MTBE including an evaporation and
21 concentration step to be superior over others tested in this study.

22 **Introduction**

23 Measurement of metabolomics in human stool samples is of great interest for a broad range of
24 applications in biomedical research including early detection of colorectal cancer (CRC) as a non-
25 invasive alternative to colonoscopy. Metabolomics might be promising for this purpose as metabolites
26 are closely related to the phenotype and depict current metabolic processes happening in an organism.
27 Stool is directly associated with the gut and may reflect changes in metabolism very early through its
28 transit in the gut (1).

29 Fecal mass consists to a great proportion of water and bacteria but also food components or
30 metabolites (2). Fecal samples show great variability in their material content and characteristics,
31 which makes it difficult to standardize the process from collection to processing and analysis including
32 the analytical platform (3). The latter defines also the sensitivity of the analysis and the type of
33 analytes available for analyses. Apart from the afore mentioned differences of stool samples,
34 metabolic changes in stool might be derived directly from the development of cancer or precancerous
35 cells or from a change in the gut microbiota which both result in a distinct metabolic phenotype that
36 might be characteristic for the disease (1). The metabolic profile of CRC or its precursors may help in
37 the understanding of disease development, progression and early detection (4).

38 Some studies have already found fecal metabolomics biomarkers for early detection of CRC but
39 metabolite selection strongly varied (1, 5-7) and as different studies were using different processing
40 methods, no direct comparison is possible. There is no consensus how to process stool samples for
41 metabolomics measurements to get reliable and reproducible results. A review by Deda et al. focused
42 on the existing various stool preparation protocols and found the metabolites to be dependent on the
43 extraction method (8). In this study we used the MxP® Quant 500 Kit (Biocrates Life Sciences AG,
44 Innsbruck, Austria) to determine and quantify a very broad range of metabolite classes in human feces.
45 In total, we used 8 different stool preparation protocols to assess the best coverage for stool metabolite
46 profiles. The protocol yielding the highest multitude of extracted metabolites was used to analyze and
47 compare additional stool samples of healthy study participants with different dietary habits.

48 **Subjects and Methods**

49 **Study design**

50 The GEKKO study (Gebt dem Krebs keine Chance – Onkocheck) is an ongoing study in southwest
51 Germany including people participating in screening colonoscopy (Arm A) or with diagnosed primary
52 cancer (Arm B). The study was approved by the ethics committees of the Medical Faculty Heidelberg
53 and of the physicians' boards of Baden-Württemberg and Rhineland Palatinate. The GEKKO study
54 Arm A was designed to evaluate novel early detection markers of CRC. People undergoing a
55 screening colonoscopy at a gastroenterology practice in or around Heidelberg, Germany, who are over
56 the age of 30 years, with no history of CRC, no inflammatory bowel disease, no colonoscopy within
57 the last 5 years and speaking and understanding the German language are invited to participate.

58 After written informed consent was received, participants are asked to fill in a questionnaire regarding
59 lifestyle and demographic data and to provide blood, stool, saliva and urine samples for research
60 purposes prior to colonoscopy. Biosamples are processed and then stored at -80°C until needed.
61 Colonoscopy reports are provided from the physicians to the study center. Participants are grouped
62 according to their most advanced finding at colonoscopy. From the individuals with no polyps or any
63 findings at colonoscopy, those with incomplete colonoscopy (Coecum not reached) or poor bowel
64 preparation were excluded. For this analysis, 3 participants of the GEKKO study Arm A were selected
65 between the age of 50 and 79 years with no polyps or any findings at colonoscopy to test 8 different
66 stool preparation protocols and to define the best analytical outcome.

67 In a further step, stool samples from additional healthy GEKKO participants (n = 18) of approximately
68 the same age (50 – 65 years) with different dietary habits (vegetarians, semi-vegetarians, and non-
69 vegetarians) were processed with the protocol that performed best with respect to numbers of detected
70 metabolites and sample handling, and results were compared between groups. Information on dietary
71 habits of the study participants was extracted from the questionnaire. Vegetarians were defined as
72 never eating meat, processed meat and poultry. Semi-vegetarians were defined as eating meat,
73 processed meat and poultry less than once a week. Other participants reporting to consume either meat
74 or processed meat or poultry more often were categorized as non-vegetarians (9).

75 **Sample collection and handling**

76 Native stool samples were collected by the participants at home from a normal bowel movement prior
77 to bowel preparation for colonoscopy with standard stool collection tubes provided with a small spoon
78 for collecting the stool. The stool samples were then directly frozen by the participants at -20°C at
79 home. The participants were asked to document date and time of sampling and the storage
80 temperature. The stool samples were taken by the participant in a freeze-cool transport container and
81 in an isolated envelope to the gastroenterologists' practices, directly frozen again at -20°C and within
82 the week of receipt delivered by a transport service on dry ice to the GEKKO study laboratory at the
83 National Center for Tumor Diseases (NCT) in Heidelberg where they were immediately frozen
84 at -80°C.

85 **Processing of the samples**

86 For this analysis, we used stool samples from 3 individuals of the GEKKO study testing 8 different
87 stool processing protocols each in triplicate which results in a total of 72 measurements. Every
88 protocol was performed with a frozen biomass of 50mg wet stool weight. Used protocols were adapted
89 in part from previously published work (protocol 1 (10), protocol 3 (11) and protocol 7 (12)) and from
90 the current recommended SOP provided from Biocrates (protocol 5 (13)). Details of the protocols are
91 shown in **Table 1**. In brief, feces was cut and weighed in frozen state and kept frozen until processing.
92 The samples were thawed and prepared according to the specific protocol and frozen again at -80°C
93 until further processing via MxP® Quant 500 Kit (Biocrates Life Sciences AG, Innsbruck, Austria).
94 Liquid sample extracts were processed according to the vendor's instructions for human plasma
95 samples. Samples dried after extraction had to be resolved before measurement (for protocol 6 and 7).
96 Therefore, 50µl of 100% isopropanol were added into the vial and the mixture was vortexed for 3 min
97 at room temperature. Additionally, 50µl of 30% isopropanol were added and again vortexed for 3min
98 at room temperature. Short centrifugation (5sec) separated the solid substances from the liquid phase
99 which was further used for analysis.

100 For metabolite measurements, a QTRAP6500+ (Sciex, Germany) MS/MS connected to an UPLC I-
101 class Plus (Waters, Germany) chromatography system was used. Conditions for LC separation and

102 FIA analyses as well as individual MRM parameters for each metabolite and respective internal
103 standards were provided by the vendor of the kit (Biocrates). The software MetIDQ (version Oxygen;
104 Biocrates Life Sciences AG, Innsbruck, Austria) was used for processing of the data.

105 In total, 630 metabolites can be measured via this kit. The limit of detection (LOD) for each
106 compound is defined as three times the background noise. The lower limit of quantitation (LLOQ) is
107 at least ten times the background noise. At LLOQ measured metabolite concentrations can be
108 considered as reliable. Data are normalized with a tissue factor for quantification under the following
109 assumption: 1mg tissue equals 1µl tissue or stool. Concentrations are given in pmol of the metabolite
110 /mg stool mass.

111 The following metabolites are measurable using the aforementioned kit: 1 alkaloid (trigonelline), 1
112 amine oxide (trimethylamine N-oxide), 20 amino acids, 30 other amino acid related metabolites, 14
113 bile acids, 9 biogenic amines, sugars (hexoses including glucose), 7 carboxylic acids, 1 cresol (p-
114 cresol sulfate), 12 fatty acids, 4 hormone and related metabolites, 4 indoles and derivatives, 2
115 nucleobases and related molecules, 1 metabolite from the group of vitamins and cofactors, 40
116 acylcarnitines, 14 lysophosphatidylcholines, 76 phosphatidylcholines, 15 sphingomyelins, 28
117 ceramides, 8 dihydroceramides, 19 hexosylceramides, 9 dihexosylceramides, 6 trihexosylceramides,
118 22 cholesteryl esters, 44 diglycerides, and 242 triglycerides. All the related isobaric and isomeric lipid
119 species can be measured but cannot be distinguished by this method.

120 **Statistical analyses**

121 We measured 630 metabolites and calculated a range of sums and ratios of metabolites indicating
122 metabolic pathways and syntheses. Those compounds with the mean below LOD were excluded and
123 described as not measured. Metabolism indicators were calculated with the MetIDQ software and
124 those with more than half of the values below LOD were marked as below LOD in the following. We
125 calculated means and standard deviations and assessed the number of metabolites and their respective
126 classes for each processing method.

127 We described the study population that was used to apply the best protocol and used ANOVA to
128 detect differences in metabolite concentrations between people with different dietary habits
129 (vegetarians, semi-vegetarians, non-vegetarians).

130 A p-value <0.05 (two-sided testing) was considered to indicate statistical significance. Statistical
131 analyses were conducted using SAS Enterprise Guide 7.1 (SAS Institute Inc., Cary, NC, USA).

132 **Results**

133 We measured metabolites with the MxP® Quant 500 kit using 8 different protocols for stool
134 processing from 3 healthy participants of the GEKKO study (free of neoplasms). We were able to
135 extract metabolites with each protocol but the number of detectable metabolites varied (**Table 2**).
136 Most metabolites were extracted (a) using isopropanol (protocol 2) and (b) when we dried the liquid
137 extracts after extraction and reconstituted the samples in a smaller volume to increase metabolite
138 concentrations (protocols 6 and 7).

139 The solvents differ in their polarity and therefore in their extraction efficacy to solve metabolites of the
140 different chemical classes studied. With the MxP® Quant 500 kit, 630 metabolites can be determined
141 of 14 classes of small molecules and 12 lipid classes (**Supplementary Table 1**). In addition, a range
142 of sum and ratios that describe certain pathways and syntheses in the organism is calculated via the
143 MetIDQ software from the obtained data (**Supplementary Table 2**). Amino acids and amino acid
144 related products were detected with concentrations above limit of detection in the analyzed stool
145 samples by all evaluated extraction protocols. Concentrations for amine oxides or carbohydrates and
146 related products were always below the LOD. Major differences were observed for triglycerides as
147 none of the compounds were above the LOD using PBS for preparation whereas more than 50 could
148 be measured using isopropanol or ethanol or methanol in combination with concentrating the liquid
149 extract.

150 Protocol 6 was favorable in terms of sample handling and measured numbers of metabolites.

151 Therefore, we further measured 18 stool samples according to this protocol from participants with
152 different dietary habits (**Table 3**). We analyzed stool samples from 18 vegetarians, semi-vegetarians
153 and non-vegetarians and found a range of metabolites that were significantly different (**Table 4**). Most
154 of the metabolites that distinguished the dietary habits were from lipid classes such as ceramides and
155 phosphatidylcholines. Some metabolism indicators were also found to be different between
156 vegetarians, semi-vegetarians and non-vegetarians. Most of the statistical significant ceramides were
157 higher abundant in non-vegetarians and the sum of ceramides was increasing from vegetarians to non-
158 vegetarians.

159 **Discussion**

160 There is no consensus so far how stool samples should be prepared for comparable, standardized
161 metabolomics measurements (8, 14). We were able to extract a broad range of metabolites with each
162 of the 8 defined protocols tested within this study. However, some methods should be preferred over
163 others in regards to the solvents used dependent on the aim of the study and if a broad or a specific
164 metabolic coverage is aimed for in a particular stool sample analysis. In this study the largest numbers
165 of metabolites could be measured after extraction with isopropanol and ethanol or methanol following
166 a drying step. We have seen differences in the measured metabolite concentration dependent on the
167 extraction method.

168 We observed that the stool processing methods differ, are not interchangeable and that metabolite
169 extraction efficiency varies. The metabolomics panels found by various research groups looking into
170 metabolomics stool sample analysis differ in metabolite composition which might be caused, amongst
171 other reasons, by the different stool processing methods. Studies that have focused on metabolomics in
172 stool samples used either PBS/D₂O buffer (6, 7), acetonitrile (5), methanol (15) or methanol/water
173 mixture (1) for metabolite extraction and each buffer/ solvent results in specific biomarker panel for
174 the specific reagent applied. In this analysis, we used PBS, acetonitrile, methanol and additionally
175 other more complex biphasic (polar/ apolar) solvent combinations for metabolite extraction. The
176 principal procedure is similar for all protocols. A solvent or mixture of solvents is added to the thawed
177 stool samples and this mixture is homogenized and centrifuged so that that supernatant can be used for
178 metabolite analysis (14). We were able to extract a broad range of metabolites with each procedure but
179 the concentrations and the type of metabolites extracted, differ.

180 A total of 630 metabolites and various sums and ratios can be measured or calculated. The typical
181 number of metabolites that can be measured with this kit in human stool is 117 using an ethanol
182 phosphate buffer based protocol (16). When we used a protocol based on PBS, we were able to
183 measure only 88 compounds. Using different methods, we were able to extract and quantify up to 340
184 metabolites from human stool samples. Amino acids, amino acid related metabolites, bile acids, fatty
185 acids, nucleobases and related metabolites amongst others can be reliably measured with almost all

186 solvents. In contrast, the numbers of acylcarnitines, glycerophospholipids or triglycerides are low in
187 human stool which was also found by Wolf et al. (16).

188 In agreement with previously reported differences between fecal metabolomics of omnivores,
189 vegetarians and vegans (17), we observed metabolite differences between vegetarians and non-
190 vegetarians. In particular, we found major differences between stool samples from vegetarians and
191 non-vegetarians in the lipid classes. Non-vegetarians were shown to have higher intake of total fat
192 compared to vegetarians (18). The significant differences in amount of metabolites of lipid classes in
193 non-vegetarians compared to vegetarians found in our study might reflect the difference in fat intake.
194 In meat-eaters, higher blood levels of glycerophospholipids or sphingolipids were found compared to
195 vegetarians or vegans as the most important sources for those metabolites are animal products (19).
196 Furthermore, it was found by different studies that meat intake is associated with the TMAO
197 metabolism as meat and meat products are rich in substances needed for the synthesis of TMAO (20).
198 In contrast, we did not find any differences in fecal TMAO. Other studies found higher amounts of
199 amino acid metabolites and bile acids excretion in urine from meat eaters as they have higher intake of
200 proteins compared to people with high vegetable intake (21). In stool samples, we did not find
201 different amounts of amino acids or bile acids between vegetarians and non-vegetarians.

202 There are some limitations of this study. First, various solvents have different abilities to dissolve and
203 extract the metabolites of different classes and therefore there will be no comparable results when
204 using different processing methods within one study or across several studies. There is a very broad
205 range of chemical classes in metabolomics and not all metabolites can be extracted equally well with
206 the same methods. Second, stool composition varies greatly depending on antibiotic use (22), diet or
207 the water content that can be in a range from 63 to 85% (14). Drying the original stool samples taken
208 out of the freezer before processing offers the advantage of referring to similar actual weights,
209 avoiding bias due to difference in the water content among different samples (23). It is difficult though
210 to imagine study participants to dry their stool samples and then freeze them in small amounts with
211 equal weights for standardization and avoiding freeze-thaw cycles. One would also need to collect
212 more wet weight stool to get enough dry weight stool for a standardized metabolomics analyses. In
213 addition, it is still an obstacle for many people to collect their stool samples for study purposes, freeze

214 them at home and finally transport them in a freezing device to their physician. And even for
215 laboratories, processing stool samples in a consistent way is a challenge also due to its varying
216 consistency. Another point are contaminations and variations introduced into the stool samples that
217 can result from toilet water or from urine, which we tried to minimize using a stool collection aid (24).
218 Further limitations are the small samples size and inclusion of healthy individuals only.

219 A major strength of this study is that we have tested various metabolite extraction protocols on the
220 same analytical platform. Stool samples are very promising in metabolomics research for CRC as they
221 directly represent the microbial activities and the cellular environment in the gut (4). The stool
222 samples were only frozen once which should ensure good metabolite stability: Composition might
223 locally differ since the 50mg were cut off as frozen biomass from the total stool sample without
224 thawing and mixing the complete stool sample.

225 In conclusion, we found a broad range of metabolites measurable in human stool samples. Some
226 chemical classes can be measured equally well with all protocols whereas others are highly dependent
227 on the extraction method. The extraction methods using (a) isopropanol or (b) ethanol or methanol and
228 MTBE including drying of the supernatant seem to be preferable over others for further metabolomics
229 analyses. To our knowledge this is the first study in stool metabolomics comparing 8 different
230 protocols for metabolite extraction with a novel highly standardized and quality controlled,
231 quantitative and reproducible assay and evaluating one methodology, the for our purpose most
232 favorable protocol, to evaluate metabolites in stool samples of participants with different dietary
233 habits.

234 There is urgent need for a consensus on standard procedures for stool processing for metabolomics
235 and for quantitative and reproducible assays to get comparable results across different studies and
236 laboratories.

Acknowledgement

The authors' responsibilities were as follows – HB: planned and designed the GEKKO study; VE, PSK, HB: designed this specific analysis; GP: designed the protocols, VE, GP: processed the samples; HB, PSK: conducted the study; VE: analyzed the data; VE, PSK, HB: drafted the manuscript; all authors critically reviewed the manuscript and approved the final draft. The authors disclose no potential conflict of interests. We received the MxP® Quant 500 Kit (Biocrates Life Sciences AG, Innsbruck, Austria) used in this study free of charge and thank Biocrates for their kind support.

References

1. Phua LC, Chue XP, Koh PK, Cheah PY, Ho HK, Chan EC. Non-invasive fecal metabonomic detection of colorectal cancer. *Cancer biology & therapy* 2014;15(4):389-97. doi: 10.4161/cbt.27625.
2. Rose C, Parker A, Jefferson B, Cartmell E. The Characterization of Feces and Urine: A Review of the Literature to Inform Advanced Treatment Technology. *Crit Rev Environ Sci Technol* 2015;45(17):1827-79. doi: 10.1080/10643389.2014.1000761.
3. Zhgun ES, Iлина EN. Fecal Metabolites As Non-Invasive Biomarkers of Gut Diseases. *Acta naturae* 2020;12(2):4-14. doi: 10.32607/actanaturae.10954.
4. Chetwynd AJ, Ogilvie LA, Nzakizwanayo J, Pazdirek F, Hoch J, Dedi C, Gilbert D, Abdul-Sada A, Jones BV, Hill EM. The potential of nanoflow liquid chromatography-nano electrospray ionisation-mass spectrometry for global profiling the faecal metabolome. *Journal of chromatography A* 2019;1600:127-36. doi: 10.1016/j.chroma.2019.04.028.
5. Amiot A, Dona AC, Wijeyesekera A, Tournigand C, Baumgaertner I, Lebaleur Y, Sobhani I, Holmes E. (1)H NMR Spectroscopy of Fecal Extracts Enables Detection of Advanced Colorectal Neoplasia. *Journal of proteome research* 2015;14(9):3871-81. doi: 10.1021/acs.jproteome.5b00277.
6. Bezabeh T, Somorjai R, Dolenko B, Bryskina N, Levin B, Bernstein CN, Jeyarajah E, Steinhart AH, Rubin DT, Smith IC. Detecting colorectal cancer by 1H magnetic resonance spectroscopy of fecal extracts. *NMR in biomedicine* 2009;22(6):593-600. doi: 10.1002/nbm.1372.
7. Lin Y, Ma C, Liu C, Wang Z, Yang J, Liu X, Shen Z, Wu R. NMR-based fecal metabolomics fingerprinting as predictors of earlier diagnosis in patients with colorectal cancer. *Oncotarget* 2016;7(20):29454-64. doi: 10.18632/oncotarget.8762.
8. Deda O, Gika HG, Wilson ID, Theodoridis GA. An overview of fecal sample preparation for global metabolic profiling. *Journal of pharmaceutical and biomedical analysis* 2015;113:137-50. doi: 10.1016/j.jpba.2015.02.006.
9. Orlich MJ, Singh PN, Sabate J, Fan J, Svein L, Bennett H, Knutsen SF, Beeson WL, Jaceldo-Siegl K, Butler TL, et al. Vegetarian dietary patterns and the risk of colorectal cancers. *JAMA internal medicine* 2015;175(5):767-76. doi: 10.1001/jamainternmed.2015.59.
10. Bjerrum JT, Wang Y, Hao F, Coskun M, Ludwig C, Günther U, Nielsen OH. Metabonomics of human fecal extracts characterize ulcerative colitis, Crohn's disease and healthy individuals. *Metabolomics* 2015;11:122-33. doi: 10.1007/s11306-014-0677-3.
11. Ivanisevic J, Zhu ZJ, Plate L, Tautenhahn R, Chen S, O'Brien PJ, Johnson CH, Marletta MA, Patti GJ, Siuzdak G. Toward 'omic scale metabolite profiling: a dual separation-mass spectrometry approach for coverage of lipid and central carbon metabolism. *Analytical chemistry* 2013;85(14):6876-84. doi: 10.1021/ac401140h.
12. Lee DY, Kind T, Yoon YR, Fiehn O, Liu KH. Comparative evaluation of extraction methods for simultaneous mass-spectrometric analysis of complex lipids and primary metabolites from human blood plasma. *Analytical and bioanalytical chemistry* 2014;406(28):7275-86. doi: 10.1007/s00216-014-8124-x.
13. Biocrates Life Sciences AG. Analysis of Human Fecal Samples with the MxP(R) Quant 500 Kit. 2019.

14. Karu N, Deng L, Slae M, Guo AC, Sajed T, Huynh H, Wine E, Wishart DS. A review on human fecal metabolomics: Methods, applications and the human fecal metabolome database. *Analytica chimica acta* 2018;1030:1-24. doi: 10.1016/j.aca.2018.05.031.
15. Goedert JJ, Sampson JN, Moore SC, Xiao Q, Xiong X, Hayes RB, Ahn J, Shi J, Sinha R. Fecal metabolomics: assay performance and association with colorectal cancer. *Carcinogenesis* 2014;35(9):2089-96. doi: 10.1093/carcin/bgu131.
16. Wolf B, Heischmann S, Dearth S, Koal T. Spotlight. The MxP Quant 500 Kit. *MetaboNews* 2019:4-9.
17. De Filippis F, Pellegrini N, Vannini L, Jeffery IB, La Stora A, Laghi L, Serrazanetti DI, Di Cagno R, Ferrocino I, Lazzi C, et al. High-level adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated metabolome. *Gut* 2016;65(11):1812-21. doi: 10.1136/gutjnl-2015-309957.
18. Rizzo NS, Jaceldo-Siegl K, Sabate J, Fraser GE. Nutrient profiles of vegetarian and nonvegetarian dietary patterns. *J Acad Nutr Diet* 2013;113(12):1610-9. doi: 10.1016/j.jand.2013.06.349.
19. Schmidt JA, Rinaldi S, Ferrari P, Carayol M, Achaintre D, Scalbert A, Cross AJ, Gunter MJ, Fensom GK, Appleby PN, et al. Metabolic profiles of male meat eaters, fish eaters, vegetarians, and vegans from the EPIC-Oxford cohort. *Am J Clin Nutr* 2015;102(6):1518-26. doi: 10.3945/ajcn.115.111989.
20. Guasch-Ferré M, Bhupathiraju SN, Hu FB. Use of Metabolomics in Improving Assessment of Dietary Intake. *Clinical chemistry* 2018;64(1):82-98. doi: 10.1373/clinchem.2017.272344.
21. Wei R, Ross AB, Su M, Wang J, Guiraud S-P, Draper CF, Beaumont M, Jia W, Martin F-P. Metabotypes Related to Meat and Vegetable Intake Reflect Microbial, Lipid and Amino Acid Metabolism in Healthy People. *Molecular Nutrition & Food Research* 2018;62(21):1800583. doi: 10.1002/mnfr.201800583.
22. Pérez-Cobas AE, Gosalbes MJ, Friedrichs A, Knecht H, Artacho A, Eismann K, Otto W, Rojo D, Bargiela R, von Bergen M, et al. Gut microbiota disturbance during antibiotic therapy: a multi-omic approach. *Gut* 2013;62(11):1591-601. doi: 10.1136/gutjnl-2012-303184.
23. Deda O, Chatziioannou AC, Fasoula S, Palachanis D, Raikos N, Theodoridis GA, Gika HG. Sample preparation optimization in fecal metabolic profiling. *Journal of chromatography B, Analytical technologies in the biomedical and life sciences* 2017;1047:115-23. doi: 10.1016/j.jchromb.2016.06.047.
24. Wu WK, Chen CC, Panyod S, Chen RA, Wu MS, Sheen LY, Chang SC. Optimization of fecal sample processing for microbiome study - The journey from bathroom to bench. *Journal of the Formosan Medical Association = Taiwan yi zhi* 2019;118(2):545-55. doi: 10.1016/j.jfma.2018.02.005.

Tables

Table 1. Overview on the tested stool protocols

Protocol 1	Protocol 2	Protocol 3	Protocol 4	Protocol 5	Protocol 6	Protocol 7	Protocol 8
50 mg stool	50 mg stool	50 mg stool	50 mg stool	50 mg stool	50 mg stool	50 mg stool	50 mg stool
150µL PBS	200µl isopropanol	200µl Methanol/ acetonitril/ H ₂ O ¹ (2/2/1)	200µl Methanol/ Acetonitril (1/1)	150µl 85% Ethanol/ 15% 20mM phosphate buffer	200µl 75% Ethanol	225µl 100% Methanol	150µl 100% methanol
vortex 2 min							
						Freeze in liquid N ₂ (1min); thaw	
5 min sonification on ice							
					500µl MTBE	750µl MTBE	
					shake 1 h at RT	shake 1h at 4°C	
						188µl H ₂ O ¹ + 0.1% ammonium acetate	
						vortex 2 min, incubate 5 min at RT	
				Centrifugation at full speed (5 min)		Centrifugation at full speed (10 min)	Centrifugation at full speed (5 min)
				supernatant		Upper supernatant	supernatant
				150µl 20% Ethanol/ 80% 20mM phosphate buffer	125µl H ₂ O ¹	~2000µl 100% methanol	150µl 20% methanol
					vortex 2 min	vortex 1 min	
					incubate 10 min at RT	Incubate 1h at -20°C	
Centrifugation at full speed at 4°C (15 min) + supernatant in extra tube							
Freeze in liquid nitrogen	Freeze in liquid nitrogen	Freeze in liquid nitrogen	Freeze in liquid nitrogen	Freeze in liquid nitrogen	Dry complete in SpeedVac	Dry complete in SpeedVac	Freeze in liquid nitrogen

¹ DNase and RNase free water

Abbreviations: MTBE, methyl tert-butyl ether; PBS, phosphate buffered saline; RT, room temperature.

Table 2. Number of metabolites that were measured with each protocol according to chemical class

Protocol	Total	1	2	3	4	5	6	7	8
analyte class									
Total	630	131	251	100	149	132	303	342	137
Alkaloids	1	1	1	1	1	1	1	1	1
Amine Oxides	1	0	0	0	0	0	0	0	0
Amino Acids	20	20	19	19	19	19	20	20	19
Amino acid related	30	25	22	23	23	22	26	25	23
Bile Acids	14	11	14	14	14	14	14	14	14
Biogenic Amines	9	8	6	6	6	6	7	7	6
Carbohydrates and related	1	1	0	0	0	0	1	0	0
Carboxylic Acids	7	4	2	3	2	2	3	4	2
Cresols	1	1	1	1	1	1	1	1	1
Fatty Acids	12	3	9	7	9	7	10	10	7
Hormones and related	4	1	2	2	2	1	2	2	2
Indoles and Derivatives	4	4	3	4	4	2	4	3	4
Nucleobases and related	2	2	2	2	2	2	2	2	2
Vitamins and Cofactors	1	1	1	1	1	1	1	1	1
Acylcarnitines	40	6	2	2	2	2	4	6	2
Glycerophospholipide (Lysophosphatidylcholines and Phosphatidylcholines)	90	5	25	7	18	10	27	38	10
Sphingomyelins	15	3	5	1	3	3	6	7	3
Cholesteryl Esters	22	3	3	0	0	0	4	6	0
Ceramides	28	0	22	0	15	10	24	26	6
Dihydroceramides	8	0	2	0	0	0	1	2	0
Glycosylceramides(Mono-, Di-, and Trihexosylceramides)	34	1	17	0	7	7	24	22	4
Diglycerides	44	6	14	4	11	5	17	18	4
Triglycerides	242	25	79	3	9	17	104	127	26
Metabolism indicators	232	66	72	85	78	86	99	96	84

Table 3. Population characteristics for the GEKKO participants analyzed with protocol 6 by dietary habits

Characteristics	Vegetarians N=6	Semi-vegetarians N=6	Non-vegetarians N=6
Sex, n (%)²			
Female	3	3	3
Male	3	3	3
Age, n (%)²			
50-54 years	1	2	2
55-59 years	4	2	4
60-64 years	1	2	0
Mean, (SD)	56.7 (\pm 2.9)	56.5 (\pm 3.6)	54.8 (\pm 2.6)

Table 4. Metabolite concentrations (and SD) and differences between vegetarians, semi-vegetarians and non-vegetarians.

Metabolite	Class	vegetarians		semi-vegetarians		non-vegetarians		ANOVA
		Mean	SD	Mean	SD	Mean	SD	p value
Sum of Cer	Metabolism Indicators	9.05	7.85	15.44	4.88	26.67	7.49	0.0017
Cer(d16:1/23:0)	Ceramides	0.18	0.21	0.34	0.06	0.56	0.14	0.0019
Sum of LCFA-Cer	Metabolism Indicators	4.82	4.69	8.62	2.57	16.16	5.74	0.0019
Cer(d18:2/18:0)	Ceramides	0.01	0.03	0.12	0.07	0.30	0.19	0.0028
Cer(d16:1/22:0)	Ceramides	0.37	0.32	0.53	0.15	0.96	0.25	0.0031
Cer(d18:1/18:1)	Ceramides	0.23	0.17	0.16	0.07	1.06	0.73	0.0046
Cer(d18:0/22:0)	Dihydroceramides	0.14	0.20	0.15	0.17	0.57	0.26	0.0049
1-Met-His Synthesis	Metabolism Indicators	6.29	3.76	0.83	1.37	0.39	0.31	0.0058
Ratio of Short-Chain to Long-Chain ACs	Metabolism Indicators	0.95	0.32	0.80	0.29	0.46	0.07	0.0058
Cer(d18:2/24:0)	Ceramides	0.04	0.06	0.10	0.11	0.21	0.05	0.0061
Sum of VLCFA-Cer	Metabolism Indicators	4.24	3.32	6.81	2.66	10.53	2.69	0.0067
PC ae C36:5	Phosphatidylcholines	0.03	0.04	0.06	0.05	0.21	0.14	0.0071
Sum of VLCFA-DH-Cer	Metabolism Indicators	0.65	0.24	1.11	0.55	2.35	0.95	0.0075
SM C26:0	Sphingomyelins	0.02	0.02	0.01	0.01	0.05	0.02	0.0080
PEA	Biogenic Amines	0.15	0.14	0.18	0.19	0.97	0.74	0.0096
CE(20:1)	Cholesteryl Esters	1.14	0.71	0.12	0.30	0.22	0.54	0.0098
Cer(d18:1/22:0)	Ceramides	0.56	0.37	0.80	0.34	1.30	0.41	0.0115
CE(14:1)	Cholesteryl Esters	0.07	0.17	0.00	0.00	0.44	0.39	0.0147
PEA Synthesis	Metabolism Indicators	0.00	0.00	0.00	0.00	0.01	0.01	0.0157
Cer(d18:0/24:1)	Dihydroceramides	0.18	0.23	0.49	0.23	0.73	0.36	0.0157
TG(17:1_32:1)	Triacylglycerides	0.00	0.00	0.05	0.12	0.23	0.18	0.0158
HexCer(d18:1/24:1)	Hexosylceramides	2.45	1.16	5.48	2.63	7.24	3.29	0.0158
Cer(d16:1/24:0)	Ceramides	0.13	0.23	0.25	0.10	0.43	0.11	0.0178
Sum of VLCFA-Glycosyl-Cer	Metabolism Indicators	6.54	3.48	14.91	6.92	18.24	8.21	0.0200
Cer(d18:1/24:1)	Ceramides	0.89	0.72	1.31	0.59	2.12	0.71	0.0200
Cer(d18:1/18:0(OH))	Ceramides	0.10	0.18	0.19	0.24	0.52	0.29	0.0213
Sum of Glycosyl-Cer	Metabolism Indicators	9.94	5.55	21.58	9.26	26.38	11.98	0.0224
Sum of HexCer	Metabolism Indicators	7.26	4.18	16.20	7.72	20.00	8.90	0.0227
HexCer(d18:1/26:0)	Hexosylceramides	0.21	0.14	0.41	0.17	0.52	0.22	0.0291
TG(18:0_32:0)	Triacylglycerides	1.79	0.21	1.46	0.27	1.34	0.32	0.0302
Cer(d18:2/16:0)	Ceramides	0.17	0.09	0.28	0.15	0.39	0.15	0.0308
DG(17:0_18:1)	Diglycerides	1.59	0.76	1.37	1.20	2.93	0.96	0.0316
Cer(d18:1/20:0)	Ceramides	0.11	0.08	0.10	0.06	0.21	0.08	0.0353
C12-DC	Acylcarnitines	1.20	0.36	0.90	0.15	1.42	0.37	0.0355
PC ae C38:6	Phosphatidylcholines	0.03	0.02	0.07	0.04	0.14	0.11	0.0365
Sum of LCFA-Glycosyl-Cer	Metabolism Indicators	3.36	2.16	6.53	2.47	8.05	3.74	0.0366
Cer(d18:1/24:0)	Ceramides	0.54	0.44	0.97	0.52	1.37	0.52	0.0368
Cer(d18:1/18:0)	Ceramides	0.38	0.27	0.55	0.31	2.76	2.76	0.0378
TG(18:1_32:2)	Triacylglycerides	1.37	1.49	3.03	1.71	3.58	0.78	0.0381
PC aa C34:4	Phosphatidylcholines	0.07	0.01	0.03	0.03	0.04	0.04	0.0395
Cer(d18:2/18:1)	Ceramides	0.03	0.02	0.04	0.02	0.09	0.05	0.0401
PC ae C38:5	Phosphatidylcholines	0.07	0.03	0.13	0.06	0.17	0.08	0.0405
PC ae C34:3	Phosphatidylcholines	0.02	0.04	0.18	0.20	0.54	0.53	0.0438
CE(18:0)	Cholesteryl Esters	1.13	0.57	0.92	0.41	3.02	2.41	0.0455
Cer(d18:1/25:0)	Ceramides	0.44	0.30	0.75	0.48	1.16	0.55	0.0472
PC ae C44:4	Phosphatidylcholines	0.11	0.01	0.12	0.01	0.13	0.02	0.0498

Abbreviations: C12-DC, Dodecanedioylcarnitine; CE, Cholesteryl ester; Cer, Ceramide, DG, Diglyceride; LCFA, Long-chain fatty acid; PC, Phosphatidylcholine; PEA, Phenylethylamine; SM, Sphingomyelin; TG, Triglyceride.