Integration of Kalman filter in the epidemiological model: a robust approach to predict COVID-19 outbreak in Bangladesh

Md. Shariful Islam¹, Mohammad Ruhul Amin², Md. Enamul Hoque^{3*},

1 Department of Mathematics and Physics, North South University, Dhaka, Bangladesh

2 Computer and Information Science, Fordham University, New York, USA

3 Computational Physics Group, Department of Physics, Shahjalal University of Science and Technology, Sylhet - 3114, Bangladesh

* mjonyh-phy@sust.edu

Abstract

As one of the most densely populated countries in the world, Bangladesh have been trying to contain the impact of a pandemic like COVID-19 since March, 2020. Although government announced an array of restricted measures to slow down the diffusion in the beginning of the pandemic, the lockdown has been lifted gradually by reopening all the industries, markets and offices with a notable exception of educational institutes. As the physical geography of Bangladesh is highly variable across the largest delta, the population of different region and their life style also differ. Thus, to get the real scenario of the current pandemic across Bangladesh, it is essential to analyze the transmission dynamics over the individual districts. In this article, we propose to integrate the Unscented Kalman Filter (UKF) with classic SIRD model to explain the epidemic evolution of individual districts in the country. We show that UKF-SIRD model results in a robust prediction of the transmission dynamics for 1-4 months. Then we apply the robust UKF-SIRD model over different regions in Bangladesh to estimates the course of the epidemic. Our analysis demonstrate that in addition to the densely populated areas, industrial areas and popular tourist spots are in the risk of higher COVID-19 transmission. In the light of these outcomes, we provide a set of suggestions to contain the pandemic in Bangladesh. All the data and relevant codebase is available at https://mjonyh.github.io.

1 Introduction

The world has been braving the COVID-19 crisis by which more than 22 million people across 113 countries over the globe got infected as of *August* 30, 2020. The Corona virus disease was first identified in Wuhan, the capital of Central China's Hubei province, on December 2019. The World Health Organization (WHO) declared it as the public health emergency on January 20, 2020 and later as Pandemic on March 11, 2020 [1,2]. Although China, Italy, Spain, USA and UK were severely affected by COVID-19 at the beginning, beginning from April 2020 onward, countries from Global South are also impacted by this devastating Corona virus. As of *August* 30, 2020, the lower middle income country Bangladesh has the second lowest number of tests conducted per confirmed cased of COVID-19.

The novel Corona virus can be transmitted via three commonly accepted modes, namely direct contact with an infected person, contaminated objects, and larger

2

10

11

12

> respiratory droplets or small airborne droplets [3]. Usually the virus infected person shows symptoms like fever, shortness in breath, cough, diarrhea, pneumonia and loss of smell, whereas the corona positive patient with premedical conditions such as diabetics, pneumonia and high blood pressure might experience the fatal death if proper medical treatment cannot be sought for in time [4,5]. Since there are no recommended medicines or remedies available for treating COVID-19 disease as of now, many countries are setting up an array of measures including wearing face mask, early detection of infected people, contact tracing, isolation of infected people, increasing the number of testing per day, and imposing lockdown to curb the COVID-19 outbreak [6,7]. Recent studies also support the effectiveness of wearing face mask, social distancing, and imposing lockdown on the declining of both infection and death rate [8].

> Bangladesh is one of the most densely populated (ranked 8^{th} worldwide) and under developing countries in the world. The recent economic development of this country makes it very competent in the South Asian region, especially in terms of foreign exchange reserves, reaching as high as second position in South Asia [9]. The first COVID-19 positive case was identified in Bangladesh on March 8, 2020. The Government of Bangladesh imposed a set of strict measures and closed all educational institutions, government, non-government offices including garments and factories [10] on March 26th, 2020. As a result, the number of infection and fatality rates are less compared to India and Pakistan. In Bangladesh, the contamination numbers spilled over 275,000 confirmed cases with death toll of more than 3,500 and it is still growing each day [11].

> To understand the dynamics of a pandemic, mathematical model can play a major role. The classic mean-field Susceptible-Infected-Recovery-Death (SIRD) model by Kermack and McKendrick [10,12], is frequently used to illustrate a quantitative picture of COVID-19 outbreak for many countries [6]. However, spread of an epidemic is a sophisticated process that depends largely on the mutated strains of the virus and its principle vectors [13,14]. Moreover, the recorded data of infected, recovered and death cases might miss significant amount of information due to many kinds of biases [15]. For instance, time series of recovered numbers are heavily unreliable as it is solely based on how a country traces its asymptomatic or mildly symptomatic patients. Therefore, usual mean-field, compartment-based models and stochastic spatial epidemic models might estimate the dynamics of an epidemic with higher error margin [16–19].

> The SIRD model is associated with a set of differential equations which compute values for an instantaneous event, whereas the time series of COVID-19 cases that we observe everyday is a discrete event. Therefore, all the daily events of COVID-19 cases lack the instantaneous effect in the differential equation based SIRD model. Here, we use the prediction based Unscented Kalman Filter (UKF) model to derive the dynamics. UKF is a classic non-linear estimation algorithm that accurately and timely forecasts the dynamic state in a nonlinear system. UKF has been used in various areas such as, navigation, target tracking, structural dynamics and vehicle positioning due to its high accuracy and rapid convergence merits [20, 21].

In this study, we integrate the UKF with classic SIRD model to examine the underlying process of COVID-19 transmission in a more precise way. As Bangladesh has 64 districts which are uneven in terms of population density, economical significance and geographical locations, we estimate the transmission trend of 64 districts to evaluate the complete picture of the outbreak in Bangladesh.

2 Method

We used the standard SIRD model to estimate the number of COVID-19 active cases, recovered cases and death cases in Bangladesh. A homogeneous immunity has been

61

62

63

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

> considered for the whole country with no transmission from animals and no significant difference between natural death and birth. We segmented the total population size, N, into four stages of disease: S, susceptible; I, infected; R, recovered; D, death and can be written by N = S + I + R + D [10, 12, 22, 23]. The interaction between these four stages is illustrated in the Figure 1.



Fig 1. Schematic diagram of the compartmental SIRD model. It represents the total populations (N) into four different health status, such as, susceptible (S), infected (I), recovered (R) and death (D). The β, γ and μ represents the rate of infection, rate of recovery and rate of death, respectively.

The differential equations of SIRD model can be written as:

$$\frac{dS(t)}{dt} = -\frac{\beta}{N}S(t)I(t) \tag{1}$$

$$\frac{dI(t)}{dt} = \frac{\beta}{N}S(t)I(t) - \gamma I(t) - \mu I(t)$$
(2)

$$\frac{dR(t)}{dt} = \gamma I(t) \tag{3}$$

$$\frac{dD(t)}{dt} = \mu I(t) \tag{4}$$

Where t denotes the time duration, S(t) represents the number of susceptible people at time t, I(t) shows the number of people infected at time t, R(t) stands for the number of people who have recovered from the infection and D(t) indicates the number of deaths. The population, N = S(t) + I(t) + R(t) + D(t), is a conserved quantity for every time step [24, 25]. The constant β , γ and μ illustrate the rate of infection, recovery and deaths, respectively.

The Eq. 1 - 4 have been discretized by forward Euler method as,

$$S_{t+1} = (1 - \beta I_t \Delta t/N) S_t \tag{5}$$

$$I_{t+1} = (1 - \gamma \Delta t + \beta S_t \Delta t/N)I_t - \mu D_t \Delta t \tag{6}$$

$$R_{t+1} = \gamma I_t \Delta t \tag{7}$$

$$D_{t+1} = \mu I_t \Delta t \tag{8}$$

An augmented state vector X is presented for simplicity as,

$$X_t = \begin{bmatrix} S_t \\ I_t \\ R_t \\ D_t \end{bmatrix}$$
(9)

69

70

71

72

73

74

75

76

and the discrete-time augmented SIRD model, Eq. 5 - 8, can be written as,

$$X_{t+1} = F(X_t) + W_t$$
 (10)

where F is the nonlinear term and W is the zero mean Gaussian uncertainty with covariance Q_f . All types of reported cases, the cumulative confirmed, recovered, deaths and active cases, are incorporated with the model using the vector as,

$$Y_{t+1} = CX_t + \nu_t \tag{11}$$

where ν is the uncertainties due to the SERS-CoV-2 test results. The considered zero mean Gaussian uncertainty has the known covariance R_f .

Let us consider an estimated vector state \hat{X}_t for the Unscented Kalman Filter (UKF), then,

$$F(X_t) = F(\hat{X}_t) + J_F(\hat{X}_t)(X_t - \hat{X}_t)$$
(12)

where $J_F(\hat{X}_t)$ is the Jacobian matrix of $F(X_t)$, given by,

$$J_F(\hat{X}_t) = \begin{bmatrix} 1 + (\beta S_t / N - \gamma - \mu) \Delta_t & 0 & 0 & 0 & 0 & 0 & 0 \\ \gamma \Delta_t & 1 & 0 & 0 & 0 & 0 & 0 \\ \mu \Delta_t & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$
(13)

Hence the algorithm of UKF is given by,

Predict:

$$\hat{X}(t+1|t) = F(\hat{X}(t|t))$$
(14)

$$P(t+1|t) = J_F(\hat{X}(t|t))P(t|t)J_F(\hat{X}(t|t))^T + Q_F(t)$$
(15)

Update:

$$\hat{Y}(t+1) = Y(t+1) - C\hat{X}(t+1|t)$$
(16)

$$K(t+1) = P(t+1|t)C^{T}(CP(t+1|t)C^{T} + R_{F}(t))^{-1}$$
(17)

$$\hat{X}(t+1|t+1) = \hat{X}(t+1|t) + K(t+1)\hat{Y}(t+1)$$
(18)

$$P(t+1|t+1) = (I - K(t+1)C)P(t+1|t)$$
(19)

Here, P(t|t) and C represent the posterior estimate covariance matrix and the data augmented matrix respectively. We integrated this updated formula of UKF in SIRD model to estimate the dynamics of the SERS-CoV-2 in Bangladesh.

The dynamical behaviour of the class of infected people is described by the basic (R_0) , effective (R_e) and time-varying (R_t) reproduction number. In the case of SIRD model, this are defined as,

$$R_0 = \frac{\beta}{\mu + \gamma} \tag{20}$$

$$R_e(t) = R_0 \frac{S(t)}{N} \tag{21}$$

Using Eq. 2, one can find that $S(t) = N/R_0$ and $R_e = 1$, at $I = I_{max}$. Besides, one can estimate the onset confirmed (C') cases as, $C' = Ce^{\gamma'(R_t-1)} = Ce^{\theta}$, where $\theta = \gamma'(R_t - 1)$ observes random walk and γ' vary independently [26]. Thus,

$$R_t = \frac{\theta}{\gamma'} + 1 \tag{22}$$

86

78

79

80

81

82

83

84

85

88

96 97

98

99

93

If this reproduction number becomes larger than the ratio between the total population and initial susceptible people, there will be a proper epidemic outbreak [27]. Otherwise, the disease will never lead to a proper outbreak.

3 Results



Fig 2. The estimation of corona active cases in Bangladesh by UKF-SIRD and SIRD model. Dynamics of the active cases population has been shown as a function of time. The real data are fitted from March 8 to July 28, 2020.

As a state of art, we first evaluate whether UKF-SIRD model provides more robust 107 estimation of COVID-19 active cases in Bangladesh compared to classic SIRD model 108 prediction. To illustrate the active cases of the outbreak, the UKF-SIRD and SIRD 109 model simulations were performed in the GNU octave (Version 5.2.0) and Python 110 programming language environment. It is clearly noticeable in figure 2 that UKF-SIRD 111 model delivers more accurate data estimation compared to normal SIRD model 112 prediction. For instance, the UKF-SIRD model predicts around 85,000 corona active 113 cases at 100 days (third week of June, 2020) relative to 90,000 real cases, whereas SIRD 114 model estimates 120,000 active cases at the same time in Bangladesh. Therefore, these 115 results illustrate the fact that integrated UKF in SIRD model is a robust method which 116 is capable of exclusive prediction of COVID-19 transmission in Bangladesh. 117

Bangladesh has 64 districts that are heterogeneous in the context of different population densities, socioeconomic activities and geographical parameters as shown in figure 3. For example, Dhaka, the capital of Bangladesh, has a population density of 23,234 people/sq. km compared to Bandarban (87 people/sq. km) district in Bangladesh. On the other hand, Narayanganj has more industries than Bogra district. Recently, Nadjat and colleagues showed the positive impact of population density on COVID-19 diffusion in Algeria [32]. Thus, considering the heterogeneous density in Bangladesh, general estimation of the COVID-19 dynamics will never match with the real outbreak scenario unless it is done on the district level. To evaluate the local transmission in Bangladesh, next we cluster whole Bangladesh into three groups according to its population density, industrial area and geographical variance.



123

124

125

127

128

118

5/13

106

103

104



Fig 3. Classification of Bangladesh map based on population density, industrial regions, agricultural land, high land, Beel and Haor

3.1 Estimation of COVID-19 diffusion in highly dense cities of 129 Bangladesh 130

In this article, we consider 12 districts (high to low dense areas) of Bangladesh to 131 examine the spread of the Corona virus infection. Fig. 3.1 clearly shows that densely 132 populated area, Dhaka is severely affected by COVID-19 outbreak as the confirmed 133 cases reach $\sim 100,000$, whereas low dense area, Bandarban has only $\sim 1,000$ positive 134 cases. In addition, the daily growth rate also follows a high trajectory trend for Dhaka 135 compared to Narsingdi, Shariatpur and Bagerhat, as shown in the Fig. 3.1. The R_t 136 values of Dhaka were over 1.0 starting from April, 2020 to end of July, 2020, whereas 137 Bagerhat had the R_t values > 1.0 for only 30 days (June, 2020). Interestingly, we notice 138 that the district of Cox's Bazar has similar number of corona positive cases $\sim 10,000$ 139 like highly dense district of Gazipur despite the fact that Cox's Bazar is a moderately 140 dense area in Bangladesh. Since Cox's Bazar is the most popular tourists spot in 141 Bangladesh, we hypothesize that high mobility in this area might cause the high 142 number of corona positive cases. Strikingly, at the end of July, 2020 R_t value is 143 approaching to < 1.0 for all twelve districts in Fig. 3.1, which could lead to the end of 144 this pandemic if the current trend continues for the next 3-4 months. 145



Fig 4. Epidemic evolution of COVID-19 based on confirmed cases, daily growth and R_t values in Bangladesh. The UKF-SIRD model illustrates the positive impact of population densities on COVID-19 confirmed cases and daily growth in major districts such as, Dhaka, Gazipur and Narayanganj.

3.2 COVID-19 transmission trend in the industrial areas of Bangladesh

The year 2020 has seen the unprecedented downfall of the economy for both the national and global market, trade, and investment to reduce the infection rate of the novel Corona virus by ceasing the means of transportation and forcing the lockdown measures that shutdown many business organizations. Moreover, many garment factories were temporarily closed to avoid the risk of Corona virus outbreak. Despite the government-announced shutdown to prevent the COVID-19 diffusion in Bangladesh, industrialists reopened the factories for the economic salvation.

However, health experts have shown their grave concerns about the proper guidelines for social distancing inside the local factories, as several ready made garment (RMG) workers have been tested Corona virus positive in the recent days. Therefore, it is important to determine the COVID-19 transmission trend in these industrial districts such as, Narayanganj, Chattogram, Khulna and Rajshahi. Our systematic UKF-SIRD

156

157

158

159



Fig 5. Epidemic evolution of COVID-19 based on confirmed cases, daily growth and R_t values in Bangladesh. The UKF-SIRD model illustrates the positive impact of industrial zones on COVID-19 confirmed cases and daily growth. High Industrial districts such as Dhaka, Chattogram are at high risk for COVID-19 diffusion compared to low industrial area like, Bogra.

model analysis illustrates a sharp daily growth of corona positive cases for the industrial area, Chattogram, compared to another industrial zone, Narayanganj. Moreover, small industrial region like Rajshahi has less number of cases ($\sim 1,000$) relative to Narayanganj ($\sim 10,000$) and Dhaka ($\sim 100,000$). Furthermore, we observe the uncertain values of R_t for Gazipur district, whereas Khulna show a growing trend of R_t from August, 2020. These analysis clearly forecast the fact that high industrial areas are at greater risk of Corona virus infection in Bangladesh.

3.3 COVID-19 outbreak in the crop land, hilly areas and low land of Bangladesh

Bangladesh is known as a predominant agricultural country where more than 50% of the population rely directly or indirectly on employment in the agricultural and fisheries sectors, as indicated by the Food and Agricultural Organization of United Nation, 2020.

168

160

161

> These sectors are playing significant role in national economy by contributing 18% to 172 country's GDP as mentioned in the Bangladesh Economic Review. Thus, next we 173 estimate the COVID-19 cases in several districts of Bangladesh that are highly 174 associated with agricultural and fisheries sectors. The UKF-SIRD model estimates 175 $\sim 1,500$ confirmed cases at the end of October for Noakhali district compared to 176 $\sim 1,000$ cases for Patuakhali and Dinajpur region. In addition, haors (large shallow 177 water body or backswamp) and wetlands area, such as Sunamganj has approximately 178 $\sim 1,200$ cases, whereas Satkhira has ~ 700 Corona virus positive cases in October, 2020. 179 Notably, the model predicts lower number of confirmed cases (< 1,000) and a smaller 180 growth rate for high lands areas namely, Rangamati and Khagrachhari districts. Since 181 these districts also have low R_t values (< 0.9) at the moment, there is a high chance 182 that Rangamati and Khagrachhari will wipe out the COVID-19 outbreak if the current 183 transmission trend continues for next 3-4 months. Taken together, this study represents 184 that the COVID-19 pandemic is less serious in agricultural and high land areas 185 compared to industrial areas in Bangladesh. 186



Fig 6. Epidemic evolution of COVID-19 based on confirmed cases, daily growth and R_t values in Bangladesh. The UKF-SIRD model illustrates that Beel, Haor areas (Sunamganj, Satkhira), Highland districts (Rangamati, Khagrachhari) and argicultural zones (Noakhali, Dinajpur) are at low risk for COVID-19 transmission in Bangladesh.

Discussion

Epidemiological models are the powerful tools that are designed for the mathematical representations of the disease transmission and their associated dynamics [28, 29]. In addition, models can be used to estimate the trend of an epidemic outbreak under certain conditions so that necessary actions can be taken at various scales to restrain the disease diffusion in a community [30]. However, to make a series of policy decisions, it is a critical prerequisite to have a robust epidemiological model that is suitable for the intended purpose and shows accurate and precise prediction of a pandemic [31]. In this study, we integrated UKF with classic SIRD model and estimate COVID-19 diffusion in different parts of Bangladesh.

Dhaka is the 6^th largest densely populated city in the World, and the key financial hub which holds the major portions of the Bangladesh's economy [32,33]. Unfortunately, we observe that Dhaka is the epicenter of the COVID-19 pandemic with the highest number of infected cases due to its high population density in Bangladesh. Although the government has implemented zone-coded lockdown in small areas within the Dhaka city [34], we observe that the mobility between zones could not be maintained strictly due to economic activities. If the government had the capacity to test and isolate each positive cases, only then the zone-coded lockdown system in Bangladesh could have been successful. Several reports show the association of population density with higher rates of transmission, infection, and mortality from COVID-19 [35,36]. Strikingly, our nationwide analysis of COVID-19 dynamics also demonstrate the relationship between population density and the COVID-19 diffusion rates for Bangladesh.

Industrial zones are at greater risk of transmitting the Corona virus as large groups of workers enter and leave the industrial area without maintaining social distance and proper hygiene in most of the cases. As a result, Narayanganj, Dhaka and Chattogram have been braving with COVID-19 for last couple of months. Notably, our model speculates the dire consequences for Chattogram if the COVID-19 diffusion continues with the current trend. Therefore, industrialists from Chattogram must ensure proper guideline for social distancing and the hygiene rules at the industrial areas to save their workers. Moreover, to reduce the public mobility, we suggest that public transportation between high risk areas should also be suspended for a time until the situation gets under control. To support this claim we would like to mention that the high rate of COVID-19 positive cases in the New York city of USA could have been controlled by implementing the proper lockdown and public mobility suspension [37].

According to our analysis, districts with the hilly areas and the wet land areas in Bangladesh show lower rate of infection than elsewhere. Besides the low population density in areas of Bandarban, Khagracchari and Rangamati, the government also control the transportation facilities in the area with some restrictions. On the other hand, the Cox's Bazar, a popular tourist destination in Bangladesh has seen rise in COVID-19 positive cases despite having less population density. So it is evident that lower population density and restricted movements are the keys to slow down the diffusion rate in any areas. Therefore, we suggest to suspend or restrict the mass transportation between densely populated area, and industrial zone in a bit to contain the pandemic in Bangladesh.

Since Bangladesh is in emergency situation, we deal with incomplete and raw data of the current outbreak. For instance, if a person from a family becomes COVID-19 positive, the whole family might turn out to be positive. But, in many families only one of the members were tested for COVID-19 and if positive then counted as one despite the necessity to test for the entire family. Moreover, there were many cases who died with COVID-19 symptoms, but could not be tested, and conversely all the death cases counted in the analysis due to COVID-19 might have died by other diseases. In addition, the recovered cases count only considered those who went through two

10/13

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

negative COVID-19 test results. Therefore, our UKF-SIRD model might not exemplify239the pandemic situation accurately. However, we are optimistic that our data-driven240mathematical modeling can serve as a base model for the policy speculation and241controlling the disease transmission in a heterogeneous region by adding more242sophisticated factors under different environmental and political conditions in243Bangladesh.244

References

- 1. Organization WH, et al. Laboratory testing for coronavirus disease 2019 (COVID-19) in suspected human cases: interim guidance, 2 March 2020. World Health Organization; 2020.
- 2. Organization WH, et al.. Statement on the second meeting of the International Health Regulations (2005) Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV); 2020.
- 3. Organization WH, et al. Transmission of SARS-CoV-2: implications for infection prevention precautions: scientific brief, 09 July 2020. World Health Organization; 2020.
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. The lancet. 2020;395(10223):497–506.
- 5. Figueiredo A, Daponte Codina A, Figueiredo M, Saez M, Cabrera León A. Impact of lockdown on COVID-19 incidence and mortality in China: an interrupted time series study. Bull World Health Organ. 2020;.
- 6. Chatterjee S, Sarkar A, Chatterjee S, Karmakar M, Paul R. Studying the progress of COVID-19 outbreak in India using SIRD model. medRxiv. 2020;.
- Lau H, Khosrawipour V, Kocbach P, Mikolajczyk A, Schubert J, Bania J, et al. The positive impact of lockdown in Wuhan on containing the COVID-19 outbreak in China. Journal of travel medicine. 2020;27(3):taaa037.
- 8. Start TD. Dealing with foreign exchange reserves;.
- 9. Worldometer CC. Worldometer.(2020) 1–22;.
- 10. Kermack WO, McKendrick AG. A contribution to the mathematical theory of epidemics. Proceedings of the royal society of london Series A, Containing papers of a mathematical and physical character. 1927;115(772):700–721.
- 11. Standard TB. Educational institutions to remain closed till March 31;.
- Kermack WO, McKendrick AG. Contributions to the mathematical theory of epidemics. II.—The problem of endemicity. Proceedings of the Royal Society of London Series A, containing papers of a mathematical and physical character. 1932;138(834):55–83.
- 13. Wang Y, Wang Y, Chen Y, Qin Q. Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID-19) implicate special control measures. Journal of medical virology. 2020;92(6):568–576.
- 14. Kupferschmidt K. Why do some COVID-19 patients infect many others, whereas most don't spread the virus at all. Science. 2020;10.

- 15. Carletti T, Fanelli D, Piazza F. COVID-19: The unreasonable effectiveness of simple models. arXiv preprint arXiv:200511085. 2020;.
- Gatto M, Bertuzzo E, Mari L, Miccoli S, Carraro L, Casagrandi R, et al. Spread and dynamics of the COVID-19 epidemic in Italy: Effects of emergency containment measures. Proceedings of the National Academy of Sciences. 2020;117(19):10484–10491.
- Giordano G, Blanchini F, Bruno R, Colaneri P, Di Filippo A, Di Matteo A, et al. Modelling the COVID-19 epidemic and implementation of population-wide interventions in Italy. Nature Medicine. 2020; p. 1–6.
- Chinazzi M, Davis JT, Ajelli M, Gioannini C, Litvinova M, Merler S, et al. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. Science. 2020;368(6489):395–400.
- Balcan D, Colizza V, Gonçalves B, Hu H, Ramasco JJ, Vespignani A. Multiscale mobility networks and the spatial spreading of infectious diseases. Proceedings of the National Academy of Sciences. 2009;106(51):21484–21489.
- 20. Julier SJ, Uhlmann JK. Unscented filtering and nonlinear estimation. Proceedings of the IEEE. 2004;92(3):401–422.
- Zheng B, Fu P, Li B, Yuan X. A robust adaptive unscented Kalman filter for nonlinear estimation with uncertain noise covariance. Sensors. 2018;18(3):808.
- Brauer F, Castillo-Chavez C, Feng Z. Challenges, Opportunities and Theoretical Epidemiology. In: Mathematical Models in Epidemiology. Springer; 2019. p. 507–531.
- Daley DJ, Gani J. Epidemic modelling: an introduction. vol. 15. Cambridge University Press; 2001.
- Anderson RM, May RM. Population biology of infectious diseases: Part I. Nature. 1979;280(5721):361–367.
- May RM, Anderson RM. Population biology of infectious diseases: Part II. Nature. 1979;280(5722):455–461.
- 26. k sys. covid-19: Estimating COVID-19's R_t in Real-Time with PYMC3; 2020. Available from: https://github.com/k-sys/covid-19.
- Hethcote HW. The mathematics of infectious diseases. SIAM review. 2000;42(4):599–653.
- Perez AM, Ward MP, Charmandarián A, Ritacco V. Simulation model of within-herd transmission of bovine tuberculosis in Argentine dairy herds. Preventive veterinary medicine. 2002;54(4):361–372.
- Smith G, Grenfell B, et al. Population biology of pseudorabies in swine. American journal of veterinary research. 1990;51(1):148–155.
- Hoque E, Islam MS, Amin MR, Das SK, Mitra DK. Adjusted Dynamics of COVID-19 Pandemic due to Herd Immunity in Bangladesh. medRxiv. 2020;.
- Sargent RG. Verification and validation of simulation models. In: Proceedings of the 2010 winter simulation conference. IEEE; 2010. p. 166–183.
- 32. Data NS, (SEDAC) AC. Population Statistics;.

- 33. Start TD. Dhaka's economic activities unplanned: analysts;.
- 34. Standard TB. Enforcing zone-based lockdown up in the air;
- 35. Post TW. The United States might have a secret weapon against coronavirus;.
- 36. Times TNY. Density Is New York City's Big 'Enemy' in the Coronavirus Fight;.
- 37. Glaeser EL, Gorback CS, Redding SJ. How much does covid-19 increase with mobility? evidence from new york and four other us cities. National Bureau of Economic Research; 2020.