

1 Gender-affirming care, mental health, and economic stability in
2 the time of COVID-19: a global cross-sectional study of
3 transgender and non-binary people
4
5

6 Brooke A. Jarrett¹, Sarah M. Peitzmeier², Arjee Restar¹, Tyler Adamson³, Sean Howell⁴,
7 Stefan Baral¹, S. Wilson Beckham⁵
8

9 ¹ Department of Epidemiology; Bloomberg School of Public Health,
10 Johns Hopkins University

11 ² Department of Health Behavior and Biological Sciences; School of Nursing;
12 University of Michigan

13 ³ Department of Health, Policy, and Management; Bloomberg School of Public Health;
14 Johns Hopkins University

15 ⁴ Hornet; San Francisco, CA, USA

16 ⁵ Department of Health, Behavior, and Society; Department of International Health;
17 Bloomberg School of Public Health; Johns Hopkins University
18
19
20

21 **CORRESPONDENCE**

22 Brooke A. Jarrett: brooke [at] jhmi [dot] edu
23

24 **FUNDING**

25 BAJ was supported by the National Institute of Mental Health (grant F31MH121128).
26 AJR was supported by the National Institute of Allergy and Infectious Diseases
27 (grant T32AI102623). SWB was supported by the National Institute of Mental Health
28 (grant K01MH114715) and also receives funding from Viiv Healthcare. The funders had no role
29 in study design, data collection and analysis, decision to publish, or preparation of the
30 manuscript.
31

32 **DISCLOSURES**

33 The authors have declared that no competing interests exist.
34

35 **KEYWORDS**

36 Transgender persons; health services for transgender persons; mental health services; social
37 determinants of health; pandemics; COVID-19
38

39 **WORD COUNTS**

40 Abstract: 300

41 Text: 4,517

42 Tables: 5

43 **ABSTRACT**

44

45 **Background**

46 Transgender and non-binary people are disproportionately burdened by barriers to quality
47 healthcare, mental health challenges, and economic hardship. This study examined the impact
48 of the novel coronavirus disease (COVID-19) pandemic and subsequent control measures on
49 gender-affirming care, mental health, and economic stability among transgender and non-binary
50 people globally.

51

52 **Methods**

53 We collected global cross-sectional data from 964 transgender and non-binary adult users of
54 the Hornet and Her apps from April to August 2020 to characterize changes in gender-affirming
55 care, mental health, and economic stability as a result of the COVID-19 pandemic. We
56 conducted Poisson regression models to assess if access to gender-affirming care and ability to
57 live according to one's gender were related to depressive symptoms, anxiety, and changes in
58 suicidal ideation.

59

60 **Results**

61 Individuals resided in 76 countries, including Turkey (27.4%,n=264/964) and Thailand
62 (20.6%,n=205). A majority were non-binary (66.8%,n=644) or transfeminine (29.4%,n=283).
63 Due to the COVID-19 pandemic, 55.0% (n=320/582) reported reduced access to gender-
64 affirming resources, and 38.0% (n=327/860) reported reduced time lived according to their
65 gender. About half screened positive for depression (50.4%,442/877) and anxiety
66 (45.8%,n=392/856). One in six (17.0%,n=112/659) expected losses of health insurance, and
67 77.0% (n=724/940) expected income reductions. The prevalence of depressive symptoms,
68 anxiety, and increased suicidal ideation were 1.63 (95% CI: 1.36-1.97), 1.61 (95% CI: 1.31-

69 1.97), and 1.74 (95% CI: 1.07-2.82) times higher for individuals whose access to gender-
70 affirming resources was reduced versus not.

71

72 **Discussion**

73 The COVID-19 pandemic has reduced access to gender-affirming resources and the ability of
74 transgender and non-binary people to live according to their gender worldwide. These
75 reductions may drive the increased depressive symptoms, anxiety, and suicidal ideation
76 reported in this sample. To improve transgender and non-binary health globally, increased
77 access to gender-affirming resources should be achieved through policies (e.g., digital
78 prescriptions), flexible interventions (e.g., telehealth), and support for existing transgender
79 health initiatives.

80 **INTRODUCTION**

81 The global pandemic caused by the SARS-CoV-2 virus has resulted in more than 37 million
82 cases of novel coronavirus disease (COVID-19) and over one million deaths.¹ In response,
83 countries have implemented a wide range of measures to quell its spread — shelter-in-place
84 orders, closures of business and schools, and the cancellation of surgeries perceived to be
85 elective.² While these interventions are focused on reducing COVID-19 cases and increasing
86 healthcare capacity, they have also negatively affected healthcare access, mental health, and
87 economic stability worldwide.³⁻⁶

88
89 A growing body of literature describes global interruptions to prescriptions for diseases like HIV,
90 increases in depression, and significant losses of job, insurance, and food security as a result of
91 the COVID-19 pandemic.⁷⁻¹¹ However, recent work has demonstrated that these effects
92 disproportionately burden groups that have been historically marginalized. Specifically,
93 pandemic control measures have exacerbated existing health disparities and social inequities
94 along lines of poverty and occupation, race and ethnicity, and sexual orientation.¹²⁻¹⁶ The
95 COVID-19 pandemic may also uniquely affect transgender and non-binary people.

96
97 Prior to the COVID-19 pandemic, transgender and non-binary individuals experienced barriers
98 to care, greater mental health challenges, and economic vulnerabilities caused by stigma,
99 discrimination, and minority stress.¹⁷⁻¹⁹ Transgender and non-binary populations face a scarcity
100 of clinicians trained in gender-affirming practices and widespread transphobia among healthcare
101 staff, both of which make healthcare less accessible.²⁰⁻²² Yet access to gender-affirming
102 healthcare (e.g., chest/breast surgery), services (e.g., hair removal), and goods (e.g., binders
103 and packers) can substantially improve the quality of life and mental health of transgender and
104 non-binary populations, who frequently have elevated levels of depression, anxiety, and suicidal
105 ideation.²³⁻²⁶ A disproportionate number of transgender and non-binary individuals also

106 experience structural vulnerabilities, such as economic, food, and housing insecurity, that can
107 reinforce or worsen barriers to gender-affirming resources and mental health counseling.^{27,28}

108

109 The COVID-19 pandemic is exacerbating these risks among transgender and non-binary
110 individuals. For example, there have been documented cancellations and delays in gender-
111 affirming surgeries;²⁹ such delays and cancellations have previously been connected to
112 negative mental health consequences.^{23,30} Furthermore, many transgender and non-binary
113 individuals who were living according to their gender prior to the COVID-19 pandemic have had
114 to return to living according to their sex assigned at birth upon moving in with relatives.³¹

115 Researchers have also reported on gendered policies from Panama, Peru, and Columbia that
116 attempted to reduce the density of crowds in public places by requiring women and men to
117 access essential services on alternating days — a policy that, like other gender-based laws,
118 would likely result in violence against transgender communities.³² In these situations,
119 transgender and non-binary individuals may be limited in their ability to live safely and
120 comfortably as themselves. The impact of the COVID-19 pandemic is likely to be especially
121 adverse for those who are already economically marginalized, occupying other marginalized
122 identities (e.g., people who are racial or ethnic minorities, living with HIV, or living with
123 disabilities), or both.

124

125 Of the studies to date about the impact of COVID-19 on transgender and non-binary individuals,
126 a majority have been conducted in a single country like the United States and focused on
127 measuring a narrow spectrum of indicators. The objective of this study was to describe the
128 global impact of the COVID-19 pandemic and subsequent control measures on gender-affirming
129 care, mental health, and economic stability among transgender and non-binary individuals. We
130 also examined the association between reduced access to gender-affirming care and ability to
131 live according to one's gender with multiple mental health indicators.

132 **METHODS**

133 **Study Design and Participants**

134 For this cross-sectional study, we used data from the global COVID-19 Disparities Survey
135 distributed between April 16 and August 3, 2020 via Hornet and Her — social networking apps
136 marketed to sexual minorities, both cisgender and transgender. We sent survey invitations to
137 the app-specific inboxes of all users who had been active on their app for at least one year. To
138 take the survey, individuals had to report being 18 years or older, view a consent form, and
139 indicate their informed consent by clicking a button to begin.

140
141 A total of 24,618 individuals began the survey. For these analyses, we included 1,285
142 transgender and non-binary adults, which we defined as people 18 years or older who self-
143 reported being transgender, non-binary, or a gender different than their sex assigned at birth.
144 We excluded all women assigned female or intersex at birth (n=161), all men assigned male or
145 intersex at birth (n=12,740), individuals who did not report a gender (n=9,751), individuals who
146 only reported not knowing, not wishing to disclose, or being unable to disclose their gender
147 (n=654), and non-transgender identifying men and women who did not report an assigned sex
148 at birth (n=27).

149
150 To ensure the quality of our study population, we screened for duplicate survey responses
151 based on IP address, and again by searching for identical responses to twenty randomly
152 selected variables but found none. We also excluded individuals who completed 89% or less of
153 the survey (n=271), who finished in less than the minimum piloted time of seven minutes (n=47),
154 or provided conflicting responses for multiple questions (n=3) for a final sample size of 964. The
155 Johns Hopkins School of Public Health Institutional Review Board provided a Category 4
156 exemption to the larger survey prior to distribution.

157

158 **Demographic Measures**

159 Individuals self-reported gender by choosing any number of the following options: woman, man,
160 transgender man, transgender woman, or non-binary (including gender-diverse, genderqueer,
161 gender nonconforming, gender expansive, and agender). They also self-reported their country
162 of residence, age, socioeconomic status, years of education, ethnic minority and immigration
163 status, access to masks during the COVID-19 pandemic, and whether the government in their
164 area had ever imposed confinement orders (e.g., mobility restrictions to stay-at-home). We
165 categorized countries according to regions defined by the World Health Organization.

166
167 We used eight mutually exclusive categories to describe reported genders. To increase the
168 power for our analyses, we then collapsed individuals into three groups, building on
169 recommendations from Reisner et al: (1) transmasculine, i.e., people who were assigned female
170 at birth (AFAB) or intersex who self-reported being transgender or being a man; (2)
171 transfeminine, i.e., people who were assigned male at birth (AMAB) or intersex who self-
172 reported being transgender or being a woman; and (3) non-binary, i.e., individuals who reported
173 being either solely non-binary, both a man and a woman, or a transgender man and
174 transgender woman.

175
176 We chose to operationalize three gender categories for multiple reasons. First, while some non-
177 binary individuals explicitly reported also being transgender, the majority did not. We wanted to
178 honor this distinction while also allowing for individuals who reported being both men and
179 women to transcend the transfeminine versus transmasculine binary. Second, our survey
180 presented a limited number of gender options to a global audience in which being a third gender
181 (e.g., two-spirit, bissu, fa'afafine, qariwarmi) is distinct from many Western concepts of being
182 transgender³³. Lastly, in line with recommendations from Restar et al., we saw statistically
183 significant differences when comparing non-binary individuals with transmasculine and

184 transfeminine individuals³⁴. Therefore, presenting results stratified by gender (i.e., a “gender-
185 specific” approach) was more appropriate than conducting analyses on all individuals together
186 and presenting them as a single population (i.e., a “gender-inclusive” approach).

187

188 **COVID-19 Impact Measures**

189 Individuals answered questions about the impact of the COVID-19 pandemic and the resulting
190 response on their access to gender-affirming resources, their mental health, and their economic
191 stability. For indicators related to gender-affirming resources, we asked those who self-identified
192 as transgender or non-binary (n=865) whether the COVID-19 pandemic limited their access to
193 the following: hormone therapy and/or medications; surgical aftercare materials (e.g., vaginal
194 dilators); cosmetic supplies and services (e.g. makeup, wigs, and hair removal); mental health
195 counseling and therapy services; and body modifiers (e.g., binders and packing materials); to
196 which they could respond, “Yes,” “No,” or “Not Applicable.” We characterized the severity of
197 interruptions to gender-affirming care by whether individuals reported that more than one
198 resource had been impacted. We also asked whether the COVID-19 pandemic had changed the
199 amount of time that the individual could live according to their gender (“Compared to before the
200 COVID-19 crisis, how often are you able to live according to your gender identity?”). We
201 categorized the five-point Likert scale into three categories: More than before (i.e., more or a lot
202 more as compared to before COVID-19), about the same as before COVID-19, and less than
203 before (i.e., less than or not at all as compared to before COVID-19).

204

205 For mental health indicators, we used the 4-item patient health questionnaire (PHQ-4) to screen
206 for common symptoms of depression and anxiety, which we dichotomized with a score of three
207 or more being considered positive³⁵. We assessed the impact of COVID-19 on loneliness
208 (“Have you been feeling lonely since the COVID crisis began?”) using a four-point Likert Scale,
209 which we dichotomized into a positive sentiment (“not lonely” or “not much lonely”) and negative

210 sentiment (“very much lonely” or “a little lonely”). We also asked how often they had thought
211 about taking their own life presently and in the six months prior to the COVID-19 pandemic with
212 the following answer options: “never,” “seldom,” “quite often,” “very often,” and “all the time.” We
213 created four categories to describe changes from pre- to mid-pandemic: was and remains rare
214 (i.e., “never” or “seldom”), was and remains frequent (i.e., “quite often,” “very often,” or “all the
215 time”), decreased, or increased. To characterize resiliency, we asked about the following in the
216 face of the COVID-19 pandemic: if they had “sources of hope, strength, comfort, and peace”; if
217 they were “intent on finding emotional support and therapy for themselves”; and if they believed
218 that they were able to “live a happy, full life despite the crisis.”

219

220 **Statistical Analyses**

221 We conducted descriptive analyses of demographic measures and presented descriptions of
222 the COVID-19 impact measures stratified by gender, using chi-squared and Fisher's exact tests
223 as appropriate. We also stratified results by country in the supplementary materials. We used
224 Poisson regression models with complete case analyses to assess for differences in the
225 prevalence of screening positive for depression and anxiety among individuals with reduced
226 (versus continued) access to gender-affirming resources and individuals who reported being
227 able to live according to their gender less (versus more) since the COVID-19 pandemic started.
228 We reported these comparisons as prevalence ratios. We used the same approach to assess
229 the impact of reductions in access to care and changes in ability to live according to one's
230 gender on changes in suicidal ideation, stratified on baseline levels of suicidal ideation. We
231 conducted all analyses in Stata version 14.³⁶

232 **RESULTS**

233 These analyses primarily consisted of non-binary (66.8%, n=644/964) and transfeminine
234 (29.4%, n=283) individuals (**Table 1**). About 47% (n=451) were from the European region and
235 25.1% (n=242) were from the South-East Asia region. There were 76 countries represented in
236 the sample; a majority of individuals were residents of Turkey (27.4%, n=264), Thailand (20.6%,
237 n=199), and Russia (11.5%, n=117). No other single country accounted for more than 5% of the
238 sample. Individuals were young and highly educated, with 50.5% (n=487) being between 18 and
239 29 years old and 42.6% (n=410) having at least a university degree. Few (12.7%, n=122) had
240 ever lacked access to a mask during the COVID-19 pandemic and 75.6% (n=729) lived in a
241 country that had issued COVID-related confinement or “stay-at-home” orders.

242
243 More than half (55%, n=320/582) of the sample reported that the COVID-19 pandemic had
244 limited their access to one or more gender-affirming resource (**Table 2**). Mental health
245 counseling and therapy was the most commonly cited resource to be affected (42.9%,
246 n=192/448), with a somewhat greater proportion of transmasculine individuals reporting reduced
247 access to counseling (61.9%, n=13/21) than non-binary (43.0%, n=122/284) and transfeminine
248 (39.9%, n=57/143) individuals (p-value=0.16). Transmasculine and transfeminine individuals
249 were more likely than non-binary individuals to report that the COVID-19 pandemic limited their
250 access to gender-affirming hormones and medications (55.0% [n=11/20] and 42.1% [n=61/145]
251 vs. 30.1% [n=71/236], p-value=0.01) as well as surgical aftercare materials (42.9% [n=6/14]
252 and 40.2% [n=51/127] vs. 28.8% [n=62/215], p-value=0.08). All geographic regions reported
253 reductions in access to gender-affirming resources; at least half of the individuals in each
254 region, beside the Western Pacific, reported reduced access to one or more resource (**S1**
255 **Table**). More than a third (38.0%, n=327/860) of individuals reported that the COVID-19
256 pandemic had reduced or completely eliminated their ability to live according to their gender,
257 with more transfeminine individuals (43.1%, n=100/232) being unable to living according to their

258 gender as compared to transmasculine (28.6%, n=8/28) and non-binary (36.5%, n=219/600)
259 individuals (p-value <0.001).

260
261 About half of the sample screened positive for depression (50.4%, n=442/877) and anxiety
262 (45.8%, n=392/856), with a larger proportion of transfeminine individuals reporting these
263 outcomes than transmasculine and non-binary individuals (**Table 3**). Overall, 10.0% (n=93/928)
264 reported that suicidal ideation had increased during the COVID-19 pandemic, and 12.5%
265 (n=116) reported that it had decreased. Transfeminine individuals were more likely to report
266 increases in suicidal ideation (11.6%, n=31/263) while being less likely to agree with statements
267 of resiliency, such as having sources of hope, strength, comfort, and peace (47.9%, n=100/209)
268 when contrasted with transmasculine (8.3%, n=3/36; 71.0%, n=22/31) and non-binary (9.5%,
269 n=59/624; 69.5%, n=367/528) individuals (p-value=0.98; p-value < 0.001). Sixteen percent
270 (n=136/851) of individuals reported that they were not intent on finding emotional support and
271 therapy for themselves during the COVID-19 pandemic. Seventy-seven percent (n=724/940) of
272 the sample expected a reduction in their income, 17% (n=112/659) expected to lose health
273 insurance, and 53.4% (n=428/801) reported not having received financial aid, despite need
274 (**Table 4**). Though 40% (n=361/900) of individuals overall reported cutting or reducing their
275 meals, fewer non-binary individuals had done so (34.9% [n=212/607] versus 51.4% [n=18/35] of
276 transmasculine individuals and 50.8% [n=131/258] of transfeminine individuals).

277
278 A positive screen for depression was 1.63 (95% confidence interval [CI]: 1.36-1.97) times more
279 common among those who had lost access to one or more gender-affirming resource during the
280 COVID-19 pandemic as compared to those without reductions in access (**Table 5**). Similarly, a
281 positive screen for anxiety was 1.61 (95% CI: 1.31-1.97) times more common among those who
282 had lost access to one or more gender-affirming resource compared to those without reductions
283 in access. Trends were similar for suicidal ideation. For example, among those with rare or no

284 suicidal ideation at the beginning of the COVID-19 pandemic, individuals who had reduced
285 access to gender-affirming resources were 1.74 (95% CI: 1.07, 2.82) times more likely to report
286 increased suicidal ideation. Screening positive for depression and anxiety were also 1.21 (95%
287 CI: 0.92, 1.58) and 1.48 (95% CI: 1.04, 2.10) times more prevalent among those reporting that
288 the COVID-19 pandemic had decreased the amount of time they could live according to their
289 gender, versus those who had increased that time. However, among individuals who had no or
290 rare suicidal ideation prior to the COVID-19 pandemic, a smaller proportion of those living less
291 according to their gender during the COVID-19 pandemic had increased suicidal ideation
292 compared to those who lived more as their gender (prevalence ratio = 0.57 [95% CI: 0.33-
293 0.98]).
294

295 **DISCUSSION**

296 This survey provides early insights into the impacts of the COVID-19 pandemic on access to
297 gender-affirming resources, mental health, and economic stability among transgender and non-
298 binary communities globally. Roughly half of individuals reported that the COVID-19 pandemic
299 had restricted their access to gender-affirming resources, and nearly two in five reported the
300 COVID-19 pandemic had negatively impacted their ability to live according to their gender.
301 Screening positive for depression and/or anxiety and increases in suicidal ideation were
302 common, but more so for those who experienced reduced access to gender-affirming
303 resources. Of these resources, counseling and therapy were the most affected by COVID-19,
304 but most people who responded to the survey also reported resiliencies, such as having
305 sources of hope and being intent on finding emotional support. This intent to seek support,
306 however, may be dampened by the fact that many individuals expected reductions in income
307 and loss of health insurance. This is the first empirical study to examine the effect of COVID-19
308 and its impact on gender-affirming resources, mental health, and economic stability among a
309 global, multi-region sample of transgender and non-binary individuals.

310
311 Half of the individuals in our survey reported reduced access to one or more gender-affirming
312 resource. This was higher in our sample than in an online survey of 1,240 transgender and non-
313 binary individuals in Germany, Switzerland, and Austria.²⁹ A third or more of individuals from the
314 European region in our study reported reductions in access to hormone therapy and surgical
315 aftercare materials, which was higher than the 18% and 3% of individuals from the European-
316 specific survey respectively. That survey also found about a quarter of individuals had delayed
317 or cancelled aftercare for a recent surgery. Differences may be due to the populations sampled,
318 as our survey primarily drew from countries in Eastern Europe where there is substantial stigma
319 against and policing of transgender individuals. Together, though, these results signify that the

320 COVID-19 pandemic is causing decreases in access to gender-affirming resources even in
321 high-income settings.

322
323 Our study demonstrated that reduced access to gender-affirming resources due to the COVID-
324 19 pandemic were associated with poorer mental health. Screening positive for depression,
325 screening positive for anxiety, and increased suicidal ideation were more common for those
326 whose access to one or more gender affirming resource had been reduced due to the COVID-
327 19 pandemic. These results mirror anecdotal data from clinics serving transgender patients in
328 China.³⁷ Gender-affirming resources are crucial to transgender and non-binary individuals, as
329 these resources often activate and enhance the interactive process of receiving recognition for
330 their gender, sense of self, and sense of humanity.³⁸ Given the abundant literature that gender
331 affirmation results in better mental health and quality of life, these data underscore the
332 importance of ensuring that transgender and non-binary individuals have access to these
333 essential services and products.^{23,24,30,39–41} Access to these resources should also be fortified to
334 avoid negative physical health outcomes. For example, about a third of this sample reported
335 reductions in access to hormones and 17% reported that they expected to lose their health
336 insurance, which may support hormone therapy procurement. These disruptions may force
337 some transgender and non-binary individuals to discontinue hormone therapy or mete out their
338 limited doses to last longer. Sustained interruptions or sub-optimal dosing of hormone therapy
339 have been connected with symptoms of hypogonadism, such as osteoporosis and
340 cardiovascular disease.⁴² Similarly, aftercare for gender-affirming surgeries is critical for
341 avoiding negative physical outcomes like urinary tract hesitancy and needing re-operation, yet a
342 third of individuals in this study reported that they had reduced access to the surgical aftercare
343 that they needed.⁴³

344

345 A third of individuals reported decreased time lived according to their gender. In comparison, a
346 poll conducted by the Trevor Project in the United States of 600 lesbian, gay, bisexual,
347 transgender, and questioning (LGBTQ) youth (13-19 years) reported that 56% of the
348 transgender and non-binary individuals had reduced their ability to express their LGBTQ identity
349 due to the COVID-19 pandemic.⁴⁴ Youth often attributed needing to move back in with
350 unsupportive caretakers as a reason for not being able to live according to their gender,^{31,44}
351 which may explain the elevated proportion compared to adults in our study. Living less
352 according to one's gender during the COVID-19 pandemic was associated with screening
353 positive for depression and anxiety. However, a counterintuitive result was found among
354 individuals in our study who had rare suicidal ideation prior to the COVID-19 pandemic and lived
355 more according to their gender during the COVID-19 pandemic. This subset of the sample was
356 more likely to have increased suicidal ideation as compared to individuals who were living
357 according to their gender less. These results, however, may reflect a limitation of our
358 measurement tool. For example, the former group likely had a low baseline ability to live
359 according to their gender pre-pandemic and hence could only increase the amount of time lived
360 according to their gender. The worse mental health in this group may be linked to low pre-
361 pandemic levels of living according to their gender, rather than the recent increase in their ability
362 to do so. Because we did not measure pre-pandemic and current ability to live according to
363 one's gender separately, we could not control for baseline levels in the model. It is also possible
364 that people who recently began living in their gender more may be subjected to elevated levels
365 of anticipated and experienced stigma that could be driving increased suicidality; prior research
366 has shown discrimination due to gender expression, rather than gender identity itself, to be
367 associated with mental distress.⁴⁵⁻⁴⁷
368
369 Positive screens for depression and anxiety were correlated with decreases in access to
370 gender-affirming care and decreased time lived according to one's gender, and were present in

371 nearly half the sample. These data align with results from transgender and non-binary youth
372 from the Trevor Project poll and another sample of 201 young adults (19-25 years) attending
373 college, both from the U.S.^{44,48} Findings in a small study of 15 Latinx trans women in the United
374 States suggest that these poor mental health indicators represent declines as a result of the
375 COVID-19 pandemic rather than just a high baseline prevalence.⁴⁹ These findings are
376 particularly concerning when contrasted against the large proportion of individuals in this and
377 other studies who reported that COVID-19 had decreased their access to mental health therapy
378 and counseling.^{44,48} To mitigate the immediate trauma of the COVID-19 pandemic and potential
379 long-lasting effects, innovative mental health interventions — from remote video and phone
380 therapy to self-help apps — have been emerging, but additional investments are needed,
381 especially to reach those with limited to no access to Internet.^{50,51} Furthermore, our findings
382 support the notion that transmasculine, transfeminine, and non-binary populations are having
383 distinct experiences during the COVID-19 pandemic and should receive gender-specific
384 support.³⁷ For example, transmasculine individuals were more likely to report having reduced
385 access to gender-affirming resources but generally reported better mental health outcomes, as
386 well as having sources of strength and comfort. Combined with the fact that transgender and
387 non-binary youth have been more likely to reach out to friends and family than cisgender
388 lesbian, gay, and pansexual youth according to the Trevor Project data, it may be possible to
389 enhance these resiliencies by strengthening and expanding trauma-informed, online peer-to-
390 peer support efforts such as Q Chat Space ^{31,44,52}. However, different approaches to
391 transfeminine, transmasculine, and non-binary individuals may be needed.

392
393 Lastly, we found that transgender and non-binary communities worldwide are experiencing
394 strains across basic needs such as finances, food, and health insurance, all as a result of
395 COVID-19 pandemic. These strains, in addition to pre-existing economic vulnerabilities, will
396 contribute to even greater barriers to gender-affirming care, mental health counseling, services,

397 and products.²⁷ For example, approximately 10% of transgender and non-binary individuals in
398 the U.S. lacked health insurance before the COVID-19 pandemic.²⁷ Though some may have
399 health insurance through their employers, transgender and non-binary people are also more
400 likely to be employed in the industries most impacted by business shutdowns.¹² We found that
401 more than three quarters of the sample expected a reduction in their income, one in six
402 expected to lose their health insurance, and more than half reported needing and not receiving
403 financial aid. These results display the pronounced structural vulnerabilities that shape
404 experiences of transgender and non-binary communities in the current context of the COVID-19
405 pandemic, and likely contribute further as stressors to mental health. These stressors may climb
406 as transgender and non-binary communities continue to experience worsening economic
407 instability due to the COVID-19 pandemic and associated control measures. Further research is
408 necessary to examine the syndemic impact of COVID-19-related stressors on transgender and
409 non-binary communities, particularly those who are experiencing multiple, intersectional
410 stressors at the individual, interpersonal, and structural levels.

411
412 This study had some limitations. While the survey reached individuals from six continents and
413 was available in 13 languages, these data are not a representative sample of transgender and
414 non-binary individuals worldwide. As a survey distributed through a mobile app, participation
415 was limited to individuals with Internet and a smartphone. Individuals in the survey were also
416 generally highly educated. Our survey likely missed highly disadvantaged individuals, and
417 consequently, likely underestimates the true magnitude of impact of COVID-19 on this
418 community. Our study also had a methodological limitation in its categorization of genders
419 — while we aimed to offer an inclusive set of options, we did not capture the full spectrum nor
420 fluidity of genders across the cultural diversity of the countries represented. Yet, our sample
421 does present results from a large number of non-binary individuals, which is a population that
422 has generally been understudied but increasingly recognized as distinct from binary transgender

423 individuals.⁵³ Furthermore, the survey only presented questions about access to gender-
424 affirming resources to individuals who self-identified as transgender (e.g., versus those who
425 selected being a man and were AFAB but did self-report as transgender). The next wave of data
426 collection will present the module to all persons whose sex assigned at birth does not match
427 their current gender.

428
429 The findings of the current study provide insights for future directions. Namely, future research
430 should expand on this work by identifying protective factors that can be potentially leveraged to
431 buffer the impact of COVID-19 pandemic on gender-affirming resources, mental health, and
432 economic stability. There is also a need to contextualize and understand how transgender and
433 non-binary communities are currently responding to the economic instabilities due to the
434 epidemic, particularly in regions where mandatory stay-at-home orders remain. These
435 restrictions may have led or will lead some transgender and non-binary individuals to turn to
436 more dangerous work for income, unregulated and unmonitored markets for gender-affirming
437 services or goods, or the use of alcohol and other substances to cope. Lastly, given the mobile
438 and online nature of recruitment for this study, researchers should look for feasibility and
439 opportunity with Hornet and other mobile apps as a platform for outreach, programming, and
440 interventions for transgender and non-binary communities in regions where the apps are
441 utilized.

442

443 **CONCLUSION**

444 Along with these research implications, our findings suggest the need for multiple programmatic
445 interventions specific to transgender and non-binary populations. Maintaining and increasing
446 secure access to lifesaving gender-affirming resources, mental health services, and economic
447 stability will require backing from both typical and atypical sources — from nonprofit
448 organizations to for-profit companies to academic researchers — both during and after the

449 COVID-19 pandemic. This could include, for example, instrumental support according to
450 community needs (e.g., coordinating food bank deliveries or monetary support for bills) or
451 resource mapping to help transgender and non-binary individuals identify where they can seek
452 pandemic-related relief and aid without stigma or discrimination⁵². Health insurers and
453 healthcare facilities could also transparently communicate changes in policies and logistics for
454 gender-affirming services to alleviate anxieties around loss of access due to pandemic control
455 measures. To prevent detrimental mental health consequences due to inaccessibility of gender-
456 affirming resources and economic hardships, rapid policies (e.g., digital prescription refills) and
457 flexible interventions (e.g., telehealth) are needed to maintain continuity of gender-affirming
458 hormones as well as therapy and counseling. To achieve these interventions, innovative
459 partnerships will be needed to reach the most marginalized— both by supporting trans-led,
460 community-based organizations to maintain and expand their transgender health services as
461 well as by increasing the capabilities of nation- and/or state-sponsored programs and private
462 sector companies to better serve transgender and non-binary communities.

463 **Acknowledgements**

464 The authors thank the individuals who volunteered their valuable time to participate in our
465 survey. The authors also thank the people who translated the survey from English into 13
466 different languages: Ana Cara-Linda, Anna Yakusik, Carol Strong, Damiano Cerasuolo, Edward
467 Sutanto, Ezgi Ayaser, Henrique Vicentim. Howie Lim, Ibu Ketty, Jose Garcia, Junming Wu,
468 Ketty Rosenfeld, Luana Araujo, Mariano Ruiz, Maryam Motaghi, Maso, Omar Cherkaoui,
469 Panyaphon Phiphatkunarnon, Pedro Moreno, Poyao Huang, Souad Orhan, Stephane Ku, Tanat
470 Chinbunchorn, Teak Sowapruks, Top Medping, and Yalda Toofan.

Preprint

471 **REFERENCES**

- 472 1. Dong, E., Du, H. & Gardner, L. An interactive web-based dashboard to track COVID-19 in
473 real time. *Lancet Infect. Dis.* **20**, 533–534 (2020).
- 474 2. Hale, T. *et al.* Oxford COVID-19 Government Response Tracker. *Blavatnik School of*
475 *Government* (2020).
- 476 3. Czeisler, M. É. *et al.* Mental Health, Substance Use, and Suicidal Ideation During the
477 COVID-19 Pandemic — United States, June 24–30, 2020. *MMWR. Morb. Mortal. Wkly.*
478 *Rep.* **69**, 1049–1057 (2020).
- 479 4. Salari, N. *et al.* Prevalence of stress, anxiety, depression among the general population
480 during the COVID-19 pandemic: a systematic review and meta-analysis. *Global. Health*
481 **16**, 57 (2020).
- 482 5. Ahmed, S. A. K. S. *et al.* Impact of the societal response to COVID-19 on access to
483 healthcare for non-COVID-19 health issues in slum communities of Bangladesh, Kenya,
484 Nigeria and Pakistan: results of pre-COVID and COVID-19 lockdown stakeholder
485 engagements. *BMJ Glob. Heal.* **5**, e003042 (2020).
- 486 6. U.S. Centers for Disease Control and Prevention. Reduced Access to Care: Household
487 Pulse Survey. *National Center for Health Statistics* (2020). Available at:
488 <https://www.cdc.gov/nchs/covid19/pulse/reduced-access-to-care.htm>. (Accessed: 1st
489 October 2020)
- 490 7. Rao, D., Kekwaletswe, T. C., Hosek, S., Martinez, J. & Rodriguez, F. Stigma and social
491 barriers to medication adherence with urban youth living with HIV. *AIDS Care - Psychol.*
492 *Socio-Medical Asp. AIDS/HIV* **19**, 28–33 (2007).
- 493 8. Ettman, C. K. *et al.* Prevalence of Depression Symptoms in US Adults Before and During
494 the COVID-19 Pandemic. *JAMA Netw. Open* **3**, e2019686 (2020).
- 495 9. Niles, M. T. *et al.* The Early Food Insecurity Impacts of COVID-19. *Nutrients* **12**, 2096
496 (2020).
- 497 10. Nicola, M. *et al.* The socio-economic implications of the coronavirus pandemic (COVID-
498 19): A review. *Int. J. Surg.* **78**, 185–193 (2020).
- 499 11. Banthin, J., Simpson, M., Buettgens, M., Blumberg, L. J. & Wang, R. *Changes in Health*
500 *Insurance Coverage Due to the COVID-19 Recession: Preliminary Estimates Using*
501 *Microsimulation.* (2020).
- 502 12. Whittington, C., Hadfield, K. & Calderón, C. *The Lives & Livelihoods of Many in the*
503 *LGBTQ Community are at Risk amidst the COVID-19 Crisis.* (2020).
- 504 13. Aldridge, R. W. *et al.* Black, Asian and Minority Ethnic groups in England are at increased
505 risk of death from COVID-19: indirect standardisation of NHS mortality data. *Wellcome*
506 *Open Res.* **5**, 88 (2020).
- 507 14. Bui, D. P. *et al.* Racial and Ethnic Disparities Among COVID-19 Cases in Workplace
508 Outbreaks by Industry Sector — Utah, March 6–June 5, 2020. *MMWR. Morb. Mortal.*
509 *Wkly. Rep.* **69**, 1133–1138 (2020).

- 510 15. Millett, G. A. *et al.* Assessing differential impacts of COVID-19 on black communities.
511 *Ann. Epidemiol.* **47**, 37–44 (2020).
- 512 16. Santos, G.-M. *et al.* Economic, Mental Health, HIV Prevention and HIV Treatment
513 Impacts of COVID-19 and the COVID-19 Response on a Global Sample of Cisgender
514 Gay Men and Other Men Who Have Sex with Men. *AIDS Behav.* (2020).
515 doi:10.1007/s10461-020-02969-0
- 516 17. Poteat, T. C., Reisner, S. L., Miller, M. & Wirtz, A. L. Letter to the Editor: Vulnerability to
517 COVID-19-related Harms Among Transgender Women With and Without HIV Infection in
518 the Eastern and Southern U.S. *JAIDS J. Acquir. Immune Defic. Syndr.* **Publish Ah**,
519 (2020).
- 520 18. Reisner, S. L. *et al.* Global health burden and needs of transgender populations: a
521 review. *Lancet* **388**, 412–436 (2016).
- 522 19. Newcomb, M. E. *et al.* High Burden of Mental Health Problems, Substance Use,
523 Violence, and Related Psychosocial Factors in Transgender, Non-Binary, and Gender
524 Diverse Youth and Young Adults. *Arch. Sex. Behav.* **49**, 645–659 (2020).
- 525 20. Casey, L. S. *et al.* Discrimination in the United States: Experiences of lesbian, gay,
526 bisexual, transgender, and queer Americans. *Health Serv. Res.* **54**, 1454–1466 (2019).
- 527 21. Jarrett, B. A., Corbet, A. L., Gardner, I. H., Weinand, J. D. & Peitzmeier, S. M. Chest
528 Binding and Care Seeking Among Transmasculine Adults: A Cross-Sectional Study.
529 *Transgender Heal.* **3**, 170–178 (2018).
- 530 22. Peitzmeier, S. M. *et al.* “It Can Promote an Existential Crisis”: Factors Influencing Pap
531 Test Acceptability and Utilization Among Transmasculine Individuals. *Qual. Health Res.*
532 **27**, 2138–2149 (2017).
- 533 23. Wernick, J. A., Busa, S., Matouk, K., Nicholson, J. & Janssen, A. A Systematic Review of
534 the Psychological Benefits of Gender-Affirming Surgery. *Urol. Clin. North Am.* **46**, 475–
535 486 (2019).
- 536 24. Murad, M. H. *et al.* Hormonal therapy and sex reassignment: a systematic review and
537 meta-analysis of quality of life and psychosocial outcomes. *Clin. Endocrinol. (Oxf)*. **72**,
538 214–231 (2010).
- 539 25. Peitzmeier, S., Gardner, I., Weinand, J., Corbet, A. & Acevedo, K. Health impact of chest
540 binding among transgender adults: a community-engaged, cross-sectional study. *Cult.*
541 *Health Sex.* **19**, 64–75 (2017).
- 542 26. Valentine, S. E. & Shipherd, J. C. A systematic review of social stress and mental health
543 among transgender and gender non-conforming people in the United States. *Clin.*
544 *Psychol. Rev.* **66**, 24–38 (2018).
- 545 27. Herman, J. L. & O’Neill, K. *Vulnerabilities to COVID-19 among Transgender Adults in the*
546 *U.S.* (2020).
- 547 28. Snow, A., Cerel, J., Loeffler, D. N. & Flaherty, C. Barriers to Mental Health Care for
548 Transgender and Gender-Nonconforming Adults: A Systematic Literature Review. *Health*
549 *Soc. Work* **44**, 149–155 (2019).

- 550 29. Koehler, A., Motmans, J., te Gueldenning, A. & Nieder, T. O. *The Impact of COVID-19 on*
551 *Transgender Health in German Speaking Countries.*
- 552 30. Bränström, R. & Pachankis, J. E. Reduction in Mental Health Treatment Utilization
553 Among Transgender Individuals After Gender-Affirming Surgeries: A Total Population
554 Study. *Am. J. Psychiatry* **177**, 727–734 (2020).
- 555 31. Fish, J. N. *et al.* “I’m Kinda Stuck at Home With Unsupportive Parents Right Now”:
556 LGBTQ Youths’ Experiences With COVID-19 and the Importance of Online Support. *J.*
557 *Adolesc. Heal.* **67**, 450–452 (2020).
- 558 32. Perez-Brumer, A. & Silva-Santisteban, A. COVID-19 Policies can Perpetuate Violence
559 Against Transgender Communities: Insights from Peru. *AIDS Behav.* **24**, 2477–2479
560 (2020).
- 561 33. Cicero, E. C., Reisner, S. L., Merwin, E. I., Humphreys, J. C. & Silva, S. G. The health
562 status of transgender and gender nonbinary adults in the United States. *PLoS One* **15**,
563 e0228765 (2020).
- 564 34. Restar, A., Jin, H. & Operario, D. Gender-inclusive and gender-specific approaches in
565 trans health research. *Transgender Heal.* (2020).
- 566 35. Kroenke, K., Spitzer, R. L., Williams, J. B. W. & Löwe, B. An Ultra-Brief Screening Scale
567 for Anxiety and Depression: The PHQ–4. *Psychosomatics* **50**, 613–621 (2009).
- 568 36. StataCorp. Stata Statistical Software: Release 14. (2015).
- 569 37. Wang, Y. *et al.* Health care and mental health challenges for transgender individuals
570 during the COVID-19 pandemic. *Lancet Diabetes Endocrinol.* **8**, 564–565 (2020).
- 571 38. Sevelius, J. M. Gender Affirmation: A Framework for Conceptualizing Risk Behavior
572 Among Transgender Women of Color. *Sex Roles* **68**, 675–689 (2013).
- 573 39. Bouman, W. P., Davey, A., Meyer, C., Witcomb, G. L. & Arcelus, J. Predictors of
574 psychological well-being among treatment seeking transgender individuals. *Sex. Relatsh.*
575 *Ther.* 1–17 (2016). doi:10.1080/14681994.2016.1184754
- 576 40. Colizzi, M. & Costa, R. The effect of cross-sex hormonal treatment on gender dysphoria
577 individuals’ mental health: a systematic review. *Neuropsychiatr. Dis. Treat.* **12**, 1953–
578 1966 (2016).
- 579 41. Nguyen, H. B. *et al.* Gender-Affirming Hormone Use in Transgender Individuals: Impact
580 on Behavioral Health and Cognition. *Curr. Psychiatry Rep.* **20**, 110 (2018).
- 581 42. Defreyne, J., Van de Bruaene, L. D. L., Rietzschel, E., Van Schuylenbergh, J. & T’Sjoen,
582 G. G. R. Effects of Gender-Affirming Hormones on Lipid, Metabolic, and Cardiac
583 Surrogate Blood Markers in Transgender Persons. *Clin. Chem.* **65**, 119–134 (2019).
- 584 43. van der Miesen, A. I. R., Raaijmakers, D. & van de Grift, T. C. “You Have to Wait a Little
585 Longer”: Transgender (Mental) Health at Risk as a Consequence of Deferring Gender-
586 Affirming Treatments During COVID-19. *Arch. Sex. Behav.* **49**, 1395–1399 (2020).
- 587 44. Morning Consult. *How COVID-19 is Impacting LGBTQ Youth.* (2020).

- 588 45. Cook, S. H., Sandfort, T. G. M., Nel, J. A. & Rich, E. P. Exploring the Relationship
589 Between Gender Nonconformity and Mental Health Among Black South African Gay and
590 Bisexual Men. *Arch. Sex. Behav.* **42**, 327–330 (2013).
- 591 46. Martin-Storey, A. & August, E. G. Harassment Due to Gender Nonconformity Mediates
592 the Association Between Sexual Minority Identity and Depressive Symptoms. *J. Sex Res.*
593 **53**, 85–97 (2016).
- 594 47. White Hughto, J. M., Reisner, S. L. & Pachankis, J. E. Transgender stigma and health: A
595 critical review of stigma determinants, mechanisms, and interventions. *Soc. Sci. Med.*
596 **147**, 222–231 (2015).
- 597 48. Gonzales, G., Loret de Mola, E., Gavulic, K. A., McKay, T. & Purcell, C. Mental Health
598 Needs Among Lesbian, Gay, Bisexual, and Transgender College Students During the
599 COVID-19 Pandemic. *J. Adolesc. Heal.* (2020). doi:10.1016/j.jadohealth.2020.08.006
- 600 49. Linnemayr, S. *et al.* Longitudinal Assessment of Changes in Mental and Sexual Health
601 Outcomes Due to COVID-19 among Latinx Sexual Minority Men and Transgender
602 Women. *JAIDS J. Acquir. Immune Defic. Syndr.* (2020).
603 doi:10.1097/QAI.0000000000002507
- 604 50. United Nations. *United Nations Policy Brief: COVID-19 and the need for action on mental*
605 *health.* (2020).
- 606 51. Moreno, C. *et al.* How mental health care should change as a consequence of the
607 COVID-19 pandemic. *The Lancet Psychiatry* **7**, 813–824 (2020).
- 608 52. Sevelius, J. M. *et al.* Research with Marginalized Communities: Challenges to Continuity
609 During the COVID-19 Pandemic. *AIDS Behav.* **24**, 2009–2012 (2020).
- 610 53. Motmans, J., Nieder, T. O. & Bouman, W. P. Transforming the paradigm of nonbinary
611 transgender health: A field in transition. *Int. J. Transgenderism* **20**, 119–125 (2019).
- 612

613 **Table 1: Demographics for transgender and non-binary individuals from the COVID-19 Disparities Survey**
 614 **distributed globally by the Hornet and Her apps between April 16 and August 3, 2020 (N=964)**
 615

	Overall (%) ^a
Self-Reported Transgender Identity	
Transgender man	95 (9.9%)
Transgender woman	201 (20.9%)
Non-binary (NB) only	594 (61.6%)
NB transgender woman	17 (1.8%)
NB transgender man	7 (0.7%)
NB, transgender woman, and transgender man	2 (0.2%)
Transgender man and transgender woman	7 (0.7%)
Man and woman	41 (4.3%)
Researcher-Generated Gender Categories	
Transmasculine (assigned female sex or intersex at birth)	37 (3.8%)
Transfeminine (assigned male sex or intersex at birth)	283 (29.4%)
Non-binary ^b	644 (66.8%)
World Health Organization Region	
Europe	451 (46.8%)
South-East Asia	242 (25.1%)
Americas	86 (8.9%)
Eastern Mediterranean	85 (8.8%)
Western Pacific	40 (4.2%)
Africa	35 (3.6%)
Age	
18-29 years	487 (50.5%)
30 - 39 years	287 (29.8%)
40 - 49 years	132 (13.7%)
50+ years	58 (6.0%)
Socioeconomic Status	
Lower	151 (15.7%)
Lower Middle	472 (49.0%)
Upper middle	290 (30.1%)
Upper	46 (4.8%)
Years of Education	
Less than 6 years	54 (5.6%)
6-12 years	176 (18.3%)
Some university, no degree	193 (20.0%)
Trade school	120 (12.5%)
University degree or more	410 (42.6%)
Ethnic Minority	
Yes	250 (26.0%)
No	493 (51.2%)
Don't know / can't answer	211 (21.9%)
Immigrant	
Yes	143 (14.8%)
No	719 (74.6%)
Not sure	85 (8.8%)
COVID-19 Pandemic Environment	
Ever lacked access to a mask	122 (12.7%)
In a location that ever issued "stay-at-home" confinement orders	729 (75.6%)

medRxiv preprint doi: <https://doi.org/10.1101/2020.11.02.20224709>; this version posted November 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

617 **Table 2: Access to and actualization of gender-affirming resources among self-identified transgender and non-binary**
 618 **individuals from the COVID-19 Disparities Survey between April 16, 2020 and August 3, 2020 (N=964)**
 619

	Overall (%)	Transmasculine	Transfeminine	Non-Binary	p-value ^b
Experienced reduced access to one or more gender-affirming resource below^a	320 / 582 (55.0%)	17 / 26 (65.4%)	110 / 189 (58.2%)	193 / 367 (52.6%)	0.25
Hormone therapy and/or gender-affirming medication	143 / 401 (35.7%)	11 / 20 (55.0%)	61 / 145 (42.1%)	71 / 236 (30.1%)	0.01
Surgical aftercare	119 / 356 (33.4%)	6 / 14 (42.9%)	51 / 127 (40.2%)	62 / 215 (28.8%)	0.08
Cosmetic supplies and services, e.g., makeup, wigs, and hair removal	189 / 500 (37.8%)	9 / 21 (42.9%)	65 / 162 (40.1%)	115 / 317 (36.3%)	0.63
Mental health counseling and therapy*	192 / 448 (42.9%)	13 / 21 (61.9%)	57 / 143 (39.9%)	122 / 284 (43.0%)	0.16
Body modifiers, e.g., binders and packing material	160 / 443 (36.1%)	8 / 18 (44.4%)	57 / 148 (38.5%)	95 / 277 (34.3%)	0.52
Compared to before the COVID-19 pandemic, how often able to live according to their gender					
More or a lot more	67 / 860 (7.8%)	7 / 28 (25.0%)	22 / 232 (9.5%)	38 / 600 (7.1%)	< 0.001
About the same	466 (54.2%)	13 (46.4%)	110 (47.4%)	343 (57.2%)	
Less or not at all	327 (38.0%)	8 (28.6%)	100 (43.1%)	219 (36.5%)	

medRxiv preprint doi: <https://doi.org/10.1101/2020.11.02.20224709>; this version posted November 16, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

^a Denominators excluded participants who were not presented with these questions, did not respond, or said that the resource was not applicable to them
^b p-values were calculated using chi-squared and Fischer's exact tests as appropriate

620
621

Preprint

622 **Table 3: Mental health and resiliency indicators among transgender and non-binary individuals from the**
 623 **COVID-19 Disparities Survey between April 16, 2020 and August 3, 2020 (N=964)^a**
 624

	Overall (%)	Transmasculine	Transfeminine	Non-Binary	p-value ^b
Screened positive (PHQ-4 ≥3)					
Depression	442 / 877 (50.4%)	17 / 35 (48.6%)	144 / 245 (58.8%)	281 / 597 (47.1%)	0.01
Anxiety	392 / 856 (45.8%)	17 / 33 (51.5%)	125 / 237 (52.7%)	250 / 586 (42.7%)	0.03
Felt lonely since COVID-19 began					
Yes	685 / 957 (71.6%)	28 / 37 (75.7%)	216 / 281 (76.9%)	441 / 639 (69.0%)	0.04
Frequency of suicidal ideation since COVID-19 vs. 6 months prior					
Was and remains rare	648 / 928 (69.8%)	26 / 36 (72.2%)	183 / 268 (68.3%)	439 / 624 (70.4%)	0.98
Decreased from frequent to rare	116 (12.5%)	4 (11.1%)	33 (12.3%)	79 (12.7%)	
Increased from rare to frequent	93 (10.0%)	3 (8.3%)	31 (11.6%)	59 (9.5%)	
Was and remains frequent	71 (7.6%)	3 (8.3%)	21 (7.8%)	47 (7.5%)	
Reported having sources of hope, strength, comfort, and peace					
Yes	489 / 768 (63.7%)	22 / 31 (71.0%)	100 / 209 (47.9%)	367 / 528 (69.5%)	< 0.001
Reported being intent on finding emotional support and therapy					
Agree	467 / 851 (54.9%)	20 / 35 (57.1%)	129 / 237 (54.4%)	318 / 579 (54.9%)	0.85
Disagree	136 (16.0%)	5 (14.3%)	43 (18.1%)	88 (15.2%)	
Reported believing they could live a happy, full life despite the pandemic^c					
Agree	516 / 847 (60.9%)	26 / 34 (76.5%)	137 / 231 (59.3%)	353 / 582 (60.6%)	0.22
Disagree	120 (14.2%)	2 (5.9%)	40 (17.3%)	78 (13.4%)	

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.11.2224709>; this version posted November 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

^a Denominators excluded individuals who did not respond or reported not knowing their answer unless otherwise noted

^b p-values were calculated using chi-squared and Fischer's exact tests as appropriate

^c Denominator includes those who stated "neither agree nor disagree"

625

626

Preprint

627 **Table 4: Economic indicators among transgender and non-binary individuals from the COVID-19 Disparities Survey**
 628 **between April 16, 2020 and August 3, 2020 (N=964)^a**
 629

	Overall (%)	Transmasculine	Transfeminine	Non-Binary	p-value ^b
Lost job due to COVID-19	151 / 953 (15.8%)	5 / 37 (13.5%)	53 / 278 (19.1%)	93 / 638 (14.6%)	0.21
Expected reduction in income					
0%	216 / 940 (23.0%)	13 / 37 (35.1%)	57 / 270 (21.1%)	10 / 633 (23.1%)	0.33
1-39%	265 (28.2%)	11 (29.7%)	70 (25.9%)	184 (29.1%)	
40-99%	337 (35.8%)	9 (24.3%)	101 (37.4%)	227 (35.9%)	
100%	122 (13.0%)	4 (10.8%)	42 (15.6%)	76 (12.0%)	
Expected to lose health insurance^c					
Yes	112 / 659 (17.0%)	7 / 23 (30.4%)	40 / 179 (22.4%)	65 / 457 (14.2%)	0.05
No	408 (61.9%)	11 (47.8%)	104 (58.1%)	293 (64.1%)	
Financial Aid					
Received and not needed	22 / 801 (2.8%)	3 / 30 (10.0%)	8 / 229 (3.5%)	11 / 542 (2.0%)	0.08
Not received but not needed	159 (19.8%)	9 (60.0%)	37 (13.2%)	113 (20.8%)	
Received and needed	92 (24.0%)	5 (16.7%)	56 (24.4%)	131 (24.2%)	
Not received and needed	428 (53.4%)	13 (43.3%)	128 (55.9%)	287 (53.0%)	
Had cut or reduced meals	361 / 900 (40.1%)	18 / 35 (51.4%)	131 / 258 (50.8%)	212 / 607 (34.9%)	< 0.001
Among those employed, ability to miss work					
Was telecommuting or on paid leave	163 / 473 (34.5%)	8 / 23 (34.8%)	32 / 128 (25.0%)	123 / 322 (38.2%)	0.06
Cannot afford to miss work but was following confinement orders	146 (30.9%)	7 (30.4%)	40 (31.2%)	99 (30.8%)	
Cannot afford to stay home and must work to survive	164 (34.7%)	8 (34.8%)	56 (43.8%)	100 (31.1%)	

medRxiv preprint doi: <https://doi.org/10.1101/2020.11.13.20211709>; this version posted November 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

^a Denominators excluded individuals who did not respond or reported not knowing their answer unless otherwise noted
^b p-values were calculated using chi-squared and Fischer's exact tests as appropriate
^c Denominator includes those who reported they "might or might not" lose their health insurance

630
631



632 **Table 5: Bivariate prevalence ratios of screening positive for depression, screening positive for anxiety, and changes in**
 633 **suicidal ideation among transgender and non-binary individuals from the COVID-19 Disparities Survey between April 16,**
 634 **2020 and August 3, 2020**
 635

	Screening positive for depression PrR (95% CI) ^a	Screening positive for anxiety PrR (95% CI)	Increased frequency of suicidal thoughts (vs. remained low) ^b	Retained frequent suicidal thoughts (vs. reduced frequency) ^c
Reported reduction in access to more than one (vs. 0) gender affirming resource	1.63 (1.36, 1.97) n = 532	1.61 (1.31, 1.97) n = 523	1.74 (1.07, 2.82) n = 449	1.37 (0.87, 2.15) n = 112
Reported decreased time (vs. increased) lived according to one's gender	1.21 (0.92, 1.58) n = 346	1.48 (1.04, 2.10) n = 331	0.57 (0.33, 0.98) n = 310	1.12 (0.50, 2.54) n = 68

^a Prevalence ratio (PrR) and 95% confidence interval (95% CI)

^b Among individuals who reported never or seldom having suicidal ideation in the six months prior to the pandemic beginning

^c Among individuals who reported having suicidal ideation quite often, very often, and all the time in the six months prior to the pandemic beginning

636

medRxiv preprint doi: <https://doi.org/10.1101/2020.11.02.20224709>; this version posted November 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Preprint

637 **SUPPORTING INFORMATION**

638 **S1 Table. Access to and actualization of gender-affirming resources among self-identified transgender and non-binary**
 639 **individuals who participated in the COVID-19 Disparities Survey, stratified by country (April 16 – August 3, 2020, N=964)**
 640

	European Region	South-East Asia Region	Region of the Americas	Eastern Mediterranean Region	Western Pacific Region	African Region	p-value ^a
Experienced reduced access to one or more gender affirming resource below^b	116 / 207 (56.0%)	109 / 204 (53.4%)	32 / 53 (60.4%)	30 / 56 (53.6%)	6 / 25 (24.0%)	15 / 23 (65.2%)	0.043
Hormone therapy and/or gender affirming medication	47 / 116 (40.5%)	55 / 186 (29.6%)	10 / 25 (40.0%)	11 / 32 (34.4%)	4 / 19 (21.1%)	8 / 13 (61.5%)	0.081
Surgical aftercare	37 / 93 (39.8%)	51 / 175 (29.1%)	7 / 15 (46.7%)	8 / 33 (24.2%)	2 / 18 (11.1%)	8 / 12 (66.7%)	0.008
Cosmetic supplies and services, e.g., makeup, wigs, and hair removal	68 / 176 (38.6%)	62 / 181 (34.3%)	21 / 46 (45.7%)	21 / 47 (44.7%)	3 / 23 (13.0%)	9 / 15 (60.0%)	0.025
Mental health counseling and therapy*	62 / 128 (48.4%)	73 / 186 (39.2%)	25 / 42 (59.5%)	12 / 42 (28.6%)	3 / 19 (15.8%)	9 / 18 (50.0%)	0.004
Body modifiers, e.g., binders and packing material	56 / 134 (41.8%)	58 / 179 (32.4%)	17 / 40 (42.5%)	14 / 46 (30.4%)	3 / 21 (14.3%)	8 / 13 (61.5%)	0.030
Compared to before the COVID-19 pandemic, how often able to live according to their gender							
More or a lot more	25 / 381 (6.6%)	20 / 236 (8.5%)	7 / 84 (8.3%)	6 / 71 (8.5%)	0 / 34 (0.0%)	6 / 33 (18.2%)	0.000
About the same	181 (47.5%)	156 (66.1%)	53 (63.1%)	26 (36.6%)	24 (70.6%)	13 (39.4%)	
Less or not at all	175 (45.9%)	60 (25.4%)	24 (28.6%)	39 (54.9%)	10 (29.4%)	14 (42.4%)	

medRxiv preprint doi: <https://doi.org/10.1101/2020.11.02.20254738>; this version posted November 24, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

^a p-values were calculated using chi-squared and Fischer's exact tests as appropriate

^b Denominators excluded participants who were not presented with these questions, did not respond, or said that the resource was not applicable to them

644
645
646
647

648 **S2 Table. Mental health and resiliency indicators among transgender and non-binary individuals who participated in the**
 649 **COVID-19 Disparities Survey, stratified by country (April 16 – August 3, 2020, N=964^a)**

	European Region	South-East Asia Region	Region of the Americas	Eastern Mediterranean Region	Western Pacific Region	African Region	p-value ^b
Screened positive per PHQ-4 (score ≥3)							
Depression	232 / 400 (58.0%)	66 / 222 (29.7%)	46 / 82 (56.1%)	48 / 76 (63.2%)	21 / 40 (52.5%)	18 / 32 (56.3%)	≤ 0.001
Anxiety	392 / 856 (45.8%)	392 / 856 (45.8%)	392 / 856 (45.8%)	392 / 856 (45.8%)	392 / 856 (45.8%)	392 / 856 (45.8%)	≤ 0.001
Felt lonely since COVID-19 began							
Yes	685 / 957 (71.6%)	685 / 957 (71.6%)	685 / 957 (71.6%)	685 / 957 (71.6%)	685 / 957 (71.6%)	685 / 957 (71.6%)	0.063
Frequency of suicidal ideation since COVID-19 vs. 6 months prior							
Was and remains rare	648 / 928 (69.8%)	648 / 928 (69.8%)	648 / 928 (69.8%)	648 / 928 (69.8%)	648 / 928 (69.8%)	648 / 928 (69.8%)	0.020
Decreased from frequent to rare	116 (12.5%)	116 (12.5%)	116 (12.5%)	116 (12.5%)	116 (12.5%)	116 (12.5%)	
Increased from rare to frequent	93 (10.0%)	93 (10.0%)	93 (10.0%)	93 (10.0%)	93 (10.0%)	93 (10.0%)	
Was and remains frequent	71 (7.6%)	71 (7.6%)	71 (7.6%)	71 (7.6%)	71 (7.6%)	71 (7.6%)	
Reported having sources of hope, strength, comfort, and peace							
Yes	489 / 768 (63.7%)	489 / 768 (63.7%)	489 / 768 (63.7%)	489 / 768 (63.7%)	489 / 768 (63.7%)	489 / 768 (63.7%)	≤ 0.001
Reported being intent on finding emotional support and therapy^c							
Agree	467 / 851 (54.9%)	467 / 851 (54.9%)	467 / 851 (54.9%)	467 / 851 (54.9%)	467 / 851 (54.9%)	467 / 851 (54.9%)	≤ 0.001
Disagree	136 (16.0%)	136 (16.0%)	136 (16.0%)	136 (16.0%)	136 (16.0%)	136 (16.0%)	
Reported believing they could live a happy, full life despite the pandemic^c							
Agree	516 / 847 (60.9%)	516 / 847 (60.9%)	516 / 847 (60.9%)	516 / 847 (60.9%)	516 / 847 (60.9%)	516 / 847 (60.9%)	≤ 0.001
Disagree	120 (14.2%)	120 (14.2%)	120 (14.2%)	120 (14.2%)	120 (14.2%)	120 (14.2%)	

medRxiv preprint doi: <https://doi.org/10.1101/2020.11.02.20224709>; this version posted November 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

^a Denominators excluded individuals who did not respond or reported not knowing their answer unless otherwise noted
^b p-values were calculated using chi-squared and Fischer's exact tests as appropriate
^c Denominator includes those who stated "neither agree nor disagree"

652
 654
 655
 656
 657

658
659
660

S3 Table: Socioeconomic indicators among transgender and non-binary individuals who participated in the COVID-19 Disparities Survey, stratified by country (April 16 – August 3, 2020, N=964)^a

	European Region	South-East Asia Region	Region of the Americas	Eastern Mediterranean Region	Western Pacific Region	African Region	p-value ^b
Lost job due to COVID-19	74 / 446 (16.6%)	38 / 238 (16.0%)	13 / 86 (15.1%)	15 / 83 (18.1%)	5 / 40 (12.5%)	2 / 35 (5.7%)	0.614
Expected reduction in income							
0%	113 / 436 (25.9%)	39 / 340 (16.3%)	25 / 86 (29.1%)	19 / 79 (24.1%)	9 / 40 (22.5%)	9 / 35 (25.7%)	0.055
1-39%	123 (28.2%)	66 (27.5%)	20 (23.3%)	20 (25.3%)	14 (35.0%)	11 (31.4%)	
40-99%	140 (32.1%)	110 (45.8%)	34 (39.5%)	25 (31.7%)	13 (32.5%)	10 (28.6%)	
100%	60 (13.8%)	25 (10.4%)	7 (8.1%)	15 (19.0%)	4 (10.0%)	5 (14.3%)	
Expected to lose health insurance^c							
Yes	32 / 307 (10.4%)	42 / 165 (25.5%)	11 / 67 (16.4%)	16 / 51 (31.4%)	7 / 35 (20.0%)	1 / 21 (4.8%)	< 0.001
No	203 (66.1%)	94 (57.0%)	48 (71.6%)	27 (52.9%)	18 (51.4%)	12 (57.1%)	
Receipt and need of financial aid							
Received and not needed	2 / 75 (2.7%)	6 / 208 (2.9%)	2 / 75 (2.7%)	3 / 64 (4.7%)	0 / 32 (0.0%)	0 / 30 (0.0%)	<0.001
Not received but not needed	86 (22.9%)	29 (13.9%)	19 (25.3%)	13 (20.3%)	6 (18.8%)	5 (16.7%)	
Received and needed	53 (14.1%)	92 (44.2%)	23 (30.7%)	8 (12.5%)	8 (25.0%)	5 (16.7%)	
Not received and needed	226 (60.3%)	81 (38.9%)	31 (41.3%)	40 (62.5%)	18 (56.3%)	20 (66.7%)	
Had cut or reduced meals	167 / 420 (39.8%)	82 / 230 (35.7%)	28 / 82 (34.2%)	45 / 75 (60.0%)	12 / 37 (32.4%)	16 / 34 (47.1%)	0.004
Among those employed, ability to miss work							
Was telecommuting or on paid leave	81 / 214 (37.9%)	40 / 127 (31.5%)	17 / 42 (40.5%)	11 / 34 (32.4%)	6 / 21 (28.6%)	6 / 20 (30.0%)	0.165
Cannot afford to miss work but was following confinement orders	72 (33.6%)	34 (26.8%)	9 (21.4%)	9 (26.5%)	5 (23.8%)	10 (50.0%)	
Cannot afford to stay home and must work to survive	61 (28.5%)	53 (41.7%)	16 (38.1%)	14 (41.2%)	10 (47.6%)	4 (20.0%)	

medRxiv preprint doi: <https://doi.org/10.1101/2020.11.02.20224709>; this version posted November 4, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC 4.0 International license.

^a Denominators excluded individuals who did not respond or reported not knowing their answer unless otherwise noted
^b p-values were calculated using chi-squared and Fischer's exact tests as appropriate
^c Denominator includes those who reported they "might or might not" lose their health insurance

661
662
663
664
665
666