

1 Factors influencing the COVID-19 daily deaths peak across 2 European countries

3 Katarzyna Jabłońska¹, Samuel Aballéa², Mondher Toumi³

4

5 ¹ MSc, Creativ-Ceutical, ul. Przemysłowa 12, 30-701 Kraków, Poland

6 ² MSc, PhD, Creativ-Ceutical, Westblaak 90, 3012KM Rotterdam, Netherlands

7 ³ MD, MSc, PhD, Professor of Public Health, Faculty of Medicine, Public Health Department,
8 Aix-Marseille University, 27 Boulevard Jean Moulin, Marseille 13005, France

9 Corresponding author: Katarzyna Jabłońska, katarzyna.jablonska@creativ-ceutical.com

10

11 Abstract

12 **OBJECTIVES:** The purpose of this study was to determine predictors of the height of COVID-
13 19 daily deaths peak and time to the peak, in order to explain their variability across European
14 countries.

15 **STUDY DESIGN:** For 34 European countries, publicly available data were collected on daily
16 numbers of COVID-19 deaths, population size, healthcare capacity, government restrictions
17 and their timing, tourism and change in mobility during the pandemic.

18 **METHODS:** Univariate and multivariate generalised linear models using different selection
19 algorithms (forward, backward, stepwise and genetic algorithm) were analysed with height of
20 COVID-19 daily deaths peak and time to the peak as dependent variables.

21 **RESULTS:** The proportion of the population living in urban areas, mobility at the day of first
22 reported death and number of infections when borders were closed were assessed as
23 significant predictors of the height of COVID-19 daily deaths peak. Testing the model with
24 variety of selection algorithms provided consistent results. Total hospital bed capacity,
25 population size, number of foreign travellers and day of border closure, were found as
26 significant predictors of time to COVID-19 daily deaths peak.

27 **CONCLUSIONS:** Our analysis demonstrated that countries with higher proportions of the
28 population living in urban areas, with lower reduction in mobility at the beginning of the
29 pandemic, and countries which closed borders having more infected people experienced
30 higher peak of COVID-19 deaths. Greater bed capacity, bigger population size and later
31 border closure could result in delaying time to reach the deaths peak, whereas a high number
32 of foreign travellers could accelerate it.

33 **Keywords:** COVID-19, mortality, healthcare capacity, modelling.

34

35

36 Introduction

37 The coronavirus infectious disease 2019 (COVID-19) outbreak was announced as a pandemic
38 by the World Health Organization (WHO) on 11 March 2020.¹ By the end of March, Europe
39 exceeded Asia and became the region experiencing the highest percentage mortality from the
40 virus across the world,² until 10 June 2020, when the rapidly growing COVID-19 mortality rate
41 in the Americas exceeded all other continents.³ According to the WHO report from 5 July 2020,
42 38% of the global mortality due to COVID-19 was from Europe.⁴

43 Since incidence and mortality rates varied between countries, numerous studies have recently
44 been published investigating factors associated with COVID-19 infection and death rate
45 across countries. A variety of potential predictors have been assessed in the literature, such
46 as country-specific demographic and health characteristics, economic and social indicators,⁵⁻
47 ⁸ mobility scores and social-distancing measures,^{9, 10} as well as ecological and environmental
48 perspectives.^{8, 11} To assess the relationship between covariates and COVID-19 incidence or
49 mortality, the most common approach used by authors was to analyse multivariate regression
50 models of total number of infections or deaths up to a given time point, using log-transformed
51 data or without any transformation, as well as daily data on infections or deaths as outcomes.

52 In this study, we used data on numbers of deaths, not infections, since the former has a much
53 higher degree of reliability than the latter, being better monitored and less dependent on the
54 number of tests done. Since all European countries seem to already reach the peak of deaths
55 by 3 June 2020 from the first wave of COVID-19 disease, our idea was to use height of daily
56 deaths peak as a primary outcome of interest, and time to the peak as a secondary outcome.

57 To the best of our knowledge, this perspective has not been investigated so far. When raw
58 and cumulative daily numbers of infections and deaths are subject to deviations from between-
59 country differences in reporting and depend on the date up to which the analysis is performed,
60 maximum number of daily deaths can be assessed as an interesting new indicator of disease
61 mortality magnitude. In addition, analysis of the height of daily deaths peak enables us to
62 assess the overall capacity of healthcare systems.

63 This study aims to detect significant drivers of COVID-19 mortality with the use of multivariate
64 generalised linear models (GLM) and distinct selection algorithms, to explain the variability of
65 height of and time to the deaths peak among European countries. This will enable us to draw
66 conclusions about how governments and societies can improve the future response on similar
67 pandemic or global life-threatening situations.

68

69 Methods

70 Data collection

71 A total of 34 European countries were included in the analysis. The height of COVID-19 daily
72 deaths peak was the primary outcome of interest, defined as maximum daily reported number
73 of people who died due to COVID-19 per country up to 3 June 2020. The value was then
74 divided by the number of inhabitants of a given country reported in January 2020 and
75 multiplied by 1 million to obtain the deaths peak height per 1 million inhabitants.

76 The secondary outcome of interest was time to COVID-19 daily deaths peak, defined as
77 number of days from the day when the first death was reported in a given country, up to the
78 day of reaching the COVID-19 daily deaths peak.

79 A set of explanatory variables used to predict outcomes consisted of the following:

- 80 • Healthcare capacity:
 - 81 ○ All beds capacity (number of hospital bed units)
 - 82 ○ Intensive care unit (ICU) beds capacity (number of ICU)
 - 83 ○ Number of tests conducted up to the time of the peak
- 84 • Government restrictions and associated factors:
 - 85 ○ 'Stay at home' order date
 - 86 ○ Educational facilities closure date
 - 87 ○ Gathering restriction date
 - 88 ○ Businesses closure date
 - 89 ○ Border closure date
 - 90 ○ Total number of COVID-19 infected people when borders were closed
 - 91 ○ Total number of COVID-19 deaths when borders were closed
- 92 • Indicators of the population size:
 - 93 ○ Country population size (in January 2020)
 - 94 ○ Percentage of population living in urban areas
 - 95 ○ Percentage of population living in metropolitan cities with more than 1 million
96 inhabitants
- 97 • Median age
- 98 • Tourism:
 - 99 ○ Number of travellers that arrived at airports in 2018
 - 100 ○ Number of foreign tourists in 2018 (arrivals at any touristic accommodation)
- 101 • Mean mobility score at the day of first reported COVID-19 death (as a difference from
102 the average mobility reported before the pandemic), calculated across mobility scores
103 at:
 - 104 ○ Retail and recreation places
 - 105 ○ Workplaces
 - 106 ○ Transit stations

107 All dates were considered as number of days after the day of first reported COVID-19 death
108 in each country. Variables indicating healthcare capacity and tourism were considered in
109 relation to the population size of a given country (per inhabitant or 1 million inhabitants).

110 *Data sources*

111 Data on COVID-19 deaths, infections, number of tests, all and ICU beds capacity, dates of
112 government restrictions, population size and urban population size were taken from
113 Worldometer.com.¹² Missing dates of government restrictions, if officially issued, were found
114 on Wikipedia.¹³

115 Mobility scores were uploaded from Google COVID-19 Community Mobility Reports.¹⁴ Scores
116 were reported as percentage changes from a usual mobility calculated before the pandemic
117 in Europe, between 3 January and 6 February 2020. Mobility scores were subject to
118 oscillations due to daily reporting and presence of weekends and holidays; therefore, we used
119 smoothed mobility scores produced with a nonparametric technique that uses local weighted
120 regression and fits a smooth curve through points in a scatter plot, called *loess regression*.¹⁵

121 Data on number of passengers arrived at airports, tourism and population living in metropolitan
122 areas were downloaded from Eurostat,¹⁶ except for Russia, Turkey and Ukraine, for which
123 data were gathered from other sources.¹⁷⁻²⁰

124 *Missing data and imputation*

125 If government restrictions were not officially set, the date was imputed with date of the peak
126 in each country. Missing data on mobility for Cyprus and Iceland were imputed with average
127 scores of remaining countries.

128

129 **Statistical analysis**

130 Descriptive statistics on outcomes and explanatory variables were produced. Since the
131 distribution of deaths peak height per population size is right-skewed, a logarithmic
132 transformation was applied.

133 Factors influencing the height of deaths peak were analysed using univariate and multivariate
134 GLM with a normal distribution function and logarithmic link function. The base case scenario
135 analysis was performed using data available for 34 countries and explanatory variables with
136 a p-value <0.1 in univariate GLM models. To avoid multicollinearity issues between
137 independent variables, Pearson correlations were investigated to detect highly correlated
138 pairs of variables. The correlation was assessed as high if the absolute value exceeded 0.7
139 and as moderate if the absolute value was contained in a range 0.5–0.7.²¹

140 Due to relatively low sample size, a risk of bias could appear for models with too high number
141 of variables included. To overcome the problem and limit the number of covariates, sensitivity
142 analyses were performed using variety of selection algorithms: stepwise, backward, forward
143 and the genetic algorithm. For the backward, forward and stepwise algorithms, a criterion of
144 having p-value lower than 0.1 was applied for each variable to stay in a model (backward and
145 stepwise) and to enter a model (forward and stepwise). For the genetic algorithm, the best
146 model was fitted based on the value of Akaike's Information Criterion (AIC) corrected for small
147 sample sizes (AICC). Models with only main effects were considered. Additional sensitivity
148 analysis was performed removing countries for which imputation of government restriction
149 dates was needed.

150 Similar methods were used to analyse time to deaths peak but without any transformation
151 since it seemed to follow a normal distribution. GLM models with a normal distribution function
152 and identity link function were analysed.

153 For all analyses, a p-value lower than 0.05 was considered as statistically significant. For each
154 GLM model, fit statistics such as AIC and its equivalent AICC were produced with lower values
155 indicating better fit.

156 Analyses were performed using SAS 9.4 software. R 3.6.2 software was used to apply the
157 genetic algorithm with the package *glmulti*.²²

158

159 **Results**

160 **Descriptive analysis**

161 Characteristics of the countries included in the study are presented in Table 1, and a histogram
162 of the height of COVID-19 deaths peak is depicted in Figure 1. The graph presenting deaths
163 peak height by country is provided in Figure 2.

164 Median height of the peak per 1 million inhabitants per day was equal to 3.48 deaths ([lower
165 quartile; upper quartile] = [1.68; 12.78]), with Belgium reaching outstandingly higher peak than
166 other countries. As can be seen from these statistics and the histogram, the height of the
167 deaths peak does not follow a normal distribution, but it seems it can be well approximated
168 using a log-normal distribution. Median time to the peak equalled 31 days from the first death
169 reported in each country with a comparable mean (31.32, SD=13.94). Average numbers of
170 days when closing schools/universities and when gatherings were banned were both negative
171 (-0.26 and -1.76, respectively).

172

173 **Height of the deaths peak**

174 Univariate GLM models of height of the peak were analysed (Table 2). Eight factors turned
175 out to be significant: educational facilities closure day, gathering restrictions day, businesses
176 closure day, proportion living in urban areas, mobility score at the day of first reported death,
177 border closure day, number of COVID-19 infections when borders were closed, and number
178 of COVID-19 deaths when borders were closed. All of them have positive estimates. Stay-at-
179 home order day, as well as the proportion living in metropolitan cities >1 million inhabitants
180 were close to reaching the significance level ($p < 0.1$). Pearson correlations between variables
181 reaching or close to reaching significance in univariate models were verified. Results can be
182 found in Supplementary Materials.

183 A base case multivariate GLM model included six covariates. The proportion living in urban
184 areas was found significant ($p < 0.001$) and the mobility score at the day of first reported death
185 was very close to reaching significance ($p = 0.052$). Results are presented in Table 2.
186 Considering small sample size ($N = 34$), selection algorithms were applied to the base case
187 model to limit the number of covariates, increase model precision and improve the model fit.
188 All selection algorithms were consistent and selected the model with three significant
189 parameters ("final" model), indicating its best fit. The proportion of the population living in urban
190 areas (6.848, $p < 0.001$), mobility score at the day of first reported death (0.049 $p < 0.001$) and
191 number of infections when borders were closed per 1 million inhabitants (0.0002, $p = 0.016$)
192 were all significantly associated with the deaths peak height (Table 2).

193 Analysis of data removing countries with imputed dates of businesses or borders closure
194 ($N = 29$) provided consistent results regarding the significance of all three covariates from the
195 final models in relation to the deaths peak height (details are provided in Supplementary
196 Materials).

197

198 **Time to the deaths peak**

199 Univariate GLM models of time to the peak were analysed (Table 3). Nine factors turned out
200 to be potentially significant (with $p < 0.1$): all beds capacity, ICU beds capacity, educational
201 facilities closure day, gathering restrictions day, population size, arrivals at airports per

202 inhabitant, number of foreign tourists per inhabitant, mobility score at the day of first reported
203 death, as well as day of border closure. The majority of these factors have positive estimates,
204 except the number of arrivals and tourists. As for the next step, linear correlations between
205 them were verified. Results can be found in Supplementary Materials.

206 A base case multivariate GLM model was run using six explanatory variables. All beds
207 capacity, population size in millions and the number of foreign tourists per inhabitant were
208 found significantly related with time to COVID-19 deaths peak. Results are presented in Table
209 3. Considering the small sample size (N=34), selection algorithms were applied to the above
210 multivariate model. The “final” model had four covariates and was selected by the backward
211 and the genetic algorithm (details in Supplementary Materials). All beds capacity per 1 million
212 inhabitants (0.003, $p=0.004$), population size in millions (0.142, $p=0.008$), the number of
213 foreign tourists per inhabitant (- 2.651, $p=0.037$) and border closure day (0.297, $p=0.008$) were
214 all significantly associated with time to the deaths peak (Table 3).

215 Analysis of data removing countries with imputed dates of borders closure (N=31) provided
216 consistent results regarding the significance of all beds capacity and population size in relation
217 to time to deaths peak, whereas number of foreign tourists and borders closure day were close
218 in reaching significance, both with $p<0.1$ (details provided in Supplementary Materials).

219

220 Discussion

221 This study provides some evidence about factors associated with COVID-19 mortality peak
222 and time to the peak. One of the strongest predictors of the COVID-19 mortality peak identified
223 in our analysis was the proportion of population living in urban areas. The relationship between
224 urbanisation and COVID-19 infections ratio was earlier outlined by the United Nations
225 Association,²³ estimating that 90% of all reported COVID-19 cases (by July 2020) came from
226 urban areas, becoming the epicentre of the pandemic. High population density increases the
227 propensity of viruses spread by increasing the contact rates of individuals. However, some
228 authors²⁴ warn readers against putting too much weight on urban density, arguing that large
229 cities just faced the coronavirus earlier (due to the higher number of incoming people) and that
230 the timing of epidemic start was of bigger interest than population density itself. This
231 observation seems to be on the contrary to our study, since neither proportion of population
232 living in urban areas, nor proportion living in metropolitan cities were related with the time to
233 deaths peak.

234 Number of infections when borders were closed was found to be another important factor
235 associated with the COVID-19 deaths peak height, whereas a positive association between
236 borders closure day and time to reach the peak was observed in this study. It shows that
237 stopping arrivals to the country at the earlier stage of epidemic can be crucial in reducing the
238 peak height, stopping the increase of daily number of deaths earlier and, consequently, to
239 flatten the deaths curve. These findings are on the contrary with Chaudhry who showed no
240 association between rapid border closures and COVID-19 mortality per million people, using
241 cumulative data available as of 1 April 2020.²⁵ However, the peak height and mortality may
242 not be strongly correlated. For example some countries experience a point peak and dropping
243 fast, (France, United Kingdom, Spain, Italy) while other countries may experience a flattened
244 peak with a mortality staying high over a long period of time (US, Brazil, Mexico).¹²

245 The difference in average mobility before the pandemic and at the day when first death was
246 reported in a country was found to be associated with deaths peak height, which suggests that
247 rapid reduction in mobility was important in terms of reducing overall mortality. These findings
248 are consistent with other authors opinion that changes in social-distancing covariates can
249 flatten the curve by changing the peak death rate.^{9, 10}

250 Tourism indicators were found not to be associated with deaths peak height, but to be
251 significant drivers of time to reach the peak. It suggests that the magnitude of inbound tourism
252 can have an impact on accelerating the moment of reaching the peak. However, Aldibasi⁵ and
253 Garcia de Alcaniz⁸ found tourism to be a significant predictor of COVID-19 mortality and
254 infection rates. Ostig and Askin²⁶ also found a significant positive relationship between number
255 of airline passengers and number of COVID-19 infections.

256 Another conclusion resulting from the study is that timing of government restrictions, especially
257 border closure, can be found as an important factor in terms of COVID-19 mortality. What can
258 be observed for many countries is that closing borders (and other government restrictions as
259 well) took place several days before any death was reported in a country. Therefore, increase
260 or decrease in mortality can be seen as a result of undertaking preventive action by the
261 government.

262 There is no general conclusion on the association between COVID-19 fatality and hospital
263 beds capacity in literature. Our study showed lack of association between beds capacity and
264 deaths peak height, but higher beds capacity was related with longer time to peak. Garcia de
265 Alcaniz showed no association between hospital beds density and both the number of
266 infections and number of deaths at any moment of the pandemic.⁸ On the contrary, Sorci
267 showed that case fatality rate was negatively associated with number of hospital beds per
268 inhabitants.⁷ High bed capacity may be associated with more intensive treatment delaying
269 time to death for a number of patients thus delaying time to peak but does not affect the peak
270 height.

271

272 **Limitations**

273 Our study has several limitations. Data on a limited number of countries were used. We
274 decided not to include countries for which reliable and comparative evidence could not be
275 found, nor countries with very low mortality. Our idea was to focus only on Europe, not to
276 merge it with countries from other continents and avoid data incomparability issues and
277 eventual difference in strain as the virus mutate rapidly.

278 However, the number of observations can be assessed as high enough to draw conclusions
279 based on multivariate GLM models by using selection algorithms. The stability of results was
280 tested with models based on different variable selection algorithms, being consistent with base
281 case findings.

282 The set of included explanatory variables can be viewed as non-exhaustive. We decided to
283 focus on variables important from a social perspective, assuming that mobility factors, tourism,
284 urbanisation, and government decisions are more of interest to inform upcoming decisions
285 than other approaches observed in the literature; for example, ecological or environmental
286 perspectives.^{8, 11}

287

288 **Conclusions**

289 This study demonstrated the significant drivers of COVID-19 mortality magnitude in Europe
290 and shed a light on reasons behind the variability between different countries. Countries with
291 a higher proportion of population living in urban areas, without a rapid reduction in mobility
292 and with delayed government restrictions, correlated with a higher COVID-19 deaths peak.
293 These findings can help improve future response in similar situations or in case of a second
294 wave.

295

296

297 **Acknowledgements:** The authors wish to thank Saima Khan (Apothecom) for proof reading
298 and editing support.

299 **Contributors:** MT conceived the study, KJ collected and analysed the data, KJ, MT and SA
300 interpreted the results, KJ wrote the first draft of the manuscript, KJ, MT and SA revised the
301 manuscript and approved the final version.

302 **Competing interests:** The authors declare no competing interests.

303

304

305 **References**

- 306 1. World Health Organisation. Coronavirus disease 2019 (COVID-19) Situation Report – 51 (11
307 March 2020) 11 March 2020.
- 308 2. World Health Organisation. Coronavirus disease 2019 (COVID-19) Situation Report – 71 (31
309 March 2020) 31 March 2020.
- 310 3. World Health Organisation. Coronavirus disease (COVID-19) Situation Report – 142 (10 June
311 2020) 10 June 2020.
- 312 4. World Health Organisation. Coronavirus disease (COVID-19) Situation Report – 167 (5 July
313 2020) 5 July 2020.
- 314 5. Aldibasi OS, Alharbi NK, Alkelya M, Zowawi H, Alghnam S. The Association of Country-Level
315 Factors with Outcomes of COVID-19: Analysis of the pandemic after one million cases. *Research*
316 *Square*. 2020.
- 317 6. Burden SJ, Rademaker J, Weedon BD, Whaymand L, Dawes H, Jones A. Associations of
318 Global Country Profiles and Modifiable Risk Factors with COVID-19 Cases and Deaths. *medRxiv*.
319 2020:2020.06.17.20133454.
- 320 7. Sorci G, Faivre B, Morand S. Why Does COVID-19 Case Fatality Rate Vary Among Countries?
321 *The Lancet*. 2020.
- 322 8. Garcia de Alcaniz JG, Romero-Lopez J, Martinez RP, Lopez-Rodas V, Costas E. What
323 variables can better predict the number of infections and deaths worldwide by SARS-CoV-2? Variation
324 through time. *medRxiv*. 2020:2020.06.04.20122176.

- 325 9. Woody S, Garcia Tec M, Dahan M, Gaither K, Lachmann M, Fox S, et al. Projections for first-
326 wave COVID-19 deaths across the US using social-distancing measures derived from mobile phones.
327 medRxiv. 2020:2020.04.16.20068163.
- 328 10. Courtemanche C, Garuccio J, Le A, Pinkston J, Yelowitz A. Strong Social Distancing Measures
329 In The United States Reduced The COVID-19 Growth Rate. Health Affairs.
330 0:10.1377/hlthaff.2020.00608.
- 331 11. Bray I, Gibson A, White J. Coronavirus disease 2019 mortality: a multivariate ecological analysis
332 in relation to ethnicity, population density, obesity, deprivation and pollution. Public Health. 2020;
333 185:261-3.
- 334 12. Worldometer. Covid-19 coronavirus pandemic. [cited 2020, 25 June 2020]; Available from:
335 <https://www.worldometers.info/coronavirus/>.
- 336 13. Wikipedia. National responses to the COVID-19 pandemic. [cited 2020, 25 June 2020];
337 Available from: https://en.wikipedia.org/wiki/National_responses_to_the_COVID-19_pandemic.
- 338 14. Google. COVID-19 Community Mobility Reports. [cited 2020, 25 June 2020]; Available from:
339 <https://www.google.com/covid19/mobility/>.
- 340 15. Wicklin R. What is loess regression? SAS Blogs: SAS; 2016 [cited 2020 29 June 2020];
341 Available from: <https://blogs.sas.com/content/iml/2016/10/17/what-is-loess-regression.html>.
- 342 16. Eurostat. Eurostat Database. [cited 2020 25 June 2020]; Available from:
343 <https://ec.europa.eu/eurostat/data/database>.
- 344 17. CEIC data. Visitor Arrivals. [cited 2020, 25 June 2020]; Available from:
345 <https://www.ceicdata.com/en/indicator/russia/visitor-arrivals>,
346 <https://www.ceicdata.com/en/indicator/turkey/visitor-arrival>,
347 <https://www.ceicdata.com/en/indicator/ukraine/visitor-arrivals>.
- 348 18. Analytical Credit Rating Agency ACRA. Air traffic increases, finances in crisis [cited 2020, 25
349 June 2020]; Available from: <https://www.acra-ratings.cn/research/1282>.
- 350 19. Hurriyet Daily News. Air passenger traffic in Turkish airports rose 8.8 percent in 2018. [cited
351 2020, 25 June 2020]; Available from: <https://www.hurriyetdailynews.com/air-passenger-traffic-in-turkish-airports-rose-8-8-percent-in-2018-140322>.
- 353 20. Kyiv Post. Record-breaking 20.5 million passengers at Ukrainian airports in 2018. [cited 2020,
354 25 June 2020]; Available from: <https://www.kyivpost.com/ukraine-politics/record-breaking-20-5-million-passengers-passed-through-ukrainian-airports-in-2018.html>.
- 356 21. Mukaka MM. Statistics corner: A guide to appropriate use of correlation coefficient in medical
357 research. Malawi Med J. 2012; 24:69-71.
- 358 22. Calcagno V, de Mazancourt C. glmulti: An R Package for Easy Automated Model Selection with
359 (Generalized) Linear Models. Journal Of Statistical Software. 2010; 34.
- 360 23. Guterres A. COVID-19 in an Urban World. United Nations.; 2020 [cited 2020 13th August 2020];
361 Available from: <https://www.un.org/en/coronavirus/covid-19-urban-world>.
- 362 24. Barr J, Tassier T. Are Crowded Cities the Reason for the COVID-19 Pandemic? Placing too
363 much blame on urban density is a mistake. Scientific American; 2020 [cited 2020 13th August 2020];
364 Available from: <https://blogs.scientificamerican.com/observations/are-crowded-cities-the-reason-for-the-covid-19-pandemic/>.
- 365

366 25. Chaudhry R, Dranitsaris G, Mubashir T, Bartoszko J, Riaz S. A country level analysis measuring
367 the impact of government actions, country preparedness and socioeconomic factors on COVID-19
368 mortality and related health outcomes. *EClinicalMedicine*. 2020; 25.

369 26. Oztig LI, Askin OE. Human mobility and coronavirus disease 2019 (COVID-19): a negative
370 binomial regression analysis. *Public Health*. 2020; 185:364-7.

371

372

373 **Table 1.** Characteristics of countries included in the analysis.

Variable	N	Mean	SD	Median	Lower quartile	Upper quartile	Min	Max
Peak height [no. of deaths per 1 mln inhabitants]	34	7.22	8.65	3.48	1.68	12.78	0.48	42.80
Peak height [raw no. of deaths]	34	180.50	322.67	32.50	8.00	185.00	2.00	1172.00
Peak time [no. of days after first death reported]	34	31.32	13.94	31.00	22.00	40.00	2.00	71.00
All beds capacity [per 1 mln inhabitants]	34	4940.25	1744.89	4634.17	3427.67	6623.84	2512.45	8248.80
ICU beds capacity [per 1 mln inhabitants]	34	141.14	82.61	110.94	86.50	191.75	20.72	349.20
Total no. of tests up to time of the peak [per 1 mln inhabitants]	34	763.09	743.98	526.54	443.77	824.78	148.92	4166.11
Stay-at-home order day	34	8.74	10.33	8.00	1.00	14.00	-6.00	41.00
Educational facilities closure day	34	-0.26	8.31	0.50	-3.00	4.00	-22.00	17.00
Gathering restrictions day	34	-1.76	8.67	-1.00	-6.00	2.00	-21.00	17.00
Businesses closure day	34	3.32	10.14	2.00	-1.00	7.00	-16.00	41.00
Population size [Jan 2020]	34	23.82	33.02	9.33	4.94	37.85	0.34	145.93
Proportion living in urban areas	34	0.74	0.12	0.74	0.66	0.83	0.54	0.98
Proportion living in metropolitan cities with more than 1 mln inhabitants	34	0.28	0.19	0.31	0.12	0.44	0.00	0.56
Median age	34	41.85	2.96	42.50	40.50	44.00	32.00	46.70
Arrivals at airports in 2018 [per 1 inhabitant]	34	2.27	2.63	1.52	0.84	2.79	0.24	14.86
No. of foreign tourists in 2018 [per 1 inhabitant]	34	1.25	1.32	0.79	0.47	1.46	0.14	6.73
Mobility score at the day of first reported death	34	-23.45	18.93	-20.02	-44.09	-5.28	-56.42	0.16
Border closure day	34	6.71	15.53	2.50	-1.00	11.00	-19.00	46.00
No. of COVID-19 infections when borders were closed [per 1 mln inhabitants]	34	490.54	1116.56	83.39	21.53	245.88	0.07	5223.83
No. of COVID-19 deaths when borders were closed [per 1 mln inhabitants]	34	28.56	71.73	0.27	0.00	3.17	0.00	297.90

374 Abbreviations: ICU, intensive care unit; SD, standard deviation; mln, million.

375

376

377 **Table 2.** Results of univariate and multivariate GLM of COVID-19 deaths peak height, with normal distribution and logit link function

Variable	Univariate analysis				Multivariate analysis								
	Estimate	Wald 95% confidence limit		p-value	Base case GLM (Full model)				Base case GLM + selection algorithms (Final model)				
		Wald 95% confidence limit			Estimate	Wald 95% confidence limit	p-value	Estimate	Wald 95% confidence limit		p-value		
		Lower	Upper						Lower	Upper			
All beds capacity [per 1 mln inhabitants]	-0.0002	-0.001	0.0001	0.109	-	-	-	-	-	-	-	-	-
ICU beds capacity [per 1 mln inhabitants]	-0.002	-0.008	0.003	0.404	-	-	-	-	-	-	-	-	-
Total no. of tests up to time of the peak [per 1 mln inhabitants]	0.000	-0.001	0.001	0.961	-	-	-	-	-	-	-	-	-
Stay-at-home order day	0.027	-0.001	0.053	0.057**	-	-	-	-	-	-	-	-	-
Educational facilities closure day	0.067	0.028	0.107	0.001*	-	-	-	-	-	-	-	-	-
Gathering restrictions day	0.045	0.006	0.084	0.023*	-	-	-	-	-	-	-	-	-
Businesses closure day	0.029	0.009	0.049	0.005*	-0.001	-0.022	0.020	0.944	-	-	-	-	-
Population size [mln]	0.001	-0.009	0.011	0.877	-	-	-	-	-	-	-	-	-
Proportion living in urban areas	8.117	3.695	12.539	<0.001*	7.127	3.642	10.611	<0.001*	6.848	4.016	9.680	<0.001*	-
Proportion living in metropolitan cities with more than 1 mln inhabitants	2.428	-0.464	5.319	0.099**	1.063	-1.331	3.457	0.384	-	-	-	-	-
Median age	-0.0003	-0.120	0.120	0.996	-	-	-	-	-	-	-	-	-
Arrivals at airports in 2018 [per 1 inhabitant]	0.016	-0.097	0.129	0.784	-	-	-	-	-	-	-	-	-
No. of foreign tourists in 2018 [per 1 inhabitant]	-0.076	-0.421	0.270	0.668	-	-	-	-	-	-	-	-	-
Mobility score at the day of first reported death	0.048	0.001	0.096	0.046*	0.041	-0.0004	0.083	0.052*	0.049	0.022	0.077	<0.001*	-
Borders closure day	0.025	0.010	0.041	0.002*	-0.005	-0.030	0.021	0.729	-	-	-	-	-
No. of COVID-19 infections when borders were closed [per 1 mln inhabitants]	0.0002	0.0000	0.004	0.022*	0.0003	-0.0001	0.001	0.113	0.0002	0.0001	0.0003	0.016*	-
No. of COVID-19 deaths when borders were closed [per 1 mln inhabitants]	0.004	0.001	0.007	0.002*	-	-	-	-	-	-	-	-	-
Multivariate models statistics													
Scale	-	-	-	-	5.020	3.958	6.367	-	5.092	4.015	6.458	-	-
AIC	-	-	-	-	222.203	-	-	-	217.169	-	-	-	-
AICC	-	-	-	-	227.963	-	-	-	219.312	-	-	-	-

378 *p-value <0.05; **p-value < 0.1.

379 Abbreviations: AIC, Akaike's Information Criterion; AICC, AIC corrected for small sample sizes; GLM, generalised linear models; ICU, intensive care unit; mln, million.

380 GLM univariate and multivariate models with normal distribution and logit link function were used to explore factors associated with COVID-19 deaths peak height as of 3rd June 2020. Each model
 381 was run using 34 observations. Variables significant in univariate models were included into the multivariate base case model, avoiding highly correlated pairs. The final multivariate model was
 382 selected based on the use of selection algorithms (backward, forward, stepwise and the genetic algorithm).
 383
 384
 385

386
387

Table 3. Results of univariate and multivariate GLM of time to COVID-19 deaths peak, with normal distribution and identity link function

Variable	Univariate analysis				Multivariate analysis							
	Estimate	Wald 95% confidence limit		p-value	Base case GLM (Full model)				Base case GLM + selection algorithms (Final model)			
		Wald 95% confidence limit			Estimate	Wald 95% confidence limit		p-value	Estimate	Wald 95% confidence limit		p-value
		Lower	Upper			Lower	Upper			Lower	Upper	
All beds capacity [per 1 mln inhabitants]	0.003	0.001	0.006	0.012*	0.003	0.001	0.005	0.006*	0.003	0.001	0.005	0.004*
ICU beds capacity [per 1 mln inhabitants]	0.046	-0.008	0.101	0.095**	-0.007	-0.049	0.034	0.731	-	-	-	-
Total no. of tests up to time of the peak [per 1 mln inhabitants]	-0.002	-0.008	0.005	0.598	-	-	-	-	-	-	-	-
Stay-at-home order day	0.201	-0.248	0.649	0.381	-	-	-	-	-	-	-	-
Educational facilities closure day	0.680	0.164	1.195	0.010*	-	-	-	-	-	-	-	-
Gathering restrictions day	0.530	0.019	1.040	0.042*	-	-	-	-	-	-	-	-
Businesses closure day	0.365	-0.081	0.810	0.108	-	-	-	-	-	-	-	-
Population size [mln]	0.250	0.135	0.364	<0.001*	0.117	0.002	0.231	0.047*	0.142	0.037	0.246	0.008*
Proportion living in urban areas	0.508	-37.667	38.683	0.979	-	-	-	-	-	-	-	-
Proportion living in metropolitan cities with more than 1 mln inhabitants	13.812	-10.931	38.554	0.274	-	-	-	-	-	-	-	-
Median age	-0.278	-1.860	1.304	0.730	-	-	-	-	-	-	-	-
Arrivals at airports in 2018 [per 1 inhabitant]	-1.908	-3.573	-0.243	0.025*	-	-	-	-	-	-	-	-
No. of foreign tourists in 2018 [per 1 inhabitant]	-5.099	-8.203	-1.995	0.001*	-2.872	-5.341	-0.403	0.023*	-2.651	-5.137	-0.165	0.037*
Mobility score at the day of first reported death	0.337	0.117	0.557	0.003*	0.132	-0.092	0.356	0.249	-	-	-	-
Borders closure day	0.326	0.045	0.607	0.023*	0.215	-0.047	0.477	0.108	0.297	0.079	0.514	0.008*
No. of COVID-19 infections when borders were closed [per 1 mln inhabitants]	0.001	-0.004	0.005	0.739	-	-	-	-	-	-	-	-
No. of COVID-19 deaths when borders were closed [per 1 mln inhabitants]	0.038	-0.026	0.102	0.244	-	-	-	-	-	-	-	-
Multivariate models statistics												
Scale	-	-	-	-	8.713	6.870	11.050	-	8.901	7.018	11.289	-
AIC	-	-	-	-	259.693	-	-	-	257.146	-	-	-
AICC	-	-	-	-	265.453	-	-	-	260.257	-	-	-


*p-value < 0.05; **p-value < 0.1.

Abbreviations: AIC, Akaike's Information Criterion; AICC, AIC corrected for small sample sizes; GLM, generalised linear models; ICU, intensive care unit; mln, million.

GLM univariate and multivariate models with normal distribution and identity link function were used to explore factors associated with time to COVID-19 deaths peak (starting from the day when the first death was reported in a given country), as of 3rd June 2020. Each model was run using 34 observations. Variables significant in univariate models were included into the multivariate base case model, avoiding highly correlated pairs. The final multivariate model was selected based on the use of selection algorithms (backward, forward, stepwise and the genetic algorithm).

388
389
390
391
392
393


394 **Figure 1.** Histogram of the height of COVID-19 daily deaths peak per 1 million inhabitants
395 with a fitted log-normal curve



396
397 Histogram presents the distribution of height of COVID-19 deaths peak per 1 million inhabitants across 34 countries. A log-
398 normal curve was fitted to the distribution with scale parameter $\sigma=1.12$ and location parameter $\zeta=1.38$, assuming
399 logarithm of height of the deaths peak is normally distributed [Normal(ζ , σ)].
400

401

402 **Figure 2.** Height of COVID-19 daily deaths peak per 1 million inhabitants across countries



403
404 The plot presents the height of COVID-19 deaths peak per 1 million inhabitants across 34 countries, with Belgium having
405 outstandingly highest peak per population size (>40 deaths per 1 million inhabitants).

406

407