

Fatigue outcomes following an influenza virus pandemic: A systematic review and meta-analysis

Kim Poole-Wright¹, Fiona Gaughran^{2,3}, Rachael Evans⁴, Trudie Chalder¹

¹ Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK

² Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK

³ National Psychosis Unit, South London and Maudsley NHS Foundation Trust, London UK

⁴ University of Leicester, Leicester, UK

Author for correspondence

T. Chalder E-mail Trudie.chalder@kcl.ac.uk

ABSTRACT

Objectives: Fatigue is a pervasive clinical symptom for many infected with an influenza virus. Prior evidence from influenza pandemics suggest that fatigue symptomology may continue beyond the acute phase, lasting for several months to several years post-discharge. This living systematic review aimed to examine long-term fatigue prevalence among survivors and among communities, as well as investigate the current evidence for associated factors.

Design: Systematic review and meta-analysis.

Setting: Hospitalised and community samples.

Participants: Patient populations with a confirmed diagnosis of a named influenza virus.

Main outcomes measured: Fatigue, fatigue syndromes

Results: Ten studies met the inclusion criteria for a pooled prevalence analysis and five studies were identified as eligible for a means differences analysis. A fatigue prevalence of 41% (95% CI 0.299-0.488) was found among a total population of 1,310. Using the 'vitality'

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

lower mean vitality score for survivors compared to population norms (M -1.523, CI -13.53 – 10.48), although this was not significant ($p = 0.803$). The most common associations with fatigue were PTSD, depression and anxiety, ICU admission, female gender and higher age.

Conclusions: This study reveals that a significant proportion of survivors (41%) experienced fatigue following their recovery from an influenza virus such as SARS or MERS and that this fatigue can be long-lasting. Also, that some factors such as female gender and psychological factors may contribute to continuing fatigue outcomes for this population.

Strengths and limitations: (a) this study provides support for long-term fatigue outcomes in people with a confirmed coronavirus or other influenza virus pandemic, (b) the study suggests an aetiological role for some psychological and social factors, (c) findings are limited by the availability of fatigue data and lack of pre-morbid fatigue information; (d) a meta-analysis on the associations was prohibited by the small number of studies investigating long-term fatigue correlates and (e) the heterogeneity of the studies (>75%) suggests the pooled estimates should be interpreted with caution.

Keywords: fatigue, chronic fatigue, survivors, pandemic, long-term, coronavirus, living review

Funding Statement

This research received no specific grant from any funding agency, commercial or not-for-profit sectors. TC receives salary support from the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed in this article are those of the authors and not necessarily those of the NIHR or the NHS.

Author Declarations

T.C. is the author of several self-help books on chronic fatigue for which she has received royalties. T.C. (King's College London, KCL) has received ad hoc payments for workshops carried out in long-term conditions. KCL have received payments for T.C.'s editor role in the Journal of Mental Health. T.C. acknowledges financial support from NIHR, Health Technology Assessment, Guy's & St Thomas' Charity, Muscle Disease UK. She has a patent background IP with a software company. No fees received to date.

INTRODUCTION

Coronavirus family such as SARS-CoV Severe Acute Respiratory Syndrome (SARS), Middle East Respiratory Syndrome (MERS) and SARS-CoV-2 are single-strand RNA viruses, characterised by severe upper respiratory tract infections. A novel A(H1N1)pdm09 influenza emerged in 2009 and before this other strains, originating in animals or birds, including the influenza virus A subtype H2N2 and HPAI A(H5N1). Each has spread rapidly worldwide with considerable impact on mortality. The case fatality rate for SARS during the course of the outbreak (2002 - 2004) was 9.7% (Petersen et al., 2020). For MERS an estimated 2,494 cases were detected since its identification in 2012, with a fatality rate of 34.4% (Bradley & Bryan, 2019). Wide-ranging clinical indicators have been observed for infectious viruses, from asymptomatic or mild to the most severe potentially resulting in respiratory distress or viral pneumonia. Outcomes are dependent upon variables such as age and physical comorbidities. However, for most infectious diseases, fatigue is among the most common presenting symptom alongside high fever, myalgia, cough and breathlessness. Clinical observations for patients infected with worldwide pandemics such as SARS have indicated fatigue in 78% of patients admitted to hospital (Han et al., 2003). Meta-analyses of the current SARS-CoV-2 (COVID-19) virus indicates prevalence of fatigue to be between 22-61.9% (Qiu et al., 2020; Wang et al., 2020) and has been found to be associated with exacerbated outcomes, including acute respiratory distress and mortality. A study investigating coronavirus found that, compared with survivors, more severe fatigue was associated with mortality in a small number of patients admitted to an ICU (Wang, Shu, Ran, Xie, & Zhang, 2020). Similarly, a meta-analysis by Qiu et al. (2020) compared SARS-CoV-2 patients admitted to ICU with non-ICU patients and found an increased risk of worse pneumonia symptoms associated with fatigue severity.

Fatigue may be characterised as tiredness or exhaustion as a result of physical or mental exertion or as a result of an illness or disease (Dittner, Wessely, & Brown, 2004). The experience of fatigue is common and is usually short-lived but, for a small number of people, it can become long-lasting, associated with a number of impairments in daily living and reduced quality of life (Dittner et al., 2004).

Post-viral fatigue has been linked to viruses including Epstein-Barr and Q Fever, while chronic fatigue has been investigated in survivors of SARS, MERS and H1N1. Chronic fatigue syndrome has been found to be associated with the H1N1 virus for instance (Magnus et al., 2015) and, in 233 SARS survivors, 43% had chronic fatigue, with around 27% fulfilling the criteria for chronic fatigue syndrome, 3 years post-infection (Lam et al., 2009). A meta-analysis comparing SARS, MERS and COVID-19 found a post-illness fatigue prevalence of

19.3% (Rogers et al., 2020). Other research indicates the development of a post-SARS syndrome for some survivors (Chrousos & Kaltsas, 2005; Moldofsky & Patcai, 2011). There is emerging evidence for lasting fatigue as a consequence of COVID-19. A recent study found moderate to severe levels of fatigue in up to 54% of 100 patients treated for coronavirus, 4 weeks post-discharge (Halpin et al., 2020) and at a median follow-up stage of 10 weeks, up to 50% reporting severe fatigue (Townsend et al., 2020).

Factors affecting post-illness fatigue have been investigated in previous viral influenza outbreaks. Studies indicate that for many SARS survivors, lung function was well preserved with significant improvements between 3-12 months post-infection, although remaining lower than healthy controls (Hui et al., 2005; Ong, 2004). Similar results have been found for exercise capacity. For example, several studies have found poorer exercise performance on the six-minute walking test (6MWT) at 3, 6 and 12 months (Hui et al., 2005; Ong, 2004). These exercise impairments, however, were not correlated with lung functions. For example, in a Singapore study of 44 SARS patients, 41% had reduced exercise capacity but mild pulmonary function defects (Ong, 2004). These findings have been replicated in other SARS survivors (Hui et al., 2009; Hui et al., 2005; Ngai et al., 2010; Tansey et al., 2007) and H1N1 patients (Hsieh et al., 2018). It is these physical functioning limitations, independent of pulmonary function, which are thought to contribute to reduced quality of life and fatigue outcomes found in numerous studies of recovered patients (Hsieh et al., 2018; Tansey et al., 2007)

These results suggest that, alongside possible physical deconditioning and corticosteroid myopathy (Dekhuijzen & Decramer, 1992; Tansey et al., 2007), other psychological, social or cognitive contributions may explain the persistence of fatigue and lower QoL. For example, long periods of social isolation or stigma (Wing & Leung, 2012). Time spent in ICU (Halpin et al., 2020) and hospital length of stay (Chen et al., 2020) have been associated with fatigue for instance and psychiatric disorders such as anxiety, post-traumatic stress disorder and depression are high in post-SARS. Further, chronic fatigue has been associated with a risk of psychiatric disorder in recovered SARS patients (Wing & Leung, 2012) and with post-traumatic stress symptoms in MERS survivors (Lee et al., 2019). Social factors have been found to be a factor in post-discharge fatigue. For instance, fatigue was more likely where patients were involved in 'medicolegal' cases (Wing & Leung, 2012) and, controlling for psychiatric morbidity, being an applicant to the survivors' fund in Hong Kong (Lam et al., 2009). Furthermore, while around 76-80% of survivors achieve a return to work (Lau et al., 2005; Rogers et al., 2020), some patients reported fatigue as one of the reasons for not returning to work (Guo et al., 2019).

Summary

Fatigue for the majority of patients usually dissipates during the course of a virus, but some experience longer lasting symptoms, independent of their pulmonary functions and exercise capacity. These symptoms may be associated with a number of psychosocial factors such as psychiatric morbidity and isolation, which have been indicated with fatigue. Long-term consequences of an epidemic such as SARS, MERS or H1N1 revealed by prior research, may be pertinent to survivors of the current COVID-19 pandemic. Therefore, the aim of this systematic review was to investigate the prevalence of persistent fatigue among survivors of a viral epidemic and consolidate current evidence for factors associated with fatigue outcomes.

METHODS

Search strategy

The protocol and PICO framework for this study was developed utilising The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA). The databases Embase, PsycINFO, Medline, CINAHL, Cochrane Database of Systematic Reviews and Open Grey were searched from inception dates to (open date), using defined search terms: H1N1 OR “swine flu” OR “swine influenza OR SARS OR “severe acute respiratory syndrome” OR coronavirus OR COVID-19 OR H5N1 OR “avian flu” OR “avian influenza” OR MERS OR “Middle East respiratory syndrome” OR H3N2 OR “Hong Kong flu” OR “Hong Kong influenza” OR H2N2 OR “Asian flu” OR “Asian influenza” OR pandemic OR epidemic AND “chronic fatigue” OR fatigue OR exhaustion OR “quality of life” OR HRQoL. MeSH (Medical Subject Headings) were included in order to maximise the search. Reference lists of the selected studies were manually searched for additional articles.

Eligibility criteria

- Studies were considered eligible for inclusion if they met the following criteria:
 - Original articles available in English;
 - Studies with primary data;
 - Studies reporting fatigue or fatigue syndromes using a fatigue measure or a fatigue/tiredness related subscale;
 - Studies investigating fatigue occurring post-discharge;
 - Studies investigating patient populations diagnosed with either COVID-19, severe acute respiratory syndrome (SARS), middle east respiratory syndrome (MERS), A/H1N1pdm09 (Swine), HPAI H5N1 (Avian), H2N2 (Asian), A/H3N2 (Hong Kong) or other sub-virus;

- Any study design including cohort, case-control, cross-sectional, randomised control trials, meta-analysis.

Studies were excluded if they (1) measured compassion or pandemic fatigue (defined as 'worn out' by pandemic warnings, by government safety instructions, or with media coverage, or with compliance requirements; (2) populations without a confirmed diagnosis of one of the named viruses; (3) reported fatigue associated with physical disorders (e.g. thyroiditis, Parkinson's disease, cancer); (4) measured fatigue as a clinical symptom during the acute phase (defined as the period of hospitalisation); (5) were protocols, vaccination studies, animals; (6) did not report fatigue outcome data; (7) were newspaper articles, conference papers, commentaries or editorials. We also excluded qualitative studies and studies investigating fatigue among healthcare workers.

Data extraction

Titles and abstracts were screened by the first researcher (KPW). Full texts were screened by KPW against the eligibility criteria. A data spreadsheet was created to include extracted data from the included studies. An additional researcher independently reviewed a proportion of the included articles and the extracted data. The senior researchers (TC, FG) reviewed a number of the final included studies and data. PRISMA Flow Diagram available in Appendix A. Discrepancies were resolved via discussion and consensus with TC. The variables obtained for the data synthesis spreadsheet were: citation, population, number of participants, control group, location, virus type, follow-up period, study design, outcome variable of interest (e.g., fatigue, vitality), associated variables (e.g., PTSD, depression, anxiety, stress), scales or measures employed, power calculation (Y/N).

Quality Assessments

Risk of bias was assessed by the Critical Appraisal Skills Programme (CASP) (2018). Each study design has an appropriate checklist (e.g., cohort) comprising 12 items designed to systematically assess a study. The items do not require a score but rather a "yes", "no" or "can't tell". The randomised and systematic review checklists were developed and adapted from JAMA's guides to medical literature. An overall assessment was made by assigning a grade of 'excellent', 'good', 'moderate' or 'low' to each included article.

Statistical analysis

We computed pooled prevalence for binary fatigue outcomes with 95% confidence intervals for 10 studies. A number of studies investigated fatigue outcomes across multiple time points. Therefore, in order to maintain the independence of observations, only one time

period was included in the analysis as follows: (a) studies investigating fatigue in the weeks following an infection, we selected the period closest to 3 months and (b) studies with 12 months or longer time periods, we selected the period closest to 12 months. This was to demonstrate that the fatigue assessed was 'long-term'. Where studies investigated both 'fatigue' and CFS outcomes, we incorporated the fatigue data only in the meta-analysis. This was because the participants had not received a confirmed diagnosis of CFS in all studies. Additional meta-analysis was conducted to compute mean differences from five included studies to estimate the 'severity' of fatigue compared to population norms. For this computation, the 'Vitality' subscale of the Medical Outcomes Study 36-item Short-Form health survey (SF-36) (Ware & Sherbourne, 1992) was utilised as a proxy for fatigue. This 'vitality' subscale is scored from 0-100, with higher scores representing better functioning/less disability. For the purposes of this study lower scores represent lower 'energy' and higher fatigue. We utilised this instrument because it the most frequently used measure assessing HRQoL among virus pandemic research. The absence of data from control groups precluded other analyses except population norms. Effect sizes were first calculated from each included study to compute the mean effect. Meta-analyses were conducted using R Studio, Version 1.3.1073 (2020). Heterogeneity was assessed using Cochran Q statistic. We obtained the I^2 statistic with the degree of heterogeneity categorised as 'not important' (0-40%), 'moderate' (30-60%), 'substantial' (50-90%) and 'considerable' (75-100%) (Higgins, 2003). We conducted an Egger's test to explore potential publication bias for our proportional analysis: bias = 1.657 (95% CI -7.19-10.5), $p = 0.723$ suggesting an absence of funnel plot asymmetry. Due to the small number of studies further analysis of the origins of heterogeneity via a meta-regression or the production of funnel plots was not performed on the mean difference analysis.

Multiple studies investigated a wide-range of post-virus outcomes including psychological, physical and/or social effects. A number of these investigated the relationship between an exposure virus and fatigue outcomes. However, each used a diverse number of outcome measures or utilised non-validated scales, which disqualified the viability of a meta-analysis. Consequently, associations which had available data were arranged in tabular form illustrating the direction of the association with fatigue or vitality (Table 2). A positive symbol (+) indicated a positive association, a negative symbol (-) indicated a negative association and a zero (0) indicated no significant association between the investigated variable and fatigue (Matcham, Ali, Hotopf, & Chalder, 2015). Associations with fatigue or vitality, measured in prospective cohort designs were demonstrated by a superscript figure representing the period the relationships were examined.

RESULTS

Search results

A total of 2,557 articles were identified using the database search protocols (Figure 1). Following the removal of duplicates, 2,108 articles remained for title and abstract screening. Of these a total of 170 articles were selected for full text screening producing a final total of 32 studies meeting the eligibility criteria and 15 were deemed eligible for a quantitative analysis. A summary of the included articles is presented in Table 1. We tabulated the studies according to the fatigue and 'vitality' outcome measures.

Table 1. Summary of included studies

Author (year), country	Setting	Study Design	Virus type	No. of participants	Time (Post-illness)	Fatigue outcome	Total Fatigue prevalence no. (%)	Risk Factors for Fatigue	Risk Factors no. (%) Other results	Prevalence of Risk Factor Controls	p
Ahmed et al (2020)	-	Meta-analysis	SARS MERS	3 studies	3, 6 months	Fatigue CFS	MERS 30 (48.1) SARS 94 (40.3)	-	-	-	-
Arnold et al. (2020) UK	Outpatients	Prospective-cohort	COVID-19	110	8-12 weeks	Fatigue	43 (39)	-	-	-	-
Halpin et al. (2020) UK	Outpatient	Cross-sectional	COVID-19	100	4-8 weeks	Fatigue	64(64)	New fatigue Ward 41 (60.3) ICU 23 (72) Fatigue Severity Severe Ward 10 (14.7) ICU 4 (12.5) Fatigue severity moderate Ward 14 (20.6) ICU 13 (40.6) Fatigue Severity mild Ward 17 (25) ICU 6 (18.8) Gender Moderate/Severe fatigue Women 46 (61) Men 54 (26.6) PTSD Severe fatigue (43.9) No fatigue (18.6) Cognitive problems Moderate/Severe fatigue (41.4) Less severe fatigue (18.6) Breathlessness Moderate severe fatigue (65.9) Less severe fatigue (39) BMI NS Age ICU v. NS Ward NS Ethnicity NS			

Author (year), country	Setting	Study Design	Virus type	No. of participants	Time (Post-illness)	Fatigue outcome	Total Fatigue prevalence no. (%)	Risk Factors for Fatigue	Risk Factors no. (%) Other results	Prevalence of Risk Factor Controls	p
Lam et al. (2009) Hong Kong	Outpatient	Cohort Retrospective	SARS	124	4 years	Chronic fatigue	73 (40.3)	Multivariate Survivors' Fund Shorter duration of follow-up Univariate Psychiatric morbidity Applied to SARS survivors' fund SARS Litigation Stigma	OR 2.64 OR 0.89 39 (73) 40 (55.6) 13 (17.8) 33 (45.2)	12 (23.5) 11 (21.6) 1 (2) 7 (13.7)	<.04 .003 <.01 <.01 <.01 <.01
Lee et al. (2018) Korea	Outpatient	Prospective cohort	MERS	52	12, 18 months	CFS	12 months 25 (48.8) 18 months 17 (32.6)	Fatigue between 12 & 18 months 12 months fatigue Depression @ 12 months Depression @ 18 months PTSD @ 12 months PTSD @ 18 months 18 months fatigue Depression @ 12 months Depression @ 18 months PTSD @ 12 months PTSD @ 18 months Multivariate FSST1 (DV) Depression T1	Pearson's r = .677 r = .742 r = .630 r = .712 r = .577 r = .579 r = .698 r = .700 r = .765		<.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001
Qi et al. (2020) China	Outpatients	Cross-sectional	COVID-19	41	27 days	Fatigue Chronic fatigue	22 (53.6)	-	-	-	-
Rogers et al (2020)		Meta-analysis	SARS MERS COVID-19	4 studies	Post-illness	Fatigue	Pooled % 61 (19.3)	-	-	-	-
Tansey et al. (2007) Canada	Outpatient	Cohort	SARS	117	3, 6, 12 months	Fatigue	3 months 53 (64) 6 months 54 (54) 12 months 70 (60)	-	-	-	-
Tenforde et al. (2020) USA	Outpatient	Cross-sectional	COVID-19	192	2-3 weeks	Fatigue	68 (35)	-	-	-	-

Author (year), country	Setting	Study Design	Virus type	No. of participants	Time (Post-illness)	Fatigue outcome	Total Fatigue prevalence no. (%)	Risk Factors for Fatigue	Risk Factors no. (%) Other results	Prevalence of Risk Factor Controls	p
Townsend et al. (2020) Ireland	Outpatient	Prospective cohort	COVID-19	128	10 weeks	CFS	67 (52.3)	Total fatigue 15.8 (5.9) Physical fatigue 11.38 (4.22) Psychological fatigue 4.72 (1.99) Severe fatigue group: Female 45 (52.3) Other factors Anxiety/Depression/anti-depressant history - Days since onset - Critical care - Length of stay - BMI - Lab tests (NLR, LDH, CRP) - COVID severity -	-	-	.002
Wang et al. (2020) China	Outpatient	Prospective cohort	COVID-19	131	1, 2 3,4 weeks	Fatigue	Total @ discharge 10 (7.63) 1-2 weeks 7 (5.34) 3-4 weeks 0 (0)	@ discharge (severity of COVID) Severe 6 (9.68) Non-severe 4 (5.8) 1-2 weeks Severe 2 (3.32) Non-severe 5 (7.25) 3-4 weeks Severe 0 Non-severe 0	-	-	0.516 0.445 - -
Wing et al. (2012) Hong Kong	Outpatient	Prospective cohort	SARS	233	3 years	Chronic Fatigue CFS	41 (45.2) 16 (17.7)	With psychiatric illness (lifetime or current) Chronic fatigue CFS 67 (75) 49 (55.1) Current psychiatric disorder Chronic fatigue CFS 58 (76.5) 45 (59.3)	-	-	-

Author (year), country	Setting	Study Design	Virus type	No. of participants	Time (Post-illness)	Fatigue outcome	Total Fatigue M (SD)	P	Risk Factors for Fatigue	Risk Factor for Fatigue M (SD) Pearson's r	p
Batawi et al (2019) Saudi Arabia	Outpatient	Cross-sectional	MERS	78	1 year	Vitality	MERS 65.96 (26.47) Non-MERS 53.60 (28.80)	< .01	ICU Ward	58.47 (31.60) 72.38 (19.29)	<.03
Chen et al (2017) China	Outpatient	Longitudinal cross-sectional	H7N9	54	3, 6, 12 months 2 years	Vitality	(Chinese Norms) 3 months 73.2 (15.8) 6 months 69.7 (16.8) 12 months 72.2 (20.9) 24 months 72.8 (21.3)	<.000 <.000 <.000 .001	-	-	-
Chen et al (2020) China	Outpatient	Cross-sectional	COVID-19	361	1 month	Vitality	(Chinese norms) NR	<.05	Gender Women Men Multivariate Length of stay Age	81.80 (16.32) 83.25 (16.13) β .113 β .128	<.001 .040 .04
Gardner et al. (2015)	-	Systematic Review	SARS	20 studies	1-6, 3-36, <54 months	Vitality	NR	-	NR	NR	NR
Guo et al (2019) China	Outpatient	Cohort	SARS	67	12 months	Vitality	With Oseltamivir 64.06 Without Oseltamivir 66.11	.723	Return to work	NR	-
Hong et al (2009) China	Outpatient	Longitudinal	SARS	70	2, 7, 10, 20, 46 months	Vitality	NR	-	@ 10 months PTSD No PTSD Follow-up time	41.32 (21.53) 69.26 (4.46) NR	<.001 .0135

Author (year), country	Setting	Study Design	Virus type	No. of participants	Time (Post-illness)	Fatigue outcome	Total Fatigue M (SD)	P	Risk Factors for Fatigue	Risk Factor for Fatigue M (SD) Pearson's r	p
Lau et al (2005) Hong Kong	Outpatient	Cohort	SARS	171	2 months	Vitality	v. HK norms 49.91 (18.88)	<.001	Gender Women Men	45.74 (17.43) 58.09 (19.09)	<.001 <.001
Lau, Ng et al (2005) Hong Kong	Outpatient	RCT	SARS	133	2 – 6 weeks	Vitality	Baseline Controls 47.5 (17.6) Exercise 51.2 (19.5)	0.25	Mean change in VT Controls 2.50 (14.2) Exercise 2.50 (14.2) Return to work	0.06 (R ²)	NS .006
Li et al. (2006) Hong Kong	Outpatient	Prospective Cohort	SARS	59	3, 6, 12 Months	Vitality			Pulmonary Function FVC FEV RV TLC DLCO KCO Exercise Capacity (6MWT) Age	r = 0.38 r = 0.27 r = -0.18 r = 0.14 r = 0.25 r = -0.23 r = 0.52 NR	NS NS NS NS NS NS <0.01 NR
Liu et al (2020)	Outpatient	RCT	COVID-19	72	6 weeks	Vitality	Pulmonary rehabilitation v. none	< .05	Intervention Group Pre-rehab Post-rehab Control Group Pre-rehab Post-rehab	60.6 (6.9) 75.6 (7.1) 60.5 (7.1) 61.2 (6.3)	< .05 NS
Lo et al (2005) Singapore	Outpatient	Cross-sectional	SARS	14	6 months	Fatigue Question	SARS 4.7 (0.9) Healthy controls 2.2. (1.5)	<.005	-	-	-
Luyt et al. (2012) France	Outpatient	Cross-sectional	H1N1	37	1 year	Vitality	NR		Population norms Non-ECLA ECLA	NR NR	.006 .058
Mak et al. (2009) Hong Kong	Outpatient	Cohort	SARS	90	30 months	Vitality	SARS 48.82 (22.32) HK Norms 60.27 (18.65)	<.001	-	-	-

Author (year), country	Setting	Study Design	Virus type	No. of participants	Time (Post-illness)	Fatigue outcome	Total Fatigue M (SD)	P	Risk Factors for Fatigue	Risk Factor for Fatigue M (SD) Pearson's r	p
Moldofsky et al. (2011) Canada	Outpatient	Cohort	SARS	22	13-66 months	CFS	NR		Compared to FMA Pre-sleep fatigue Post sleep fatigue	4.57 (1.57) 4.30 (1.87)	NS
Ngai et al. (2010) Hong Kong	Outpatient	Prospective cohort	SARS	123	3, 6, 12, 18, 24 months	Vitality	NR	<.001	Age 6MWT Intubated	NR NR NR	<.01 NR NS
Skinner et al. (2015) Australia	Outpatient	Prospective cohort	H1N1	62	6, 12 months	Vitality	6 months ICU 44.5 Ward 42.1 H1N1 ICU 49.7 Non-H1N1 ARDS ICU 41.0 12 months H1N1 61.5 Non-H1N1 ARDS 50.2	NR NR 0.93	-	-	-

Study characteristics

A total of 3,262 participants were represented in the included studies. There were 13 (40.6%) studies investigating fatigue/chronic fatigue using a fatigue scale and 19 (59.4%) studies using the 'vitality' subscale of the SF-36. The most common country of origin was Hong Kong with 10 studies, followed by China (7). Singapore, Canada and the UK (2 each). France, Saudi Arabia, Ireland, Korea, Australia and USA each had 1 study. SARS was the most frequently investigated virus (21 articles), COVID-19 (9), MERS (4) and H1N1 (3). Zero studies investigated fatigue succeeding the other named viruses. There were 15 prospective-cohort designs, 7 cross-sectional, 1 case-control, 5 cohort, 2 randomised-control and 3 systematic reviews. Time periods under study ranged from 2 weeks to 12 years post-discharge or infection (for community cases) with the most frequent time point being fatigue <3 months and 12 months post-discharge. For the purposes of the meta-analysis, time periods ranged from 2 weeks to 4 years (Appendix B). CASP quality assessments resulted in most studies receiving a 'low' or 'moderate' assessments indicated bias. Lower grades were assigned for multiple reasons including selection bias, lack of adequate control groups, small sample sizes and methodological bias (employment of unvalidated/unreliable scales).

Factors associated with fatigue

Not all 32 studies investigated or reported factors associated with fatigue following a virus infection. Also, the available data for each risk factor were too few to conduct a quantified analysis, with a diverse number of outcome measures. However, associations that were available are illustrated in Table 2. Results are stratified by study design and univariate or multivariate results. Only available data was included in the table. In summary, four studies reported gender differences. ICU admission was reported in 8 studies and 6 studies reported age/fatigue relationships. Anxiety, PTSD and exercise capacity, depression and, pulmonary functions were investigated in 3 studies. Three studies investigated pulmonary function, exercise capacity and vitality outcomes. Factors such as length of stay, medicolegal involvement and shorter response to follow-up were examined in 2 studies. The remainder of risk factors were analysed by 1 study each.

Table 2. Variables associated with fatigue

Factor	Cross-sectional <i>Bivariate</i>	<i>Multivariate</i>	Prospective Cohort <i>Bivariate</i>	<i>Multivariate</i>
PTSD ↑	+, + ³		+ ¹⁰ , + ¹² , + ¹⁸	
Anxiety symptoms ↑	+		+	
Depression ↑	+, 0		+ ¹² , + ¹⁸	
Psychiatric morbidity ↑	0		+	
Pulmonary function ↓			+ ⁶ , + ¹² , 0 ¹²	
Age ↑	0, 0, -	+	0, ³ + ⁶ , + ¹² , + ³ , + ⁶ , + ¹² , + ³ , + ⁶ , + ³ , + ⁶ , + ¹² , + ¹⁸ , + ²⁴	+
ICU Admission	+, +, +, 0		0, 0 ⁶ , 0 ¹² , 0	
Medicolegal involvement	+		+	+
Female gender	+, 0, 0		+	
Ethnicity	0			
BMI	0			
Stigma	0, 0			
Returned to work	+	+		
Exercise capacity ↓			+, + ¹² , + ¹²	
Intubated			0 ⁶	
Response to follow-up <	-		-, +	+
Length of stay <		+	0	+
Education ↓	0			
Physical health ↓	0			
Steroid treatment	+			
Days since onset ↑			0	
Perceived social support ↓	0			
Cognitive problems ↑	+			
Breathlessness ↑	+			
Personal coping ↓	+			

+ = positive association with fatigue ($p < 0.05$); - = negative association with fatigue ($p < 0.05$); 0 = no association with fatigue ($p < 0.05$), ^{3,6,12} months; ↑ = higher; ↓ = lower; > = longer duration; < = shorter duration

Meta-analyses

Ten studies with a total of 1,310 participants were included for the proportional analysis using a random-effects model. A pooled prevalence from 10 studies was found to be 41% (95% CI 0.31-0.52). I^2 statistic was 93% indicating 'considerable' heterogeneity. Details of this analysis are represented by a forest plot (Figure 2). Five studies reported means and standard deviations on SF-36 'vitality' outcomes and were included for the means differences analysis. These results indicate vitality scores were -1.52 points lower for the

participants compared to population norms (95% CI 13.53-10.49). This was non-significant $p = 0.803$. A forest plot of the results is available in Figure 3.

Figure 2. Forest plot for pooled prevalence for fatigue





Figure 3. Forest plot for mean differences in vitality between survivors and population norms



DISCUSSION

This review investigated the long-term fatigue outcomes in populations who had a confirmed coronavirus or other named virus diagnosis. We found 32 studies that provided data on fatigue and chronic fatigue, CFS and 'vitality'. The majority of investigated viruses were SARS and COVID-19. MERS, H1N1 and 1 study examining H7N9 ('avian') made up the balance. No fatigue data was found among the remaining viruses. We found the pooled prevalence of fatigue to be considerable at 41% (95% CI 0.31-0.52) among 1,310 survivors and this fatigue was present between < 1 month (Tenforde et al. 2020; Wang et al., 2020) to 4 years post-infection (Lam et al., 2009). Our results are higher than one other study, which found a pooled estimate of 19.3% for fatigue in 316 survivors of SARS, MERS or COVID-19 (Rogers et al., 2020). To our knowledge no other studies have investigated fatigue systematically. The vitality subscale of the SF-36 (Ware & Sherborne, 1992) was used as a proxy for fatigue, with lower vitality scores representing higher fatigue. Our results indicated an overall lower mean for survivors compared to population norms. Although the results were not significant ($p = 0.803$), they suggest impaired functioning up to 2.5 years following exposure (Mak, Chu, Pan, Yiu, & Chan, 2009). Our results compare to previous systematic reviews examining vitality in recovered patients, which found lower mean vitality in SARS and MERS survivors compared to normative values (Ahmed et al., 2020). Similarly, in a literature review by Gardner et al. (2015), all domains of health-related quality of life measures, including vitality, were found to be lower than local population norms.

Fatigue and vitality changes over time were mixed, with some studies indicating improved scores between time points (Hong et al. 2009; Kwek et al. 2006; Lee et al. 2019) and other studies reporting no differences in fatigue for each follow-up period (Hui, Wong, et al. 2005, Hui, Joynt et al. 2005). Nevertheless, most reported lower vitality or higher fatigue compared to population norms at all time points, up to 4 years post-infection (Lam et al. 2009; Li et al. 2006). We were not able to quantify the differences in fatigue between those who were admitted to ICU compared to ward patients, however, fatigue for ICU groups was generally found to be higher compared to population norms (Hui, Wong et al. 2005; Hui, Joynt et al. 2005). Where studies compared ICU with ward groups, results were mixed, either finding a difference (Batawi et al. 2019; Halpin et al. 2020; Kwek et al, 2006) or no difference in vitality scores (Hui, Wong et al. 2005; Hui, Joynt et al. 2005).

Our study did not calculate risk factors for fatigue due to the small number of studies investigating potential correlates, diverse outcome measures and aggregated SF-36 scores. Associations between fatigue and particular lung functions, however, were reported in 2 studies. For example, FVC and VC were found to be positively correlated with vitality at 3, 6 and 12 months (Hui et al. 2005) and FEV₁ with higher vitality at 6 months (Hui et al. 2005).

Both suggest lower fatigue is associated with better pulmonary performance. Contrary evidence from an Australian H1N1 study, found that lung functions and fatigue for 62 survivors had improved to within normal range at 1-year post-discharge (Skinner et al., 2015). Exercise capacity, measured by the 6MWT, was lower than norms in most studies and correlated with fatigue in 3 studies (Hui et al., 2005; Li et al., 2006; Ngai et al., 2010). A systematic review found that mean walking distance was lower for all survivors compared to norms (Ahmed et al., 2020) with a 'levelling' of test results at 12 and 18 months (Ngai et al., 2010). Such limitations resulting from critical illness hospitalisation, are thought to contribute to reduced quality of life and fatigue outcomes in recovered patients (Herridge et al., 2003). However, these effects were observed independent of physical functioning. For example, in a study of 9 H1N1 recovered patients given pulmonary rehabilitation, there was a significant decrease in quality of life between 3 to 6 months, despite improved lung capacities and exercise capacity (Hsieh et al., 2018). In a study examining 1-year outcomes of SARS patients, Tansey et al. (2007) found that, while lung functions were within normal ranges, many reported persistent fatigue and shortness of breath.

Psychological factors were found to be significant factors for SARS and MERS survivors in systematic reviews (Gardner & Moallem, 2015; Rogers et al., 2020). Ahmed et al. (2020) found a pooled estimate for PTSD, depression and anxiety among SARS and MERS survivors to be 39%, 33% and 30% respectively. Among the included studies in our study chronic fatigue had a 'reciprocal' association with psychiatric diagnosis (Wing & Leung, 2012), while lower vitality was associated with PTSD symptomology (Halpin et al., 2020; Hong et al., 2009; Kwek et al., 2006), anxiety and depression (Kwek et al., 2006). Recovered MERS patients with chronic fatigue 1-year post exposure, had persistent PTSD symptoms mediated via depression 18 months post-infection (Lee et al., 2019). Moreover, those with psychiatric diagnosis were more likely to experience chronic fatigue 1 year later (Lam et al., 2009). In contrast, Qi et al., (2020) found no association between psychiatric morbidity and chronic fatigue in recovered COVID-19 patients.

Our study suggests that survivors of a serious influenza virus experienced considerable and long-lasting distress, physical impairments and psychological effects, including fatigue. The persistence of physical and psychological symptoms has been identified as a post-SARS sickness syndrome (Chrousos & Kaltsas, 2005). A syndrome distinguished by fatigue, myalgia, general weakness, psychological distress and sleep disorders. In respect of the current epidemic, there is potential for longer-term fatigue outcomes following a COVID-19 infection and current evidence points to a syndrome-like collection of symptoms. However, there are few studies examining probable correlates between fatigue or vitality outcomes

with physical and pulmonary functioning; psychological or social functioning in recovering patients. Future studies should investigate such risk factors to help inform clinical interventions to address 'long COVID' symptomology. The generalisability of our results should be applied with caution due to the diversity of measurements, distance between investigation and exposure (i.e., < 1 month or > 2 years), variability in populations, different admission and discharge protocols and lung function reference ranges between countries (Chan, 2005).

References

- Ahmed, H., Patel, K., Greenwood, D. C., Halpin, S., Lewthwaite, P., Salawu, A., ... Sivan, M. (2020). Long-term clinical outcomes in survivors of severe acute respiratory syndrome and Middle East respiratory syndrome coronavirus outbreaks after hospitalisation or ICU admission: A systematic review and meta-analysis [Article]. *Journal of Rehabilitation Medicine*, 52(5), 0. <https://doi.org/10.2340/16501977-2694>
- Bradley, B. T., & Bryan, A. (2019). Emerging respiratory infections: The infectious disease pathology of SARS, MERS, pandemic influenza, and Legionella. *Seminars in Diagnostic Pathology*, 36(3), 152–159. <https://doi.org/10.1053/j.semmdp.2019.04.006>
- Chen, K. Y., Li, T., Gong, F. H., Zhang, J. S., Li, X. K., Chen, K.-Y., ... Zhang, J.-S. (2020). Predictors of Health-Related Quality of Life and Influencing Factors for COVID-19 Patients, a Follow-Up at One Month. *Frontiers in Psychiatry*, 11, 668. <https://doi.org/http://dx.doi.org/10.3389/fpsy.2020.00668>
- Chrousos, G. P., & Kaltsas, G. (2005). Post-SARS sickness syndrome manifestations and endocrinopathy: How, why, and so what? [Article]. *Clinical Endocrinology*, 63(4), 363–365. <https://doi.org/http://dx.doi.org/10.1111/j.1365-2265.2005.02361.x>
- Dekhuijzen, P. N., & Decramer, M. (1992). Steroid-induced myopathy and its significance to respiratory disease: a known disease rediscovered. *The European Respiratory Journal*, 5(8), 997–1003. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/1426209>
- Dittner, A. J., Wessely, S. C., & Brown, R. G. (2004). The assessment of fatigue: a practical guide for clinicians and researchers. *Journal of Psychosomatic Research*, 56(2), 157–170. [https://doi.org/10.1016/S0022-3999\(03\)00371-4](https://doi.org/10.1016/S0022-3999(03)00371-4)
- Gardner, P. J., & Moallem, P. (2015). Psychological impact on SARS survivors: Critical review of the English language literature. [Article]. *Canadian Psychology/Psychologie Canadienne*, 56(1), 123–135. <https://doi.org/10.1037/a0037973>
- Guo, L., Han, Y., Li, J., Chen, Q., Ren, Y., Wu, Q., ... Zhang, M. (2019). Long-term outcomes in patients with severe acute respiratory syndrome treated with oseltamivir: a

- 12-year longitudinal study. *International Journal of Clinical and Experimental Medicine*, 12(10), 12464–12471. Retrieved from www.ijcem.com/
- Halpin, S J, Mclvor, C., Whyatt, G., Adams, A., Harvey, O., McLean, L., ... Sivan, M. (2020). Postdischarge symptoms and rehabilitation needs in survivors of COVID-19 infection: A cross-sectional evaluation. *Journal of Medical Virology*, 1–10.
<https://doi.org/http://dx.doi.org/10.1002/jmv.26368>
- Halpin, Stephen J., Mclvor, C., Whyatt, G., Adams, A., Harvey, O., McLean, L., ... Sivan, M. (2020). Postdischarge symptoms and rehabilitation needs in survivors of COVID-19 infection: A cross-sectional evaluation. *Journal of Medical Virology*, jmv.26368.
<https://doi.org/10.1002/jmv.26368>
- Han, Y., Geng, H., Feng, W., Tang, X., Ou, A., Lao, Y., ... Li, Y. (2003). A follow-up study of 69 discharged SARS patients. *Journal of Traditional Chinese Medicine*, 23(3), 214–217.
- Higgins, J. P. T. (2003). Measuring inconsistency in meta-analyses. *BMJ*, 327(7414), 557–560. <https://doi.org/10.1136/bmj.327.7414.557>
- Hsieh, M-J., Lee, W-C., Cho, H-Y., Wu, M-F., Hu, H-C., ... Huang, C-C. (2018). Recovery of pulmonary functions, exercise capacity, and quality of life after pulmonary rehabilitation in survivors of ARDS due to severe influenza A (H1N1) pneumonitis. *Influenza and Other Respiratory Viruses*, 12(5), 643–648. <https://doi.org/10.1111/irv.12566>
- Hui, D. S. C., Wong, K. T., Antonio, G. E., Tong, M., Chan, D. P., & Sung, J. J. Y. (2009). Long-term sequelae of SARS: physical, neuropsychiatric, and quality-of-life assessment. [Article]. *Hong Kong Medical Journal = Xianggang Yi Xue Za Zhi*, 15 Suppl 8, 21–23. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20393208>
- Hui, D. S., Joynt, G. M., Wong, K. T., Gomersall, C. D., Li, T. S., Antonio, G., ... Sung, J. J. Y. (2005). Impact of severe acute respiratory syndrome (SARS) on pulmonary function, functional capacity and quality of life in a cohort of survivors. *Thorax*, 60(5), 401–409.
<https://doi.org/http://dx.doi.org/10.1136/thx.2004.030205>
- Hui, David S, Wong, K. T., Ko, F. W., Tam, L. S., Chan, D. P., Woo, J., & Sung, J. J. Y. (2005). The 1-year impact of severe acute respiratory syndrome on pulmonary function, exercise capacity, and quality of life in a cohort of survivors. *CHEST*, 128(4), 2247–2261. <https://doi.org/10.1378/chest.128.4.2247>
- Lam, M. H.-B., Wing, Y. K., Yu, M. W. M., Leung, C. M., Ma, R. C. W., Kong, A., ... Lam, S. P. (2009). Mental morbidities and chronic fatigue in severe acute respiratory syndrome survivors: Long-term follow-up. *Archives of Internal Medicine*, 169(22), 2142–2147.
<https://doi.org/10.1001/archinternmed.2009.384>
- Lam, M., Wing, Y., Yu, M., Leung, C., Ma, R., Kong, A., ... Lam, S. (2009). Mental morbidities and chronic fatigue in severe acute respiratory syndrome survivors: Long-term follow-up. *Archives of Internal Medicine*, 169(22), 2142–2147.


- <https://doi.org/10.1001/archinternmed.2009.384>
- Lau, H. M.-C. C., Ng, G. Y.-F. F., Jones, A. Y.-M. M., Lee, E. W.-C. C., Siu, E. H.-K. K., & Hui, D. S.-C. C. (2005). A randomised controlled trial of the effectiveness of an exercise training program in patients recovering from severe acute respiratory syndrome. *Australian Journal of Physiotherapy*, *51*(4), 213–219. [https://doi.org/10.1016/S0004-9514\(05\)70002-7](https://doi.org/10.1016/S0004-9514(05)70002-7)
- Lee, S. H., Shin, H.-S., Park, H. Y., Kim, J. L., Lee, J. J., Lee, H., ... Han, W. (2019). Depression as a mediator of chronic fatigue and post-traumatic stress symptoms in middle east respiratory syndrome survivors. *Psychiatry Investig*, *16*(1), 59–64. <https://doi.org/http://dx.doi.org/10.30773/pi.2018.10.22.3>
- Magnus, P., Gunnes, N., Tveito, K., Bakken, I. J., Ghaderi, S., Stoltenberg, C., ... L., T. (2015). Chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME) is associated with pandemic influenza infection, but not with an adjuvanted pandemic influenza vaccine. *Vaccine*, *33*(46), 6173–6177. <https://doi.org/http://dx.doi.org/10.1016/j.vaccine.2015.10.018>
- Mak, I. W. C., Chu, C. M., Pan, P. C., Yiu, M. G. C., & Chan, V. L. (2009). Long-term psychiatric morbidities among SARS survivors. *General Hospital Psychiatry*, *31*(4), 318–326. <https://doi.org/10.1016/j.genhosppsych.2009.03.001>
- Matcham, F., Ali, S., Hotopf, M., & Chalder, T. (2015, July 1). Psychological correlates of fatigue in rheumatoid arthritis: A systematic review. *Clinical Psychology Review*, Vol. 39, pp. 16–29. <https://doi.org/10.1016/j.cpr.2015.03.004>
- Moldofsky, H., & Patcai, J. (2011). Chronic widespread musculoskeletal pain, fatigue, depression and disordered sleep in chronic post-SARS syndrome; a case-controlled study [Article]. *BMC Neurology*, *11*(1), 37. <https://doi.org/10.1186/1471-2377-11-37>
- Ngai, J. C., Ko, F. W., Ng, S. S., To, K.-W. W., Tong, M., & Hui, D. S. (2010). The long-term impact of severe acute respiratory syndrome on pulmonary function, exercise capacity and health status. *Respirology (Carlton, Vic.)*, *15*(3), 543–550. <https://doi.org/10.1111/j.1440-1843.2010.01720.x>
- Ong, K.-C. (2004). Pulmonary function and exercise capacity in survivors of severe acute respiratory syndrome. *European Respiratory Journal*, *24*(3), 436–442. <https://doi.org/10.1183/09031936.04.00007104>
- Petersen, E., Koopmans, M., Go, U., Hamer, D. H., Petrosillo, N., Castelli, F., ... Simonsen, L. (2020). Comparing SARS-CoV-2 with SARS-CoV and influenza pandemics. *The Lancet Infectious Diseases*, *20*(9), e238–e244. [https://doi.org/10.1016/S1473-3099\(20\)30484-9](https://doi.org/10.1016/S1473-3099(20)30484-9)
- Qiu, P., Zhou, Y., Wang, F., Wang, H., Zhang, M., Pan, X., ... Liu, J. (2020). Clinical characteristics, laboratory outcome characteristics, comorbidities, and complications of

- related COVID-19 deceased: a systematic review and meta-analysis. *Aging Clinical and Experimental Research*. <https://doi.org/10.1007/s40520-020-01664-3>
- Rogers, J. P., Chesney, E., Oliver, D., Pollak, T. A., McGuire, P., Fusar-Poli, P., ... G., L. (2020). Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic [Article]. *The Lancet Psychiatry*, 7(7), 611–627. [https://doi.org/10.1016/S2215-0366\(20\)30203-0](https://doi.org/10.1016/S2215-0366(20)30203-0)
- Tansey, C. M., Louie, M., Loeb, M., Gold, W. L., Muller, M. P., de Jager, J., ... A.S., S. (2007). One-year outcomes and health care utilization in survivors of severe acute respiratory syndrome. *Archives of Internal Medicine*, 167(12), 1312–1320. <https://doi.org/http://dx.doi.org/10.1001/archinte.167.12.1312>
- Townsend, L., Dyer, A. H., Jones, K., Dunne, J., Kiersey, R., Gaffney, F., ... Bergin, C. (2020). Full Title: Persistent fatigue following SARS-CoV-2 infection is common and independent of severity of initial infection Short title: Fatigue following SARS-CoV-2 infection. *MedRxiv*, 2020.07.29.20164293. <https://doi.org/10.1101/2020.07.29.20164293>
- Wang, L., Shen, Y., Li, M., Chuang, H., Ye, Y., Zhao, H., & Wang, H. (2020). Clinical manifestations and evidence of neurological involvement in 2019 novel coronavirus SARS-CoV-2: a systematic review and meta-analysis. *Journal of Neurology*, 1, 3. <https://doi.org/10.1007/s00415-020-09974-2>
- Wang, Z.-H., Shu, C., ... Zhang, L. (2020). Critically Ill Patients with Coronavirus Disease 2019 in a Designated ICU: Clinical Features and Predictors for Mortality. *Risk Management and Healthcare Policy*, Volume 13, 833–845. <https://doi.org/10.2147/RMHP.S263095>
- Ware, J. E., & Sherbourne, C. D. (1992). *The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection* (Vol. 30, pp. 473–483). Vol. 30, pp. 473–483. Retrieved from <https://doi.org/10.1097/00005650-199206000-00002>
- Wing, Y. K., & Leung, C. M. (2012). Mental health impact of severe acute respiratory syndrome: a prospective study. *Hong Kong Medical Journal = Xianggang Yi Xue Za Zhi*, 18 Suppl 3(4), 24–27. Retrieved from <https://pubmed.ncbi.nlm.nih.gov/22865219/>

APPENDIX A



PRISMA 2009 Flow Diagram



APPENDIX B

Studies included in pooled prevalence meta-analysis

Study	Virus Type	Scale	Post-discharge
Lam et al. 2009	SARS	Chalder Fatigue Scale	4 years
Lee et al. 2019	MERS	Fatigue Severity Scale	12 months
Tansey et al. 2007	SARS	Fatigue question	12 months
Townsend et al. 2020	COVID-19	Chalder Fatigue Scale	2.5 months
Qi et al. 2020	COVID-19	Chalder Fatigue Scale	1 month
Wing et al. 2012	SARS	Fatigue question	3 years
Halpin et al. 2020	COVID-19	Interview question	1 month
Tenforde et al. 2020	COVID-19	Fatigue questionnaire	< 1 month
Wang, Xu et. al. 2020	COVID-19	Fatigue question	< 1 month
Arnold et al. 2020	COVID-19	Fatigue question	3 months

Studies included in means differences meta- analysis

Study	Virus Type	Scale	Post-discharge
Chen, Wu et al. 2017	H7N9	SF-36	3 months
Lam et al. 2006	SARS	SF-36	< 3 months
Lau, Lee et al. 2005	SARS	SF-36	< 3 months
Lo et al. 2005	SARS	ADQ	6 months
Mak et al. 2009	SARS	SF-36	30 months