

# 1 Risk of thrombotic complications in influenza 2 versus COVID-19 hospitalized patients

3 Dutch COVID & Thrombosis Coalition (DCTC)

4

5 Corresponding author: MV Huisman MD, PhD FESC; Department of Thrombosis and Hemostasis,  
6 Leiden University Medical Center, Leiden, the Netherlands; Albinusdreef 2, 2300RC, Leiden, the  
7 Netherlands; Phone: +31-715269111; E-mail: [m.v.huisman@lumc.nl](mailto:m.v.huisman@lumc.nl)

8

## 9 **Abstract**

10 **Background:** Whereas accumulating studies on COVID-19 patients report high incidences of  
11 thrombotic complications, large studies on clinically relevant thrombosis in patients with other  
12 respiratory tract infections are lacking. How this high risk in COVID-19 patients compares to those  
13 observed in hospitalized patients with other viral pneumonias such as influenza is unknown.

14 **Objectives:** To assess the incidence of venous and arterial thrombotic complications in hospitalized  
15 influenza patients as opposed to that observed in hospitalized COVID-19 patients.

16 **Methods:** Retrospective cohort study; we used data from Statistics Netherlands (study period: 2018)  
17 on thrombotic complications in hospitalized influenza patients. In parallel, we assessed the  
18 cumulative incidence of thrombotic complications – adjusted for competing risk of death - in patients  
19 with COVID-19 in three Dutch hospitals (February 24th - April 26th 2020).

20 **Results:** Of the 13.217 hospitalized influenza patients, 437 (3.3%) were diagnosed with thrombotic  
21 complications, versus 66 (11%) of the 579 hospitalized COVID-19 patients. The 30-day cumulative  
22 incidence of any thrombotic complication in influenza was 11% (95%CI 9.4-12) versus 25% (95%CI 18-  
23 32) in COVID-19. For venous thrombotic complications (VTE) and arterial thrombotic complications  
24 alone, these numbers were respectively 3.6% (95%CI 2.7-4.6) and 7.5% (95%CI 6.3-8.8) in influenza  
25 versus 23% (95%CI 16-29) and 4.4% (95%CI 1.9-8.8) in COVID-19.

26 **Conclusions:** The incidence of thrombotic complications in hospitalized influenza patients was lower  
27 than in hospitalized COVID-19 patients. This difference was mainly driven by a high risk of VTE  
28 complications in the COVID-19 patients admitted to ICU. Remarkably, influenza patients were more  
29 often diagnosed with arterial thrombotic complications.

30

**NOTE:** This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

1    **Essentials:**

- 2           • It is unknown how COVID-19 compares to patients with other virus infections regarding
- 3           thrombosis.
- 4           • Hospitalized patients with influenza and COVID-19 were evaluated and compared to each
- 5           other.
- 6           • 30-day cumulative incidence of thrombosis was lower in influenza (11%) than in COVID-19
- 7           (25%).
- 8           • Difference was mainly driven by a high risk of VTE in COVID-19 patients admitted to the ICU.

9

10

11

12    **Keywords:**

13    COVID-19

14    Incidence

15    Influenza, Human

16    Pneumonia, Viral

17    Thrombosis

18

19

20

## 1 **Introduction**

2 The clinical course of acute infections may become complicated by venous and arterial thrombotic  
3 disease.[1, 2] Precipitating factors for thrombotic complications in this setting include inflammation,  
4 activation of the coagulation system, immobilization, and diffuse intravascular coagulation (DIC).[1,  
5 2] In light of this, international guidelines recommend pharmacological thromboprophylaxis in  
6 patients with infectious diseases if they are hospitalized.[3]  
7 Respiratory viruses, including influenza, were already known to lead to a procoagulant state.[4-6]  
8 Moreover, previous studies showed an transient increase in the incidence of thrombotic vascular  
9 complications after respiratory tract infections.[1, 7] Still, to date it is largely unknown how often  
10 influenza infection leads to thrombotic complications, since only small case series yielding conflicting  
11 results have been published.[4, 8-10] Contrarywise, the COVID-19 pandemic has led to a large  
12 number of cohort studies on thrombotic complications in hospitalized COVID-19 patients. These  
13 studies showed that COVID-19 infections were characterized by a sometimes excessive activation of  
14 blood coagulation [11-13] and high incidences of thrombotic complications were reported in patients  
15 with COVID-19, aggravated by admittance to the Intensive Care Unit (ICU).[14-19] How this high  
16 incidence of thrombotic complications in COVID-19 patients compares to those observed in  
17 hospitalized patients with non-COVID-19 virus infections such as influenza is largely unknown.[20]  
18 This is especially relevant to understand the etiology of COVID-19 associated thrombosis.[21, 22]  
19 We evaluated the incidence of thrombotic complications in hospitalized patients with influenza as  
20 opposed to that in hospitalized patients with COVID-19, in an effort to better understand the  
21 epidemiology of COVID-19 associated thrombotic complications.

22

## 23 **Methods**

### 24 *Setting and study population*

25 We used anonymous information on registered diagnoses during hospital admissions in Dutch  
26 hospitals from the National Basic Register of Hospital Care of Dutch Hospital Data (DHD), which

1 includes all general and academic Dutch hospitals, provided via Statistics Netherlands (“Centraal  
2 Bureau voor de Statistiek”, CBS ), for information on hospitalized influenza patients. Patients  
3 hospitalized with an admission date between January 1st 2018 and November 30th 2018 with a  
4 diagnosis of influenza were included for the primary analysis and data from January 1<sup>st</sup> 2013 until  
5 December 31<sup>st</sup> 2018 were used to collect information on comorbidities. Information on the use of  
6 pharmacological thromboprophylaxis was not available in DHD.  
7 We also used data from adult COVID-19 patients admitted to the wards and ICUs of one university  
8 hospital (Leiden University Medical Center, Leiden, the Netherlands) and two non-university teaching  
9 hospitals (Amphia Hospital Breda and Alrijne Hospital Leiderdorp, both in the Netherlands) between  
10 February 24<sup>th</sup> and April 26<sup>th</sup> 2020. COVID-19 was confirmed by a positive polymerase chain reaction  
11 (PCR) test or considered positive in patients with a negative PCR but highly suggestive symptoms and  
12 typical COVID-19 abnormalities on CT-scan of the chest (CO-RADS 4 or 5 following Dutch Radiology  
13 Society[23]) with no alternative diagnosis. All COVID-19 patients had received pharmacological  
14 thromboprophylaxis from admission on, according to local hospital protocols, which changed over  
15 time (Appendix 3). Informed consent was obtained by an opt-out approach. This study was approved  
16 by the Institutional Review Boards of the LUMC for observational studies.

17

### 18 *Objectives*

19 The primary objective was to assess the incidence of venous and arterial thrombotic complications in  
20 hospitalized influenza patients (general ward and ICU combined) and compare this with that  
21 observed in our cohort of hospitalized COVID-19 patients.

22

### 23 *Data collection*

24 A diagnosis of influenza in the allocated time frame was identified in the DHD data based on  
25 International Classification of Diseases (ICD) codes (Appendix 1). For those patients who had more  
26 than one influenza related hospitalization, only the first hospitalization was included. In these data, a

1 differentiation is made between admission to a normal ward only or (temporarily) admission to an  
2 ICU, but it was not possible to distinguish exact days of admittance on the general ward versus ICU.  
3 Therefore, influenza patients in the DHD database were categorized as hospitalized at the ward or  
4 ICU by the following definition: 1. Ward: 'hospitalizations that did not involve ICU admission'  
5 (includes patients that were only admitted to the wards) and 2. ICU: 'hospitalizations that did involve  
6 ICU admission' (includes patients that were admitted to the ICU at some timepoint during  
7 hospitalization). Patient characteristics of included influenza patients as age, sex, and comorbidities  
8 were collected. A history of comorbidities as malignancy and prior history of venous  
9 thromboembolism (VTE) were screened for within 8 years before the influenza related  
10 hospitalization. Prior to 2018, categorization of the hospitalization as ward and/or ICU was not  
11 possible as this information was not available in the CBS database. Therefore, we studied the  
12 incidence of thrombotic complications in influenza related hospitalizations during influenza seasons  
13 between 2013 and 2017 as a sensitivity analysis, rather than as a main analysis. Influenza seasons  
14 started from the 40<sup>th</sup> week of one year to the 20<sup>th</sup> week of the next year.  
15 The patient charts from COVID-19 patients were retrospectively scrutinized for baseline  
16 characteristics and outcomes of interest. For COVID-19 we were able to strictly distinguish general  
17 ward from ICU admission by exact days of admittance, but we used the influenza categorization of  
18 ward and ICU in order to be consistent.

19

## 20 *Outcomes*

21 Information on thrombotic complications in influenza was identified by ICD codes (Appendix 1). Any  
22 VTE (DVT and/or PE) or arterial thrombotic complication (ATE: ischemic stroke, myocardial infarction  
23 and/or systemic arterial embolism) that occurred during the index hospitalization was regarded an  
24 outcome event. The exact date of the thrombotic complication during admission was unavailable in  
25 DHD. Therefore, the date halfway the hospitalization period was chosen as the date of the  
26 thrombotic complication in influenza patients.

1 In COVID-19 patients, information on thrombotic complications was identified based on scrutinizing  
2 patient charts. Thrombotic complications consisted of acute pulmonary embolism (PE), deep-vein  
3 thrombosis (DVT), ischemic stroke, myocardial infarction and systemic arterial embolism. No VTE  
4 screening strategies were applied during this study in COVID-19 patients. Upon clinical suspicion of  
5 one of the thrombotic complications, appropriate diagnostic tests were applied, i.e. computed  
6 tomography pulmonary angiography (CTPA) for suspected PE and compression ultrasonography  
7 (CUS) for suspected DVT, cardiac enzymes including troponin, electrocardiogram and  
8 echocardiography for suspected acute coronary syndrome, and CT scan of the brain and CT  
9 angiography of the carotid and intracerebral arteries for suspected ischemic stroke.

10

#### 11 *Statistical analysis*

12 Patient characteristics were described using standard descriptive statistics. The analysis on the  
13 comparison between influenza and COVID-19 patients, comprised the full time of hospitalization  
14 (ward and ICU combined). Hospitalized influenza patients were followed from the day of hospital  
15 admission (i.e., the starting date) until day of discharge, day of death or end of data collection (31th  
16 December 2018), whichever came first. COVID-19 patients were followed from the day of hospital  
17 admission until discharge, transfer to another hospital, until they died, or until end of data collection  
18 (between April 17 and 26, 2020), whichever came first.

19 The incidence of thrombotic complications was estimated for all ICU and non-ICU patients combined,  
20 patients admitted to general wards only and for patients who had been admitted to the ICU at some  
21 timepoint during hospitalization (according to the DHD categorization of ward and ICU), and for all  
22 thrombotic complications as well as for venous and arterial complications separately. Cumulative  
23 incidences were estimated using the Kaplan-Meier method and the cumulative incidence competing  
24 risk (CICR) method, to adjust for the competing risk of death. In addition, incidence rates were  
25 calculated over the total follow-up period.

26 SPSS Statistics version 25.0 and R version 4.0.2 served for data analysis.

## 1 **Results**

### 2 *Patients*

3 A total of 13,217 hospitalized influenza patients and 579 hospitalized COVID-19 patients were  
4 included. Characteristics of both patient cohorts are summarized in Table 1. Patients with influenza  
5 and COVID-19 were comparable with regard to age, history of cancer, and length of hospital stay  
6 (median and IQR: five days (3-10) in influenza and seven days (4-11) in COVID-19). There were more  
7 male patients with COVID-19 than with influenza.

8

### 9 *Thrombotic complications in influenza patients*

10 Of the influenza patients, 437/13,217 patients (3.3%) were diagnosed with thrombotic complications,  
11 of whom 126 with venous and 319 with arterial complications. The 30-day cumulative incidence of all  
12 thrombotic complications, adjusted for competing risk of death, was 11% (95%CI 9.4-12) (Table 2).

13 The 30-day adjusted cumulative incidence for venous thrombotic complications was 3.6% (95%CI 2.7-  
14 4.6) and for arterial thrombotic complications 7.5% (95%CI 6.3-8.8). The incidence-rates in  
15 hospitalized influenza patients were 1.6/patient-year (95%CI 1.4-1.7) for any thrombotic  
16 complication, 0.44/patient-year (95%CI 0.37-0.53) for venous thrombotic complications and  
17 1.1/patient-year (95%CI 1.0-1.3) for arterial thrombotic complications. The incidences calculated for  
18 the year 2018 were comparable with previous years (2013-2017; Appendix 2).

19 For influenza patients that had only been admitted to the ward, the adjusted 30-day cumulative  
20 incidences for all thrombotic complications, venous thrombotic complications and arterial  
21 thrombotic complications were: 8.9% (95%CI 7.5-10), 2.4% (95%CI 1.8-3.2) and 6.6% (95%CI 5.4-8.0),  
22 respectively. From these ward admitted patients, 766 out of 12,412 died during hospitalization  
23 (6.2%). In influenza patients that had been admitted to the ICU, the adjusted 30-day cumulative  
24 incidences for all thrombotic complications, venous thrombotic complications and arterial  
25 thrombotic complications were: 18% (95%CI 14-23), 7.7% (95%CI 5.0-11) and 11% (95%CI 7.9-15),  
26 respectively (Table 3). In this group of patients, 171 out of 805 died during hospital admission (21%).

1

## 2 *Thrombotic complications in COVID-19 patients*

3 Of the 579 COVID-19 patients, 66 patients (11%) were diagnosed with 70 thrombotic complications.

4 PE was the most frequent thrombotic complication of all (n=54, 77%). The 30-day cumulative

5 adjusted incidence of all thrombotic complications was 25% (95%CI 18-32) (Table 2). The 30-day

6 adjusted cumulative incidence for venous thrombotic complications was 23% (95%CI 16-29) and for

7 arterial thrombotic complications 4.4% (95%CI 1.9-8.8). Incidence-rates in COVID-19 patients were

8 5.1/patient-year (95%CI 3.9-6.4) for any thrombotic complication, 4.5/patient-year (95%CI 3.4-5.8)

9 for venous thrombotic complications and 0.78/patient-year (95%CI 0.39-1.4) for arterial thrombotic

10 complications.

11 For the COVID-19 patients that had only been admitted to the ward, the adjusted 30-day cumulative

12 incidences for all thrombotic complications, VTE alone and ATE alone were: 6.1% (95%CI 3.1-11),

13 4.4% (95%CI 1.8-8.8) and 1.7% (95%CI 0.61-3.9), respectively. From these ward admitted patients, 66

14 out of 401 died during hospitalization (17%). In COVID-19 patients that had been admitted to the ICU

15 at any moment during the in-hospital stay, the adjusted 30-day cumulative incidences for all

16 thrombotic complications, VTE alone and ATE alone were: 36% (95%CI 28-45), 34% (95%CI 26-43)

17 and 5.3% (95%CI 2.0-11), respectively (Table 3). In this group of patients, 45 out of 178 died during

18 hospital admission (25%).

19

## 20 **Discussion**

21 In this study we observed that, compared to hospitalized influenza patients, patients admitted with

22 COVID-19 had a distinctly increased risk for getting thrombotic complications by day 30. This risk

23 was driven by a difference in venous thrombotic complications (3.6% in influenza and 23% in COVID-

24 19) and was particularly observed in patients admitted to the ICU.

25 How often influenza infection led to clinically relevant thrombotic disease was hitherto

26 underreported, since only small case series have been published, yielding conflicting results.[4, 8-10]



1 Nevertheless, respiratory viruses, including influenza, were already known to lead to a procoagulant  
2 state.[4-6] Previous studies for instance showed a transient increase in the risk of vascular  
3 complications after respiratory tract infections, with a twofold increase for venous thrombotic  
4 disease and an even fivefold increase for ischemic heart disease.[1, 7] Of note, in our study arterial  
5 thrombotic complications, and in particular myocardial infarction, occurred more in influenza  
6 patients than in COVID-19 patients. A recent study showed stroke rates quite similar to our  
7 study.[24] Strokes occurred in 0.2% in the influenza patients and in 1.6% in the COVID-19 patients  
8 [24], in our study this was 0.8% in the influenza patients and 1.7% in the COVID-19 patients.  
9 Furthermore, another Chinese study was recently published, which compared the incidence of VTE in  
10 patients hospitalized with COVID-19 versus those hospitalized with community-acquired pneumonia  
11 (CAP).[20] Surprisingly, this study found no difference in VTE rates between CAP and COVID-19.  
12 However, COVID-19 patients in this study were overall younger and had fewer comorbidities.[20] In  
13 addition, arterial thrombotic complications were not included in this study and VTE risk is known to  
14 be lower in East Asians than in white Caucasians, which limits generalizing these findings to other  
15 populations. A recent large cross-sectional study on 80,261 influenza patients illustrated that the  
16 high risk of arterial thrombotic complications coincides with influenza virus infection.[25] Therefore,  
17 it is reasonable to believe that viral pneumonias such as influenza virus and SARS-CoV-2 virus both  
18 interact with the coagulation system.[4]

19 Virus specific features could play an important role and maybe explain the observed difference in  
20 thrombotic complications between influenza and COVID-19 patients.[4, 26, 27] Recent studies  
21 suggest the possibility of in-situ immunothrombosis in COVID-19 patients, due to a SARS-CoV-2  
22 specific effect. Following this hypothesis, COVID-19 associated alveolar injury and an extreme  
23 inflammatory response contribute to small vessel thrombus formation in the lungs. Autopsy studies  
24 support this hypothesis by reporting the presence of platelet-fibrin thrombi in small arterial vessels  
25 of the lungs.[22, 26, 28] On the other hand, other studies observed a high incidence of DVT upon  
26 screening as well, which is compatible with the conventional thromboembolic origin of PE.[29, 30]

1 Therefore, both mechanisms likely play a role in the pathophysiology of COVID-19 associated  
2 thrombosis and further research in this field is needed.

3 Methodological strengths of our study include the large sample size and multicenter study design. In  
4 addition, VTE screening strategies have not been applied during the course of this study and  
5 diagnostic tests were only performed upon clinical suspicion. Limitations of our study regarding  
6 influenza data are that not all baseline patient characteristics were available, since these could only  
7 be determined during a specific time period before hospital admission, and that exact dates of  
8 thrombotic complications were not available. Furthermore, we do not know whether, and if so,  
9 which dose thrombosis prophylaxis was given to the hospitalized influenza patients. National  
10 guidelines in the Netherlands recommend pharmacological thromboprophylaxis with low molecular  
11 weight heparin (LMWH) in ward admitted patients in the case of a PADUA Prediction Score  $\geq 4$  and in  
12 the absence of contra-indications. All patients admitted to the ICUs receive pharmacological  
13 thromboprophylaxis, irrespective of the PADUA Prediction Score.[31] Standard practice in the LUMC,  
14 Amphia Hospital and Alrijne Hospital is the administration of nadroparin 2850 IU sc. per day. It is  
15 likely that these recommendations were followed for the hospitalized influenza patients, but still we  
16 do not know for sure. In addition, outcome classification differed between influenza and COVID-19  
17 patients, as information on thrombotic complications was identified in influenza based on ICD codes  
18 and in COVID-19 by patient chart review. Besides, it is important to recall that thrombotic risks in  
19 influenza may differ between the seasonal influenza virus – from which patients were included in this  
20 study - and the H1N1 influenza virus pandemic from the year 2009/2010. For COVID-19 data,  
21 limitations include the different doses of thrombosis prophylaxis being prescribed over time and the  
22 lowering threshold to suspect VTE over the course of time in COVID-19 patients. In addition,  
23 thrombosis in COVID-19 due to in-situ immunothrombosis could have been underestimated, since  
24 this diagnosis is often only established after death in pathology reports. Still we were interested in  
25 clinically relevant confirmed thrombosis and believe that this underestimation will likely be minimal,  
26 especially given this lowering threshold to suspect VTE in COVID-19. Furthermore, compared to

1 influenza patients, we observed a higher mortality rate in the COVID-19 ward admitted patients,  
2 which indicates that the COVID-19 ward admitted patients had more severe illness than the influenza  
3 ward admitted patients. Although this could (partially) explain the observed difference in thrombotic  
4 risk between the ward admitted patients in both groups, this cannot explain differences found in  
5 thrombotic complications between patients that had been admitted to ICU, as mortality rates in  
6 these patients were quite comparable, but a large difference was found in (especially venous)  
7 thrombotic complications. Altogether, the comparison between influenza and COVID-19 patients  
8 could have been biased because of the difference in outcome classification and the lowered  
9 threshold for diagnostic testing upon suspicion of VTE in COVID-19 patients was probably lower than  
10 in influenza patients after the high incidence of thrombotic complications in COVID-19 was  
11 elucidated.

12

13 In conclusion, this study suggests that the incidence of thrombotic complications in hospitalized  
14 influenza patients is lower than in hospitalized COVID-19 patients. The difference in thrombotic  
15 complications in our study was particularly driven by an exceptionally high risk for VTE in the COVID-  
16 19 patients that had been admitted to ICU. Remarkably, influenza patients were more often  
17 diagnosed with arterial thrombotic complications than with VTE. Further studies are needed to  
18 substantiate our findings and to explore explanations for this difference.

19

20

21 **The Dutch COVID & Thrombosis Coalition study group**

22 *Authors:*

23 MAM Stals, MD: Department of Thrombosis and Hemostasis, Leiden University Medical Center,  
24 Leiden, the Netherlands. Gathered and verified data, performed the analyses and primarily drafted  
25 the first version of the manuscript.

- 1 MJJH Grootenboers, MD: Department of Pulmonology, Amphia Hospital Breda, the Netherlands.
- 2 Gathered data and revised the manuscript critically for important intellectual content.
- 3 C van Guldener, MD: Department of Internal Medicine, Amphia Hospital Breda, the Netherlands.
- 4 Revised the manuscript critically for important intellectual content.
- 5 FHJ Kaptein, MD: Department of Thrombosis and Hemostasis, Leiden University Medical Center,
- 6 Leiden, the Netherlands. Gathered data and revised the manuscript critically for important
- 7 intellectual content.
- 8 SJE Braken: Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden,
- 9 the Netherlands. Gathered data and revised the manuscript critically for important intellectual
- 10 content.
- 11 Q Chen, MD: Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, the
- 12 Netherlands. Analysed the DHD Influenza data and revised the manuscript critically for important
- 13 intellectual content.
- 14 G Chu, MD: Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden,
- 15 the Netherlands. Gathered data and revised the manuscript critically for important intellectual
- 16 content.
- 17 EM van Driel, MD: Department of Intensive Care Medicine, Alrijne Hospital Leiderdorp, the
- 18 Netherlands. Revised the manuscript critically for important intellectual content.
- 19 A Iglesias del Sol, MD: Department of Internal Medicine, Alrijne Hospital Leiderdorp, the Netherlands.
- 20 Revised the manuscript critically for important intellectual content.
- 21 E de Jonge, MD: Department of Intensive Care Medicine, Leiden University Medical Center, Leiden,
- 22 the Netherlands. Revised the manuscript critically for important intellectual content.
- 23 KM Kant, MD: Department of Intensive Care Medicine, Amphia Hospital Breda, the Netherlands.
- 24 Revised the manuscript critically for important intellectual content.
- 25 F Pals: Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden, the
- 26 Netherlands. Gathered data and revised the manuscript critically for important intellectual content.

1 MMA Toorop, MD: Department of Clinical Epidemiology, Leiden University Medical Center, Leiden,  
2 the Netherlands. Analysed the DHD Influenza data and revised the manuscript critically for important  
3 intellectual content.

4 SC Cannegieter, PhD: Department of Thrombosis and Hemostasis, Leiden University Medical Center,  
5 Leiden, the Netherlands; Department of Clinical Epidemiology, Leiden University Medical Center,  
6 Leiden, the Netherlands. Designed the study and provided methodological input. In addition, revised  
7 the manuscript critically for important intellectual content.

8 FA Klok, MD: Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden,  
9 the Netherlands. Designed the study, performed the analyses and drafted the first version of the  
10 manuscript and revised the manuscript critically for important intellectual content.

11 MV Huisman, MD: Department of Thrombosis and Hemostasis, Leiden University Medical Center,  
12 Leiden, the Netherlands. Designed the study, verified data, performed the analyses and drafted the  
13 first version of the manuscript and revised the manuscript critically for important intellectual  
14 content.

15

16 All authors agree with the final version.

17

18 **Contributors:**

19 See **Appendix A**.

20

21 **Acknowledgements**

22 The authors thank Statistics Netherlands for providing data from the Dutch Hospital Data registry.

23

24 **Disclosures**

25 Frederikus Klok reports research grants from Bayer, Bristol-Myers Squibb, Boehringer-Ingelheim,

26 Daiichi-Sankyo, MSD and Actelion, the Dutch Heart foundation, ZonMW Dutch Healthcare Fund and

1 the Dutch Thrombosis association, all outside the submitted work. Menno Huisman reports grants  
2 from ZonMW Dutch Healthcare Fund, and grants and personal fees from Boehringer-Ingelheim,  
3 Pfizer-BMS, Bayer Health Care, Aspen, Daiichi-Sankyo, Leo Pharma, all outside the submitted work.  
4 The other authors having nothing to disclose.

5

#### 6 **Role of the funding source**

7 This study was funded by unrestricted grants of the participating hospitals. The steering committee,  
8 consisting of the authors, had final responsibility for the study design, oversight, and data verification  
9 and analyses. The sponsor was not involved in the study. All members of the steering committee  
10 contributed to the interpretation of the results, approved the final version of the manuscript, and  
11 vouch for the accuracy and completeness of the data reported. The corresponding author (MV  
12 Huisman) had full access to all the data in the study and takes responsibility for the integrity of the  
13 data and the accuracy of the data analysis. The final decision to submit the manuscript was made by  
14 the corresponding author on behalf of all co-authors.

15

#### 16 **Data sharing**

17 Anonymous data on hospitalized COVID-19 patients as collected in this study can be made available  
18 to others, on request to the corresponding author of this manuscript (MV Huisman). The request will  
19 be granted after approval of a methodologically sound proposal and a signed data sharing agreement  
20 is required. Data will be available after publication of the article and until five years following article  
21 publication. Unfortunately, data on hospitalized influenza patients cannot be shared with third  
22 parties as Statistics Netherlands does not permit this. No other relevant documents are available.

23

24

25

## 1 References

- 2 1 Smeeth L, Cook C, Thomas S, Hall AJ, Hubbard R, Vallance P. Risk of deep vein thrombosis and  
3 pulmonary embolism after acute infection in a community setting. *Lancet*. 2006; **367**: 1075-9.
- 4 2 Tichelaar YI, Kluin-Nelemans HJ, Meijer K. Infections and inflammatory diseases as risk  
5 factors for venous thrombosis. A systematic review. *Thromb Haemost*. 2012; **107**: 827-37.
- 6 3 Kahn SR, Lim W, Dunn AS, Cushman M, Dentali F, Akl EA, et al. Prevention of VTE in  
7 nonsurgical patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American  
8 College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012; **141**: e195S-  
9 e226S.
- 10 4 Goeijenbier M, van Wissen M, van de Weg C, Jong E, Gerdes VE, Meijers JC, et al. Review:  
11 Viral infections and mechanisms of thrombosis and bleeding. *J Med Virol*. 2012; **84**: 1680-96.
- 12 5 Wong RS, Wu A, To KF, Lee N, Lam CW, Wong CK, et al. Haematological manifestations in  
13 patients with severe acute respiratory syndrome: retrospective analysis. *Bmj*. 2003; **326**: 1358-62.
- 14 6 Giannis D, Ziogas IA, Gianni P. Coagulation disorders in coronavirus infected patients: COVID-  
15 19, SARS-CoV-1, MERS-CoV and lessons from the past. *J Clin Virol*. 2020; **127**: 104362.
- 16 7 Smeeth L, Thomas SL, Hall AJ, Hubbard R, Farrington P, Vallance P. Risk of myocardial  
17 infarction and stroke after acute infection or vaccination. *N Engl J Med*. 2004; **351**: 2611-8.
- 18 8 van Wissen M, Keller TT, Ronkes B, Gerdes VE, Zaaier HL, van Gorp EC, et al. Influenza  
19 infection and risk of acute pulmonary embolism. *Thromb J*. 2007; **5**: 16.
- 20 9 Obi AT, Tignanelli CJ, Jacobs BN, Arya S, Park PK, Wakefield TW, et al. Empirical systemic  
21 anticoagulation is associated with decreased venous thromboembolism in critically ill influenza A  
22 H1N1 acute respiratory distress syndrome patients. *J Vasc Surg Venous Lymphat Disord*. 2019; **7**: 317-  
23 24.
- 24 10 Bunce PE, High SM, Nadjafi M, Stanley K, Liles WC, Christian MD. Pandemic H1N1 Influenza  
25 Infection and Vascular Thrombosis. *Clin Infect Dis*. 2011; **52**: e14-e7.
- 26 11 Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor  
27 prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost*. 2020; **18**: 844-7.
- 28 12 Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized  
29 Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *Jama*. 2020; **323**: 1061-  
30 9.
- 31 13 Helms J, Tacquard C, Severac F, Leonard-Lorant I, Ohana M, Delabranche X, et al. High risk of  
32 thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study.  
33 *Intensive Care Med*. 2020. doi: 10.1007/s00134-020-06062-x.
- 34 14 Lodigiani C, Iapichino G, Carenzo L, Cecconi M, Ferrazzi P, Sebastian T, et al. Venous and  
35 arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in  
36 Milan, Italy. *Thromb Res*. 2020; **191**: 9-14.
- 37 15 Poissy J, Goutay J, Caplan M, Parmentier E, Duburcq T, Lassalle F, et al. Pulmonary Embolism  
38 in COVID-19 Patients: Awareness of an Increased Prevalence. *Circulation*. 2020; **142**: 184-6.
- 39 16 Middeldorp S, Coppens M, van Haaps TF, Foppen M, Vlaar AP, Müller MCA, et al. Incidence of  
40 venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost*. 2020; **18(8)**:  
41 1995-2002.
- 42 17 Klok FA, Kruip M, van der Meer NJM, Arbous MS, Gommers D, Kant KM, et al. Incidence of  
43 thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res*. 2020; **191**: 145-7.
- 44 18 Klok FA, Kruip M, van der Meer NJM, Arbous MS, Gommers D, Kant KM, et al. Confirmation  
45 of the high cumulative incidence of thrombotic complications in critically ill ICU patients with COVID-  
46 19: An updated analysis. *Thromb Res*. 2020; **191**: 148-50.
- 47 19 Thomas W, Varley J, Johnston A, Symington E, Robinson M, Sheares K, et al. Thrombotic  
48 complications of patients admitted to intensive care with COVID-19 at a teaching hospital in the  
49 United Kingdom. *Thromb Res*. 2020; **191**: 148-50.

- 1 20 Mei F, Fan J, Yuan J, Liang Z, Wang K, Sun J, et al. Comparison of Venous Thromboembolism  
2 Risks Between COVID-19 Pneumonia and Community-Acquired Pneumonia Patients. *Arterioscler*  
3 *Thromb Vasc Biol.* 2020; **40(9)**: 2332-7.
- 4 21 Varga Z, Flammer AJ, Steiger P, Haberecker M, Andermatt R, Zinkernagel AS, et al.  
5 Endothelial cell infection and endotheliitis in COVID-19. *Lancet.* 2020; **395**: 1417-8.
- 6 22 Ackermann M, Verleden SE, Kuehnel M, Haverich A, Welte T, Laenger F, et al. Pulmonary  
7 Vascular Endothelialitis, Thrombosis, and Angiogenesis in Covid-19. *N Engl J Med.* 2020;**383**: 120-8.
- 8 23 Prokop M, Everdingen Wv, Vellinga TvR, Ufford JQv, Stöger L, Beenen L, et al. CO-RADS – A  
9 categorical CT assessment scheme for patients with suspected COVID-19: definition and evaluation.  
10 *Radiology.* 2020;**296(2)**: E97-104.
- 11 24 Merkler AE, Parikh NS, Mir S, Gupta A, Kamel H, Lin E, et al. Risk of Ischemic Stroke in  
12 Patients With Coronavirus Disease 2019 (COVID-19) vs Patients With Influenza. *JAMA Neurology.*  
13 2020. 10.1001/jamaneurol.2020.2730.
- 14 25 Chow EJ, Rolfes MA, O'Halloran A, Anderson EJ, Bennett NM, Billing L, et al. Acute  
15 Cardiovascular Events Associated With Influenza in Hospitalized Adults. *Ann Intern Med.* 2020. doi:  
16 10.7326/M20-1509.
- 17 26 Carsana L, Sonzogni A, Nasr A, Rossi RS, Pellegrinelli A, Zerbi P, et al. Pulmonary post-mortem  
18 findings in a series of COVID-19 cases from northern Italy: a two-centre descriptive study. *Lancet*  
19 *Infect Dis.* doi: 10.1016/S1473-3099(20)30434-5.
- 20 27 Wichmann D, Sperhake JP, Lutgehetmann M, Steurer S, Edler C, Heinemann A, et al. Autopsy  
21 Findings and Venous Thromboembolism in Patients With COVID-19: A Prospective Cohort Study. *Ann*  
22 *Intern Med.* 2020. 10.7326/m20-2003.
- 23 28 Dolhnikoff M, Duarte-Neto AN, de Almeida Monteiro RA, da Silva LFF, de Oliveira EP, Saldiva  
24 PHN, et al. Pathological evidence of pulmonary thrombotic phenomena in severe COVID-19. *J Thromb*  
25 *Haemost.* 2020; **18**: 1517-9. 10.1111/jth.14844.
- 26 29 Llitjos J-F, Leclerc M, Chochois C, Monsallier J-M, Ramakers M, Auvray M, et al. High  
27 incidence of venous thromboembolic events in anticoagulated severe COVID-19 patients. *J Thromb*  
28 *Haemost.* 2020; **18**: 1743-6.
- 29 30 Santoliquido A, Porfida A, Nesci A, De Matteis G, Marrone G, Porceddu E, et al. Incidence of  
30 Deep Vein Thrombosis among non-ICU Patients Hospitalized for COVID-19 Despite Pharmacological  
31 Thromboprophylaxis. *J Thromb Haemost.* 2020. doi:10.1111/jth.14992.
- 32 31 NIV. Richtlijn Antitrombotisch beleid: Nederlandse Internisten Vereniging; [updated  
33 December 2015]. Available from:  
34 [https://internisten.nl/files/Richtlijn%20Antitrombotisch%20beleid\\_def.pdf](https://internisten.nl/files/Richtlijn%20Antitrombotisch%20beleid_def.pdf)



1 **Tables and figures**

2 **Table 1:** Characteristics of hospitalized influenza patients and hospitalized COVID-19 patients

	Influenza (N=13217)*	COVID-19 (N=579)*
Age (mean, SD)	69 (19)	67 (13)
Male sex (number, %)	6306 (48)	380 (66)
Body weight in kg (median, IQR)	NA	84 (73-95)
Cancer history (number, %) <sup>~</sup>	2343 (18)	86 (15)
Prior history VTE (number, %) <sup>~</sup>	404 (3.1)	NA
Therapeutic anticoagulation at admission (number, %)	NA	77 (13)
Length of hospital stay in days (median, IQR)	5 (3-10)	7 (4-11)

3 Note: SD: standard deviation; IQR: interquartile range; VTE: venous thromboembolism; n: number; NA : not  
4 available

5 \* Ward and Intensive Care Unit patients combined

6 <sup>~</sup> Within influenza this could only be determined within 8 years before date of admission

1 **Table 2:** Cumulative incidences of thrombotic complications in hospitalized influenza patients versus  
2 hospitalized COVID-19 patients

	<b>Influenza~</b> <b>Crude</b> % (95% CI)	<b>Influenza~</b> <b>Adjusted^</b> % (95% CI)	<b>COVID-19*</b> <b>Crude</b> % (95% CI)	<b>COVID-19*</b> <b>Adjusted^</b> % (95% CI)
All thrombotic complications				
10 days	4.8% (4.3-5.4)	4.6% (4.1-5.2)	13% (9.3-17)	12% (8.9-15)
20 days	8.9% (7.8-10)	8.2% (7.3-9.2)	30% (21-38)	24% (18-30)
30 days	12% (10-14)	11% (9.4-12)	32% (22-41)	25% (18-32)
VTE				
10 days	1.3% (0.97-1.6)	1.2% (0.95-1.5)	12% (8.2-15)	11% (7.7-14)
20 days	2.8% (2.2-3.5)	2.6% (2.1-3.2)	27% (19-36)	21% (16-28)
30 days	4.1% (2.9-5.2)	3.6% (2.7-4.6)	29% (20-38)	23% (16-29)
ATE				
10 days	3.6% (3.2-4.1)	3.5% (3.1-4.0)	2.4% (0.83-4.0)	2.2% (1.1-4.1)
20 days	6.3% (5.4-7.2)	5.8% (5.0-6.7)	3.9% (0.57-7.2)	3.2% (1.4-6.4)
30 days	8.4% (6.9-10)	7.5% (6.3-8.8)	5.7% (0.8-11)	4.4% (1.9-8.8)

3 Note: All thrombotic complications: VTE and ATE together; VTE: venous thrombotic complications; ATE: arterial

4 thrombotic complications; CI: confidence interval

5 ~13217 influenza patients (ward and Intensive Care Unit combined)

6 \*579 COVID-19 patients (ward and Intensive Care Unit combined)

7 ^Adjusted: cumulative incidence adjusted for competing risk of death

8

1 **Table 3:** Adjusted cumulative incidences of thrombotic complications in ward and ICU Influenza and COVID-19  
2 patients\*

	<b>Influenza ward</b> <b>N=12412</b> % (95% CI)	<b>COVID-19 ward</b> <b>N=401</b> % (95% CI)	<b>Influenza ICU</b> <b>N=805</b> % (95% CI)	<b>COVID-19 ICU</b> <b>N=178</b> % (95% CI)
All thrombotic complications				
10 days	4.3% (3.8-4.9)	4.7% (2.6-7.9)	6.8% (5.1-8.9)	21% (15-28)
20 days	7.2% (6.2-8.2)	6.1% (3.1-11)	13% (10-17)	35% (27-44)
30 days	8.9% (7.5-10)	6.1% (3.1-11)	18% (14-23)	36% (28-45)
VTE				
10 days	1.1% (0.80-1.4)	3.1% (1.4-5.9)	2.5% (1.5-3.9)	21% (15-27)
20 days	2.2% (1.7-2.8)	4.4% (1.8-8.8)	4.9% (3.2-7.1)	33% (25-42)
30 days	2.4% (1.8-3.2)	4.4% (1.8-8.8)	7.7% (5.0-11)	34% (26-43)
ATE				
10 days	3.4% (2.9-3.9)	1.7% (0.61-3.9)	4.4% (3.0-6.1)	2.5% (0.83-5.9)
20 days	5.2% (4.4-6.1)	1.7% (0.61-3.9)	8.7% (6.4-11)	3.8% (1.3-8.5)
30 days	6.6% (5.4-8.0)	1.7% (0.61-3.9)	11% (7.9-15)	5.3% (2.0-11)

3 \*Following the classification of ward (hospitalizations that did not involve ICU) and ICU (hospitalizations that

4 did involve ICU) as is done in the CBS hospitalization data

5 Note: All thrombotic complications: VTE and ATE together; VTE: venous thrombotic complications; ATE: arterial

6 thrombotic complications; N= number; CI: confidence interval

7 Adjusted: cumulative incidence adjusted for competing risk of death

8