

- 1 Knowledge, attitude and practice toward COVID-19 among
- 2 healthcare workers in public health facilities, Eastern Ethiopia.
- 3 Alinoor Mohamed Farah¹
- 4 Email: alinuriana@yahoo.com
- 5 Tahir Yousuf Nour¹
- 6 Email: daahir571@gmail.com
- 7 Muse Obsiye
- 8 Email: museibnuobsiye22@gmail.com
- 9 Mowlid Aqil Adan¹
- 10 Email: mowlidakil@gmail.com
- 11 Omar Moeline Ali¹
- 12 Email: yaanyo571@gmail.com
- 13 Muktar Arab Hussein¹
- 14 Email: muktararab59@gmail.com
- 15 Abdullahi Bedel Budul¹
- 16 Email: abdulahi.bbh@gmail.com
- 17 Muktar Omer¹
- 18 Email: muktaomer82@gmail.com
- 19 Fentabil Getnet¹
- 20 Email: b.infen4ever@gmail.com
- 21 Authors' Information:
- 22 *¹Corresponding Author
- 23 *¹Correspondence and reprint requests should be addressed to:
- 24 Alinoor Mohamed Farah
- 25 College of Medicine and Health Sciences,
- 26 Jigjiga University, Jigjiga, Ethiopia
- 27 Lecturer at Jigjiga University
- 28 Phone: +251911053913
- 29 Email: alinuriana@yahoo.com
- 30 ¹ Department of Public Health, College of Medicine and Health
- 31 Sciences, Jigjiga University, Ethiopia

32 **Abstract**

33 On 13 March 2020, Ethiopia reported the first confirmed case of
34 COVID-19 in Addis Ababa. COVID-19 is likely to overwhelm an
35 already fragile health- care delivery system and reduce the
36 availability of services for endemic health concerns such as
37 malaria and diarrheal diseases.

38 Cross sectional study was conducted on health care workers in
39 three public health facilities in Somali region to assess knowledge,
40 attitude and practice towards COVID-19. T-test and ANOVA were
41 used to analyze the relationship between the dependent, and
42 independent variables. Spearman's correlation was used to assess
43 the relationship between mean knowledge and attitude scores.

44 A vast majority of the participants were male (n = 293, 67.5%),
45 with a mean age of 27.6 (SD: 5.3) years. The mean knowledge
46 score was 13.7 (SD: 2.6) and the mean attitude score 10.5 (SD:
47 4.1). Only 45.2 % (n = 196) of the participants had a good attitude
48 toward COVID-19. There was a negative correlation between
49 knowledge scores, attitude scores ($r=-0.295$, $P<0.001$) and practice
50 ($r=-0.298$, $P<0.001$).

51 The overall level of knowledge was good. However, the attitude
52 and practice were relatively low. We recommend strategies for
53 enhancing the capacity of healthcare workers to develop positive
54 attitude and practice.

55 **Key words:** covid-19, knowledge, attitude, practice, health care
56 workers, Eastern Ethiopia

57

58

59

60 **Background**

61 A cluster of pneumonia cases in Wuhan, China, was reported to the
62 World Health Organization (WHO) on 31 December 2019. The
63 cause of the pneumonia cases was identified as the 2019 novel
64 coronavirus. On 30 January 2020, in response to this serious
65 situation, the World Health Organization (WHO) declared it a
66 public health emergency of international concern on January 30
67 and called for collaborative efforts of all countries to prevent the
68 rapid spread of COVID-19 (Organization, 2005) and on 11 March
69 2020, declared COVID-19 a pandemic(Organization, 2020d). In a
70 period of just 11 weeks from January to mid-March 2020, COVID-
71 19 has spread rapidly from its epicenter in Wuhan City to the rest
72 of the world. At the time of writing of current work, there are more
73 than 25 million confirmed covid-19 cases globally and almost
74 950,000 associated deaths(Organization, 2020b).

75 COVID-19 is highly infectious, and its main clinical symptoms
76 include fever, dry cough, fatigue, myalgia, and dyspnea. The most
77 commonly reported symptoms are fever, cough, myalgia or
78 fatigue, pneumonia, and complicated dyspnea, whereas less
79 common reported symptoms include headache, diarrhea,
80 hemoptysis, runny nose, and phlegm-producing cough(Chen et al.,
81 2020; Control & Prevention, 2019; Huang et al., 2020; Novel,
82 2020). Fatality rate was higher among middle-aged and elderly
83 patients with pre-existing medical condition (tumor surgery,
84 cirrhosis, hypertension, coronary heart disease, diabetes, and
85 Parkinson's disease). The virus is also affecting the health of
86 younger adults(Li et al., 2020).

87 The pandemic is adding to the burden of endemic infectious
88 diseases that prevail in many countries with an ongoing
89 humanitarian response, such as cholera, measles, malaria, HIV and

90 tuberculosis. Pre-existing poor hygiene practices, poor coverage in
91 water and sanitation services and overcrowded living conditions
92 also augment the incidence and spread of contamination by the
93 virus.

94 Ethiopia reported the first confirmed case of COVID-19 in Addis
95 Ababa on 15 March 2020. As of 3rd of August, there were a total
96 of 18,706 COVID-19 confirmed cases and 310 associated deaths. In
97 Somali region, the first case was identified on 24 of April 2020 and
98 as of 3rd of August, a total of 704 cases were identified with 18
99 associated deaths. There are already a limited number of health
100 workers in the country and in particular Somali region, putting
101 extreme strain on capacity to serve patients, especially for non-
102 emergency care (EPHI, 2020).

103 Thus, COVID-19 is likely to overwhelm an already fragile health-
104 care delivery system and reduce the availability of services for
105 endemic health concerns such as malaria and diarrheal diseases
106 including cholera (Poole et al., 2020). Ethiopia's safe water supply
107 is 30 per cent, serving as the leading cause of communicable
108 diseases in the country.

109 To date, no specific antiviral treatment has been confirmed to be
110 effective against COVID-19, and there is no vaccine so far
111 (Cascella et al., 2020). Regarding infected patients with COVID-
112 19, it has been recommended to apply appropriate symptomatic
113 treatment and supportive care (Control & Prevention, 2019; Huang
114 et al., 2020). Therefore, applying preventive measures to reduce
115 the spread of disease is of utmost importance.

116 In this regard, airborne precautions and other protective measures
117 have been discussed and proposed for prevention. Recommended
118 preventive measures include regular hand washing with soap,
119 social/physical distancing, and respiratory hygiene (covering

120 mouth and nose while coughing or sneezing) (Control &
121 Prevention, 2019; Organization, 2020c).

122 The WHO also issued guidelines on the use of face masks in
123 different settings including in the community, home based care,
124 and in the health care settings of COVID-19(Organization, 2020a).
125 In this guideline, health care workers are recommended to use
126 facemask such as those certified N95 when performing aerosol-
127 generating procedures and to use medical masks when providing
128 care to suspected or confirmed COVID-19 cases. However, these
129 effective prevention and control practices depends on awareness
130 and compliance among health workers at all levels.

131 A poor level of knowledge has been implicated in the rapid spread
132 of the infection in health facilities (McCloskey & Heymann,
133 2020; Selvaraj et al., 2018) and delay of treatment(Hoffman &
134 Silverberg, 2018), and may put patients' lives at risk. Therefore,
135 this study is aimed to investigate the knowledge, attitude and
136 practice of health workers towards COVID-19 infection. The
137 findings may be useful in recommending any remedial measures
138 and additional interventions in the study area to improve
139 awareness, attitude and practice among health care workers.

140

141

142

143

144

145

146

147

148 **Method and materials**

149 **Study design, setting and period**

150 An institution based cross-sectional study conducted from June to
151 August 2020 at one public hospital and two health centers in
152 Jigjiga town. The referral hospital is a multidisciplinary specialized
153 teaching hospital with 351 inpatient beds and the other one is
154 district hospital with 193 inpatient beds. Currently, it provides
155 health services to more than one million inhabitants in its
156 catchment area. During data collection, Karamada hospital which
157 one of the facilities selected, was turned into covid-19 treatment
158 center and was not accessible. Most of their staff were reassigned
159 to other health faculties to support the provision of basic health
160 services.

161 **Sample size determination**

162 Employees that were invited to participate in this study were all
163 health care workers employed by the public health facilities in
164 Jigjiga city council, Somali region of Ethiopia. 686 respondents
165 believed to have patient care or specimen contact such as
166 physicians, nurses, midwives, health officers, laboratory
167 professions, x-ray professionals, pharmacists/druggists,
168 anesthetists and biomedical professionals.

169 **Sampling technique**

170 All public health facilities in Jigjiga city council of Somali region
171 were identified and all health facilities were selected. The sample
172 size of 686 was allocated to each facility proportional to the size of
173 health professionals who were working during data collection. The
174 health professionals were stratified by their professions in each

175 selected facility and all health workers were selected from each
176 stratum.

177 **Inclusion and exclusion criteria**

178 Only full-time employee (health care workers) who are potentially
179 at high-risk (physicians, medical laboratory technologists, nurses,
180 and midwives), available during the data collection period and who
181 are ready to take part in the study were included.

182 **Data collection and quality control**

183 A structured self-administered questionnaire was used to collect
184 the data. The questionnaire was designed as per Ahmed M. Asaad'
185 study towards the Middle East Respiratory Syndrome Coronavirus
186 (MERs CoV) and also adapted from the current interim guidance
187 and information for healthcare workers published by the CDC,
188 updated on March 7, 2020(CDC, 2020; Giao et al., 2020). The self-
189 administered questionnaire consisting of socio-demographic
190 questions, and 35 questions based on knowledge, attitude and
191 infection control practices related to COVID-19 disease in the
192 healthcare setting.

193 Information sheet and consent forms was included in the first part
194 of the questionnaire. The study participants were requested to read
195 the informed-consent statements before they proceeded to the next
196 section. Study participants who unwilling to take part in the study
197 were not forced to proceed with the questions. The questionnaires
198 were collected daily, checked for completeness and any incomplete
199 questionnaire the respondent were contacted for completion. In
200 addition, timely supervision of the data collection process was
201 done by the investigators. During collection of the questionnaires
202 precautionary measures such as wearing of masks, physical
203 distancing was observed.

204

205

206

207 **Methods of data analysis**

208 Data was coded and entered into Epi info version 3.5.1 software
209 and exported into STATA version 14.1 for analysis. Knowledge
210 and practice questions were scored as 1 or 0 for correct and
211 incorrect responses, respectively. The total knowledge score varied
212 between 0 (with no correct answer) and 15 (for all correct
213 answers), and a cut off level of ≤ 12 was considered as poor
214 knowledge, and >12 indicated good knowledge. The question
215 regarding the practice was fourteen (with minimum score 14 and
216 maximum score 42). The score of the practice based on 3 points, in
217 which the score of 1 to 3 was given from always to never. A mean
218 score >18.8 (answering for always) was considered as having good
219 practice and a score of ≤ 18.5 indicated a poor practice (answering
220 never or occasionally).

221 Whereas, attitude responses were provided with 1, 2, 3, 4 or 5 for
222 “Strongly Agree”, “Agree”, “Neutral”, “Disagree” and “Strongly
223 Disagree”, respectively. The question regarding the attitude was
224 six (with minimum score of 6 and maximum score of 30). The
225 score of the attitude was based on 5 points Likert scale, in which
226 the score of 1 to 5 was given from strongly disagree to strongly
227 agree. A mean score >10.5 (answering for strongly agree or agree)
228 was considered as a favorable attitude and a score of ≤ 10.5 was
229 considered negative attitude (answering strongly disagree or
230 disagree or neutral).

231 Summary statistics such as frequencies and proportions were
232 computed as appropriate. Chi square test was used to compare
233 categorical variables and ratios. T-test and ANOVA were used to

234 analyze the relationship between the dependent (knowledge,
235 attitude and practice), and independent variables (demographic
236 characteristics of the participants). Spearman's correlation was
237 used to assess the relationship between mean knowledge and
238 attitude scores. All the differences of estimated variables were
239 considered statistically significant if $P < 0.05$.

240

241 **Ethical considerations**

242 Ethical clearance and support letters were obtained from the
243 Ethical Review Committee (ERC) of the College of Medicine and
244 Health Sciences, Jigjiga University. The support letter then
245 submitted to the public health facilities. Then, permission was
246 obtained from the health facilities director and department/section
247 heads. Study participants were informed about the purpose and
248 importance of the study through written informed consent before
249 the data collection process. In addition, participants who are
250 unwilling to take part in the study and those who need to quit their
251 participation at any stage were informed to do so without any
252 restriction.

253

254

255

256

257

258

259

260 **Result**

261 **Socio-demographic characteristics**

262 A total of 434 health workers completed the survey questionnaire.
263 Of the 686 HCWs approached, total of 434 HCWs responded
264 (response rate = 63%). A vast majority of the participants were
265 male (n = 293, 67.5%), with a mean age of 27.6 (SD: 5.3) years
266 and below 40 years of age (n = 423, 97.5%). Majority of the
267 participants were working in Referral Hospital (n = 345, 79.5%).
268 Three hundred twenty-two (74.2%) participants were nurses, 36
269 (8.3%) were physician, 31 (7.1%) were laboratory
270 technician/technologist, 26(6.0%) were pharmacist, 12(2.8%) were
271 x-ray technicians and 7 (1.6%) were dentist. Of the 434
272 participants, 307 (70.7%) had more than 5 years' experience. The
273 main sources of information about COVID-19 among participants
274 were information from international health organizations like the
275 CDC and WHO, Ministry of Health, Ethiopia media sites, News
276 Media and social media such as WhatsApp and Facebook. Table 1
277 summarizes the sociodemographic characteristics of the
278 participants.

279

280

281

282

283

284

285

286

287 **Table 1: Sociodemographic characteristics of the participants**

Variable	Frequency (n)	%
Sex		
Male	293	67.5
Female	141	32.5
Age (Mean, SD)		
18-39	423	97.5
>40	11	2.5
Qualification		
Nurse (including midwives)	322	74.2
Physician	36	8.3
Pharmacist	26	6.0
Dentist	7	1.6
Laboratory technologist	31	7.1
X-ray physician	12	2.8
Place of work		
Referral hospital	345	79.5
Karamara hospital	33	7.6
Ablele HC	44	10.1
Ayrdage HC	12	2.8
Work experience (Mean, SD)		
0-4 years	127	29.3
>5 years	307	70.7
Source of information on covid-19		
WHO	241	55.5
Government site and media	105	24.2
Social media	52	12.0
Other news media	30	6.9
Other	6	1.4

288

289

290

291 **Knowledge**

292 Almost all HCWs were able to correctly identify COVID-19 key
293 symptoms. Data from current study revealed that 93% of
294 respondents fully understood that isolation and treatment of people
295 who are infected with the COVID-19 virus are effective ways to
296 reduce the spread of the virus. Almost 90% were also aware of
297 factors likely to be associated with severity of the disease.

298 Majority of the respondents understood dynamics of COVID-19
299 infectiousness: 91.5% respondents were aware of
300 possibilities to infect before the onset of symptoms;
301 90.8% of HCWs responded true on droplets as a major
302 transmission route. Considerable proportion of respondents
303 (93.1%) were informed that the incubation period is not constant
304 and could vary from 2-14 days.

305

306

307

308

309

310

311

312

313

314

315

318 **Table 2: Knowledge of HCWs toward the COVID-19, 2020**

Questions	Response (N=434)		
Knowledge	True (n, %)	False (n, %)	I don't know (n, %)
The common symptoms of COVID-19 are fever fatigue and dry cough	414(95.4)	13(3.0)	7(1.6)
The causative agent of COVID-19 is corona virus	410(94.4)	12(2.8)	12(2.8)
The incubation period of COVID-19 is 2-14 days	404(93.1)	21(4.8)	9(2.1)
Not all persons with COVID-2019 will develop to severe cases. Only those who are elderly, have chronic illnesses are more likely to be severe cases.	390(89.9)	33(7.6)	11(2.5)
COVID-19 can be transmitted through direct contact of respiratory droplets when infected persons coughs or sneezes.	394(90.8)	28(6.4)	12(2.8)
The disease can be transmitted from asymptomatic patient	397(91.5)	26(6.0)	11(2.5)
Training and observation of standard precautionary measures are required by care-giving personnel in suspected and probable cases of COVID-19 infection	392(90.3)	30(6.9)	12(2.8)
Visitors to patients with suspected, probable and confirmed cases of COVID-19 infection should be limited both in hospital and at home	406(93.6)	22(5.1)	6(1.3)
The disease can be spread by touching a surface or object that has the virus on it and then touching their own mouth, nose, or possibly their eyes.	401(92.4)	22(5.1)	11(2.5)
Isolation and treatment of people who are infected with the COVID-19 virus are effective ways to reduce the spread of the virus.	404(93.1)	23(5.3)	7(1.6)
People who have contact with someone infected with the COVID-19 virus should be immediately isolated in a proper place. In general, the observation period is 14 days.	400(92.2)	24(5.5)	10(2.3)

A person with mild symptoms of COVID-19 must remain at home until resolution of clinical symptoms	398(91.7)	30(6.9)	6(1.4)
Standard precautions should be followed by health care providers in dealing with suspected, probable and confirmed cases of COVID-19 infection	395(91.0)	29(6.7)	10(2.3)
Oxygen therapy should be given to all cases of severe COVID-19 with acute respiratory infection	369(85.0)	54(12.5)	11(2.5)
Ventilation with an endotracheal tube must be carried out in patient with confirmed and or suspected COVID-19 with clinical manifestations of acute respiratory distress symptoms.	379(87.3)	40(9.2)	15(3.5)

319 **Attitude**

320 The mean attitude score 10.5 (SD: 4.1). Overall, there was poor attitude among HCWs toward COVID-
 321 19. Only 45.2 % (n = 196) of the participants had a good attitude toward COVID-19. Only 48.8% (n =
 322 212) were confident that COVID-19 can be controlled by public health institute (Table 3). When asked
 323 about the using mask, up to 49.3% (n = 214) believed that It is imperative to use surgical mask when
 324 working with patient with COVID-19.

325

326 **Table 3: Attitude of HCWs toward the COVID-19, 2020**

Questions	Response (N=434)				
	SA	A	N	D	SD
Caring for patients with COVID-19 infection may be a threat to health care personnel	226(52.1)	147(33.9)	28(6.4)	24(5.5)	9(2.1)
Public health agencies like EPHI can control Outbreak of COVID-19	212(48.8)	121(27.9)	42(9.7)	50(11.5)	9(2.1)
COVID-19 can have a negative effect on the economies of the countries involved	231(53.2)	135(31.1)	24(5.5)	36(8.3)	8(1.9)
It is important to report suspected cases to health authorities	301(69.3)	104(24.0)	11(2.5)	13(3.0)	5(1.2)
COVID-19 is preventable	226(52.1)	136(31.3)	33(7.6)	23(5.3)	16(3.7)
It is imperative to use surgical mask when working with patient with COVID-19	214(49.3)	136(31.3)	23(5.3)	26(6.0)	35(8.1)

327 **Practice**

328 About practical measures put in place by HCWs in order to protect themselves and their families, 72%
329 of health care workers reported to always wore personal protective equipment when coming into contact
330 with the patients and up to 67.5% washed their hands before and after touching each patient. Other
331 measures included: avoiding going to crowded places and avoiding shaking hands, hugging or
332 kissing. Unfortunately, as high as 72.6% (n = 315) of the participants had avoided patients with
333 symptoms suggestive to those of COVID-19 (Table 4).

334

335

336

337

338

339

340

341

342

343

344

345 **Table 4: Practice of HCWs toward the COVID-19, 2020**

Questions	Response (N=434)		
Practice	Always (n, %)	Occasional (n, %)	Never (n, %)
I always wear the personal protective equipment when handling a patient	314(72.3)	101(23.3)	19(4.4)
I wash my hands with water and soap before putting on the gloves	302(69.6)	105(24.2)	27(6.2)
For residual protection, I use hand sanitizer after washing hands with soap and water	303(69.8)	103(23.7)	26(6.5)
I wash hand between patients (before and after examining)	293(67.5)	105(24.2)	36(8.3)
I wash my hands whenever they are soiled	298(68.7)	113(26.0)	23(5.3)
I wash my hands after blowing nose or covering a sneeze	289(66.6)	114(26.3)	31(7.1)
I wash my hands with soap and water and apply with hand sanitizer when leaving the health facility	208(71.0)	107(24.6)	19(4.4)
When I sneezing or coughing, I cover my nose or mouth with a tissue/clean cloth or in to my elbow	308(71.0)	106(24.4)	20(4.6)
I keep at list one miter distance from a sick person	311(71.7)	108(24.9)	15(3.4)
I avoid shaking hands, hugging or kissing with colleagues/patients	310(71.4)	110(25.4)	14(3.2)
I avoid touching my mouth, nose and eyes with my hand if I have not washed them with water and soap	307(70.7)	117(27.0)	10(2.3)
I avoid going to crowded places	316(72.8)	107(24.7)	11(2.5)
I safely dispose the used PPE items when I finish the service	315(72.6)	100(23.0)	19(4.4)
I avoid patients with signs and symptoms suggestive of COVID-19.	315(72.6)	91(21.0)	28(6.4)

346

347 **Comparison of KAP between demographic characteristics of**
348 **HCWs**

349 Almost ninety percent (n = 381) of the participants scored 12 or
350 more and were considered to have sufficient knowledge. Of the
351 total participants, more male participants had sufficient knowledge
352 than those of female participants (89.8 vs. 83.7%), however this
353 difference was not statistically significant (p = 0.07). More
354 participants working at the referral hospital had sufficient
355 knowledge than other health facilities (p<0.001).

356 Only 45% of the participants had positive attitude (n=196).
357 Bivariate analysis revealed that more HCWs working at Ayardage
358 HC had good attitude compared to those HCWs working at other
359 health facilities (p=0.02). More HCWs who used government
360 media like television to access information on COVID-19 had
361 better attitude (p = 0.009). There was no statistically significant
362 difference between attitude and other sociodemographic variables
363 (Table 5).

364 Overall, up to 38.5% (n = 167) of the participants had good
365 practices (mean score \geq 18.8). HCWs \leq 40 (p=0.04) and those
366 working at Ayardage HC practiced more than their counter parts (p
367 <0.001) (Table 5).

368

369

370

371

372

373 **Table 5: Comparison of COVID-19 knowledge, attitude and**
 374 **practice between different groups**

Variable	Sufficient Knowledge (n=381)	<i>P</i> ¹	Positive attitude (n=196)	<i>P</i> ¹	Good practice (n=167)	<i>P</i> ¹
	Freq (%)		Freq (%)		Freq (%)	
Overall	381(87.8)		196(45.2)		167(38.5)	
Sex						
Male	263(89.8)	0.07	130(44.4)	0.63	110(37.5)	0.56
Female	118(83.7)		66(46.8)		57(40.4)	
Age						
18-39	372(87.9)		190(44.9)	0.52	166(39.2)	0.04
>40	9(81.2)		6(54.6)		1(9.1)	
Qualification						
Nurse (including midwives)	287(89.1)	0.45	137(42.5)	0.44	129(40.1)	0.23
Physician	29(80.6)		18(50.0)		17(47.2)	
Pharmacist	23(88.5)		14(53.9)		5(19.2)	
Dentist	5(71.4)		4(57.1)		3(42.9)	
Laboratory technologist	26(83.9)		15(48.4)		9(29.0)	
X-ray physician	11(91.0)		8(66.7)		4(33.3)	
Place of work						
Referral hospital	317(91.9)	<0.001	140(40.6)	0.02	129(37.4)	0.001
Karamara hospital	23(69.7)		20(60.6)		10(30.3)	
Ablele HC	32(72.7)		27(61.4)		27(61.4)	
Ayrdage HC	9(75.0)		9(75.0)		1 (8.30)	
Work experience						
0-4 years	113(89.0)	0.62	53(41.7)	0.36	51(40.2)	0.64
>5 years	268(87.8)		143(46.6)		116(37.8)	
Source of information on covid-19						
WHO	213 (88.4)	0.28	100(41.5)	0.009	142(40.7)	0.17
Government site and media	95(90.5)		61(58.1)		74(28.6)	
Social media	45(86.4)		16(30.8)		27(46.2)	

Other news media	24(80.0)	16(53.3)	18(40.0)
Other	4(66.7)	3(50.0)	3(50.0)

375 | Pearson chi-squared test to compare KAP between demographic
376 | characteristics of HCWs

377

378

379 | **Distribution of KAP**

380 | The mean knowledge, attitude and practice score was 13.7 (SD:
381 | 2.6), 10.5 (SD:4.1) and 18.8 (SD:5.8) respectively. Male
382 | participants had higher knowledge score than female (p=0.013).
383 | Similarly, nurses had higher knowledge score compared to other
384 | health professionals (p=0.04). The level of knowledge among the
385 | healthcare workers working at Referral hospital (p<0.001) and
386 | those who seek covid-19 related information from world health
387 | organization (p<0.001). (Table 6).

388 | Attitude score was higher among health workers working at Ablele
389 | HC and among HCWs who seek information from other sources.

390 | Likewise, practice score was higher among younger professional
391 | (p=0.04), those working at Ablele hospital (p=0.01) and those who
392 | seek covid-19 information from other sources (p=0.02). (Table 6

394 **Table 6: Distribution of knowledge and attitude scores among healthcare workers, 2020**

Variable	Knowledge (Max=15)	<i>t-value</i>	<i>F-value</i>	<i>P-value</i>	Attitude (Max=30)	<i>t-value</i>	<i>F-value</i>	<i>P-value</i>	Practice (Max=42)	<i>t-value</i>	<i>F-value</i>	<i>P-value</i>
	Mean ± SD				Mean ± SD				Mean ± SD			
Overall	13.7 ± 2.6				10.5 ± 4.1				18.8 ± 5.8			
Sex												
Male	13.9 ± 2.3	2.49	-	0.013*	10.3 ± 4.1	-1.02	-	0.31	18.6 ± 5.8	-1.18	-	0.24
Female	13.3 ± 3.1				10.8 ± 4.1				19.3 ± 6.3			
Age												
18-39	13.7 ± 2.6	0.014	-	0.99	10.5 ± 4.1	0.24	-	0.81	18.9 ± 6.0	2.01	-	0.04*
>40	13.7 ± 1.8				10.2 ± 2.9				15.3 ± 2.7			
Qualification												
Nurse (including midwives)	13.9 ± 2.2	-	2.39	0.04*	10.3 ± 3.8	-	1.45	0.21	19.2 ± 6.1	-	1.61	0.16
Physician	12.9 ± 3.9				11.0 ± 4.4				19.3 ± 5.9			
Pharmacist	13.2 ± 2.4				10.7 ± 3.8				16.5 ± 3.5			
Dentist	12.2 ± 3.5				9.9 ± 2.3				18.3 ± 4.5			
Laboratory technologist	13.0 ± 3.5				10.9 ± 5.6				17.3 ± 5.9			
X-ray physician	13.6 ± 3.1				13.2 ± 6.7				17.4 ± 4.4			
Place of work												
Referral hospital	14.3 ± 1.8	-	40.5	<0.001**	10.1 ± 4.1	-	5.93	<0.01*	18.8 ± 6.0	-	3.87	0.01*
Karamara hospital	12.3 ± 3.4				12.0 ± 4.2				17.6 ± 4.4			
Ablele HC	10.6 ± 4.0				12.3 ± 3.8				20.9 ± 6.6			

Ayrdage HC	12.6 ± 2.4				11.5 ± 2.4				15.3 ± 1.8			
Work experience												
0-4 years	13.8 ± 2.3	0.41	-	0.68	9.9 ± 3.4	-1.78	-	0.08	18.9 ± 6.0	0.54	-	0.59
>5 years	13.7 ± 2.7				10.7 ± 4.4				15.3 ± 2.7			
Source of information on covid-19												
WHO	14.0 ± 2.0	-	5.94	<0.01	10.4 ± 3.9	-	2.96	0.02*	19.2 ± 6.1	-	2.88	0.02*
Government site and media	13.9 ± 2.2				11.1 ± 4.4				17.2 ± 4.5			
Social media	13.3 ± 3.1				9.1 ± 3.3				19.1 ± 6.1			
Other news media	12.6 ± 3.9				11.6 ± 4.4				19.7 ± 7.0			
Other	9.8 ± 6.9				12.2 ± 7.0				22.2 ± 11.2			

395 Data were expressed as mean ± SD. *t*-test and ANOVA to compare the score between KAP between demographic
396 characteristics of HCWs. * p<0.05, **P<0.01

397 **Correlation between knowledge, attitude and practice**

398 Table 5 presents the correlation coefficient between whole grains
399 knowledge, attitude and practice. It was found that attitude and
400 practice domain were inversely associated knowledge score ($r=-$
401 0.295 , $p < 0.01$) and ($r=-0.298$, $p < 0.01$) for attituded and practice
402 respectively. Significant positive correlations were found between
403 attitude with attitude ($r=0.173$, $p < 0.0003$).

404 **Table 6: Correlation between knowledge, attituded and**
405 **practice**

	Knowledge	Attitude	Practice
Knowledge	-		
Attitude	-0.295**		
Practice	-0.298**	0.173*	

406 *** significant at level of < 0.001 , ** significant at level < 0.05

407 **Discussion**

408 The current work assessed the knowledge, attitude and practice
409 toward the COVID-19 outbreaks among HCWs working at public
410 health facilities. In this study, the socio-demographic, knowledge
411 level, attitude, and infection prevention practical level response of
412 434 HCWs were analyzed.

413 Our study found that most of the HCWs were well informed with
414 COVID-19 related knowledge, during the outbreak. Among these
415 HCWs, the level of knowledge about COVID-19 was higher male
416 participants and nurse. From our study, 87.8% of HCW had
417 sufficient knowledge about COVID-19 which is higher than values
418 reported by studies conducted in Uganda and Northern
419 Ethiopia(Olum et al., 2020; Tadesse et al., 2020) where 69% and
420 74% of HCWs had sufficient knowledge. The proportion of HCW
421 with sufficient knowledge in the current study is similar to those
422 studies conducted in China(Huynh et al., 2020; Zhou et al., 2020).

423 In our study, most of the participants used information from
424 international and governmental media (websites and verified social
425 media pages). Our study suggests that majority of HCWs with
426 sufficient knowledge on COVID-19 used news media such
427 television. This suggests that such media should be frequently used
428 to disseminate information on COVID-19 by the stakeholders
429 involved in covid-19 response.

430 About 52% of HCW believed that caring for COVID-19 patients
431 may be a threat to health care workers which is similar to findings
432 by study conducted in China(Zhou et al., 2020) which showed
433 around 85% of the surveyed HCWs were afraid of getting infected
434 while caring for their patients. HCWs help patients during their
435 routine tasks such as patient consultation, infusion, dressing
436 changes and surgery. They must also deal with many other

437 emergency situations, and they may become infected with the virus
438 if they are not cautious.

439 About 14% of HCW believed that wearing general medical masks
440 may not protect the spread COVID-19 contrary to findings by Ng
441 et al. which showed adequate protection(Ng et al., 2020).

442 Our study reveals that majority of HCWs have a negative attitude
443 toward COVID-19 which is in congruence with a KAPs study on
444 COVID in Uganda among HCWs(Olum et al., 2020) but in
445 contrast to study in China on COVID-19 and Ethiopia (Jemal et
446 al., 2020; Zhou et al., 2020). However, Spearman's analysis found
447 that a significant negative correlation between the mean knowledge
448 and attitude scores of HCWs about COVID-19 ($r=-0.295$, $P<0.001$)
449 which is in consistent with a study conducted in Vietnam(Huynh et
450 al., 2020). In other words, the lower the attitude scores were, the
451 higher the probability of positive attitudes; while the higher the
452 knowledge scores were, the higher the probability of good
453 knowledge. Therefore, a good knowledge COVID-19 was directly
454 associated with a positive attitude.

455 Our study shows that majority of HCWs have low score of
456 COVID-19 prevention practices contrary to findings by other
457 studies on coronaviruses(Olum et al., 2020; Tadesse et al., 2020;
458 Zhou et al., 2020). Though majority of the HCWs are following
459 key infection prevention and control practices recommended by the
460 Ministry of Health Ethiopia and WHO. These include regular hand
461 hygiene, social distancing and wearing personal protective
462 equipment when in high risk situations. Seventy two percent and
463 67.5% of HCWs reported wearing personal protective equipment
464 when in contact with patients and washing hands before/after
465 handling patients. These are very vital practices to prevent transfer
466 of COVID-19 from patients to patients and to the HCWs

467 themselves. However, up to 73% of HCWs admitted having
468 avoided patients with symptoms suggestive of COVID-19. This
469 can be attributed to shortage of personal protective equipment
470 globally (Bauchner et al., 2020; Ranney et al., 2020).

471 Our study has some limitations. Firstly, no standardized tool for
472 assessing KAPs on COVID-19 has been previously validated. We
473 have however adapted and modified a previously published tool for
474 assessment of KAP toward prevention of respiratory tract
475 infections. Secondly, only HCWs in public health facilities in parts
476 of Somali region were surveyed and the results of this study may
477 not reflect the KAPs of HCWs in the private sectors. A similar
478 study may be extended to the community.

479 In conclusion, we found that more than 85% of HCWs have
480 sufficient knowledge on the transmission, diagnosis and prevention
481 of the transmission of COVID-19. Knowledge on COVID-19 was
482 significantly higher among HCWs who are working at referral
483 hospital. There was statistically significant difference in the level
484 of knowledge about COVID19 among health care workers. About
485 55% of the respondents had poor attitude toward COVID-19 and
486 just over 38% of the HCWs had good practices toward COVID-19
487 especially those younger than 40 years. In nutshell, the overall
488 level of knowledge was good. However, the attitude and practice
489 were relatively low. We therefore, recommend strategies for
490 enhancing the capacity of healthcare workers to develop positive
491 attitude and practice.

492

493

494

495

496 **List of abbreviation**

497

498 CDC: Centre of Disease Control

499 CI: Confidence Interval

500 EOC: Emergency Operating Centre

501 EPHI: Ethiopia Public Health Institute

502 ERC: Ethical Review Committee

503 HCW: Health Care Worker

504 IPC: Infection Prevention Control

505 PPE: Personal Protective Equipment

506 WHO: World Health Organization

507

508 **Declarations**

509 **Acknowledgment**

510 We would like to extend a special thanks to Jigjiga University for
511 funding researches on covid-19 which may be used in planning
512 health programs about the emerging viral disease.

513 **Funding**

514 This research was funded by Jigjiga University

515

516 **Availability of data and materials**

517 The dataset for this study is available from the corresponding
518 author on reasonable request.

519 **Ethical approval**

520 Ethical clearance and support letters were obtained from the
521 Ethical Review Committee (ERC) of the College of Medicine and
522 Health Sciences, Jigjiga University. The support letter then
523 submitted to the five public health facilities. Then, permission
524 obtained from the health facilities director and department/section
525 heads. Study participants were informed about the purpose and
526 importance of the study through written informed consent before
527 the data collection process. In addition, participants who are
528 unwilling to take part in the study and those who need to quit their
529 participation at any stage were informed to do so without any
530 restriction.

531 **Consent for publication**

532 Not applicable

533

534 **Conflict of interest:**

535 The authors declare that they have no competing of interest.

536 **Authorship:**

537 AM conceived the study, prepared the proposal, analyzed the data,
538 interpreted the findings and wrote the manuscript. TY, MO, MA,
539 OM, MA and AB were involved in developing the study proposal,
540 data analysis and reviewing the manuscript.

541

542

543

544

545

546 **References**

- 547 Bauchner, H., Fontanarosa, P. B., & Livingston, E. H. (2020).
548 Conserving supply of personal protective equipment—a
549 call for ideas. *Jama*, 323(19), 1911-1911.
- 550 Cascella, M., Rajnik, M., Cuomo, A., Dulebohn, S. C., & Di
551 Napoli, R. (2020). Features, evaluation and treatment
552 coronavirus (COVID-19) (*StatPearls [Internet]*). StatPearls
553 Publishing.
- 554 CDC. (2020). *Information for health care professionals* Retrieved
555 April 15, 2020 from
556 [https://www.cdc.gov/coronavirus/2019-](https://www.cdc.gov/coronavirus/2019-ncov/hcp/index.html)
557 [ncov/hcp/index.html](https://www.cdc.gov/coronavirus/2019-ncov/hcp/index.html)
- 558 Chen, N., Zhou, M., Dong, X., Qu, J., Gong, F., Han, Y., Qiu, Y.,
559 Wang, J., Liu, Y., & Wei, Y. (2020). Epidemiological and
560 clinical characteristics of 99 cases of 2019 novel
561 coronavirus pneumonia in Wuhan, China: a descriptive
562 study. *The lancet*, 395(10223), 507-513.
- 563 Control, C. f. D., & Prevention. (2019). Novel coronavirus,
564 Wuhan. *China. Information for Healthcare Professionals*.
565 [https://www.cdc.gov/coronavirus/2019-nCoV/hcp/index.](https://www.cdc.gov/coronavirus/2019-nCoV/hcp/index.html)
566 [html](https://www.cdc.gov/coronavirus/2019-nCoV/hcp/index.html) (Accessed on February 14, 2020).
- 567 EPHI. (2020). Notification Note on COVID-19 Situational Update.
- 568 Giao, H., Han, N. T. N., Van Khanh, T., Ngan, V. K., Van Tam,
569 V., & Le An, P. (2020). Knowledge and attitude toward
570 COVID-19 among healthcare workers at District 2
571 Hospital, Ho Chi Minh City. *Asian Pacific Journal of*
572 *Tropical Medicine*, 13.
- 573 Hoffman, S. J., & Silverberg, S. L. (2018). Delays in global
574 disease outbreak responses: lessons from H1N1, Ebola, and
575 Zika. *American journal of public health*, 108(3), 329-333.
- 576 Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L.,
577 Fan, G., Xu, J., & Gu, X. (2020). Clinical features of
578 patients infected with 2019 novel coronavirus in Wuhan,
579 China. *The lancet*, 395(10223), 497-506.
- 580 Huynh, G., Nguyen, T. N. H., Vo, K. N., & Pham, L. A. (2020).
581 Knowledge and attitude toward COVID-19 among
582 healthcare workers at District 2 Hospital, Ho Chi Minh
583 City. *Asian Pacific Journal of Tropical Medicine*, 13(6),
584 260.
- 585 Jemal, B., Ferede, Z. A., Mola, S., Hailu, S., Abiy, S., Wolde, G.
586 D., Tilahun, A., Tesfaye, B., Lemma, D. T., & Alimaw, A.
587 A. (2020). Knowledge, attitude and practice of healthcare
588 workers towards COVID-19 and its prevention in Ethiopia:
589 a multicenter study.
- 590 Li, T., Wei, C., Li, W., Hongwei, F., & Shi, J. (2020). Beijing
591 Union Medical College Hospital on " pneumonia of novel

- 592 coronavirus infection" diagnosis and treatment proposal
593 (V2. 0). *Med J Peking Union Med Coll Hosp.*
- 594 McCloskey, B., & Heymann, D. L. (2020). SARS to novel
595 coronavirus—old lessons and new lessons. *Epidemiology &*
596 *Infection, 148.*
- 597 Ng, K., Poon, B. H., Kiat Puar, T. H., Shan Quah, J. L., Loh, W. J.,
598 Wong, Y. J., Tan, T. Y., & Raghuram, J. (2020). COVID-
599 19 and the risk to health care workers: a case report. *Annals*
600 *of internal medicine.*
- 601 Novel, C. P. E. R. E. (2020). The epidemiological characteristics
602 of an outbreak of 2019 novel coronavirus diseases
603 (COVID-19) in China. *Zhonghua liu xing bing xue za zhi=*
604 *Zhonghua liuxingbingxue zazhi, 41(2), 145.*
- 605 Olum, R., Chekwech, G., Wekha, G., Nassozi, D. R., & Bongomin,
606 F. (2020). Coronavirus Disease-2019: Knowledge,
607 Attitude, and Practices of Health Care Workers at Makerere
608 University Teaching Hospitals, Uganda. *Frontiers in Public*
609 *Health, 8, 181.*
- 610 Organization, W. H. (2005). Statement on the second meeting of
611 the International Health Regulations (2005) Emergency
612 Committee regarding the outbreak of novel coronavirus
613 (2019-nCoV).
- 614 Organization, W. H. (2020a). *Advice on the use of masks in the*
615 *community, during home care and in healthcare settings in*
616 *the context of the novel coronavirus (2019-nCoV)*
617 *outbreak: interim guidance, 29 January 2020 (*
- 618 Organization, W. H. (2020b). Coronavirus disease 2019 (COVID-
619 19): situation report, 72.
- 620 Organization, W. H. (2020c). Infection prevention and control
621 during health care when novel coronavirus (nCoV)
622 infection is suspected: interim guidance, 25 January 2020.
- 623 Organization, W. H. (2020d). WHO director-general's opening
624 remarks at the media briefing on covid-19-11 march 2020.
625 2020.
- 626 Poole, D. N., Escudero, D. J., Gostin, L. O., Leblang, D., & Talbot,
627 E. A. (2020). Responding to the COVID-19 pandemic in
628 complex humanitarian crises. *International Journal for*
629 *Equity in Health, 19(1), 1-2.*
- 630 Ranney, M. L., Griffeth, V., & Jha, A. K. (2020). Critical supply
631 shortages—the need for ventilators and personal protective
632 equipment during the Covid-19 pandemic. *New England*
633 *Journal of Medicine, 382(18), e41.*
- 634 Selvaraj, S. A., Lee, K. E., Harrell, M., Ivanov, I., & Allegranzi, B.
635 (2018). Infection rates and risk factors for infection among
636 health workers during Ebola and Marburg virus outbreaks:

637 a systematic review. *The Journal of infectious diseases*,
638 218(suppl_5), S679-S689.
639 Tadesse, D. B., Gebrewahd, G. T., & Demoz, G. T. (2020).
640 Knowledge, Attitude, Practice and Psychological response
641 toward COVID-19 among Nurses during the COVID-19
642 outbreak in Northern Ethiopia, 2020.
643 Zhou, M., Tang, F., Wang, Y., Nie, H., Zhang, L., You, G., &
644 Zhang, M. (2020). Knowledge, attitude and practice
645 regarding COVID-19 among health care workers in Henan,
646 China. *Journal of Hospital Infection*.
647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

691

692

693