1	The Association of COVID-19 Incidence with Sport and Face Mask
2	Use in United States High School Athletes
3	
4	
5	Andrew M. Watson, MD, MS, ¹ Kristin Haraldsdottir, PhD, ¹ Kevin Biese, MA, LAT, ² Leslie
6	Goodavish, PA, ¹ Bethany Stevens, ² Timothy McGuine, PhD, LAT ¹
7	
8	From the Department of Orthopedics and Rehabilitation, ¹
9	University of Wisconsin School of Medicine and Public Health, Madison, WI, and
10	The Department of Kinesiology, ² University of Wisconsin – Madison
11	
12	
13	
14	
15	Address for Correspondence:
16	Andrew Watson
17	Department of Orthopedics and Rehabilitation, Division of Sports Medicine
18	1685 Highland Avenue
19	Madison, WI 53705
20	Phone: (608) 263-6477
21	Fax: (608) 263-0503
22	Email: Watson@ortho.wisc.edu
23	

24 ABSTRACT

25 **Purpose:** To evaluate the influence of sport characteristics and face mask use on COVID-19

26 incidence among high school athletes.

27 Methods: Surveys were distributed to high school athletic directors throughout the United States

regarding sport re-initiation, COVID-19 cases, and risk reduction procedures in fall 2020.

29 Separate mixed effects Poisson regression models were developed to evaluate the associations

30 between reported COVID-19 incidence and 1) sport characteristics (contact/non-contact,

31 individual/team, indoor/outdoor) and 2) face mask use while playing (yes/no).

32 **Results:** 991 schools had restarted fall sports, representing 152,484 athletes on 5,854 teams.

2,565 cases of COVID-19 were reported, representing a case rate of 1,682 cases per 100,000

34 athletes and an incidence rate of 24.6 cases per 100,000 player-days. COVID-19 incidence was

35 lower among outdoor versus indoor sports (incidence rate ratio [IRR]=0.54, 95% CI=0.49-0.60,

36 p<0.001) and non-contact versus contact sports (IRR=0.78 [0.70-0.87], p<0.001), but not team

37 versus individual sports (IRR=0.96 [0.84-1.1], p=0.49). Face mask use was associated with a

decreased incidence in girls' volleyball (IRR=0.53 [0.37-0.73], p<0.001), boys' basketball

39 (IRR=0.53 [0.33-0.83], p=0.008) and girls' basketball (IRR=0.36 [0.19-0.63], p<0.001), and

40 approached statistical significance in football (IRR=0.79 [0.59-1.04], p=0.10) and cheer/dance

41 (IRR=0.75 [0.53-1.03], p=0.081).

42 **Conclusions:** In this nationwide survey of US high school athletic directors representing

43 152,484 athletes, lower COVID-19 incidence was independently associated with participation in

44 outdoor versus indoor and non-contact versus contact sports, but not team versus individual

45 sports. Face mask use was associated with decreased COVID-19 incidence among indoor sports,

46 and may be protective among outdoor sports with prolonged close contact between participants.

47

48 Keywords: adolescent; infection; pediatric; SARS-CoV-2

50 **INTRODUCTION**

51 In an effort to control the spread of the coronavirus disease 2019 (COVID-19) in spring 52 2020, restrictions were placed on virtually all facets of American society, including the 53 cancellation of school and interscholastic athletics. Early research suggests that school and sport 54 cancellations during the initial months of the COVID-19 pandemic were associated with 55 significant decreases in physical activity and worsening of depressive symptoms in children and athletes.^{3,6,8} It has been projected that prolonged restriction could contribute significantly to 56 long-term increases in obesity and mental health disorders.^{2,7,15} In a nationwide survey of over 57 58 13,000 adolescent athletes in May 2020, 37% reported moderate to severe symptoms of anxiety and 40% reported moderate to severe symptoms of depression.⁹ Together these results suggest 59 60 that isolation and physical inactivity during COVID-19 restrictions may represent a significant 61 threat to physical and mental health in children and adolescents.

62 Nonetheless, efforts to promote the benefits of youth sports participation are necessarily 63 balanced against the potential risk of COVID-19 transmission. While high school sports have 64 restarted in many areas of the United States, they remain shut down in other areas, with a lack of 65 specific evidence to guide decision-making. A number of organizations have developed risk 66 reduction protocols in an attempt to mitigate the spread of COVID-19 among youth sport 67 participants, but it is widely recognized that there is very little prior research within sport environments to guide these decisions.^{1,5,11,12,17} The available evidence has been limited to case 68 69 reports in adult recreational athletes, media reports of infections among adults and adolescents 70 associated with interscholastic athletics, and a single evaluation of youth soccer in a small-group, physically-distanced setting.^{4,5,13,18} 71

72	Organizations have also attempted to classify sports in terms of the risk of COVID-19
73	transmission during participation. These recommendations are based on a number of
74	characteristics but differ between organizations, as evidence derived from sport contexts is
75	lacking. ^{1,11,12,17} In fact, we are aware of no research which has evaluated the relative risks of
76	COVID-19 among athletes between different sports or between sport characteristics such as
77	indoor versus outdoor, individual versus team or contact versus non-contact.
78	Similarly, the recommendations regarding face mask use during sport participation differ
79	between public health organizations, and the American Academy of Pediatrics (AAP) recently
80	changed its recommendation to encourage their use among youth athletes during most sport
81	contexts. ^{1,10,12,17} Although there is general consensus among the scientific community that face
82	mask use in community settings can reduce transmission of COVID-19, there is no evidence
83	specifically within youth sport environments regarding their efficacy, and whether any benefit
84	differs between sports or sport characteristics. Therefore, the purpose of this study was to
85	determine the associations between COVID-19 risk and specific sports, sport characteristics, and
86	face mask use among US high school athletes.

87

88 **METHODS**

89 Study Design

All procedures performed in this study were approved by the Institutional Review Board of the University of Wisconsin-Madison. In collaboration with the National Federation of State High School Associations (NFHS), surveys were distributed to all state high school athletic associations in the United States between November 1, 2020 and November 3, 2020. Among states in which fall high school athletics had restarted, surveys were forwarded on to the athletic

95 directors of high schools within the state. In addition to school name and location, athletic 96 directors were asked whether they had restarted participation in sports since the initial COVID-97 19 restrictions in the spring of 2020. Those schools who reported reinitiating sports were asked 98 to provide the specific sports and the date of restarting, number of athletes, number of practices 99 and games, number of COVID-19 cases among athletes, and reported sources of infection (if 100 known) within each sport during the months of August, September and October 2020. Schools 101 were asked about their type of instruction during the fall (virtual, in-person) and what restarted 102 sports were using face masks for players while playing. Schools were included if they had any 103 sport that had restarted participation during August, September or October 2020.

104 Statistical Analysis

105 Data were initially evaluated using descriptive statistics, including estimates of central 106 tendency (mean, median) and variability (standard deviation, interguartile range, range) for 107 continuous variables, and counts and percentages for categorical variables. Case rates were 108 expressed as the number of reported cases per 100,000 players (cases / total number of players * 109 100,000) overall and for each sport. Duration of participation for each sport at each school was 110 determined as the difference in days between the date of restarting and October 31, 2020 and 111 player-days was determined as the product of the number of participating players and duration. 112 Incidence rates were expressed as the number of reported cases per 100,000 player-days (cases / 113 total number of player-days * 100,000) overall and for each sport, with confidence intervals 114 calculated using an exact method.

In addition, the number of cases, total population, case rate and incidence rate during
August, September, and October were determined for each state in which a respondent high
school was located, from publicly available online information aggregated from the US Centers

for Disease Control and local health authorities.¹⁹ In order to determine whether background state COVID-19 rates were associated with reported COVID-19 rates among high school athletes, the total number of athletes and COVID-19 cases were aggregated by state. For those states with >100 athletes, the relationship between COVID-19 case rates among high school athletes and the general population were evaluated with a linear regression model weighted for the number of players from each state.

124 For those sports with data from 50 or more schools, the relative risk of each sport was 125 evaluated using a mixed effects Poisson regression model to predict the number of COVID-19 126 cases for each team with local incidence, instructional delivery type (in-person, virtual), and 127 sport as fixed effects, school as a random effect, and the log of player-days as an offset, yielding 128 an incidence rate ratio (IRR) with Soccer – Boys as the reference (since it represented the median 129 unadjusted incidence rate and is typically considered a moderate risk sport). To evaluate the 130 independent relationships between COVID-19 cases and sport characteristics, a mixed effects 131 Poisson regression model was developed to predict the number of cases with local incidence, 132 instructional delivery type, sport location (indoor, outdoor), sport contact (contact, non-contact), 133 and sport type (individual, team) as fixed effects, school as a random effect and the log of player-134 days as an offset including data from all sports reported. The specific sports that were reported 135 and their respective characteristics are shown in Supplemental Table 1.

To evaluate the association between overall COVID-19 incidence and reported face mask use, a mixed effects Poisson regression model was developed to predict the number of cases for each team, with local incidence, instructional delivery type, and face mask use (yes/no) as fixed effects, school as a random effect and the log of player-days as an offset. Similar, separate models were developed for each sport characteristic (indoor, outdoor, contact, non-contact,

141	individual, team) and each specific sport with greater than 40 reported cases. Within each of
142	these sports, incidence rates and 95% confidence intervals were calculated within each sport for
143	those teams reporting face mask use or not and compared descriptively. Swimming was excluded
144	from the face mask analyses. Coefficients from Poisson models were exponentiated to represent
145	IRRs for binary variables and Wald confidence intervals were generated. Significance level was
146	determined a priori at the 0.05 level and all tests were 2-tailed. All statistical analyses were
147	performed in R. ¹⁴

148

149 **RESULTS**

150 1,508 schools submitted survey responses, of which 991 schools had restarted a fall sport. 151 These schools represented 152,484 student-athletes on 5,854 teams that had participated in 152 159,947 practices and 48,582 games. Eight hundred eighty-nine (89.7%) respondent schools 153 reported utilizing in-person instruction during the fall of 2020. Among the schools that had 154 restarted participation, 2,565 cases of COVID-19 were reported, yielding a case rate of 1,682 155 cases per 100,000 athletes and an incidence rate of 24.6 (95% CI = 23.7-25.6) cases per 100,000 156 player-days. Of the cases with a reported, known source, 870 (55%) were attributed to 157 household contact followed by community contact outside sport or school (516, 32%), school 158 contact (115, 7.3%), sport contact (69, 4.3%) and other (24, 1.5%). For those sports with greater 159 than 50 participating schools, the unadjusted COVID-19 incidence rate ranged from 10.4 (Tennis 160 - Boys) to 52.0 cases per 100,000 player-days (Basketball - Girls), as shown in Figure 1 (full 161 data for all sports is available in Supplementary Table 2). 162 When aggregated by state, the overall COVID-19 case rates for athletes were highly

163 correlated with the case rates for their respective state's general population ($\beta = 1.09 \pm 0.16$, r =

164	0.81, $p < 0.001$; see Figure 2). The COVID-19 IRRs for specific sports, adjusted for state
165	COVID-19 incidence, instruction delivery type and school repeated measures are shown in
166	Figure 3. After adjusting for state COVID-19 incidence and school instruction type, reported
167	COVID-19 incidence among high school athletes was significantly and independently lower
168	among outdoor versus indoor sports (IRR=0.54 [0.49-0.60], p<0.001) and non-contact versus
169	contact sports (0.78 [0.70-0.87], p<0.001), while no association was identified with respect to
170	team versus individual sports (0.96 [0.84-1.1], p=0.49).
171	284 schools (28%) reported face mask use by players while playing certain sports,
172	representing 1,677 (28.6%) of all teams participating during the study period. Overall, teams
173	reporting face mask use did not have a lower incidence of COVID-19 among players (IRR =
174	0.94 [95% CI = 0.75-1.16], p=0.55). However, COVID-19 incidence was lower with face mask
175	use among players participating in indoor sports (Table 1). For those sports with greater than 40
176	reported cases, differences in COVID-19 incidence between teams with and without face mask
177	use within each sport are shown in Figure 4. Finally, face mask use was associated with a
178	decreased COVID-19 incidence in girls' volleyball, girls' basketball, and boys' basketball, and
179	approached significance in football and cheer/dance, but no association was identified in other
180	sports (Table 2).

181

182 **DISCUSSION**

These findings suggest that the incidence of COVID-19 among US high school athletes in the fall of 2020 differs between sports and sport characteristics. Although only a small proportion of the cases with a reported source were attributed to sport contact, indoor location and contact were independently associated with an increased incidence rate of COVID-19. This

187 is the first evidence we are aware of that has been derived specifically from a high school sport 188 context, but it is in agreement with prior research and recommendations from various public 189 health organizations that have suggested that COVID-19 is most likely to be transmitted between 190 individuals in close proximity for prolonged periods, and may be more easily transmissible 191 indoors than outdoors.^{1,12,17} Specifically, sport participation indoors versus outdoors appeared to 192 have the strongest influence on COVID-19 incidence within our represented group of athletes, 193 while contact had an independent, yet weaker relationship. We did not find an independent 194 association between COVID-19 incidence and team versus individual sport participation, 195 suggesting that this effect is minimal after accounting for the influences of location and contact. 196 Public health organizations and sport governing bodies have attempted to classify sports based on expected risk of COVID-19 transmission,^{1,11,12,17} although we are aware of no prior 197 198 evidence that has evaluated these within sport environments. With respect to the risk categories offered for high school athletics by NFHS,¹² our findings are in agreement with the suggestion 199 200 that outdoor, non-contact sports have the lowest COVID-19 incidence. These data also align 201 with the suggestion that sports with close, sustained contact may carry a relatively increased risk, 202 but suggest that indoor location may have the strongest influence on COVID-19 risk. However, 203 it should be noted that wrestling demonstrated an intermediate risk despite being an indoor sport 204 with prolonged, close contact between participants. It is unclear why this would be the case, 205 although the sample size within this sport was relatively small and consequently the confidence 206 intervals were relatively wide, making it difficult to classify the risk associated with this specific 207 sport with confidence.

It should be recognized that not all sports reported participation during the same
timeframe and may therefore have had differing background COVID-19 incidence during their

respective seasons. Nationwide COVID-19 cases decreased through August and were relatively
stable during September, but began increasing in October.¹⁶ Although we tried to account for
differences in local COVID-19 disease burden within our adjusted models by including state
COVID-19 rates during the fall months, we cannot exclude the possibility that the higher
incidence among traditional winter sports may be partly due to higher local COVID-19 incidence
later in the study period when these sports began participation.

216 We found that face mask use was associated with a decreased incidence of COVID-19 217 among specific sports. In general, those sports with the highest incidence of COVID-19 were 218 also found to have the greatest benefit from reported face mask use. Specifically, COVID-19 219 incidence was lower among indoor sports in which face masks were reportedly used when 220 evaluated collectively, but this was also true within volleyball, girls' basketball, and boys' 221 basketball when evaluated individually. Importantly, reported COVID-19 incidence among 222 indoor team sports (volleyball, basketball) when using face masks appeared comparable to the 223 incidence among outdoor team sports, suggesting that the increased risk associated with being 224 indoors may be reduced considerably by face mask use. Face mask use also appeared to be 225 associated with a decreased COVID-19 incidence in football and cheer/dance, although this did 226 not reach statistical significance. This may be attributable to the relatively small proportion of 227 the teams in these sports that reported face mask use, and a larger sample of teams using face 228 masks in these sports may have revealed a significant association.

While face mask use was not found to be associated with COVID-19 incidence among other outdoor contact sports, "contact" as a sport characteristic surely exist along a continuum with respect to the time spent in close proximity to other players. The risk of COVID-19 transmission likely varies between sports with brief contact and relatively little time spent within

close proximity to others, to sports with prolonged periods of close contact that constitute an
increased likelihood of sufficient exposure for COVID-19 transmission between participants.
This supports the suggestion that although face mask use did not appear to have a large effect
within sports like soccer, it may still be protective in an outdoor sport with sustained close
contact.¹

238 Face mask use in the community has been widely recommended by public health 239 agencies but remains a contentious issue within the public at large. Within sports, 240 recommendations differ between organizations, and recently the AAP revised their 241 recommendations regarding face mask use in youth sports, suggesting that they be used in most sports contexts when it is safe to do so.¹ These differences likely represent the fact that there has 242 243 been no primary evidence regarding the utility of face masks to reduce COVID-19 transmission 244 during sport participation. While we cannot directly evaluate the true transmission rate within 245 the data available, our findings nonetheless suggest that face masks may have a meaningful 246 influence on COVID-19 risk among indoor sports and outdoor sports with sustained close 247 contact, but less influence among outdoor sports with less time spent in close proximity to other 248 players.

Although we were unable to identify publicly available, state-specific, adolescent case rates during the fall months for many of the represented states within our sample, we nonetheless identified a strong relationship between reported COVID-19 case rates in our high school athletes and the COVID-19 case rates among the general population in their respective states. In addition, the majority of cases with a reported source were attributed to household and community contact with a much smaller proportion attributed to school or sport contact. This may suggest that COVID-19 incidence among adolescent athletes is largely reflective of

256 background COVID-19 rates within their community. The overwhelming majority of schools 257 reported in-person instruction, making it difficult to fully evaluate the role of in-person school 258 instruction in COVID-19 incidence among high school athletes. Nonetheless, we included school 259 instruction type within our adjusted models in order to account for the potential confounding role 260 this could play in comparing different groups. Importantly, it should be recognized that this 261 study cannot comment on the incidence or transmission risk of COVID-19 among attendees at 262 high school sporting events such as fans, coaches, staff, and spectators. While this risk remains 263 undefined, it nonetheless represents a potential contribution to community COVID-19 spread and 264 risk mitigation procedures should continue to be prioritized to not only protect athletes but also 265 to help reduce the risk of infection among attendees.

266 This study has several limitations. The information is self-reported by the athletic 267 directors of each school and cannot be directly verified through medical records or another 268 independent source. Local, state-level daily COVID-19 case data was often not available for 269 adolescents or children, so our adjusted models could only account for the population-level 270 background incidence from each state. Nonetheless, we found that reported case rates from our 271 sample and the case rates from the state general populations were highly related. As mentioned 272 above, the incidence of COVID-19 was likely not stable throughout the fall in many areas, and 273 those sports that initiated play during periods of increased local incidence (winter sports in 274 October, for example) may be biased toward a higher incidence that is not directly attributable to 275 the sport itself. Reported sources of infection were provided by the schools themselves and not 276 through formal contact tracing by local health authorities. We cannot directly account for the 277 possibility of transmission between players that went unidentified. Finally, while this data

278	represents information regarding a large number of male and female high school athletes from a
279	nationwide sample, it may not be generalizable to other populations.
280	In conclusion, this study suggests that certain high school sports and sport characteristics
281	may have a greater relative risk of COVID-19 and that face mask use may help reduce the risk of
282	COVID-19 among adolescent athletes in sports with higher risk. Specifically, indoor sports

- appear to have a greater risk of COVID-19 infection among participants, while outdoor, non-
- 284 contact sports have the lowest risk. However, face mask utilization was associated with a

significantly decreased incidence of COVID-19 in indoor sports, and this appeared to mitigate a

large portion of the increased risk. Given the general lack of information regarding COVID-19

risk among youth sport participants, this information may help guide decision-making to reduce

the risk of COVID-19 transmission, while facilitating the wide-ranging physical and mental

289 health benefits of sport participation.

290 **REFERENCES**

- 291 American Academy of Pediatrics. COVID-19 Interim Guidance: Return to Sports. 1. 292 https://services.aap.org/en/pages/2019-novel-coronavirus-covid-19-293 infections/clinical-guidance/covid-19-interim-guidance-return-to-sports/. Accessed 294 December 21, 2020. 295 An R. Projecting the impact of the coronavirus disease-2019 pandemic on childhood 2. 296 obesity in the United States: A microsimulation model. J Sport Health Sci. 297 2020;9(4):302-312.
- 2983.Bates LC, Zieff G, Stanford K, et al. COVID-19 Impact on Behaviors across the 24-299Hour Day in Children and Adolescents: Physical Activity, Sedentary Behavior, and300Sleep. Children (Basel). 2020;7(9).
- commercial appeal.com. Most school-associated COVID-19 cases in Shelby County
 are sports-related, officials say.
- https://www.commercialappeal.com/story/news/education/2020/10/22/most school-associated-covid-19-cases-shelby-county-sports-related-officials say/3735873001/. Accessed December, 21, 2020.
- 306 5. Drezner JA, Drezner SM, Magner KN, Ayala JT. COVID-19 Surveillance in Youth
 307 Soccer During Small Group Training: A Safe Return to Sports Activity. Sports Health.
 308 2021;13(1):15-17.
- 309 6. Dunton GF, Do B, Wang SD. Early effects of the COVID-19 pandemic on physical
 310 activity and sedentary behavior in children living in the U.S. *BMC Public Health.*311 2020;20(1):1351.
- 3127.Golberstein E, Wen H, Miller BF. Coronavirus Disease 2019 (COVID-19) and Mental313Health for Children and Adolescents. JAMA Pediatr. 2020.
- Gouttebarge V, Ahmad I, Mountjoy M, Rice S, Kerkhoffs G. Anxiety and Depressive
 Symptoms During the COVID-19 Emergency Period: A Comparative Cross-Sectional
 Study in Professional Football. *Clin J Sport Med.* 2020.
- McGuine TA, Biese KM, Petrovska L, et al. The health of US adolescent athletes
 during Covid-19 related school closures and sport cancellations. *Journal of athletic training.* 2020.
- National Athletic Trainers Association. COVID-19 Return to Sport Considerations for
 Secondary School Athletic Trainers.
- 322 https://www.nata.org/sites/default/files/covid_19_return-to-
- 323 sport_considerations_for_secondary_school_ats_1.pdf. Accessed September 22,
 324 2020.
- 325 11. National Collegiate Athletics Association. Resocialization of Collegiate Sport:
 326 Checklist. http://www.ncaa.org/sport-science-institute/resocialization-collegiate 327 sport-checklist.
- 328 12. National Federation of State High School Associations. Guidance For Opening Up
 329 High School Athletics And Activities.
- New York Times. College Sports Has Reported at Least 6,629 Virus Cases. There Are
 Many More. https://www.nytimes.com/2020/12/11/sports/coronavirus-college sports-football.html.

- Record #133 is using an undefined reference type. If you are sure you are using the
 correct reference type, the template for that type will need to be set up in this
 output style.
- Singh S, Roy D, Sinha K, et al. Impact of COVID-19 and lockdown on mental health of
 children and adolescents: A narrative review with recommendations. *Psychiatry Res.*2020;293:113429.
- 339 16. US Centers for Disease Control and Prevention. CDC COVID Data Tracker.
 340 https://covid.cdc.gov/covid-data-tracker/#trends_dailytrendscases. Accessed
 341 December 30, 2020.
- 342 17. US Centers for Disease Control and Prevention. Considerations for Youth Sports.
 343 https://www.cdc.gov/coronavirus/2019-ncov/community/schools 344 childcare/youth-sports.html.
- 34518.US Centers for Disease Control and Prevention. An Outbreak of COVID-19 Associated346with a Recreational Hockey Game Florida, June 2020.
- 347 https://www.cdc.gov/mmwr/volumes/69/wr/mm6941a4.htm. Accessed
 348 December 21, 2020.
- 349 19. USAFACTS.org. US Coronavirus Cases and Deaths.
 350 https://usafacts.org/visualizations/coronavirus-covid-19-spread-map/. Accessed
 351 January 7, 2021.
- 352

- 354 Figure Legends
- Figure 1. Unadjusted incidence rates of COVID-19 among United States high school sports
- during fall 2020. Incidence rate is shown as reported cases per 100,000 player-days for those
- 357 sports with greater than 50 schools reporting re-initiation.
- Figure 2. Reported COVID-19 case rates for high school athletes and the general population of
- their respective states during the fall of 2020. Size of points scaled to number of players from
- ach state and dashed line represents a line of equality. Solid line and shaded area represent
- 361 regression line and 95% confidence interval from linear model weighted for number of players
- 362 from each state. r = correlation coefficient.
- 363 Figure 3. COVID-19 incidence rate ratios during fall 2020 for United States high school sports,
- adjusted for local (state) COVID-19 incidence, instructional delivery type and repeated measures
- 365 from the same school. Includes those sports with greater than 50 schools reporting participation,
- 366 with Soccer Boys as reference. p<0.05.
- 367 Figure 4. Unadjusted COVID-19 incidence rates reported among US athletes in the fall 2020,
- 368 comparing teams with or without reported face mask use, within each sport.

Table 1. The association of reported face mask use with COVID-19 incidence within each sport

Sport Characteristic	N (%)	IRR (95% CI)	р
Indoor	305 (23.5%)	0.44 (0.29-0.66)	< 0.001
Outdoor	773 (18.4%)	1.02 (0.80-1.31)	0.88
Individual	441 (17.7%)	0.89 (0.64-1.25)	0.50
Team	637 (21.2%)	0.92 (0.70-1.20)	0.52
Contact	452 (20.9%)	0.86 (0.63-1.18)	0.36
Non-Contact	626 (18.8%)	0.88 (0.66-1.16)	0.36

371 characteristic among US high school athletes during fall 2020.^a

^aIncidence rate ratios and Wald confidence intervals from separate mixed effects Poisson models

373 within each sport characteristic to predict COVID-19 cases with local incidence and face mask

374 use (yes/no) as fixed effects, school as a random effect and log(player-days) as an offset; CI =

375 Confidence Interval; IRR=Incidence Rate Ratio.

Table 2. The association of reported face mask use with COVID-19 incidence within each sport

Sport	N (%)	IRR (95% CI)	р
Baseball	39 (26.4%)	1.2 (0.44-2.93)	0.69
Basketball - Boys	74 (25.7%)	0.53 (0.33-0.83)	0.0084
Basketball - Girls	78 (22.6%)	0.36 (0.19-0.63)	<0.001
Cheer/Dance	96 (18.3%)	0.75 (0.53-1.03)	0.081
Cross Country	123 (15.7%)	0.86 (0.53-1.31)	0.5
Football	94 (14.0%)	0.79 (0.59-1.04)	0.10
Golf - Boys	69 (18.9%)	0.57 (0.17-1.42)	0.28
Soccer - Boys	74 (24.5%)	1.1 (0.65-1.71)	0.77
Soccer - Girls	70 (20.5%)	1.3 (0.76-2.18)	0.3
Softball	39 (17.3%)	0.53 (0.12-1.71)	0.34
Volleyball - Girls	120 (22.0%)	0.53 (0.37-0.73)	<0.001

among US high school athletes during fall 2020.^a

^aIncidence rate ratios and Wald confidence intervals from separate mixed effects Poisson models
within each sport to predict COVID-19 cases with local incidence, school instruction type, and
face mask use (yes/no) as fixed effects, school as a random effect and log(player-days) as an
offset; CI = Confidence Interval; IRR=Incidence Rate Ratio.



Sport





Sport

