TITLE:

"Susceptibility and risk of suffering SARS-COV-2 infection by demographic characteristics and pre-existing medical conditions among middle-aged and older adults in Tarragona, Spain: results from the *COVID19 TARRACO* Cohort Study, March-June 2020".

AUTHORS:

Satué-Gracia, EM.^{a,b}, Vila-Córcoles, A.^{a,b}, de Diego-Cabanes, C.^a, Vila-Rovira, A.^b, Torrente-Fraga, C.^c, Gómez-Bertomeu, F^d., Hospital-Guardiola, I^a., Ochoa-Gondar, O.^{a,b} and Martín-Luján, F.^{a,b}

AFFILIATIONS:

^a Primary Care Department Camp de Tarragona. Institut Català de la Salut (ICS), Tarragona, Spain

^b Fundació Institut Universitari per a la recerca a l'Atenció Primària de Salut Jordi Gol i Gurina (IDIAPJGol), Barcelona, Spain

^c Technology and Informatic Department, Institut Català de la Salut (ICS), Tarragona, Spain.

^d Microbiology Department. Hospital Universitari Joan XXIII. Institut Català de la Salut (ICS), Tarragona, Spain.

CORRESPONDENCE:

Eva M Satué Gracia Unitat de Suport a la Recerca Camp de Tarragona-Reus. Institut Català de la Salut Camí de Riudoms 53-55, 43205, Reus (Tarragona), Spain Tel: 34 977 778518

Email: esatue.tgn.ics@gencat.cat

SUMMARY

Objective: To analyse susceptibility/risk of suffering COVID19 among adults with distinct underlying medical conditions.

Methods: Cohort study (population-based) including 79,083 people ≥50 years-old in Tarragona (Southern Catalonia, Spain). At study start (01/03/2020) baseline cohort characteristics (demographic, previous comorbidities, chronic medications and vaccinations' history) were recorded. Primary outcome was laboratory-confirmed COVID19 incurred in cohort members throughout 01/03/2020-30/06/2020. Risk of suffering COVID19 was evaluated by Cox regression, estimating multivariable hazard ratios (HRs) adjusted for age/sex and previous comorbidities.

Results: Across study period, 536 laboratory-confirmed COVID19 cases were observed (mean incidence: 39.5 cases per 100,000 persons-week). In multivariableanalysis, age/years (HR: 1.01; 95% CI: 1.00-1.02; p=0.050), nursing-home (HR: 20.19; 95% CI: 15.98-25.51; p<0.001), neurological disease (HR: 1.35; 95% CI: 1.03-1.77; p=0.029), taking diuretics (HR: 1.39; 95% CI: 1.10-1.75; p=0.006), antiplatelet (HR: 1.36; 95% CI: 1.05-1.76; p=0.021) and benzodiazepines (HR: 1.24; 95% CI: 1.00-1.53; p=0.047) significantly increased risk; while smoking (HR: 0.57; 95%CI: 0.41-0.80; p=0.001), angiotensin converting enzyme inhibitors (HR: 0.78; 95% CI: 0.61-1.00; p=0.048), angiotensin II receptor blockers (HR: 0.70; 95%CI: 0.51-0.96; p=0.027) and statins (HR: 0.75; 95% CI: 0.58-0.96; p=0.025) were associated with reduced risk. Among non-institutionalised persons, cancer, renal and cardiac disease appeared also related to increased risk, whereas influenza vaccination was associated with reduced risk.

Conclusion: In a setting with relatively low incidence of COVID19 across the first wave of pandemic period, age, nursing-home residence and multiple comorbidities appear predisposing for COVID19 among middle-aged/older adults. Conversely, statins, angiotensin-receptor blockers/inhibitors and influenza vaccination were related with decreased risk.

KEYWORDS: Coronavirus; SARS-COV-2; COVID19; Incidence; Risk.

INTRODUCTION

The novel severe acute respiratory syndrome coronavirus 2 (SARS-COV-2), discovered in December 2019 following an outbreak of pneumonias in Wuhan, China, causes an infectious illness known as coronavirus disease 2019 (COVID19). The World Health Organisation (WHO), due to the alarming levels of spread of the disease and its severity, declared it to be a pandemic on March 11, 2020. [1] Currently the disease continues to spread in the world, with almost 100 million confirmed cases in later January 2021. [2]

In Spain there was a first wave in March-April that was considered contained on May 11, with a total of just over 250,000 diagnosed cases; the contagion curve grew again in June until reaching the maximum peak of second wave in mid-October and as of later January 2021, in the middle of the third wave, the Spanish Ministry of Health reports 2,705,001 cases and almost 58,000 deaths. [3]

There is much information updating COVID19 surveillance epidemiological data (number of cases and deaths by demographic characteristics, accumulated incidences, ratios for positive laboratory testing, etc.); but there is few populationbased data on specific incidence and susceptibility for SARS-COV-2 infection in population subgroups with distinct underlying -believed as at-risk- conditions. [2,3] Earlier studies regarding clinical characteristics and prevalence of comorbidities among COVID19 patients reported that cardiovascular disease (including hypertension), chronic respiratory disease, renal disease, cancer, diabetes mellitus, obesity and smoking were major risk factors for severe COVID19. However, most of these results (mainly based on hospitalised severe cases) are likely confounded by age and other unadjusted variables (including else socio-demographic, medical or health care-related factors). [4-7] Of note, COVID19 patients managed as outpatients were generally poorly represented in COVID19 research studies. [4-7] In this sense, a UK report based on a primary care cohort found that positive SARS-COV-2 testing was associated with similar risk factors (except smoking) as described for severe COVID19 in hospital settings. [8] Nevertheless, there is uncertainty about the possibility that some of these conditions, although they increase the risk of severe COVID19, may not "per se" increase the risk of SARS-COV-2 infection. [9]

In this context, we designed a large population-based retrospective-prospective cohort study (initiated in April 2020 and named as *TARRACO Cohort Study*) aimed to investigate the incidence, clinical characteristics, susceptibility and risk profile of suffering COVID19 among middle-aged and older adults (who supported the greatest burden of COVID19 at this time) in the region of Tarragona across pandemic period.

First-time analyses reporting earlier data have been published. [10,11] The present study updates population-based incidence data and risk of suffering laboratory-confirmed COVID19 among the 79,083 cohort members by distinct underlying medical conditions (previous comorbidities, vaccinations' history and common chronic medications' use), reporting final data observed at the end of the planned study period (from 1st March to 30th June, 2020).

METHODS

Design, setting and study population

The design, setting and study population have been extensively described elsewhere. [10,11] Briefly, this is a retrospective-prospective population-based cohort study (began in April 2020) that included 79,083 persons \geq 50 years old in the region of Tarragona, whose total population (all-age) is 210,672 inhabitants. The cohort incorporates all persons \geq 50 years old assigned to 12 Primary Care Centres (PCCs) managed by the *Institut Català de la Salut (ICS)* in the study area (*Tarragonés, Alt Camp and Conca Barberà* counties). The reference Hospital and Laboratory for the 12 participating PCCs were *Hospital Universitari Joan XXIII* and its Microbiological Department in the city of Tarragona. According to census data, the study cohort represents approximately 75% of inhabitants aged 50 years or older in the area. [12] Cohort members were followed since March 1, 2020 (date before the onset of the epidemic in the region), until the appearance of any study event (COVID19 diagnosis or death), or until the end of 4-month follow-up (June 30, 2020).

The study was approved by the Ethics Committee of the Foundation University Institute for Primary Health Care Research (*IDIAP Jordi Gol*, Barcelona, file 20/065-PCV) and was conducted according to the Helsinki Declaration and Spanish legislation on biomedical studies, data protection and respect for human rights. [13] Considering public health surveillance emergency, informed consent for all cohort members was waived.

Data sources

An institutional clinical research database earlier used for other cohort studies in the area (CAPAMIS), [14] was updated and became the primary data source for this COVID19 epidemiologic report; it includes administrative and clinical data (coded according to the International Classification of Diseases 10th Revision, ICD-10), extracted from the PCCs' clinical records electronic system, that operates since the 2000s. We collected socio-demographic characteristics, previous comorbidities,

> chronic medication and vaccinations' history for cohort members in order to establish their baseline conditions (at March 1, 2020).

> At the COVID19 epidemic period onset in the area, two electronic alerts, linked to COVID19 laboratory tests results and ICD-10 codes for disease suspicion (B34.2 and B97.29), were added to the electronic clinical records system of PCCs. Subsequently, both data sources were associated in the anonymised research database used for this report. Vital status for all cohort members was considered according to administrative data periodically updated in the electronic PCCs' clinical records.

Outcomes

The main outcome was laboratory-confirmed COVID19 diagnosed by reverse transcription-polymerase chain reaction (RT-PCR) or serological testing. For laboratory diagnosis of COVID19, guidelines of the Health Department of the *Generalitat de Catalunya* were implemented. [15] Of note, the availability of PCR testing was scarce during the first weeks of epidemic period, and they were prioritised for severe cases (requiring hospital admission) and nursing-home residences (where several outbreaks took place); while less PCR tests were made among suspected cases with mild symptoms managed as outpatients. We report, in descriptive analysis, presumed COVID19 cases (people who presented clinical suspicion who were no PCR-tested) and, also, COVID19-excluded cases (with negative result in PCR-test) detected among cohort members across the study period.

Covariates

Besides socio-demographic (age, sex and type of residence as communitydwelling/nursing-home), main covariates for this study were vaccinations' history (influenza vaccine in prior autumn, pneumococcal [pneumococcal polysaccharide vaccine -PPV23- and pneumococcal conjugate vaccine -PCV13] or tetanus vaccination at any time) and presence of pre-existing comorbidities. The following comorbidities/underlying conditions were considered according to ICD-10 codes data registered in the electronic PCCs clinical records on 01/03/2020: neurological disease (including dementia and stroke), cancer (solid organ or haematological neoplasia (diagnosed in past 5 years), chronic renal failure, systemic autoimmune rheumatic diseases (including rheumatoid arthritis and lupus), chronic respiratory disease (including chronic bronchitis/emphysema and/or asthma), chronic heart disease (including congestive heart failure, coronary artery disease and other

cardiopathies), atrial fibrillation, chronic liver disease (including chronic hepatitis and cirrhosis), arterial hypertension, diabetes mellitus, hypercholesterolemia, obesity, alcoholism and smoking.

In addition, baseline use of some common chronic medications, which could be hypothetically related with physiopathologic mechanism of SARS-COV-2 infection or virulence (e.g., antihypertensive, antiplatelet/anticoagulant and/or anti-inflammatory drugs), were also considered as exposure co-variables possibly related with the occurrence of COVID19 for the present study. Thus, active medication treatments in each cohort member on 01/03/2020, coded according to the Anatomical, Therapeutic, and Chemical classification system (ATC) of the World Health Organization, [16] were identified from the patient treatment plan registered in the PCC's clinical records system, including the following therapeutic groups: antihypertensive (diuretics, beta-blockers, angiotensin-converting enzyme inhibitors [ACEIs], angiotensin II receptor blockers [ARBs], calcium channel blockers), statins, anticoagulants (warfarin and new oral anticoagulant drugs), antiplatelet drugs, antidiabetic drugs (insulin, oral anti-diabetic drugs), inhaled respiratory drugs, antineoplastic agents, systemic corticosteroids, non-steroidal anti-inflammatory drugs (NSAIDs), chloroquine/hydroxychloroquine, antihistamines, proton-pump inhibitors and benzodiazepines, were registered. Criteria used to define and collect preexisting conditions and chronic medications use are extensively described in an appendix elsewhere. [11]

Statistical analyses

We calculated frequencies and percentages, means and standard deviations (SD) for qualitative and continuous variables, respectively. Incidence rates (IRs) for laboratory-confirmed COVID19 cases were calculated for persons-weeks, considering in the denominator the sum of time contributed by each cohort member during the study period (17.4 weeks). Confidence intervals (CIs) for IRs were calculated assuming a Poisson distribution for uncommon events. In bivariate analyses, baseline characteristics according to final presence/absence of study outcome (COVID19 diagnosis) were compared using Chi-squared or Fisher's test as appropriate.

Cox regression analyses were used to calculate unadjusted, age&sex-adjusted and multivariable-adjusted hazard ratios (HRs) aimed at estimating the association between baseline exposure conditions and the time to the first outcome (laboratory-confirmed COVID19) in cohort members throughout the study period (from 1st March to 30th June 2020). We included in exploratory multivariable Cox models all above

mentioned exposure and covariates (i.e., age, sex, residence, vaccinations' history, comorbidities/underlying conditions and medication use). The method to select a subset of variables to include in final models was the purposeful selection. [17] These models incorporate significant and confounder variables as well as all covariates judged clinically or epidemiologically relevant. We performed a main analysis with the total study cohort (N=79,083) and two subgroup analyses restricted to community-dwelling people (N=77,669) and nursing-home residents (N=1414). Statistical significance was set at p <0.05 (two-tailed). Analyses were performed using IBM SPSS Statistics for Windows, version 24 (IBM Corp., Armonk, N.Y., USA).

RESULTS

Study cohort included 37,626 (47.6%) men and 41,457 (52.4%) women, with a mean age of 65.8 (SD: 11.3) years. Stratifying by age, 42,684 (54%) were 50-64 years and 10,386 (13.1%) were 80 years or more; by residence, 1414 (1.8%) were nursing-home residents and 77,669 (98.2%) community-dwelling individuals.

Total cohort members were followed for 1,356,358 persons-weeks (1,335,069 persons-weeks community-dwelling and 21,289 persons-weeks nursing-home residents).

Of the 79,083 cohort members, 4084 (5.2%) were PCR tested for SARS-COV-2 infection across the study. Of them, 3577 (87.6%) had a negative result and 507 (12.4%) had a positive result (319 community-dwelling and 188 nursing-home residents). Additionally, 29 cohort members had a positive laboratory serological testing for SARS-COV-2 (twenty-six community-dwelling and three nursing-home residents). In two hundred and eighty-three cohort members COVID19 was suspected (clinically compatible) but no laboratory test was performed.

Of the 536 laboratory-confirmed COVID19 cases, 235 (43.8%) were men and 301 (56.2%) women, with a mean age of 74 years (SD: 14.4). By age groups, 177 (33.0%) occurred in those who were 50-64 years, 137 (25.6%) in those 65-79 years and 222 (41.4%) in aged 80 years or more. Three hundred and forty-five cases (64.4%) occurred in community-dwelling and 191 (35.6%) in nursing-home residents. The most prevalent comorbidities were hypertension (54.1%), cardiac disease (28.5%), obesity (25.2%), diabetes (23.5%), respiratory disease (14.9%) and neurological disease (14.7%).

Figure 1 shows weekly distribution of COVID19-confirmed cases and cumulative incidences at 7-days and 14-days across the 18 weeks of the study period.



Figure 1. Weekly distribution of COVID19-confirmed cases accross study period (March 1st to June 30th, 2020).

Mean IR across study period was 39.5 (95% CI: 36.1-43.2) cases per 100,000 persons-week (24.1 in 50-64 years vs 30.6 in 65-79 years vs 127.0 in 80 years or older; 36.4 in men vs 42.3 in women; 25.8 in community-dwelling vs 897.2 in nursing-home residents).

By underlying conditions, maximum IRs (per 100,000 persons-week) emerged among those persons with neurological diseases (206.0) followed by atrial fibrillation (109.4), chronic renal failure (85.9), chronic heart disease (66.9), inflammatory bowel disease (66.0), chronic respiratory disease (64.6), cancer (59.6), diabetes (55.4), and hypertension (48.5). Lower IRs were observed among persons with rheumatic diseases (47.0), liver disease (44.0), hypercholesterolemia (38.6), obesity (36.3), alcoholism (35.7) and smokers (18.7).

According to baseline chronic medications, maximum IRs (per 100,000p-w) appeared among those receiving diuretics (97.0), oral anticoagulants (87.3), insulin (83.1), and inhaled-respiratory therapy (72.1) (Table 1).

Table 2 shows unadjusted, age&sex-adjusted and multivariable-adjusted analyses evaluating risk of suffering laboratory-confirmed COVID19 in the whole study cohort. In the unadjusted analysis, many previous comorbidities and medications were associated with increased risk. However, after multivariable-adjustment, only age (HR: 1.01; 95% CI: 1.00-1.02; p=0.051), nursing-home residence (HR: 20.19; 95% CI: 15.98-25.51; p<0.001), neurological disease (HR: 1.35; 95% CI: 1.03-1.77; p=0.029), taking diuretics (HR: 1.39; 95% CI: 1.10-1.75; p=0.006), antiplatelet drugs (HR: 1.36; 95% CI: 1.05-1.76; p=0.021), benzodiazepines (HR: 1.24; 95% CI: 1.00-1.53; p=0.047) and history of PCV13 vaccination (HR: 1.74; 95% CI: 1.06-2.85;

p=0.028) appeared significantly associated with increased risk. Oppositely, smoking (HR: 0.57; 95%CI: 0.41-0.80; p=0.001), taking angiotensin-converting enzyme (ACE) inhibitors (HR: 0.78; 95% CI: 0.61-1.00; p=0.048), angiotensin II receptor blockers - ARBs- (HR: 0.70; 95%CI: 0.51-0.96; p=0.027) and statins (HR: 0.75; 95% CI: 0.58-0.96; p=0.025) appeared associated with a reduced risk.

Table 3 (incidence rates) and Table 4 (unadjusted and adjusted risks) show subgroup analyses restricted to community-dwelling persons (N=77,669). In the multivariable analysis focused on these individuals, chronic renal disease (HR: 1.50; 95% CI: 1.02-2.21; p=0.038), cancer (HR: 1.45; 95% CI: 1.05-2.01; p=0.023), cardiac disease (HR: 1.32; 95% CI: 0.98-1.79; p=0.070), taking diuretics (HR: 1.54; 95% CI: 1.11-2.13; p=0.010) and benzodiazepines (HR: 1.28; 95% CI: 0.97-1.68; p=0.078) were associated (significantly or near significantly) with increased risk. Oppositely, smoking (HR: 0.51; 95% CI: 0.35-0.74; p<0.001), receiving ACE-inhibitors (HR: 0.63; 95% CI: 0.45-0.89; p=0.009), statins (HR: 0.66; 95% CI: 0.49-0.90; p=0.009) and influenza vaccination in prior autumn (HR: 0.68; 95% CI: 0.51-0.92; p=0.012) were associated with decreased risk.

Table 5 shows subgroup analyses restricted to nursing-home residents (N=1414). In the multivariable analysis, increasing age (HR: 1.03; 95% CI: 1.01-1.05; p=0.001), oral anti-diabetic drugs (HR: 1.79; 95% CI: 1.04-3.08; p=0.037), antineoplastic agents (HR: 2.51; 95% CI: 1.05-6.01; p=0.039) and influenza vaccination in prior autumn (HR: 1.56; 95% CI: 1.07-2.29; p=0.022) were associated with increased risk. In contrast, ARBs (HR: 0.47; 95% CI: 0.25-0.87; p=0.017) and history of tetanus vaccination (HR: 0.63; 95% CI: 0.44-0.91; p=0.013) were associated with decreased risk.

DISCUSSION

Population-based clinical data on susceptibility and risk of SARS-COV-2 infection is limited. [9, 18] This large cohort study was aimed to assess population-based incidence of (laboratory-confirmed) COVID19 among middle-aged and older adults with distinct underlying medical conditions in a well characterised geographic area (Tarragona region) during the first wave of the COVID19 epidemic (March-June 2020); we investigated possible relationships between pre-existing medical conditions (including common comorbidities, vaccinations' history and chronic medication use) with the susceptibility/risk of suffering COVID19.

As main result, we found a mean IR of 39.5 laboratory-confirmed COVID19 cases per 100,000 persons-week thru study period, which is a relatively low incidence as compared to other Spanish or European regions during the same time period. [19,20].

During the study period (17.4 weeks since 1st March until 30th June 2020), approximately 5% of cohort members were PCR tested (with 12.4% positive results). Taking this data into account, and considering that PCR test were scarcely available in the study area for patients with less severe symptoms during the first weeks of the epidemic period, the true incidence of COVID19 was logically underestimated. Nevertheless, considering the relatively low number of presumptive cases (clinical suspicion alone without PCR performed), our data also suggests that the overall number of infected people was considerably lower than initially speculated (as showed by the national seroprevalence study). [21] Assuming this relatively little incidence, we found that increasing age, nursing-home residence, some pre-existing comorbidities (cancer, neurological, renal or cardiac disease and taking certain chronic medications (diuretics, antiplatelet drugs and benzodiazepines) were independent factors related to increased risk of suffering COVID19. In contrast, cohort members receiving statins or ACEIs/ARBs (and, in the case of communitydwelling individuals, those vaccinated against influenza in prior autumn) presented, after multivariable adjustment, a decreased risk.

Of note, in the crude analyses several covariates (comorbidities, medications and vaccinations) appeared significantly associated with increased risk/susceptibility of suffering COVID19. However, for most of them the significance disappeared after age&sex-adjustment and only a few remained independently associated with an increased risk after multivariable adjustments. This highlights the importance of reporting adjusted data and note that differences in adjustment methods may explain, in part, the heterogeneous results (when evaluating the possible association between some pre-existing conditions and risk of COVID19) reported to date. [22] At present, there is not clear evidence about possible clinical predisposing and protecting factors front SARS-COV-2 infection, and uncertainty surrounds this issue. [23]

If we consider demographic characteristics, apart from nursing-home residence that increased twenty-times the adjusted-risk of suffering laboratory-confirmed COVID19, we found that age increased approximately 1-2% for each year the adjusted-risk of suffering COVID19. While COVID19 was more frequent in women, sex did not alter significantly the risk of infection in multivariable analysis.

Considering pre-existing comorbidities, only neurological disease (basically patients with a history of dementia and/or stroke living in nursing-home residences) was associated with increased multivariable-adjusted risk of infection in the analysis of full cohort. Concomitantly, previous cancer, renal disease and cardiac disease emerged significantly associated with increased risk in subgroup analysis restricted to community-dwelling individuals.

In the present study, chronic respiratory disease, diabetes, hypertension and obesity did not emerge independently associated with significant increased risk of suffering laboratory-confirmed COVID19 after multivariable-adjusted analyses. There is general accordance considering these conditions as major risk factors for poor prognosis in hospitalised COVID19 patients, but there is few data assessing their role in predisposing for SARS-COV-2 infection. [24]

Surprisingly, smokers presented the lowest specific incidence rate of COVID19 in the present report, and smoking was associated with a statistically significant reduced risk of COVID19, both in multivariable analyses evaluating the total study cohort and in the restricted to community-dwelling individuals. Notably, active smoking has been linked with decreased odds of a positive result SARS-CoV-2-test in other studies. [8] Opposite findings about poor prognosis among smokers with COVID19 have been reported. [25,26]

Considering chronic medication use, receiving diuretics, antiplatelet drugs and benzodiazepines appeared significantly associated with increased risk of suffering COVID19. In contrast, taking statins and ACEIs or ARBs emerged associated with decreased risk. At onset of COVID19 pandemic, after discovering that the host receptor for SARS-CoV-2 cell entry was the angiotensin-converting enzyme 2 (ACE2), it was feared that receiving ACEIs/ARBs could facilitate SARS-COV-2 infection and predispose to suffer severe COVID-19. However, most recent studies have concluded that there is no clinical or experimental evidence supporting that ACEIs or ARBs augment the susceptibility to/severity of COVID-19 at present. Indeed, ACEIs/ARBs have been associated with lower incidence and/or improved outcomes in patients with lower respiratory tract infections, and with lower risk of all-cause mortality in COVID19 hospitalised patients. Our data also support this conclusion.

Considering statins, while first studies reported opposite results, our data showing a decreased risk of laboratory-confirmed COVID19 among people taking these drugs are in accordance with conclusions of a recent meta-analysis on this concern. [28] Anyway, caution is recommended until further evidence emerges surrounding statin use in COVID19 patients.

> Regarding other chronic medications that appeared associated with increased risk in the current study (i.e., diuretics, antiplatelet drugs and benzodiazepines), we have not a clear interpretation for this data. It is possibly due to an unresolved residual confounding, linked to unmeasured health-care-related factors or insufficient ageadjustment, but we also note that some of them (e.g., benzodiazepines) have already been reported associated with greater risk of pneumonia (common clinical manifestation in COVID19 patients).

> Antihistamines' use was not associated with any statistically significant change in the risk of suffering COVID-19 in the present study. We note that antihistamines' use was associated (marginally significative) with a lower risk of COVID19 in an earlier report, [12] but we also remark that it could be due to an alpha error considering that this final report has greater statistical power (since include more persons-time follow-up and greater number of events analysed).

Similar to data observed in earlier analyses, we found that influenza vaccination in prior autumn was significantly associated with a reduction risk of suffering COVID19 among community-dwelling individuals. [10] Conversely, it was associated with a greater risk in the subgroup of nursing-home residents (where a significant protective association appeared for those who had received tetanus vaccination). While these associations could be due to residual confounding related with unmeasured life-style or healthcare factors, further investigations are needed exploring a possible immunity-related mechanism explanation. [29,30]

Major strengths in this study were the representativity (due to large size) of the study cohort, that included almost 80,000 people, which represents about 75% of the entire population over 50 years in the area, [13] and the use of survival analysis to precisely asses susceptibility and risk of suffering a SARS-COV-2 infection adjusting by important potential confounders (as socio-demographic, and medical underlying conditions clinically relevant)

Major limitations were related to its partially retrospective design and poor access to PCR tests at the beginning of the epidemic period in our setting, especially considering that the most specific diagnosis for COVID19 is a positive PCR test. Moreover its reliability depends on the correct collection of nasopharyngeal swabs, moment of collection, sensitivity of tests and guidelines for testing followed throughout the study period. Due to above mentioned scarce availability of PCR tests they were mainly performed in hospitalised/severe patients and in nursing homes (after some outbreaks were detected). Thereby, patients with mild symptoms (many of them without PCR done) have been logically underestimated in this study.

Even though we did subgroup analyses (community-dwelling/nursing-home) and multivariable-adjustments, residual confounding when estimating risks (common in observational studies), tied to non-included variables such as socio-economical, labor-, lifestyle- or health care-related factors, may not be discarded.

Despite the large size of the cohort, there were relatively few events (536 laboratoryconfirmed COVID19 cases), which limits statistical power. This is especially relevant in the subgroup of nursing-home residents where, even if great incidence occurred (200 cases per 100,000 persons-week), only 191 confirmed COVID19 cases were detected, fact that largely affects the precision of estimates (confidence intervals very wide).

Notwithstanding all mentioned limitations, authors may emphasise the added value that provide adjustments (age&sex and multivariable), not present in many other studies. Their importance is clearly evident if we compare, for example in Table 3 and 5, crude with adjusted results.

Concerning external validity, we must consider that the study was conducted in a single geographic area; thus, specific incidence data cannot be directly extrapolated to other regions with distinct epidemic conditions.

Nevertheless, we also note that the adjusted-risks estimated here may be helpful to better characterize the risk profile (socio-demographic, pre-existing conditions/comorbidities, chronic medication use) for suffering COVID19 among middle-aged and older adults living in a setting with low/moderate intensity of COVID19, as in the present study.

CONCLUSIONS

In summary, in this large population-based study that followed a cohort of 79,083 individuals ≥50 years old between March-June 2020 in Tarragona area, the incidence of laboratory-confirmed SARS-COV-2 infection was relatively low (39.5 cases per 100,000 persons-week) as compared with that of other Spanish regions during the first wave of COVID19 pandemic.

In this epidemic context (low incidence), our data support that increasing age, nursing-home residence, pre-existing cancer, neurological, renal and cardiac disease are independent major predisposing conditions to suffer COVID19 among middle-aged and older adults. Patients receiving diuretics, antiplatelet drugs and benzodiazepines were also at increased risk. Oppositely, patients taking statins, rennin-angiotensin-aldosterone system inhibitors and those community-dwelling

individuals that received influenza vaccination in prior autumn presented decreased risk.

ACKNOWLEDGEMENTS

This study was supported by a grant from the Instituto de Salud Carlos III of the Spanish Health Ministry (file COV20/00852; call for the SARS-COV-2/COVID-19 disease, RDL 8/2020, March 17, 2020).

CONTRIBUTORS

ESG and AVC wrote the manuscript; CDC, CTF, FGB, and IHG obtained data; ESG and AVR assessed outcomes and did statistical analyses; AVR edited the manuscript; OOG and FML revised the final version; AVC coordinated the study.

FUNDING

This study is supported by a grant from the Instituto de Salud Carlos III of the Spanish Health Ministry (file COV20/00852; call for the SARS-COV-2/COVID19 disease, RDL 8/2020, March 17, 2020). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

COMPETING INTEREST

All authors, none declared.

REFERENCES

- 1.World Health Organisation (WHO). Listings of WHO's response to COVID-19. Available at: <u>https://www.who.int/news/item/29-06-2020-covidtimeline</u>. [Accessed January 20, 2021].
- 2. World Health Organisation (WHO). Coronavirus Disease (COVID19). Weekly epidemiological update-27 January 2021. Available at: <u>https://www.who.int/publications/m/item/weekly-epidemiological-update---27-january-2021</u>. [Accessed]
- January 28, 2021].
 3. Ministerio de Sanidad. Dirección General de Salud Pública. Centro de Coordinación de Alertas y Emergencias Sanitarias. Actualización nº 300. Enfermedad por el coronavirus (COVID-19). 28.01.2021. Available https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov/documentos/Actualizac

ion_300_COVID-19.pdf. [Accessed January 28, 2021].

- 4. Rodríguez-Molinero A, Gálvez-Barrón C, Miñarro A, et al. Association between COVID-19 prognosis and disease presentation, comorbidities and chronic treatment of hospitalized patients. PLoS One. 2020;15(10):e0239571. doi:10.1371/journal.pone.0239571.
- 5. Grasselli G, Zangrillo A, Zanella A, et al. Baseline Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV-2 Admitted to ICUs of the Lombardy Region, Italy. JAMA. 2020; 323:1574□81. doi: 10.1001/jama.2020.5394.
- 6. Yang J, Zheng Y, Gou X, et al. Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis. Int J Infect Dis. 2020; 94: 91-95. doi: 10.1016/j.ijid.2020.03.017.
- 7. Ssentongo P, Ssentongo AE, Heilbrunn ES, Ba DM, Chinchilli VM. Association of cardiovascular disease and 10 other pre-existing comorbidities with COVID-19 mortality: A systematic review and metaanalysis. PLoS One. 2020;15(8):e0238215. doi: 10.1371/journal.pone.0238215.
- 8. de Lusignan S, Dorward J, Correa A, et al. Risk factors for SARS-CoV-2 among patients in the Oxford Royal College of General Practitioners Research and Surveillance Centre primary care network: a cross-sectional study. Lancet Infect Dis. 2020;20(9):1034-42. doi: 10.1016/S1473-3099(20)30371-6.
- 9. Lipsitch M, Swerdlow DL, Finelli L. Defining the epidemiology of Covid-19—studies needed. N Engl J Med. 2020;382:1194-96. doi: 10.1056/NEJMp2002125.
- 10. Vila-Córcoles Á, Ochoa-Gondar O, Torrente-Fraga C, et al. Evaluación de la incidencia y perfil de riesgo de Covid-19 según comorbilidad previa en adultos ≥50 años del área de Tarragona [Evaluation of incidence and risk profile for suffering Covid-19 infection by underlying conditions among middle-aged and older adults in Tarragona.]. *Rev Esp Salud Publica*. 2020;94:e202006065.
- 11. Vila-Córcoles A, Ochoa-Gondar O, Satué-Gracia EM, et al. Influence of prior comorbidities and chronic medications use on the risk of COVID-19 in adults: a population-based cohort study in Tarragona, Spain. *BMJ Open*. 2020;10(12):e041577. doi:10.1136/bmjopen-2020-041577.
- 12. Generalitat de Catalunya. Statistical institute of Catalonia. IDESCAT. Available at: <u>https://www.idescat.cat/?lang=en</u>. [Accessed April 15, 2020]
- 13. World Medical Association. WMA Declaration of Helsinki Ethical Principles for Medical Research Involving Human Subjects. Available at: https://www.wma.net/policies-post/wmadeclaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/. [Accessed January 15, 2021].
- 14. Vila-Corcoles A, Hospital-Guardiola I, Ochoa-Gondar O, et al. Rationale and design of the CAPAMIS study: effectiveness of pneumococcal vaccination against community-acquired pneumonia, acute myocardial infarction and stroke. BMC Public Health. 2010;10:25. doi: 10.1186/1471-2458-10-25.
- 15. Who Collaborating Centre for Drug Statistics Methodology. ATC/DDD Index 2020. Available at: https://www.whocc.no/atc_ddd_index/. [Accessed May 12, 2020].
- 16. Generalitat de Catalunya. Subdirecció General de Vigilància i Resposta a Emergències de Salut Pública. Procediment d'actuació enfront de casos d'infecció pel nou coronavirus SARSCoV-2. Available at: https://canalsalut.gencat.cat/web/.content/_A-Z/C/coronavirus-2019-ncov/materialdivulgatiu/procediment-actuacio-coronavirus.pdf. [Accessed April 16, 2020]
- 17. Hosmer DW, Lemeshow S. Applied Survival Analysis. Regression Modelling of Time to Event Data. New York: John Wiley & Sons, 1999.
- 18. Koh WC, Naing L, Chaw L, et al. What do we know about SARS-CoV-2 transmission? A systematic review and meta-analysis of the secondary attack rate and associated risk factors. *PLoS One*. 2020;15(10):e0240205. doi:10.1371/journal.pone.0240205.
- 19. Gobierno de España. Secretaría General de Sanidad y Consumo. Dirección General de Salud Pública, Calidad e innovación. Centro de Coordinación de Alertas y Emergencias Sanitarias. Información científica-técnica. Enfermedad por coronavirus, COVID-19 Actualización 12 de noviembre, 2020. Available

https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov/documentos/ITCorona virus.pdf. [Accessed December 5, 2020].

 20. European Centers for Disease Prevention and Control. Situation updates on COVID19. Available at: <u>https://www.ecdc.europa.eu/en/covid-19/situation-updates</u>. [Accessed June 30, 2020]

- 21. Instituto de Salud Carlos III. Estudio ENE-COVID19: informe final. Estudio nacional de seroepidemiología de la infección por SARS-CoV-2 en España. July 6, 2020. Available at: <u>https://www.mscbs.gob.es/ciudadanos/ene-covid/docs/ESTUDIO_ENE-</u> <u>COVID19_INFORME_FINAL.pdf</u>. [Accessed July 30, 2020].
- 22. Yusuf E, Maiwald M. COVID-19, equipoise and observational studies: a reminder of forgotten issues [published online ahead of print, 2020 Jun 25]. *Infection.* 2020;1-3. doi:10.1007/s15010-020-01466-9.
- 23. Fakhroo AD, AI Thani AA, Yassine HM. Markers Associated with COVID-19 Susceptibility, Resistance, and Severity. *Viruses*. 2020;13(1):45. doi:10.3390/v13010045
- 24. Centres for Disease Control and Prevention. Coronavirus Disease 2019 (COVID19). People at increased risk for severe illness. Available at: https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html. [Accessed December 5, 2020].
- 25. Zhao Q, Meng M, Kumar R, et al. The impact of COPD and smoking history on the severity of COVID-19: A systemic review and meta-analysis. *J Med Virol.* 2020;92(10):1915-21. doi:10.1002/jmv.25889-
- 26. Lippi G, Henry BM. Active smoking is not associated with severity of coronavirus disease 2019 (COVID-19). Eur J Intern Med. 2020; 75:107-108. doi: 10.1016/j.ejim.2020.03.014.
- 27. Kreutz R, Algharably EAE, Azizi M, et al. Hypertension, the renin-angiotensin system, and the risk of lower respiratory tract infections and lung injury: implications for COVID-19. *Cardiovasc Res.* 2020;116(10):1688-99. doi:10.1093/cvr/cvaa097.
- 28. Kow CS, Hasan SS. Meta-analysis of Effect of Statins in Patients with COVID-19. Am J Cardiol. 2020;134:153-55. doi: 10.1016/j.amjcard.2020.08.004.
- 29. Salem ML, El-Hennawy D. The possible beneficial adjuvant effect of influenza vaccine to minimize the severity of COVID-19. Med Hypotheses 2020;140:109752. doi:10.1016/j.mehy.2020.109752.
- 30. letto G. SARS CoV-2: Reasons of epidemiology of severe ill disease cases and therapeutic approach using trivalent vaccine (tetanus, diphtheria and Bordetella pertussis). *Med Hypotheses*. 2020;141:109779. doi:10.1016/j.mehy.2020.109779.

TABLES

Table 1. Incidence of laboratory-confirmed COVID-19 cases according to baseline demographic and clinical characteristics in the total study cohort (N=79,083). Tarragona region, 01/03/2020-30/06/2020.

CHARACTERISTIC	Study population (N=79,083)	LABORATORY-CONFIRMED COVID-19 CASES (n=536)			
		Univariate analysis	Time follow-up	Incidence rate	
	n (%)	n (%) p-value	(persons-week)	IR (95% CI)	
Socio-demographic				, , , , , , , , , , , , , , , , , , ,	
Age: 50-64 years	42.684 (54.0)	177 (33.0) <0.001	734.123	24.1 (20.8-28.0)	
65-79 years	26.013 (32.9)	137 (25.6)	447.474	30.6 (25.8-36.2)	
≥80 years	10,386 (13.1)	222 (41.4)	174,761	127.0 (110.2-146.2)	
Sex Men	37,626 (47.6)	235 (43.8) 0.082	645,470	36.4 (32.1-41.3)	
Women	41,457 (52.4)	301 (56.2)	710,888	42.3 (37.7-47.4)	
Community-dwelling	77,669 (98.2)	345 (64.4) <0.001	1,335,069	25.8 (23.2-28.7)	
Nursing-home residence	1414 (1.8)	191 (35.6)	21,289	897.2 (778.8-1032.7)	
Comorbidities	· · · · ·				
Neurological disease	2317 (2.9)	79 (14.7) <0.001	38,344	206.0 (164.4-257.5)	
Renal disease	4476 (5.7)	65 (12.1) <0.001	75,629	85.9 (67.4-109.1)	
Cancer	6630 (8.4)	67 (12.5) 0.001	112,481	59.6 (46.8-75.7)	
Rheumatic disease	872 (1.1)	7 (1.3) 0.651	14,897	47.0 (18.8-96.8)	
Inflammatory bowel disease	532 (0.7)	6 (1.1) 0.204	9097	66.0 (24.2-143.9)	
Respiratory disease	7272 (9.2)	80 (14.9) <0.001	123,794	64.6 (51.6-80.8)	
Cardiac disease	13,435 (17.0)	153 (28.5) <0.001	228,716	66.9 (57.1-78.3)	
Atrial fibrillation	3786 (4.8)	70 (13.1) <0.001	63,962	109.4 (85.9-138.9)	
Liver disease	1465 (1.9)	11 (2.1) 0.731	24,998	44.0 (22.0-78.8)	
Diabetes	13,317 (16.8)	126 (23.5) <0.001	227,640	55.4 (46.1-66.5)	
Hypertension	34,945 (44.2)	290 (54.1) <0.001	597,890	48.5 (43.3-54.4)	
Hypercholesterolemia	27,314 (34.5)	181 (33.8) 0.707	468,535	38.6 (33.3-44.8)	
Obesity	21,678 (27.4)	135 (25.2) 0.247	372,086	36.3 (30.6-43.0)	
Smoking	12,750 (16.1)	41 (7.6) <0.001	219,518	18.7 (13.4-25.4)	
Alcoholism	1796 (2.3)	11 (2.1) 0.733	30,774	35.7 (17.8-63.9)	
Chronic medication use	Γ				
Diuretics	8481 (10.7)	139 (25.9) <0.001	143,357	97.0 (81.9-114.8)	
Beta blockers	9571 (12.1)	87 (16.2) 0.003	163,450	53.2 (43.0-66.0)	
ACEIs	16,419 (20.8)	114 (21.3) 0.772	281,514	40.5 (33.7-48.6)	
ARBs	8869 (11.2)	54 (10.1) 0.401	152,376	35.4 (26.3-46.7)	
Calcium channel blockers	6490 (8.2)	64 (11.9) 0.002	111,107	57.6 (44.4-74.9)	
Statins	16,134 (20.4)	91 (17.0) 0.048	277,237	32.8 (26.5-40.7)	
Oral anticoagulants	3912 (4.9)	58 (10.8) <0.001	66,424	87.3 (67.2-113.5)	
Antiplatelet drugs	9154 (11.6)	108 (20.1) <0.001	155,957	69.2 (56.6-84.4)	
Insulin	3042 (3.8)	43 (8.0) <0.001	51,741	83.1 (60.6-111.4)	
Oral anti-diabetic drugs	10,585 (13.4)	85 (15.9) 0.091	181,452	46.8 (37.9-58.0)	
Inhaled respiratory drugs	6293 (8.0)	// (14.4) <0.001	106,857	72.1 (57.5-90.1)	
Antineoplastic agents	1614 (2.0)	12 (2.2) 0.745	27,691	43.3 (22.4-75.8)	
Systemic corticosteroids	1252 (1.6)	10 (1.9) 0.599	21,030	47.6 (22.8-87.6)	
NSADIS	4321 (5.5)	21 (3.9) 0.114	74,152	28.3 (17.5-43.3)	
	168 (0.2)	1 (0.2) 0.896	2867	34.9 (0.9-194.4)	
Antinistamines	3264 (4.1)	15 (2.8) 0.121	56,243	20.7 (15.0-44.1)	
Proton-Pump Innibitors	17,931 (22.7)	183 (34.1) <0.001	305,566	59.9 (51.6-69.5)	
Vacaination's history	13,040 (16.5)	120 (23.5) <0.001	222,537	50.0 (47.1-67.9)	
Elu vaccina in prior outurer	22 606 (22 6)	250 (40 4) -0.004	205 660	66.0 (50.0.75.0)	
PIU Vaccine in prior autumn		200 (40.1) <0.001	300,000 447,070	00.9 (39.0-73.9)	
	20,103 (33.1)	212(30.7) < 0.001	447,270	00.0 (33.0-00.9)	
	50 700 (64 2)	10 (3.4) <0.001	13,272	30.4 (33.4-147.0)	
reidhus	JU,799 (04.2)	344 (04.2) 0.978	071,204	JY.J (JJ.J-43.9)	

NOTE: p-values in univariate analysis were calculated by chi-squared, or Fisher's test as appropriate, comparing percentages in the study population vs in COVID-19 cases; IR denotes incidence rates per 100,000 persons-week; CIs denotes confidence intervals for incidence rates and were calculated assuming a Poisson distribution for uncommon events.

Table 2. Cox regression analyses assessing unadjusted, age & sex adjusted and multivariable-adjusted risks to suffer laboratory-confirmed COVID-19 in the total study cohort (N=79,083). Tarragona region (Southern Catalonia, Spain), from 01/03/2020 to 30/06/2020.

	LABORATORY-CONFIRMED COVID-19 CASES (n=536)				
CHARACTERISTIC	Unadjusted	Age & sex adjusted	Multivariable-adjusted		
	HR (95% CI) p-value	HR (95% CI) p-value	HR (95% CI) p-value		
Socio-demographic					
Age (continuous years)	1.06 (1.05-1.07) <0.001	1.06 (1.05-1.07) <0.001	1.01 (1.00-1.02) 0.051		
Sex: women	1.16 (0.98-1.38) 0.083	1.00 (0.84-1.19) 0.992	0.89 (0.74-1.07) 0.214		
Nursing-home residence	33.18 (27.80-39.59) <0.001	23.80 (19.18-29.53) <0.001	20.19 (15.98-25.51)		
			<0.001		
Comorbidities					
Neurological disease	5.86 (4.62-7.44) <0.001	3.01 (2.33-3.88) <0.001	1.35 (1.03-1.77) 0.029		
Renal disease	2.33 (1.80-3.02) <0.001	1.09 (0.83-1.44) 0.518	1.01 (0.76-1.34) 0.959		
Cancer	1.57 (1.22-2.03) 0.001	1.16 (0.90-1.50) 0.264	1.18 (0.91-1.55) 0.216		
Rheumatic disease	1.19 (0.57-2.51) 0.646	1.07 (0.51-2.25) 0.866	1.29 (0.59-2.84) 0.519		
Respiratory disease	1.74 (1.38-2.21) <0.001	1.36 (1.07-1.73) 0.012	1.16 (0.83-1.62) 0.377		
Cardiac disease	1.97 (1.63-2.37) <0.001	1.21 (1.00-1.48) 0.054	0.99 (0.79-1.25) 0.950		
Atrial fibrillation	3.02 (2.35-3.89) <0.001	1.57 (1.20-2.04) 0.001	1.25 (0.83-1.88) 0.281		
Liver disease	1.12 (0.61-2.03) 0.721	1.26 (0.69-2.28) 0.453	1.13 (0.62-2.06) 0.699		
Diabetes	1.52 (1.25-1.86) <0.001	1.13 (0.92-1.38) 0.243	1.09 (0.76-1.55) 0.644		
Hypertension	1.49 (1.26-1.77) <0.001	0.83 (0.69-1.00) 0.050	0.97 (0.77-1.23) 0.823		
Obesity	0.89 (0.73-1.08) 0.245	0.81 (0.66-0.98) 0.032	0.90 (0.73-1.11) 0.328		
Smoking	0.43 (0.31-0.59) <0.001	0.69 (0.49-0.95) 0.024	0.57 (0.41-0.80) 0.001		
Alcoholism	0.90 (0.50-1.64) 0.735	1.16 (0.64-2.13) 0.623	0.96 (0.52-1.77) 0.896		
Chronic medication use					
Diuretics	2.95 (2.43-3.58) <0.001	1.71 (1.39-2.10) <0.001	1.39 (1.10-1.75) 0.006		
Beta blockers	1.41 (1.12-1.78) 0.003	1.01 (0.80-1.28) 0.920	0.94 (0.73-1.22) 0.664		
ACEIs	1.03 (0.84-1.27) 0.769	0.75 (0.61-0.92) 0.007	0.78 (0.61-1.00) 0.048		
ARBs	0.89 (0.67-1.17) 0.399	0.62 (0.47-0.82) 0.001	0.70 (0.51-0.96) 0.027		
Calcium channel blockers	1.52 (1.17-1.97) 0.002	1.04(0.80-1.35) 0.793	1.19 (0.90-1.58) 0.225		
Statins	0.80 (0.64-1.00) 0.049	0.60 (0.48-0.75) <0.001	0.75 (0.58-0.96) 0.025		
Oral anticoagulants	2.35 (1.79-3.09) <0.001	1.29 (0.98-1.71) 0.074	1.18 (0.76-1.84) 0.456		
Antiplatelet drugs	1.94 (1.57-2.39) <0.001	1.21 (0.97-1.51) 0.085	1.36 (1.05-1.76) 0.021		
Insulin	2.20 (1.61-3.00) <0.001	1.64 (1.20-2.25) 0.002	1.24 (0.85-1.81) 0.261		
Oral anti-diabetic drugs	1.22 (0.97-1.54) 0.091	0.94 (0.74-1.18) 0.584	1.02 (0.70-1.48) 0.922		
Inhaled respiratory drugs	1.96 (1.54-2.49) <0.001	1.43 (1.12-1.83) 0.004	1.23 (0.87-1.72) 0.242		
Antineoplastic agents	1.10 (0.62-1.95) 0.746	1.04 (0.59-1.84) 0.899	0.94 (0.51-1.73) 0.841		
Systemic corticosteroids	1.20 (0.64-2.24) 0.574	0.88 (0.47-1.65) 0.687	0.80 (0.42-1.53) 0.499		
NSAIDs	0.71 (0.46-1.09) 0.118	0.84 (0.54-1.30) 0.433	1.13 (0.72-1.76) 0.603		
Antihistamines	0.67 (0.40-1.12) 0.122	0.68 (0.41-1.14) 0.144	0.73 (0.43-1.22) 0.231		
Proton-Pump Inhibitors	1.78 (1.49-2.13) <0.001	1.11 (0.92-1.34) 0.276	0.92 (0.74-1.15) 0.464		
Benzodiazepines	1.56 (1.28-1.91) <0.001	1.19 (0.97-1.46) 0.092	1.24 (1.00-1.53) 0.047		
Vaccination's history					
Flu vaccine (prior autumn)	2.33 (1.97-2.76) <0.001	1.27 (1.05-1.54) 0.013	1.00 (0.80-1.25) 0.987		
PPV23	2.09 (1.77-2.48) <0.001	0.85 (0.69-1.04) 0.106	1.05 (0.82-1.35) 0.693		
PCV13	2.40 (1.50-3.84) <0.001	1.79 (1.12-2.87) 0.015	1.74 (1.06-2.85) 0.028		
Tetanus	1.00 (0.84-1.19) 0.983	0.73 (0.61-0.87) <0.001	0.84 (0.68-1.03) 0.091		

NOTE: HRs denotes Hazard ratios, and were calculated for those who had the condition as compared with those who had not the condition. In multivariable-adjusted analysis the HRs were adjusted for age (continuous years), sex, residence, comorbidities/underlying conditions and chronic medication use. Cls denote confidence intervals.

Table 3. Incidence of laboratory-confirmed COVID-19 cases according to baseline demographic and clinical characteristics (comorbidities/medications) in community-dwelling individuals (N=77,669). Tarragona region (Southern Catalonia, Spain), 01/03/2020-30/06/2020.

	Study population	LABORATORY-CONFIRMED COVID-19 CASES (n=345)		
CHARACTERISTIC	(N=77,669)	Univariate analysis	Time follow-up	Incidence rate
	n (%)	n (%) p-value	(persons-week)	IR (95% CI)
Socio-demographic				
Age: 50-64 years	42,533 (54.8)	175 (50.7) 0.004	731,553	23.9 (20.6-27.7)
65-79 years	25,712 (33.1)	108 (31.3)	442,706	24.4 (20.0-29.8)
≥80 years	9424 (12.1)	62 (18.0)	160,810	38.6 (29.7-50.2)
Sex Men	37,143 (47.8)	172 (49.9) 0.449	638,226	26.9 (23.2-31.2)
Women	40,526 (52.2)	173 (50.1)	696,843	24.8 (21.4-28.8)
Comorbidities				
Neurological disease	1950 (2.5)	14 (4.1) 0.066	33,074	42.3 (23.1-71.1)
Renal disease	4240 (5.5)	40 (11.6) <0.001	72,114	55.5 (39.6-75.5)
Cancer	6461 (8.3)	46 (13.3) 0.001	109,968	41.8 (30.5-56.0)
Rheumatic disease	860 (1.1)	6 (1.7) 0.261	14,699	40.8 (15.0-88.9)
Inflammatory bowel disease	520 (0.7)	3 (0.9) 0.648	8927	33.6 (6.9-98.1)
Respiratory disease	7074 (9.1)	61 (17.7) <0.001	120,762	50.5 (38.9-65.7)
Cardiac disease	12,923 (16.6)	90 (26.1) <0.001	221,049	40.7 (32.9-50.5)
Atrial fibrillation	3560 (4.6)	37 (10.7) <0.001	60,657	61.0 (42.5-84.8)
Liver disease	1438 (1.9)	8 (2.3) 0.519	24,570	32.6 (14.1-64.2)
Diabetes	12,925 (16.6)	68 (19.7) 0.125	221,837	30.7 (24.1-39.0)
Hypertension	33,992 (43.8)	159 (46.1) 0.384	583,619	27.2 (23.2-31.9)
Hypercholesterolemia	26,765 (34.5)	114 (33.0) 0.579	460,087	24.8 (20.7-29.8)
Obesity	21,343 (27.5)	91 (26.4) 0.646	366,948	24.8 (20.1-30.8)
Smoking	12,640 (16.3)	32 (9.3) <0.001	217,755	14.7 (9.9-21.0)
Alcoholism	1756 (2.3)	8 (2.3) 0.942	30,125	26.6 (11.5-52.4)
Chronic medication use				
Diuretics	8028 (10.3)	70 (20.3) <0.001	136,857	51.1 (40.1-64.9)
Beta blockers	9311 (12.0)	58 (16.8) 0.006	159,529	36.4 (28.0-47.3)
ACEIs	16,030 (20.6)	56 (16.2) 0.043	275,658	20.3 (15.6-26.4)
ARBs	8709 (11.2)	42 (12.2) 0.571	149,802	28.0 (20.0-38.1)
Calcium channel blockers	6315 (8.1)	35 (10.1) 0.170	108,465	32.3 (22.5-44.9)
Statins	15,910 (20.5)	66 (19.1) 0.532	273,714	24.1 (18.9-30.6)
Oral anticoagulants	3741 (4.8)	37 (10.7) <0.001	63,865	57.9(40.4-80.5)
Antiplatelet drugs	8809 (11.3)	55 (15.9) 0.007	150,896	36.4(28.0-47.3)
Insulin	2904 (3.7)	24 (7.0) 0.002	49,699	48.3 (31.0-72.0)
Oral antidiabetic drugs	10,352 (13.3)	45 (13.0) 0.876	178,039	25.3 (18.4-33.9)
Inhaled respiratory drugs	6095 (7.8)	57 (16.5) <0.001	103,830	54.9 (42.3-71.4)
Antineoplastic agents	1581 (2.0)	6 (1.7) 0.696	27,209	22.1 (8.1-48.2)
Systemic corticosteroids	1216 (1.6)	10 (2.9) 0.046	20,457	48.9 (23.5-90.0)
NSAIDs	4305 (5.5)	21 (6.1) 0.658	73,888	28.4 (17.6-43.5)
Antihistamines	3221 (4.1)	13 (3.8) 0.723	55,545	23.4 (12.4-40.0)
Proton-Pump Inhibitors	17,315 (22.3)	108 (31.3) <0.001	296,238	36.5 (29.9-44.5)
Benzodiazepines	12,654 (16.3)	74 (21.4) 0.009	216,635	34.2 (26.8-43.4)
Vaccination's history				
Flu vaccine in prior autumn	21,569 (27.8)	103 (29.9) 0.386	370,309	27.8 (22.7-33.9)
PPV23	25,222 (32.5)	142 (41.2) 0.001	432,783	32.8 (27.7-38.8)
	1115 (1.4)	13 (3.8) <0.001	18,935	68.7 (36.5-117.5)
letanus	49,817 (64.1)	231 (67.0) 0.274	856,072	27.0 (23.8-30.6)

NOTE: p-values in univariate analysis were calculated by chi-squared, or Fisher's test as appropriate, comparing percentages in the study population vs COVID-19 cases; IR denotes incidence rates per 100,000 persons-week; CIs denotes confidence intervals for incidence rates and were calculated assuming a Poisson distribution for uncommon events.

Table 4. Cox regression analyses assessing unadjusted, age & sex adjusted and multivariable-adjusted risks to suffer laboratory-confirmed COVID-19 in community-dwelling individuals (N=77,669). Tarragona region (Southern Catalonia, Spain) from 01/03/2020 to 30/06/2020.

	LABORATORY-CONFIRMED COVID-19 CASES (n=345)				
CHARACTERISTIC	Unadjusted	Age & sex adjusted	Multivariable-adjusted		
	HR (95% Cl) p-value	HR (95% CI) p-value	HR (95% CI) p-value		
Socio-demographic					
Age (continuous years)	1.02 (1.01-1.02) 0.001	1.02 (1.01-1.03) 0.001	0.99 (0.98-1.01) 0.212		
Sex: women	0.92 (0.75-1.14) 0.448	0.89 (0.72-1.11) 0.301	0.90 (0.72-1.13) 0.369		
Comorbidities					
Neurological disease	1.66 (0.97-2.83) 0.065	1.38 (0.80-2.39) 0.250	1.06 (0.60-1.86) 0.837		
Renal disease	2.29 (1.65-3.18) <0.001	2.00 (1.40-2.86) <0.001	1.50 (1.02-2.21) 0.038		
Cancer	1.71 (1.25-2.33) 0.001	1.57 (1.14-2.15) 0.005	1.45 (1.05-2.01) 0.023		
Rheumatic disease	1.59 (0.71-3.56) 0.262	1.57 (0.70-3.52) 0.276	1.50 (0.63-3.58) 0.358		
Respiratory disease	2.16 (1.64-2.84) < 0.001	2.03 (1.53-2.68) < 0.001	1.39 (0.89-2.18) 0.147		
Cardiac disease	1.78 (1.40-2.26) <0.001	1.62 (1.26-2.10) <0.001	1.32 (0.98-1.79) 0.070		
Atrial fibrillation	2.52 (1.79-3.54) <0.001	2.20 (1.53-3.17) <0.001	1.36 (0.70-2.63) 0.360		
Liver disease	1.26 (0.63-2.55) 0.512	1.28 (0.64-2.58) 0.490	1.10 (0.54-2.24) 0.796		
Diabetes	1.23 (0.95-1.61) 0.123	1.12 (0.85-1.47) 0.419	1.31 (0.81-2.13) 0.277		
Hypertension	1.10 (0.89-1.36) 0.377	0.95 (0.75-1.19) 0.631	0.98 (0.72-1.34) 0.909		
Obesity	0.95 (0.74-1.20) 0.645	0.92 (0.72-1.17) 0.473	0.83 (0.64-1.08) 0.159		
Smoking	0.53 (0.37-0.76) 0.001	0.56 (0.39-0.81) 0.002	0.51 (0.35-0.74) <0.001		
Alcoholism	1.03 (0.51-2.07) 0.939	1.03 (0.51-2.09) 0.931	1.10 (0.53-2.24) 0.804		
Chronic medication use					
Diuretics	2.22 (1.71-2.89) <0.001	2.07 (1.56-2.74) <0.001	1.54 (1.11-2.13) 0.010		
Beta blockers	1.49 (1.12-1.97) 0.006	1.36 (1.02-1.81) 0.038	1.03 (0.74-1.43) 0.869		
ACEIs	0.75 (0.56-0.99) 0.044	0.67 (0.50-0.89) 0.006	0.63 (0.45-0.89) 0.009		
ARBs	1.10 (0.80-1.52) 0.572	0.99 (0.71-1.37) 0.933	0.77 (0.52-1.13) 0.181		
Calcium channel blockers	1.28 (0.90-1.81) 0.172	1.14 (0.80-1.63) 0.477	1.07 (0.73-1.56) 0.741		
Statins	0.92 (0.70-1.20) 0.535	0.82 (0.62-1.08) 0.158	0.66 (0.49-0.90) 0.009		
Oral anticoagulants	2.39 (1.70-3.36) <0.001	2.10 (1.46-3.00) <0.001	1.33 (0.68-2.60) 0.413		
Antiplatelet drugs	1.49 (1.11-1.98) 0.007	1.31 (0.97-1.77) 0.083	1.35 (0.94-1.96) 0.106		
Insulin	1.93 (1.28-2.93) 0.002	1.77 (1.17-2.69) 0.007	1.42 (0.85-2.37) 0.179		
Oral anti-diabetic drugs	0.98 (0.71-1.34) 0.878	0.88 (0.64-1.21) 0.441	0.66 (0.40-1.09) 0.102		
Inhaled respiratory drugs	2.34 (1.76-3.11) <0.001	2.18 (1.63-2.92) <0.001	1.49 (0.93-2.37) 0.095		
Antineoplastic agents	0.85 (0.38-1.91) 0.696	0.85 (0.38-1.90) 0.685	0.52 (0.22-1.23) 0.139		
Systemic corticosteroids	1.90 (1.01-3.57) 0.045	1.76 (0.94-3.31) 0.080	1.21 (0.61-2.38) 0.592		
NSAIDs	1.11 (0.71-1.72) 0.653	1.16 (0.75-1.81) 0.509	1.20 (0.76-1.88) 0.436		
Antihistamines	0.90 (0.52-1.57) 0.721	0.91 (0.52-1.59) 0.746	0.73 (0.42-1.29) 0.282		
Proton-Pump Inhibitors	1.60 (1.27-2.00) <0.001	1.47 (1.16-1.87) 0.002	1.15 (0.86-1.52) 0.348		
Benzodiazepines	1.41 (1.09-1.82) 0.009	1.36 (1.04-1.78) 0.023	1.28 (0.97-1.68) 0.078		
Vaccination's history	1	1	1		
Flu vaccine in prior autumn	1.11 (0.88-1.40) 0.381	0.92 (0.71-1.19) 0.508	0.68 (0.51-0.92) 0.012		
PPV23	1.46 (1.18-1.81) 0.001	1.29 (0.97-1.70) 0.080	1.33 (0.95-1.88) 0.100		
PCV13	2.71 (1.56-4.72) <0.001	2.48 (1.42-4.33) 0.001	1.65 (0.92-2.98) 0.095		
Tetanus	1.13 (0.91-1.42) 0.272	1.05 (0.83-1.32) 0.703	0.99 (0.77-1.28) 0.935		

NOTE: HRs denotes Hazard ratios, and were calculated for those who had the condition as compared with those who had not the condition. In multivariable-adjusted analysis the HRs were adjusted for age (continuous years), sex, comorbidities/underlying conditions and chronic medication use. Cls denote confidence intervals.

Table 5. Univariate and multivariate analyses assessing susceptibility/risk to suffer laboratoryconfirmed Covid-19 among nursing-home residents (N=1,414). Tarragona region (Southern Catalonia, Spain), 01/03/2020-30/06/2020.

	Study population (N=1414)	LABORATORY-CONFIRMED Covid-19 CASES (n=191)		
CHARACTERISTIC		Univariate analysis	Multivariable analysis	
	n (%)	n (%) p value	HR (95% Cl) p value	
Socio-demographic				
Age: 50-64 years	151 (10.7)	2 (1.0) <0.001	Age, continuous years:	
65-79 years	301 (21.3)	29 (15.2)	1.03 (1.01-1.05) 0.001	
≥80 years	962 (68.0)	160 (83.8)		
Sex: Men	483 (34.2)	63 (33.9) 0.713	1.00 (reference) 0.001	
Women	931 (65.8)	128 (66.1)	0.78 (0.56-1.09) 0.140	
Comorbidities				
Neurological disease	367 (26.0)	65 (34.0) 0.006	1.28 (0.93-1.75) 0.127	
Renal disease	236 (16.7)	25 (13.1) 0.151	0.69 (0.44-1.08) 0.106	
Cancer	169 (12.0)	21 (11.0) 0.661	0.83 (0.51-1.35) 0.462	
Rheumatic disease	12 (0.8)	1 (0.5) 0.598	0.73 (0.10-5.42) 0.762	
Respiratory disease	198 (14.0)	19 (9.9) 0.082	0.75 (0.43-1.32) 0.315	
Cardiac disease	512 (36.2)	63 (33.0) 0.319	0.80 (0.57-1.12) 0.188	
Atrial fibrillation	226 (16.0)	33 (17.3) 0.600	1.23 (0.72-2.08) 0.447	
Liver disease	27 (1.9)	3 (16) 0.713	0.83 (0.26-2.63) 0.744	
Diabetes	392 (27.7)	58 (30.4) 0.380	1.02 (0.61-1.68) 0.955	
Hypertension	953 (67.4)	131 (68.6) 0.706	0.98 (0.68-1.41) 0.906	
Obesity	335 (23.7)	44 (23.0) 0.819	1.13 (0.79-1.61) 0.520	
Smoking	110 (7.8)	9 (4.7) 0.089	0.98 (0.47-2.02) 0.952	
Alcoholism	40 (2.8)	3 (1.6) 0.259	0.59 (0.18-1.92) 0.379	
Chronic medication use				
Diuretics	453 (32.0)	69 (36.1) 0.193	1.26 (0.91-1.75) 0.162	
Beta blockers	260 (18.4)	29 (15.2) 0.219	0.77 (0.50-1.19) 0.243	
ACEIs	389 (27.5)	58 (30.4) 0.342	0.97 (0.68-1.38) 0.867	
ARBs	160 (11.3)	12 (6.3) 0.018	0.47 (0.25-0.87) 0.017	
Calcium channel blockers	175 (12.4)	29 (15.2) 0.205	1.37 (0.89-2.10) 0.156	
Statins	224 (15.8)	25 (13.1) 0.263	0.92 (0.58-1.48) 0.739	
Oral anticoagulants	171 (12.1)	21 (11.0) 0.617	0.87 (0.46-1.65) 0.675	
Antiplatelet drugs	345 (24.4)	53 (27.7) 0.246	1.35 (0.92-1.99) 0.123	
Insulin	138 (9.8)	19 (9.9) 0.925	0.82 (0.47-1.43) 0.481	
Oral anti-diabetic drugs	233 (16.5)	40 (20.9) 0.074	1.79 (1.04-3.08) 0.037	
Inhaled respiratory drugs	198 (14.0)	20 (10.5) 0.130	0.81 (0.47-1.40) 0.447	
Antineoplastic agents	33 (2.3)	6 (3.1) 0.427	2.51 (1.05-6.01) 0.039	
Systemic corticosteroids	36 (2.5)	0 (0.0) 0.016	0.00 (-) 0.945	
NSAIDs	16 (1.1)	0 (0.0) 0.112	0.00 (-) 0.967	
Antihistamines	43 (3.0)	2 (1.0) 0.084	0.40 (0.10-1.63) 0.199	
Proton-Pump Inhibitors	616 (43.6)	75 (39.3) 0.198	0.74 (0.53-1.04) 0.083	
Benzodiazepines	392 (27.7)	52 (27.2) 0.869	0.97 (0.70-1.36) 0.879	
Vaccination's history				
Flu vaccine in prior autumn	1037 (73.3)	155 (81.2) 0.009	1.56 (1.07-2.29) 0.022	
PPV23	961 (68.0)	130 (68.1) 0.975	0.95 (0.65-1.39) 0.778	
PCV13	24 (1.7)	5 (2.6) 0.290	1.89 (0.73-4.88) 0.187	
Tetanus	982 (69.4)	113 (59.2) 0.001	0.63 (0.44-0.91) 0.013	

NOTE: p-values in univariate analysis were calculated by chi-squared, or Fisher's test as appropriate, comparing percentages in the study population vs COVID-19 cases; HRs denotes Hazard ratios, and were calculated for those who had the condition as compared with those who had not the condition, and were adjusted for age (continuous years), sex, comorbidities/underlying conditions and chronic medication use. Cls denote confidence intervals.