Impact of the COVID-19 pandemic on mental health among Greek adults: a cross-sectional survey.

Efthimios Dragotis¹, Korina Atsopardi², Anastasia Barbouni³, Konstantinos Farsalinos², Konstantinos Poulas²,

¹ Department of Business Management and Administration, University of Patras, Greece ² Laboratory of Molecular Biology and Immunology, Department of Pharmacy, University of Patras, Greece

³ Department of Public and Community Health, University of West Attica

Abstract

The COVID-19 virus characterised as a pandemic in 2020 at its outbreak around the world has been recorded to have severe effects on the global health and mental health. The current research is accomplished during the first period of the restrictive measures in Greece March – April 2020 and is aiming to evaluate the COVID-19 effect on mental health of the population. The STAI and DASS-21 questionnaires have been used in order to accomplish the aims of the study. The results (n = 527) show a moderate level on mental health status of the sample STAI-S: 45.8, STAI-T: 40.7, Depression: 4.6, Anxiety: 3.1 and Stress: 6.1. The research shows that women, younger respondents, lower income households and those living on smaller houses suffer from increased depression, anxiety and stress. Additionally, factors of the daily habits within the period of COVID-19 such as mask, gloves and use of antiseptic can drastically decrease illnesses of mental health. Such findings can be used from the Greek State in order to reduce the effects of COVID-19 in mental health of the population and protect the socially vulnerable groups.

Introduction

The coronavirus disease (COVID-19) was first reported and confirmed in Wuhan (Hubei Province, China) in late December 2019 and from then it began to spread first in China, and soon throughout all over the world. The World Health Organization (WHO) declared the COVID-19 outbreak as a Public Health Emergency of International Concern (PHEIC) on January 30, 2020 and characterized it as a pandemic on March 11, 2020. The coronavirus pandemic detected in Greece on February 26, 2020. The Government took instant measures of decongestion of the public such as the cancelation of the upcoming carnival events that compose a period of extreme crowding in many cities. In the following days, the Greek authorities imposed restrictive measures regionally and on March 23, 2020 a national lockdown was imposed as soon as the number of positive COVID-19 cases and the number of deaths started to increase. Due to the high transmissibility, Lockdown was an option for the most European countries in an effort to contain the COVID-19 outbreak. The restrictive measures disrupted social and occupational life, education, transportation and traveling, and other aspects of daily life. Moreover, restrictive measures and isolation have negative psychological effects such as confusion, anger and posttraumatic distress [1,2].

Infectious diseases, such as severe acute respiratory syndrome (SARS) [3–5], the new influenza A (HIN1) epidemic [6–8], the Middle East Respiratory Syndrome (MERS) [9,10] and Ebola virus disease (EVD) [11] have become one of the major threats to global public health recent years with impact in both physical and mental health. Epidemic diseases that are globally spread rapidly are characterized as pandemic. Pandemics usually compose a public health emergency situation that affects the community's normal functioning. Anxiety, stress, depression and post-traumatic stress are some of the psychological problems that humans can experiences through these outbreaks [3,7,10,11].

The reported symptoms of COVID-19 are primarily respiratory. Although most patients present with mild symptoms, in some cases the disease progresses to viral pneumonia,

which in severe cases can lead to acute respiratory failure and to death. The limited knowledge of the COVID-19, the overwhelming news, the increasing number of patients, suspected cases and deaths lead to anxiety and fear in the public [12,13]. In confirmed or suspected cases of COVID-19 except from the physical suffering, the patient may experience anxiety and fear due to the high contagiousness and the serious consequences of the disease [14,15]. Moreover, under restrictive measures they may also experience boredom, disappointment, irritability loneliness, denial, and depression. [1,15]. Researches in China and Turkey showed that lockdown during the COVID-19 epidemic affects the general population with the present of anxiety and depression [16–18].

Health professionals and affiliated healthcare workers in Greece and worldwide are in the frontline on the fight against COVID-19 pandemic and they are under both physical and psychological pressure [19,20]. An impact on the mental health status of university students in Greece has already been documented and shows increased anxiety and depression [21,22]. Moreover, COVID-19 pandemic and the lockdown measures had also adverse mental health impact on children and adolescents in Greece [2]. During nationwide lockdown, anxiety and depression had increased for the already stressed last year senior high school students preparing for the upcoming Panhellenic university entrance exams [23]. Patients with chronic disease have increased distress and somatization during quarantine in Greece [24].

COVID-19 outbreak and the restrictive measures raising concerns of the adverse effects, although literature that focus on the impact of the COVID-19 outbreak on mental health has not been systematically studied. The aim of the study was to investigate the effect of COVID-19 outbreak to the public mental health in Greece, during the period of the lockdown.

Methods

The study designed on April 2020 and accomplished in May 2020. The study uses quantitative methods of primary research. The procedure of the study was the online survey since from March to May 2020 the Greek State has issued laws, provisions and

Ministerial decisions that set traffic and circulation restrictions of the population. Furthermore, during this period, the university institutions' operation has been suspended and personal contacts were limited. Therefore, in the need of conducting the research at the period of the lock-down, the online distribution of the questionnaire has been used. The research conducted from May 2 to May 5, while the restrictive measures were still in charge. The final sample is consisted of 527 respondents who answered the questionnaire in more than six (00:06) and less than forty (00:40) minutes.

The questionnaire is divided into six sections of issues that the respondents are up to answer. The first section is about the demographics of the sample where the respondents are answering fourteen (14) questions: gender, age group, type of place of living, education level, citizenship, work status, if they are health professionals, if they own a type A of movement document, income level, family status, dependent members in family and m^2 of house/flat. These questions will be used as control variables in our research.

The second section of the questionnaire is about the medical history and habits of the sample. Thirteen (13) different illnesses – health status are listed where the respondents mark if they suffer from them (General medical record, Diabetes, Hypertension, Increased Cholesterol levels, Stroke, Lung diseases, Asthma, Cancer, Autoimmune illnesses, Transplant history, Immunosuppression, no problem – being healthy). Moreover, the sample is reporting the use of cigarettes, alcohol, caffeine and exercise, habits that are connected with stress, anxiety and depression among the relevant literature.

The third section contains three questions concerning the knowledge about the transmission and their personal protection measures against the covid19, in order to assess the level of medical and epidemiological knowledge of the respondents (report the most vulnerable subgroups, the ways of transmission and ways of protection). The fourth section of the questionnaire is dedicated to the symptoms, the reactions and the health status of the sample during the period of the wide spread of covid19 and the lock down period, from February to May 2020. The sample answers if from February 2020 has presented one or more symptoms that are related to covid19, among relevant literature

and medical sources (fever 370 C and above, Cough, Headache, Runny nose, Sore throat, Shortness of breath, Pain in joints and muscles, felt tired, Immune, Tasteleness, Gastrointestinal disorders and if they were full healthy), what they did with the abovementioned symptoms and state their future decisions in case they will present suspicious symptoms related with the covid19.

The fifth section of the questionnaire is concerning the Greek authorities, their decisions and the respondents' agreement to the policies that implied in Greece from February 2020. The section contains three questions where the sample is called to report its agreement to the restrictive measures, its satisfaction from authorities and its main sources of information. The final section is scheduled to measure mental health status of the respondents. Stress on work environment measured using STAI questionnaire (State-Trait Anxiety Inventory), introduced by Spielberger [25]. This part of the questionnaire consists of 40 statements to which the respondent is asked to answer. Through them, we examine transient and permanent stress, measuring the employee's emotional state at the time of completing the questionnaire and how they generally feel. The evaluation is reported in a 4 items scale, at the first 20 questions (transient stress) the answers range from "Not at all" to "Too much" while for the next 20 questions (permanent stress) the answers range from "Almost never" to "Almost always.". Furthermore, the research also uses the Depression Anxiety and Stress scale (DASS-21) [26] in order to measure these illnesses on the respondents. The questions are answered in a 4 items Likert scale where range varies from 0 – "does not apply for me" to 3- "does apply for me very much, most of the times". The STAI and DAS-21 research tools have previously used in research related to the COVID-19 pandemic [27-32].

Results

General beliefs about COVID-19

As seen on table 1, the most perceived vulnerable groups are the people with health problems and elderly people, while the least vulnerable are children and teenagers. The sample mostly believes that the virus is spread easier from "the drops from the mouth and nose when someone coughs or sneezes" and least from mosquitos or other insect bites.

Tuble 1 – General beliefs ubbal COVID-19 spread						
Question	Statement	N	Mean	Std. Deviation		
Vulnerability	Children	511	3,62	0,900		
	Teenagers	506	3,50	0,776		
	Elderly people	520	1,38	0,637		
	People with health problems	525	1,28	0,556		
	Pregnant women	516	2,68	0,961		
Way of transmission	Mosquitos or other insect bites	511	4,72	0,726		
	Sexual contacts	513	3,25	1,560		
	Infected animals	506	4,01	1,245		
	Contaminated objects	521	2,10	1,016		
	Drops from the mouth and nose when	526	1,15	0,481		
	Unhealthy cooking or poor cooking	509	3,74	1,333		
	Raw vegetables - fruits	514	3,78	1,212		

Table 1 – General beliefs about COVID-19 spread

General habits of the population

Table 2 results show that the daily practices of sample were mostly the covering of the mouth in case of cough / sneeze, washing hands with soap and water, and washing hands after contacting contaminated objects. An interesting finding is the habits that the respondents do less are wearing gloves and masks.

Table 2 – ways of protection from COVID-19					
Statement	N	Mean	Std. Deviation		
Cover my mouth when you cough or sneeze	525	1,11	0,382		
Avoid sharing utensils	518	1,93	1,115		
Wash your hands with soap and water	522	1,20	0,467		
Wash hands immediately after coughing, sneezing, or wiping nose.	524	2,00	1,059		
Wash hands after contact with contaminated objects	525	1,31	0,674		
Wash the products you buy	522	2,36	1,329		
You used a hand antiseptic	522	1,88	1,009		

Table 2 – Ways of protection from COVID-19

Wear a face mask when you go out regardless of the	524	3,35	1,353
presence or absence of symptoms			
Wear gloves when going out regardless symptoms	522	3,49	1,403

The 47,4% declared having no symptoms related to COVID-19 from period of February to May 2020. The most common indication of illness among the sample was the headaches that 26,8% of the respondents declare having (Table 3). The 37,2% of the sample declared that had done nothing from the above, taking no action against these symptoms and 9,3% reported that addressed to a doctor by phone. The 75,3% declared being health and the group that had some symptoms of the COVID-19 was the 22% of the sample.

Table 3 – COVID-19 symptoms		
Statement	N	Mean
Fever 37° C and above	67	12,7%
Cough	85	16,1%
Headache	150	28,5%
Runny nose	116	22%
Sore throat	102	19,4%
Shortness of breath	29	5,5%
Pain in joints and muscles	51	9,7%
Feeling tired	95	18 %
Immune	12	2,3%
Tasteleness	13	2,5%
Gastrointestinal disorders	50	9,5%
None of the above	246	46,7%

Table 4 – Reaction on COVID-19 symptoms				
Statement	N	Percent		
To a private doctor (family or other) by phone	49	16.8%		
To a private doctor (family or other) with a visit	23	7.9%		
In a private hospital	1	0,4%		
In a public hospital	8	2.7%		
By phone at NPHO (1135)	7	2.4%		
I made a SARS-COV-2 test	8	2.7%		
Nothing from the above	196	67,1%		

Table 5 – Health status			
Statement	Ν	Percent	
Full healthy	397	75,3%	

Had some symptoms of illness	117	22,2%
Had symptoms and I went to hospital	3	0,6%
Had symptoms and did the test	4	0,8%
Did the test and I was positive	2	0,4%

According to the mean scores of the following table, the sample is recording more positive opinions for calling doctors by phone and calling the National Public Health Organization than other options. The least positive opinion is recorded about going to a private hospital.

Statement	N	Mean	Std. Deviation	
To a private doctor (family or other) by phone	509	1,96	1,27	
To a private doctor (family or other) with a visit	499	3,19	1,32	
In a private hospital	492	4,21	1,02	
In a public hospital	499	3,02	1,40	
To a pharmacist	490	3,91	1,22	
By phone at NPHO (1135)	507	2,63	1,44	
In a family and friendly environment	499	3,25	1,59	
To anyone / I will deal with it alone	494	4,14	1,17	

_ . . . _

As table 7 shows, the most popular mean of information was the doctors' advice and the daily briefing at 18.00 that was accomplished by the Civil Protection, the National Public Health Organization and information through the Internet websites.

Table 7 = 1 requercy of using sources of information about COVID-17.					
Source of Information	N	Mean	Std. Deviation		
Newspapers and magazines	519	3,43	1,123		
The news on TV	517	3,56	1,213		
Online information (websites)	519	2,66	1,021		
Social media (e.g. Facebook)	520	3,62	1,070		
Daily government briefing at 6 p.m.	523	2,65	1,320		
Advice from Doctors	522	2,01	0,895		
Tips from family / friends	516	3,89	0,964		

Table 7 – Frequency of using sources of information about COVID-19

Public satisfaction

The sample reports highest level of agreement to the cancellation of the carnival events and the restrictive measure in churches. The higher satisfaction levels are reported for the response of the National Health System and the responsibility in the peoples' behavior. The lowest satisfaction levels are reported for the University reactivity in applying the restrictive measures within their daily life.

N	Mean	Std. Deviation
522	2,55	1,336
523	2,11	1,303
523	2,24	1,318
524	1,75	1,116
524	1,75	1,158
	N 522 523 523 524 524	N Mean 522 2,55 523 2,11 523 2,24 524 1,75 524 1,75

Table 8 – Agreement in specific decisions

5			
Statement	N	Mean	Std. Deviation
Response of the National Health System	521	2,31	1,128
Daily briefing at 18.00 from Government	520	2,58	1,324
University reactivity (distance learning platforms)	515	2,75	1,158
Remote working and work from home system	513	2,67	1,172
Government readiness regarding the scope of measures taken	520	2,70	1,268
Responsibility of peoples' behavior	520	2,42	0,938

Table 9 – Satisfaction

Anxiety, Stress and Depression

Measuring the Depression, Anxiety and Stress levels, Table 10 shows moderate level of mental health status. Normality tests performed and showed that all the six scales do now follow the normal distribution. Reliability analysis performed and the Cronbach's alpha scores were higher than 0,8 for the DAS-21 scales and higher than 0,9 for the STAI scales. The table 11 presents high and significant (a = 99%) correlation levels between variables.

Table 10 – Anxiety, Stress and Depression

	N	Min	Max	Mean	Std. Deviation
STATE	511	2,00	77,00	45,8552	12,53679
TRAIT	510	1,00	75,00	40,7471	10,41651
TOTAL	510	5,00	151,00	86,6333	20,46668
Depression	496	0,00	21,00	4,6472	4,42793
Anxiety	495	0,00	21,00	3,1192	3,71831
Stress	495	0,00	21,00	6,0848	4,66150

	State	Trait	Total	Depression	Anxiety	Stress
State	1	0,567**	0,909**	0,517**	0,487**	0,604**
Trait		1	,0849**	0,623**	0,592**	0,591**
Total			1	0,638**	0,602**	0,672**
Depression				1	0,686**	0,791**
Anxiety					1	0,723**
Stress						1

Table 11 – Correlation between scales

Depression, Anxiety and Stress non-related with COVID-19.

Demographics and medical history were tested in order to report their relations with mental health. The results (table 12) indicate that women present significantly higher mean scores on stress, depression and anxiety, the 18-24 years old group (the younger sample group) presents significantly higher mean scores on anxiety and stress and the group that works at the private sector present higher stress levels. Moreover, people with two dependent members in their family have significantly higher trait stress and depression than the other groups of the sample. Furthermore, the sub-groups with income up to 500 euros and living at an apartment up to 50m² present significantly higher anxiety symptoms. As for the medical history, results indicate that the mental health issue are presented significantly higher in only four cases: people with asthma had higher state anxiety, people with cancer had higher trait anxiety, people with immunosuppression had higher anxiety and people with hypertension had lower stress.

	State anxiety		Trait Anxiety		Total A	Total Anxiety		Depression		Anxiety		ress
	В	ť	В	ť	В	ť	в	t	В	ť	В	t
Gender												
Male	-2,16	-1,73	-1,06	-1,03	-3,07	-1,53	-0,99	-2,20*	-0,81	-2,11*	-1,37	-2,88**
Female							Reference					
Age group												
18-24	7,02	2,03*	6,16	2,16*	11,33	2,03*	2,35	1,85	1,51	1,38	2,68	1,99*
25-34	4,67	1,76	6,05	2,78*	9,04	2,12*	1,56	1,62	0,47	0,58	0,86	0,84
35-44	1,74	0,73	4,55	2,34*	4,51	1,19	1,35	1,56	0,4	0,54	1,46	1,59
45-54	0,34	0,15	1,29	0,67	0,72	0,19	1,03	1,21	0,37	0,51	0,96	1,07
55+							Reference					
Living at												
City	-2,21	-0,86	-1,68	-0,81	-3,05	-0,75	-0,25	-0,28	-0,37	-0,47	0,05	0,05
Town	-2,28	-0,79	-2,38	-1,01	-4,86	-1,05	-0,23	-0,22	-0,00	-0,00	-0,25	-0,23
Village							Reference					
Education												
High-School	-2,66	-1,37	0,68	0,43	-1,89	-0,60	0,21	0,29	-0,13	-0,21	-0,62	-0,84
Institute	-2,46	-1,02	-0,2	-0,11	-2,00	-0,53	.028	0.33	0,01	0,01	-0,70	-0,78
BSc	-1,90	-1,25	-0,496	-0,40	-1,98	-0,82	-0,21	-0,39	-0,13	-0,29	-0,83	-1,45
MSc / PhD	,	,	,	, ,	,	·	Reference	,	,	,	,	,
Citizenship												
Greek	-2.20	-0.46	3.32	0.84	1.69	0.22	-0.20	-0.12	-1.14	-0.78	-0.28	-0.16
Non-Greek	, -	-, -	- ,-	- / -	,	- /	Reference	- ,	,	- ,	- , -	- , -
Working at												
Public Sector	3 28	1.08	0.21	0.08	3.88	0 79	0.95	0.86	1 43	1.51	1.62	1 37
Private Sector	0.81	0.30	0.40	0.18	2.88	0,77	1 48	1 54	1,19	1 44	2 21	2 16*
Freelancer	0.022	0.08	-0.24	-0.10	3.18	0,07	1,10	1	1,17	1 33	1.26	1 16
Retired	-1.26	-0.33	-2 144	-0.69	-4 17	-0.68	0.89	0.64	0.63	0.53	1,20	0.77
Unemployed	-0.66	-0.23	1 12	0.48	1 78	0.38	1.0	1.00	0,89	1.01	1,12	1 41
Student	0,00	0,23	1,12	0,40	1,70	0,50	Reference	1,00	0,09	1,01	1,55	1,41
Hoalth professional							Reference					
Health worker	0.53	0.30	0.56	0.39	0.95	0.34	0.39	0.63	0.29	0.55	-0.06	-0.09
Non health	0,55	0,50	0,50	0,37	0,75	0,54	Deference	0,05	0,27	0,55	-0,00	-0,07
Moving with document							Reference					
	1 1 9	0.83	82	0.71	2 00	1.26	0.68	1 22	0.21	0.71	0.19	0.24
Non Type A	-1,10	-0,85	-,05	-0,71	-2,00	-1,20	-0,08 Poference	-1,55	0,31	0,71	-0,18	-0,34
Income							Reference					
Income No income	2 27	1.50	1.22	0.67	6 20	1 70	1.50	1 97	1.20	1 72	1.25	1.57
In to 500	5,57	1,50	1,25	0,07	0,39	1,79	1,52	1,87	1,20	1,75	1,55	1,37
Up to 500	3,04	2,41*	4,55	2,20*	7.44	2 20*	0,72	0,84	1,72	2,50*	0,77	0,80
Up to 800	4,51	2,20*	1,62	0,99	7,44	2,32*	0,9	1,28	0,72	1,17	0,14	0,19
Op to 1000	0,39	0,23	0,32	0,23	1,15	0,41	-0,060 Defense	-0,09	0,58	1,09	-0,07	-0,11
Over 1000							Reference					
Living status	1.24	0.20	2.01	1.01	2.62	0.25	1.25	0.76	0.01	0.01	0.15	0.09
Married	1,34	0,29	-3,81	-1,01	-2,62	-0,35	-1,25	-0,76	0,01	0,01	0,15	0,08
Unmarried	-1,95	-0,40	-5,39	-1,34	-7,00	-0,89	-0,57	-0,32	0,23	0,15	0,41	0,22
Divorced	1,8	0,36	-2,75	-0,66	-0,63	-0,08	0,23	0,13	0,88	0,57	1,33	0,70
Widowed							Reference					
Dependent members	1.02	0.65	0.40	0.01	2.10	0.40	1.6	1.57	1.01	1.51	1.50	1.40
0 members	-1,83	-0,65	0,48	0,21	-2,18	-0,48	-1,6	-1,57	-1,31	-1,51	-1,50	-1,40
1 member	-1,58	-0,57	,010	0,04	-1,85	-0,41	-0,90	-0,91	-1,14	-1,35	-1,70	-1,63
2 members	-1,11	-0,41	-1,77	-0,79	-2,96	-0,67	-2,02	-2,06*	-1,19	-1,43	-2,39	-2,31*
3+ members							Reference					
m ² of house						<u> </u>						
Up to 50	-0,90	-0,49	1,55	1,04*	1,58	0,54*	0,0	0,14	0,33	0,59	0,21	0,30
50 - 100	-0,59	-0,43	-0,33	-0,30	-0,20	-0,09	-0,090	-0,18	0,57	1,37	-0,05	-0,10
Over 100							Reference					

Table 12 - Association between demographics and the mental health status

Depression, Anxiety and Stress related with COVID-19.

The effect of different protection ways during the spread of COVID-19 on the three stress, depression and anxiety levels are reported on Table 11. The correlations indicate that there is a significant decrease on anxiety levels with the use of mask (trait anxiety, total anxiety and anxiety), with the use of antiseptic (state anxiety and anxiety) and with the use of gloves (anxiety). People who use more antiseptic, wearing masks and gloves more often, tend having lower levels of anxiety. Moreover, a significant higher depression levels are evaluated as the people share more often their utensils. The following table summarizes the correlations that come up in a statistically significant level.

	Table 13 – Correlation between scales									
	Cover mouth on Sneeze	Sharing utensils	Hand washing	Washing after cough	Washing after contaminated	Products washing	Antiseptic	Mask	Gloves	
State.anx							-0,09			
Trait.anx								-0,08		
Total.anx								-0,08		
Depression		0,09								
Anxiety Stress							-0,11	-0,15	-0,12	

Furthermore, in order to evaluate the effect of COVID-19 potential symptoms on mental health levels, twelve symptoms – conditions were tested. The results (table 14) indicate those who presented each symptom had significantly higher levels of stress, depression and anxiety than those who did not. Specifically, people who had fever 37o C and above, cough, headache, runny nose, sore throat, pain in joints and muscles and felt tired had significantly higher mean scores of depression, stress and anxiety. Moreover, the sample who had gastrointestinal disorders and immune presented only higher levels of anxiety. Furthermore, the sample who had shortness of breath and tasteleness presented significantly higher mean scores of depression. The same time, the sample that declared to be healthy presented significantly lower mean scores for depression, stress, and anxiety.

	State anxiety		Trait Anxiety To		Total	otal Anxiety De		Depression		Anxiety		Stress	
	В	t	В	t	В	t	В	t	В	t	В	t	
Fever 37° C plus													
No	-4,76	-2,9**	-3,75	-2,75**	-8,48	-3,17**	-1,5	-2,54*	-9,94	-2*	-1,54	-2,47*	
Yes						Re							
Cough													
No	-0,86	-0,57	-1,86	-1,49	-2,68	-1,09	-1,05	-1,97*	-1,1	-2,6**	-1,26	-2,24*	
Yes						Re	ference						
Headache													
No	-4,96	-4,1**	-4,1	-4,07**	-9,03	-4,58**	-2,3	-5,35*	-1,45	-3,97**	-2,19	-4,83**	
Yes						Re	ference						
Runny nose													
No	-3,02	-2,2*	-2,87	-2,6**	-5,86	-2,69**	-1,8	-3,74**	-1,22	-3**	-1,97	-3,91**	
Yes						Re	ference						
Sore throat													
No	-4,85	-3,5**	-3,62	-3,12**	-8,43	-3,71**	-1,63	3,27**	-1,41	-3,37**	-1,84	-3,49**	
Yes						Re	ference						
Shortness of breath													
No	-6,7	-2,76**	-3,78	-1,87	-10,4	-2,64**	-1,55	-1,77	-2,61	-3,6**	-2,2	-2,4*	
Yes						Re	ference						
Pain in joints and muscles													
No	-0,83	-0,45	-3,55	-2,32*	-4,35	-1,44	-1,43	-2,13*	-1,55	-2,76**	-1,6	-2,2*	
Yes						Re	ference						
Feeling tired													
No	-3,72	-2,61**	-4,86	-4,15**	-8,55	-3,7**	-2,38	-4,76**	-2,2	-5,23**	-2,64	-5**	
Yes						Re	ference						
Immune													
No	-1,85	-0,5	-2,82	-0,93	-4,64	-0,77	-1,05	-0,81	-3,38	-3,13**	-1,87	-1,38	
Yes						Re	ference						
Tasteleness													
No	-6,55	-1,79	-7,17	-2,36*	-13,7	-2,3*	-1,94	1,44	-5,28	-4,75**	-3,17	-2,24*	
Yes						Re	ference						
Gastrointestinal disorders													
No	-4,57	-2,46*	-3,43	-2,22*	-7,97	-2,63**	-2,11	-3,2**	-2,68	-4,89**	-1,97	-2,82	
Yes						Re	ference						
Full healthy													
No	4,28	3,91**	4,51	5**	8,86	5**	2,5	6,5**	1,94	5,98**	2,76	6,88**	
Yes						Re	ference						

Table 24 - Association between symptoms related to COVID-19 and the mental health status

The sample was consisted of 527 respondents, *p < 0,05, **p < 0,01

Discussion

The results show that the sample shows sufficient level of knowledge about the virus in terms of vulnerability and transmission ways. The people with health problems and the elderly people face increased threat in case of coronavirus contamination. Recent studies and clinical data have shown that the mortality levels are increased for people 55+ and people who suffer from an underlying disease. The sample seems to be well informed about the ways that the virus is transmitted (table 1) and their most common practices is covering mouth when cough or sneeze and wash their hands with soap and water and the use of a face mask in public places.

The symptom-related findings at table 3-5 are showing that the sample did not perceive symptoms as serious threats. The majority of the sample (54,3%) reported that from February to May have presented one or more symptoms related to COVID-19 virus. However, the majority of them (67%) did nothing about the specific symptoms in terms of visiting or calling a doctor, hospitals, calling the NPHO or making a COVID-19 test. As result, the 75,3% of the sample considered themselves as totally healthy for the specific period as not facing any serious symptoms related to COVID-19. Consequently, and given the low diagnostic test numbers that has been conducted in Greece during this period we still not know the real number of people who were positive on COVID-19.

Considering the work anxiety levels (STAI), the sample reports a STAI-S score of 45,85 and a STAI-T score of 40,74. Relevant research have evaluated either higher [30] or lower [33] STAI-S and STAI-T scores. Our study reveals medium anxiety levels (STAI-S: 32,7% low, 54,2% moderate, 12,3% severe – STAI-T: 47,2% low, 47,9% moderate, 4,1% severe) in similar levels with the relevant literature. Moderate to severe levels are reported to the 66,5% (STAI-S) and 52% (STAI-T) of the respondents which is a respectively high percentage within the relevant literature [34].

Considering the rest of mental health illnesses (DASS-21) the levels of Depression was 4,65, Anxiety 3,12 and Stress 6,08. These results represent a lower level of mental health illnesses compared to various researches within the relevant literature in different places in the same research period: According to Odriozola-Gonzáleza et al. (2020) [35] the levels of Depression is 5,52, Anxiety 3,34 and Stress 6,81 in Spain. According to Wang et al., (2020) [29] there are significant differences between DASS-21 scores in Europe (Poland) and Asia (China) but these two cases are still higher than Greece (In China Depression:

6,25, Anxiety: 6,16 and Stress: 7,76 while in Poland Depression: 10,06, Anxiety: 7,65 and Stress: 14,00). According to Verma et al. (2020) [36] the level of Depression is 8,39, Anxiety 6,53 and Stress 8,83 in India. Although the above findings signify a higher level of mental health illnesses than our results, our sample has a relatively high score in severe and extremely severe das scales. The scores of the sample are classified in five categories: normal, mild, moderate, severe and extremely severe. The majority of our sample (60-65%) demonstrate normal DASS-21 levels, however these scores are relatively lower than the current literature. The mild and moderate levels are between 20-24% of the sample which is higher than the relevant literature. Finally, the severe and extremely severe levels are between 11,7-15% which is significantly higher than the relevant literature [27,28,32,37–39].

Level / Mental health illness	Depression	Anxiety	Stress
Normal	60%	66,5%	64,8%
Mild	11,5%	13,1%	12,4%
Moderate	12,5%	6,7%	11,1%
Severe	10%	6,7%	9,3%
Extremely Severe	5%	6,7%	2,4%

Table 15 – Levels of mental health illnesses

Stress, Anxiety and Depression are strongly connected. The results of the study suggest a high and significant correlation between variables at table 11: work stress, stress, anxiety and depression are highly linked. People who suffer from a mental health illness have increased potential to suffer from others too.

The research aimed to assess potential significant differences between sub-groups of the sample in order to evaluate the factors that have a significant role in mental health status. The results show that these factors are the gender, the age, the income level and the place of living size. The female subgroup has significantly higher Depression, Stress and Anxiety levels, younger respondents have significantly higher State and Trait Anxiety levels, low-income workers have significantly lower Anxiety, State and Trait Anxiety levels and sample that lives in a place less than 50m² has significantly higher Trait Anxiety scores. The findings are similar to the relevant literature within the period of the COVID-19 outbreak in Europe: the female subgroups are reported having a significantly higher anxiety, stress and depression level [18,38–46], Another group of increased Depression, Anxiety or Stress levels within the relevant literature is the younger sub-group [27,28,30,33,40,41,46] and our findings confirm this significant difference. A third factor that is

strongly linked with higher Depression, Anxiety or Stress levels is related with the income. Our findings support the relevant literature conclusions that low-income and unemployed people have higher levels of DASS during the current period [30,33,34,39,43,46]. Some researchers also point out the significantly decreased Depression, Anxiety or Stress levels on singles / unmarried groups [32,38,43] however our research does not evaluate significant differences within different family status sub-groups. Furthermore, our study did not reveal significant differences in Depression, Anxiety or Stress (normal, state or trait) levels between healthcare professionals and other community sample in contrast to relevant literature findings that state healthcare workers are suffering from higher Depression Anxiety and Stress levels [33,39,43].

The results of our research show that the link between some specific measures of protection and mental health illnesses is strong. The use of mask, antiseptic and gloves as way of protection found to be vital the mental health of the sample. The use of each of them decreases the levels of specific mental health illnesses. These results are supported from the current literature findings. Within the relevant literature, the mask and the gloves, as way of protection, are reported to generate significantly lower scores of mental health illnesses [27,28,38,46]. Moreover, people who wash their hands more often, specifically after coughing or sneezing, are reporting lower Depression, Anxiety or Stress levels [32,38,39,46].

Another significant finding is that the specific symptoms of COVID-19 can create serious mental health issues as people who suffered from the above-mentioned symptoms had a significantly higher level of Depression, Anxiety, Stress, State and Trait Anxiety (Table 14). The results of our research support the relevant literature findings that disclose a significant relation between mental health illnesses and symptoms of COVID-19: the people who had once or more symptoms related with COVID-19 had significant higher levels of Depression, Anxiety and Stress [27–29,32,38,39,46]. During the pandemic outbreak, the people who had any symptom of illness that can be related to COVID-19 symptoms suspected that were positive to the virus. The suspicion of being ill to a severe virus such as COVID-19 constitutes a Depression, Anxiety and Stress increase factor.

Conclusions

A prevalent finding is the intense effect of COVID-19 and the restrictive measures in the social environment in mental health of specific groups. Relevant literature findings and the current study results

show adverse mental health impact during the COVID-19 outbreak and the period of the lockdown. However, the impact on specific sub-groups is significantly higher: the females, the sample living in a limited house or flat (under $50m^2$), the low-income and the younger population experience increased levels of Depression, Anxiety and Stress.

Another conclusion is based on the results of DASS-21 and specifically on the mean scores and the percentages of severe and extremely severe mental health illnesses that are increased. That means that people who have these mental health issues experience them more intensively.

Lastly, the results suggest that there are factors with a moderating role at the relation between COVID-19 outbreak, restrictive measures and mental health status. Such factors are the use of a mask for protection in public places, use of gloves and antiseptic that are significantly decreasing the mental health illnesses such as Anxiety and Stress. The results of our study show that except from the commonly accepted medical significance (protecting from the spread), these factors have a mental health significance too.

Based on the above-mentioned results and the relevant findings of the current literature, the authorities have to focus on the most vulnerable groups and take specific measures for their medical and mental health status. The COVID-19 outbreak demands enactment of targeted care and reinforcing the public health of the population, focusing on every sub-group and specifically the most vulnerable, in contrast of taking just horizontal measures that affect commonly the whole population. The restrictive measures, specifically the strictest, the prolonged and the recurrent ones, might lead to serious mental health problems.

References

- Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. Lancet 2020;395:912–20. doi:10.1016/S0140-6736(20)30460-8.
- [2] Magklara K, Lazaratou H, Barbouni A, Poulas K, Farsalinos K, Greece -Coronavirus, et al. Impact of COVID-19 pandemic and lockdown measures on mental health of children and adolescents in

Greece. MedRxiv Prepr 2020. doi:10.1101/2020.10.18.20214643.

- [3] Ko CH, Yen CF, Yen JY, Yang MJ. Psychosocial impact among the public of the severe acute respiratory syndrome epidemic in Taiwan. Psychiatry Clin Neurosci 2006;60:397–403. doi:10.1111/j.1440-1819.2006.01522.x.
- [4] Peng EYC, Lee MB, Tsai ST, Yang CC, Morisky DE, Tsai LT, et al. Population-based post-crisis psychological distress: An example from the SARS outbreak in Taiwan. J Formos Med Assoc 2010;109:524–32. doi:10.1016/S0929-6646(10)60087-3.
- [5] Mak IWC, Chu CM, Pan PC, Yiu MGC, Ho SC, Chan VL. Risk factors for chronic post-traumatic stress disorder (PTSD) in SARS survivors. Gen Hosp Psychiatry 2010;32:590–8. doi:10.1016/j.genhosppsych.2010.07.007.
- [6] Yeung NCY, Lau JTF, Choi KC, Griffiths S. Population responses during the pandemic phase of the influenza A(H1N1)pdm09 epidemic, Hong Kong, China. Emerg Infect Dis 2017;23:813–5. doi:10.3201/eid2305.160768.
- [7] Taha SA, Matheson K, Anisman H. H1N1 Was Not All That Scary: Uncertainty and stressor appraisals predict anxiety related to a coming viral threat. Stress Heal 2014;30:149–57. doi:10.1002/smi.2505.
- [8] Lau JTF, Griffiths S, Choi KC, Tsui HY. Avoidance behaviors and negative psychological responses in the general population in the initial stage of the H1N1 pandemic in Hong Kong. BMC Infect Dis 2010;10. doi:10.1186/1471-2334-10-139.
- [9] Jeong H, Yim HW, Song YJ, Ki M, Min JA, Cho J, et al. Mental health status of people isolated due to Middle East Respiratory Syndrome. Epidemiol Health 2016;38:e2016048. doi:10.4178/epih.e2016048.
- [10] Lee SM, Kang WS, Cho AR, Kim T, Park JK. Psychological impact of the 2015 MERS outbreak on hospital workers and quarantined hemodialysis patients. Compr Psychiatry 2018;87:123–7. doi:10.1016/j.comppsych.2018.10.003.
- [11] Shultz JM, Baingana F, Neria Y. The 2014 Ebola outbreak and mental health: Current status and recommended response. JAMA - J Am Med Assoc 2015;313:567–8. doi:10.1001/jama.2014.17934.

- Shigemura J, Ursano RJ, Morganstein JC, Kurosawa M, Benedek DM. Public responses to the novel 2019 coronavirus (2019-nCoV) in Japan: Mental health consequences and target populations.
 Psychiatry Clin Neurosci 2020;74:281–2. doi:10.1111/pcn.12988.
- [13] Bao Y, Sun Y, Meng S, Shi J, Lu L. 2019-nCoV epidemic: address mental health care to empower society. Lancet 2020;395:e37–8. doi:10.1016/S0140-6736(20)30309-3.
- [14] Xiang YT, Yang Y, Li W, Zhang L, Zhang Q, Cheung T, et al. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. The Lancet Psychiatry 2020;7:228–9. doi:10.1016/S2215-0366(20)30046-8.
- [15] Li W, Yang Y, Liu ZH, Zhao YJ, Zhang Q, Zhang L, et al. Progression of mental health services during the COVID-19 outbreak in China. Int J Biol Sci 2020;16:1732–8. doi:10.7150/ijbs.45120.
- [16] Lei L, Huang X, Zhang S, Yang J, Yang L, Xu M. Comparison of Prevalence and Associated Factors of Anxiety and Depression among People Affected by versus People Unaffected by Quarantine during the COVID-19 Epidemic in Southwestern China. Med Sci Monit 2020;26:e924609-1. doi:10.12659/MSM.924609.
- [17] Li J, Yang Z, Qiu H, Wang Y, Jian L, Ji J, et al. Anxiety and depression among general population in China at the peak of the COVID-19 epidemic. World Psychiatry 2020;19:249–50. doi:10.1002/wps.20758.
- [18] Özdin S, Bayrak Özdin Ş. Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish society: The importance of gender. Int J Soc Psychiatry 2020;66:504–11. doi:10.1177/0020764020927051.
- [19] Walton M, Murray E, Christian MD. Mental health care for medical staff and affiliated healthcare workers during the COVID-19 pandemic. Eur Hear J Acute Cardiovasc Care 2020;9:241–7. doi:10.1177/2048872620922795.
- [20] Tsamakis K, Rizos E, Manolis A, Chaidou S, Kympouropoulos S, Spartalis E, et al. [Comment] COVID-19 pandemic and its impact on mental health of healthcare professionals. Exp Ther Med 2020;19:3451. doi:10.3892/etm.2020.8646.
- [21] Kaparounaki CK, Patsali ME, Mousa DP V., Papadopoulou EVK, Papadopoulou KKK, Fountoulakis

KN. University students' mental health amidst the COVID-19 quarantine in Greece. Psychiatry Res 2020;290:113111. doi:10.1016/j.psychres.2020.113111.

- [22] Patsali ME, Mousa DP V., Papadopoulou EVK, Papadopoulou KKK, Kaparounaki CK, Diakogiannis I, et al. University students' changes in mental health status and determinants of behavior during the COVID-19 lockdown in Greece. Psychiatry Res 2020;292. doi:10.1016/j.psychres.2020.113298.
- [23] Giannopoulou I, Efstathiou V, Triantafyllou G, Korkoliakou P, Douzenis A. Adding stress to the stressed: Senior high school students' mental health amidst the COVID-19 nationwide lockdown in Greece. Psychiatry Res 2021;295:113560. doi:10.1016/j.psychres.2020.113560.
- [24] Louvardi M, Pelekasis P, Chrousos GP, Darviri C. Mental health in chronic disease patients during the COVID-19 quarantine in Greece. Palliat Support Care 2020;18:394–9. doi:10.1017/S1478951520000528.
- [25] Spielberger CD, Gorsuch RL, Lushene RE, Vagg PR, Jacobs GA. Manual for the State-Trait Anxiety Inventory. Palo Alto, CA Consult Psychol Press 1983.
- [26] Norton PJ. Depression Anxiety and Stress Scales (DASS-21): Psychometric analysis across four racial groups. Anxiety, Stress Coping 2007;20:253–65. doi:10.1080/10615800701309279.
- [27] Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. Int J Environ Res Public Health 2020;17. doi:10.3390/ijerph17051729.
- [28] Wang C, Pan R, Wan X, Tan Y, Xu L, McIntyre RS, et al. A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. Brain Behav Immun 2020;87:40–8. doi:10.1016/j.bbi.2020.04.028.
- [29] Wang C, Chudzicka-Czupała A, Grabowski D, Pan R, Adamus K, Wan X, et al. The Association Between Physical and Mental Health and Face Mask Use During the COVID-19 Pandemic: A Comparison of Two Countries With Different Views and Practices. Front Psychiatry 2020;11. doi:10.3389/fpsyt.2020.569981.

- [30] Lin Y, Hu Z, Alias H, Wong LP. Knowledge, Attitudes, Impact, and Anxiety Regarding COVID-19 Infection Among the Public in China. Front Public Heal 2020;8. doi:10.3389/fpubh.2020.00236.
- [31] Tan BYQ, Chew NWS, Lee GKH, Jing M, Goh Y, Yeo LLL, et al. Psychological Impact of the COVID-19 Pandemic on Health Care Workers in Singapore. Ann Intern Med 2020;173:317–20. doi:10.7326/M20-1083.
- [32] Tan W, Hao F, McIntyre RS, Jiang L, Jiang X, Zhang L, et al. Is returning to work during the COVID-19 pandemic stressful? A study on immediate mental health status and psychoneuroimmunity prevention measures of Chinese workforce. Brain Behav Immun 2020;87:84– 92. doi:10.1016/j.bbi.2020.04.055.
- [33] Chen L, Zhao H, Razin D, Song T, Wu Y, Ma X, et al. Anxiety levels during a second local COVID-19 pandemic breakout among quarantined people: A cross sectional survey in China. J Psychiatr Res 2021;135:37–46. doi:10.1016/j.jpsychires.2020.12.067.
- [34] Wong LP, Hung CC, Alias H, Lee TSH. Anxiety symptoms and preventive measures during the COVID-19 outbreak in Taiwan. BMC Psychiatry 2020;20. doi:10.1186/s12888-020-02786-8.
- [35] Odriozola-González P, Planchuelo-Gómez Á, Irurtia MJ, de Luis-García R. Psychological effects of the COVID-19 outbreak and lockdown among students and workers of a Spanish university. Psychiatry Res 2020;290. doi:10.1016/j.psychres.2020.113108.
- [36] Verma S, Mishra A. Depression, anxiety, and stress and socio-demographic correlates among general Indian public during COVID-19. Int J Soc Psychiatry 2020;66:756–62. doi:10.1177/0020764020934508.
- [37] Zhang Y, Ma ZF. Impact of the COVID-19 pandemic on mental health and quality of life among local residents in Liaoning Province, China: A cross-sectional study. Int J Environ Res Public Health 2020;17. doi:10.3390/ijerph17072381.
- [38] Tee ML, Tee CA, Anlacan JP, Aligam KJG, Reyes PWC, Kuruchittham V, et al. Psychological impact of COVID-19 pandemic in the Philippines. J Affect Disord 2020;277:379–91. doi:10.1016/j.jad.2020.08.043.
- [39] Alkhamees AA, Alrashed SA, Alzunaydi AA, Almohimeed AS, Aljohani MS. The psychological

impact of COVID-19 pandemic on the general population of Saudi Arabia. Compr Psychiatry 2020;102. doi:10.1016/j.comppsych.2020.152192.

- [40] Hyland P, Shevlin M, McBride O, Murphy J, Karatzias T, Bentall RP, et al. Anxiety and depression in the Republic of Ireland during the COVID-19 pandemic. Acta Psychiatr Scand 2020;142:249–56. doi:10.1111/acps.13219.
- [41] Bäuerle A, Teufel M, Musche V, Weismüller B, Kohler H, Hetkamp M, et al. Increased generalized anxiety, depression and distress during the COVID-19 pandemic: A cross-sectional study in Germany. J Public Heal (United Kingdom) 2020;42:672–8. doi:10.1093/pubmed/fdaa106.
- [42] Mazza MG, De Lorenzo R, Conte C, Poletti S, Vai B, Bollettini I, et al. Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors. Brain Behav Immun 2020;89:594–600. doi:10.1016/j.bbi.2020.07.037.
- [43] Hacimusalar Y, Kahve AC, Yasar AB, Aydin MS. Anxiety and hopelessness levels in COVID-19 pandemic: A comparative study of healthcare professionals and other community sample in Turkey. J Psychiatr Res 2020;129:181–8. doi:10.1016/j.jpsychires.2020.07.024.
- [44] Bigalke JA, Greenlund IM, Carter JR. Sex differences in self-report anxiety and sleep quality during COVID-19 stay-at-home orders. Biol Sex Differ 2020;11. doi:10.1186/s13293-020-00333-4.
- [45] Prete G, Fontanesi L, Porcelli P, Tommasi L. The psychological impact of COVID-19 in Italy: Worry leads to protective behavior, but at the cost of anxiety. Front Psychol 2020;11. doi:10.3389/fpsyg.2020.566659.
- [46] Traunmüller C, Stefitz R, Gaisbachgrabner K, Schwerdtfeger A. Psychological correlates of COVID-19 pandemic in the Austrian population. BMC Public Health 2020;20. doi:10.1186/s12889-020-09489-5.