A novel benchmark for COVID-19 pandemic testing effectiveness enables the

2 accurate prediction of new Intensive Care Unit admissions

3 Dimitris Nikoloudis^{1*}, Dimitrios Kountouras¹ and Asimina Hiona¹

⁴ ¹ Center for Preventive Medicine & Longevity, Bioiatriki Healthcare Group, 11525 Athens,

- 5 Greece
- 6 *Corresponding author
- 7

8 Abstract

9 The positivity rate of testing is currently used both as a benchmark of testing adequacy and for assessing the evolution of the COVID-19 pandemic. However, since the former is a prerequisite for the latter, its 10 11 interpretation is often conflicting. We propose as a benchmark for COVID-19 testing effectiveness a new 12 metric, termed 'Severity Detection Rate' (SDR), that represents the daily needs for new Intensive Care 13 Unit (ICU) admissions, per 100 cases detected (t-i) days ago, per 10,000 tests performed (t-i) days ago. 14 Based on the announced COVID-19 monitoring data in Greece from May 2020 until August 2021, we 15 show that beyond a certain threshold of daily tests, SDR reaches a plateau of very low variability that 16 begins to reflect testing adequacy. Due to the stabilization of SDR, it was possible to predict with great 17 accuracy the daily needs for new ICU admissions, 12 days ahead of each testing data point, over a period of 10 months, with Pearson r = 0.98 ($p = 10^{-197}$), RMSE = 7,16. We strongly believe that this metric will 18 19 help guide the timely decisions of both scientists and government officials to tackle pandemic spread 20 and prevent ICU overload by setting effective testing requirements for accurate pandemic monitoring. 21 We propose further study of this novel metric with data from more countries to confirm the validity of 22 the current findings.

- 24
- 25
- 26

It is made available under a CC-BY 4.0 International license .

2	7
_	

- 28
- 29

30 Introduction

31

Although no country knows at any point the true total number of COVID-19 cases, it is crucial for public health administrations to be confident that the daily testing performed is stably representative of that number. Effective testing provides health professionals and officials with a clear picture of SARS-CoV-2 spread within the community, as well as of the dynamics of COVID-19 pathology, and guides them for prompt and adequate interventions towards containment of the pandemic at the local and national levels.

38 The percentage of tests that return a positive result, also known as the "positivity rate", is an 39 important outcome of testing that is used both as a benchmark for testing adequacy and as a metric for assessing the current spread of the virus^{1,2}. However, this dual usage presents an inherent drawback in 40 41 entrusting the metric in any one of two possible ways: is a high positivity rate due to a high number of 42 infected individuals, or due to a low number of tests performed? A rule of thumb says that a positivity 43 rate of 5% is too high, and the WHO suggests that the positivity rate should rest below that threshold for 44 a length of at least two weeks before officials decide to progressively reopen professional and social activities¹. Another evidence-based perception suggests that the positivity rate must remain below 3% 45 to ensure that surveillance is broad and accurate enough². However, these rules may only cover either 46 47 the virus spread surveillance criterion or that of testing adequacy, not both. Indeed, officials often respond to a high positivity rate both with an increase in testing and with measures to restrict virus 48 49 transmission, such as social distancing and soft or hard lockdowns. But by doing so, it is expectedly hard 50 to timely assess the true rate of the virus spreading out, or being contained, as the new higher levels of 51 testing must be stabilized for a length of time before allowing again to reliably follow the pandemic 52 dynamics. In such a scenario, if health officials rely only on the positivity rate metric, the timing of the 53 response would lag and thus be almost invariably suboptimal.

54 Fundamentally, a metric that would serve as a benchmark for the effectiveness of COVID-19 55 testing should not concurrently be used for assessing the evolution of the pandemic, as the former is a

It is made available under a CC-BY 4.0 International license .

56 prerequisite for the latter and therefore the interpretation would be conflicting; indeed, the health 57 administrations of a country should be confident that a sufficient number of tests is performed to 58 effectively track the virus spread. However, if such a metric also implemented measurable outcomes of 59 the pandemic in the community (e.g., number of deaths, number of ICU admissions, etc.), they could 60 introduce by their more factual nature a link between expectation and actuality, since the outcomes of 61 COVID-19 are inherently tied to the virus's pathogenesis. Therefore, such a link could, in theory, 62 introduce a benchmarkable step of convergence towards a soft cap (threshold) that would in turn reflect 63 testing adequacy, e.g., usually a maximized or minimized value, or a state of minimized variation. In this 64 report, we present an easy-to-implement metric that we developed while independently monitoring 65 and analyzing COVID-19 pandemic evolution in Greece, which considers outcomes that are already monitored in most countries, such as the daily numbers of human losses, COVID-19 patients in the ICU 66 67 (Intensive Care Units), and patients who are being discharged from the ICU. In our example we show 68 that this metric displays remarkable output stability when a certain threshold of daily testing is reached, 69 which to our view clearly reflects testing adequacy. Furthermore, we validated its benchmarking 70 efficiency by forecasting, not only with high accuracy but also great precision, the total daily needs for 71 new ICU admissions, roughly two weeks in advance, over a period of 10 months.

72

73 Methods

74

The national monitoring data for the evolution of the COVID-19 pandemic in Greece were retrieved from the Hellenic National Public Health Organization³ and Greek Government's official daily announcements⁴. Specifically, the daily official announcements included the following parameters: (a) number of new COVID-19 cases detected, (b) number of deaths due to COVID-19, (c) total number of COVID-19 ICU patients, (d) total number of COVID-19 patients discharged from ICU, (e) total number of SARS-CoV-2 PCR tests performed⁵, and (f) total number of SARS-CoV-2 rapid antigen tests performed⁶.

81 Based on the available data, we defined *the daily needs for new COVID-19 ICU admissions* as 82 number U:

83 $U = (x_0 - x_{t-1}) + d + e$ (1)

84 where:

85	-	Today's deaths due to COVID-19:	d
86	-	Today's number of COVID-19 patients discharged from ICU:	e
87	-	Today's total number of COVID-19 ICU patients:	x 0
88	-	Yesterday's total number of COVID-19 ICU patients:	X _{t-1}

This number U represents the actual daily new COVID-19 ICU admissions, plus those patients who died in the community (not in ICU), whom we theorize to have required ICU admission, hence the definition of *the daily needs for new COVID-19 ICU admissions*.

92

93 Next, we defined as the Severity Detection Rate with a time lag (t-i) (SDR_i), a metric that 94 represents the percentage of patients who require ICU admission, per new cases, detected (t-i) days 95 ago, per 10,000 tests, performed (t-i) days ago:

96
$$SDR_i = (U^*100/c_{t-i})/(n_{t-i}/10,000) \Rightarrow SDR_i = U^*1,000,000/(c_{t-i}^* n_{t-i})$$
 (2)

97 where:

- 98 Today's rolling 7-day average of new daily needs for COVID-19 ICU: U
- 99 Rolling 7-day average of detected COVID-19 cases, (t-i) days ago: c_{t-i}
- 100 Rolling 7-day average of total number of COVID-19 tests, (t-i) days ago: n_{t-i}

101

102 Tests in Greece were performed freely by any individual who wanted to get tested, in selected 103 hospitals, or in most private diagnostic centers and clinics, or in mobile testing hubs, dispatched by the 104 public healthcare administration. Also, an individual may get tested in regular intervals (e.g., up to twice 105 per week), as requested by their employer or the administration, due to the nature of their profession. 106 To the best of our knowledge, only one swab is taken from the individual per test, in Greece. 107 Furthermore, the reported COVID-19 cases detected, and daily tests performed, are used for the official calculation of positivity rate, announced routinely by the country's healthcare administration³; if 108 109 multiple tests per individual were simply added to the total daily number, this would constitute a 110 systematic error in the calculation of positivity rate. Therefore, for the reasons explained above, for this 111 analysis, the daily number of tests reported publicly is presumed to represent unique individuals.

It is made available under a CC-BY 4.0 International license .

112	For initial data exploration, the lag of Severity Detection Rate (SDR) metrics was set to 14 days,
113	which means that the current day's critical outcomes of COVID-19 (i.e., ICU admission or death in the
114	community) were attributed to COVID-19 cases detected 14 days ago. For the identification of the
115	optimal lag point between the critical outcomes of COVID-19 and the detected cases, we searched
116	within an interval between 7 to 21 days, in the period 17/10/20 to 31/1/21 of the dataset, for the most
117	stable correlation between the numerator (number U) and the denominators ([cases _{t-i} * tests _{t-i}]) of the
118	metrics studied. The best correlation was obtained for a lag of 12 days (i=12) (see Discussion section)
119	and therefore, for consistency, all charts and tables reflect this optimal time lag (i=12).

120

Finally, for completeness of the study, we also defined as ICU admission Rate with a time lag (t-i) (henceforth "ICU Rate", IR_i), a metric that represents the percentage of patients who require ICU admission, per new cases, detected (t-i) days ago:

124
$$IR_i = U^* 100/c_{t-i}$$
 (3)

125 where:

126	-	Today's rolling 7-day average of new daily needs for COVID-19 ICU:	U
127	-	Rolling 7-day average of detected COVID-19 cases, (t-i) days ago:	C _{t-i}

128

129 IR metric is essentially a simpler form of the SDR metric, which doesn't take into account the number of 130 daily tests performed. As we wanted to also evaluate its predictive performance, we doubled every 131 piece of analysis performed on the SDR metric, on the IR metric as well. The related charts and tables 132 are not part of the Results section in favor of clarity for the main metric presented (SDR), but are, 133 nonetheless, commented upon in the Discussion section.

134

135 The dataset was locked on August 8th 2021.

136

137

It is made available under a CC-BY 4.0 International license .

139	
140	
141	
142	Results
143	For observation, the daily evolution of SDR ₁₂ , from the 7^{th} of May 2020 onwards, was traced on

144 the same chart versus the observed number of daily ICU needs, the positivity rate and the corresponding

145 number of testing samples (Figure 1).

- 146
- 147
- 148



It is made available under a CC-BY 4.0 International license .

Figure 1. Comparison of trendlines of Severity Detection Rate, Daily needs for new ICU admissions, Positivity rate, and number of Daily Tests, in the period from 7/5/2020 to 8/8/2021. The Daily needs for new ICU admissions and the number of Daily Tests represent rolling 7-day averages. Severity Detection Rates and Positivity Rates were calculated from the rolling 7-day averages of their components. All numbers were normalized by their maximum value in the examined period.

155

156 Compared to the other quantities, the SDR metric shows a remarkable stabilization past the time mark 157 on approximately 20/8/2020, which also corresponds to the attainment of an average daily testing 158 number of 10,000/day. From that point forward, the observed daily ICU needs, the positivity rate and 159 the testing rate continue to fluctuate independently and considerably, but without accordingly 160 perturbing SDR stabilization.

161

162

163 The rate of daily testing in Greece has been scaled up significantly on four occasions, 164 approximately (a) on 29/7/2020, (b) on 17/10/2020, (c) on 1/2/2021 and (d) on 11/7/2021. As the new testing levels were preserved after each scale-up, it is possible to define 5 distinct periods of testing 165 166 intensity thus far during the COVID-19 pandemic in Greece. Interestingly, a sixth distinct period is noted 167 between 1/5/2021 and 10/7/2021, where, inversely, a steady reduction in the number of daily tests is 168 observed, although the daily average number of tests is preserved from the immediately previous 169 period. We qualified this behavior as noteworthy and chose to study the respective period separately. 170 We therefore characterized the SDR number and the rates of testing for each of the following time 171 intervals: (i) 1/5/2020 - 28/7/2020, (ii) 29/7/2020 - 16/10/2020, (iii) 17/10/2020 - 31/1/2021, (iv), 1/2/2021 – 30/4/2021, (v) 1/5/2021 – 10/7/2021, and (vi) 11/7/2021 – 8/8/2021 (Table 1). 172

- 173
- 174
- 175
- 176
- 177

It is made available under a CC-BY 4.0 International license .

- 178
- 179
- 180
- **Table 1.** Characterization of the Severity Detection Rate and the number of daily tests for each of the
- 182 six time intervals of distinct testing levels in Greece.

183

Distinct periods of testing levels						
intervals 1/5/2020 - 28/7/2020		29/7/2020 -	- 16/10/2020	17/10/2020	- 31/1/2021	
	SDR ₁₂	Samples	SDR ₁₂	Samples	SDR ₁₂	Samples
max	92.0%	7309	7.0%	20310	3.2%	31602
average	20.1%	4051	2.7%	12861	2.1%	24377
median	14.3%	3992	2.6%	12453	2.0%	24743
mi n	0.1%	1400	1.1%	9706	1.3%	17315
sd	19.6%	1260	1.0%	2439	0.4%	3172
CV	0.97	0.31	0.36	0.19	0.19	0.13
intervals	1/2/2021 - 30/4/2021		1/5/2021 - 10/7/2021		11/7/2021 - 27/7/2021	
	SDR ₁₂	Samples	SDR ₁₂	Samples	SDR ₁₂	Samples
max	1.6%	58578	1.0%	57206	0.12%	76262
average	0.9%	42905	0.4%	42150	0.11%	69974
median	0.9%	41906	0.4%	41922	0.11%	73005
mi n	0.5%	29504	0.2%	30871	0.10%	58583
sd	0.3%	7510	0.2%	7190	0.01%	5919
CV	0.32	0.18	0.36	0.17	0.06	0.08

184

185

Tripling the average daily rate of testing (from 4K to 12K) in the second (ii) interval brought a 7-fold lower average value of SDR ($20.1\% / 2.7\% \sim 7.4$), with a remarkable 20-fold decrease (19.6% / 1%) in the Standard Deviation (SD) of SDR, and a concomitant 3-fold decrease in the CV (Coefficient of Variation) of SDR ($0.97/0.36 \sim 2.7$). Further doubling of the average daily number of tests (from 12 K to 24 K) in the third (iii) interval again brought an equivalent decrease in the SDR SD (1.0% / 0.4% = 2.5) although the average value of SDR was now only moderately diminished by approximately 30% ($2.7\% / 2.1\% \sim 1.29$), indicating a tendency towards stabilization of the SDR value and a continuous reduction of the Standard

It is made available under a CC-BY 4.0 International license .

193 Deviation (SD). Overall, it is noteworthy that specifically the average and SD values of SDR continued to 194 drop consistently in all 6 periods.

195

We then traced the values of SDR metric against the daily number of tests. The SDR values display a strong correlation with the daily number of tests, employing power regression (Spearman r = -0.90, p = 10^{-167} , N = 451) and suggest that beyond a threshold of daily tests performed, SDR becomes significantly stabilized (Figure 2); for Greece, this stabilization begins once the number of daily tests exceeds the mark of 10,000 per day.



201

Figure 2. Correlation between the Severity Detection Rate and rolling 7-day averages of the number of daily tests, with Spearman r = -0.90, $p = 10^{-167}$, N = 451. Numbers of daily tests derived from the period from 15/5/2020 to 8/8/2021.

205

The next step was to study the correlation between the numerator (number U) and the denominator ([cases_{t-i} * tests_{t-i}]) of SDR metric, for the period 17/10/2020 to 8/8/2021 (Figure 3). The starting period was chosen to be the same as the start of testing period (iii) (Table 1). Before that date,

It is made available under a CC-BY 4.0 International license .

- both the numbers of new daily needs for ICU and daily cases were relatively low (Figure 1) and therefore
- of smaller interest to the specific study, *i.e.*, when added to the rest of the data, the respective
- correlation is innately stronger due to the near-baseline nature of the data points prior to 17/10/2020.

212



Figure 3. Correlation of the numerator and denominator of SDR, i.e., number U versus the product (cases_{t-12}* tests_{t-12}), with a lag of 12 days. Numbers of daily tests and detected cases were derived from the period from 17/10/2020 to 8/8/2021.

217

It is made available under a CC-BY 4.0 International license .

- 219
- 220

221	Finally, we applied the linear regression equations to forecast the rolling 7-day average daily
222	needs for new ICU admissions, 12 days ahead of each data point of daily announced cases and tests, for
223	the corresponding periods, i.e., from 17/10/2020 to 8/8/2021. The forecast employing the SDR
224	regression equations (Figure 3) proved very accurate (Pearson r = 0.98, p = 10^{-197} , RMSE = 7.16; with n =
225	296, observed U[max] = 125, U[average] = 51) (Figure 4). Expectedly, as can be noticed in Figure 4B,
226	most of the few intense discrepancies in the fitted values are observed around dates of transition from
227	one regression equation to another; a rolling regression window could possibly help improve the
228	forecast of even these phases. Overall, forecasting with the use of Severity Detection Rate proved to be
229	functional as it indicated a very strong agreement between the predicted and observed values for a
230	period of nearly 10 months, which included the two major pandemic waves in the country, thus far.
231	
222	
232	
233	
234	
225	
235	
236	
237	
220	
238	
239	
240	
241	
242	
243	
244	

It is made available under a CC-BY 4.0 International license .

245

246 **A**







It is made available under a CC-BY 4.0 International license .

250

Figure 4. (A) Correlation between observed and predicted daily needs for new ICU admissions for the period between 17/10/2020 and 8/8/2021 employing Severity Detection Rate regression equations, with Pearson r = 0.98, p = 10^{-197} , RMSE = 7.16; with n = 296, observed U[max] = 125, U[average] = 51. (B) The respective time-series plots for visual inspection of the fit.

- 255
- 256

257 **Discussion**

258

259 We have shown that beyond a threshold of daily tests performed, SDR reaches a plateau that 260 displays very low variation. This threshold appears roughly around the 10,000 daily samples mark in 261 Greece, a country of approximately 11 million people, but this number is expected to vary greatly from 262 country to country depending on the total population, rural density, societal particularities, population 263 immune profile, and sampling strategies⁷. Reaching that threshold should not mean that there is no 264 need for further increase in the number of daily tests, as it is strongly suggestive that the more tests a 265 country performs, the more informative the results are about the actual viral spread in the community, 266 and consequently health administrations are in better position to respond accordingly. In terms of the 267 SDR metric, more daily tests appear to further decrease its variation (Table 1). The weaker its variation, 268 the stronger the correlation coefficient between the numerator and denominator of SDR, i.e., number U 269 versus the product (cases_{ti}* tests_{ti}), and therefore, the more accurately we can predict the number of 270 daily needs for new ICU admissions, t+i days in advance. In the studied example, predictions were highly 271 accurate with an average daily number of tests as high as 24,000 (Table 1), which resulted in a SD of the 272 SDR of 0.4%. As the SD of the SDR showed a consistent decrease over a period of 15 months in our 273 studied example (Table 1), we propose it can possibly act as an actual numerical threshold that denotes 274 the attainment of the SDR plateau.

As a direct consequence of this potential predictability, when SDR establishes a plateau, we consider that the bulk of daily tests is returning a set of positive cases that is stably representative of the current spread of the virus. Therefore, the SDR metric constitutes a benchmark of testing effectiveness.

It is made available under a CC-BY 4.0 International license .

The metric is potentially efficient at a local level as well, if cases that require delocalization, *e.g.*, due to lack of available ICU locally, are effectively tracked and taken into account. As the full segmentation of the necessary data was not available at a local level for the present study, it was not possible to evaluate the effects of viral spread uniformity across the country and, more specifically, the metric's behavior due to disproportionate testing intensities locally, e.g., higher number of tests in districts with lower viral load, and relatively lower numbers of daily tests in districts with higher true viral load. In such a case, it would be helpful to apply the SDR monitoring at a local level.

285 The metric's median value is expected to decrease monotonically and with decreasing variation 286 as daily tests increase, or due to the gradual containment of the virus, immunization of the population, 287 thanks to an efficient vaccination program, improvement of therapeutic protocols that reduce the 288 number of very severe cases, or even a significant reduction in the average age of infected individuals 289 due to the efficient protection of the elderly. Conversely, the metric's median value may increase 290 (interrupting the plateau) if the viral spread becomes greatly enhanced with time, e.g., due to the prevalence of a new more infectious variant^{8,9,10}, without the testing levels catching up. In such a case 291 292 the SDR's median will increase disproportionately and beyond its expected variability.

293 In order to comprehend the nontrivial nature of the plateau attainment and retainment in the 294 plot of SDR versus the number of daily tests (Figure 2), it is useful to look more carefully at some notable 295 boundaries of the SDR metric. For instance, if it was possible to test the entire population every day for 296 newly infected individuals (minus the individuals that are already known to be infected), then the 297 "discovery" of every new infection case would be guaranteed (assuming 100% accurate tests). With a number of daily tests as big as the entire population and with the highest possible number of detected 298 299 cases (i.e., equal to the actual cases), the SDR value becomes [U / ((actual new cases) * population)] 300 with the denominator assuming its greatest possible value, hence producing the lowest possible SDR.

301 In a different approach that hypothetically guarantees the detection of all the actual new 302 infected cases (without testing the entire population), we can consider testing all the newly infected 303 individuals, and only them, so that the number of daily tests becomes equals to the number of new 304 infections (again, assuming 100% testing accuracy). In this case the SDR value becomes [U/(actual cases * actual cases) = U/ actual cases²]. Whether the possible values of the SDR metric can be bigger or 305 306 smaller than the value obtained in this second hypothetical scenario, depends on whether the product 307 $[cases_{t-i} * test_{t-i}]$ is smaller or bigger than the square of the number of actual new infection cases (see 308 mathematical demonstration, below). Finally, as the theoretical maximum of all the possible SDR values

It is made available under a CC-BY 4.0 International license .

309 we may consider the case where the denominator $[cases_{t-i} * tests_{t-i}]$ is equal to 1, and therefore SDR 310 would be equal to number U. Specifically:

311

312	$U/(\alpha_{t-1}*population) \le SDR \le U/{\alpha_{t-1}}^2 \le SDR' \le U $ =>				
313	$U/(\alpha_{t-i}*population) \le U/(cases_{t-i}*tests_{t-i}) \le U/\alpha_{t-i}^2 \le U/(cases_{t-i}'*tests_{t-i}') \le U$				
314	$1/(\alpha_{t-i}*population) \le 1/(cases_{t-i}*tests_{t-i}) \le 1/\alpha_{t-i}^2 \le 1/(cases_{t-i}'*tests_{t-i}') \le 1$	(4)			
315	therefore:				
316	$(cases_{t-i}^*tests_{t-i}) \le \alpha_{t-i}^*$ population	(5)			
317	$(cases_{t-i} * tests_{t-i}) \ge \alpha_{t-i}^2 \implies \forall (cases_{t-i} * tests_{t-i}) \ge \alpha_{t-i}$	(6)			
318	$(cases_{t-l}'*tests_{t-l}') \le \alpha_{t-l}^{2} \qquad = > \qquad \forall (cases_{t-l}'*tests_{t-l}') \le \alpha_{t-l}$	(7)			
319	where:				
320	- Today's rolling 7-day average of new daily needs for COVID-19 ICU:	U			
321	- The actual new cases, (t-i) days ago:	α_{t-1}			
322	- Severity Detection Rate at plateau: SDR				
323	- Severity Detection Rate outside the plateau:	SDR'			
324	- Rolling 7-day average of detected COVID-19 cases, (t-i) days ago:	cases _{t-l} , cases _{t-l} ,			
325	with $1 \leq cases_{t\text{-} } \leq \alpha_{t\text{-} }$, and $1 \leq cases_{t\text{-} }' \leq \alpha_{t\text{-} }$				
326	- Rolling 7-day average of total number of COVID-19 tests, (t-i) days ago:	$tests_{t\text{-}i}, tests_{t\text{-}i}'$,			
327	with $1 \leq \text{tests}_{t-1} \leq \text{population}$, and $1 \leq \text{tests}_{t-1}' \leq \text{population}$				
328					
329					
330	Inequality (5) is trivial as the number of actual new cases (α_{t}) and the	entire population of the			
331	country, or area of interest, are by definition the highest possible values of the product (cases _{t-i} *tests _{t-i}).				
332	However, inequality (6) describes a situation where the number of tests can	only be equal or greater			

than $\alpha_{t-i}^2/cases_{t-i}$, and which may increase up to the number of the entire population, causing the reduction of the SDR value till its described minimum of U/(α_{t-i} *population). Inequality (7), inversely,

It is made available under a CC-BY 4.0 International license .

describes a situation where the number of tests can only be equal or lower than $\alpha_{t-1}^2/case_{t-1}$, and which may decrease to as low as 1 test, causing the increase of the SDR value to its maximum that equals the number U.

338 Therefore, because of this demonstrated relationship between the number of daily tests and the 339 number of actual new infections, we theorize that in a plot of SDR versus the number of daily tests, the 340 observed plateau is a consequence of the SDR starting to adopt values that are smaller than U/α^2 . Inversely we observe values outside the plateau as long as SDR adopts values greater than U/α^2 . This is 341 potentially what happened around the mark of 10,000 tests in our studied example (roughly around 342 20/8/2020), with the product (cases * tests) increasing almost 10-fold within a few days and presumably 343 becoming greater than the square of the actual new cases, thus collapsing the SDR variability into the 344 345 observed plateau (Figures 2 & 5). The importance of the plateau being, as previously explained, the 346 reduction of the metric's variability (i.e., Standard Deviation), enabling a correspondingly robust 347 forecasting of ICU needs, (t+i) days ahead of each datapoint.



It is made available under a CC-BY 4.0 International license .

Figure 5. Visualization of the domain space where the number of actual new cases are to be found

hypothetically (blue area), relatively to the square root of the product [cases $_{t-12}$ * tests $_{t-12}$] (orange line),

before and after the date of 8/20/2020, which marked the beginning of the SDR plateau.

353

354 In the context of the regression analysis of the daily needs for new ICU admissions (U) vs. the 355 product of [Detected Cases * Performed Tests] (Figure 3), significant changes in the SDR median would 356 be reflected as changes in the slope and/or the intercept of the regression line. Specifically, changes in 357 the slope most likely translate into two possibilities: (A) a change in virulence (*i.e.*, how many individuals 358 per group of 100 positive cases, per 10,000 tests, are expected to develop very severe COVID-19, given a 359 theoretical zero regression intercept), or (B) a modification in sampling parameters (e.g., testing more or 360 fewer asymptomatic persons, or testing a younger subset of the population). Accordingly, a change in the intercept will likely signify either (a) changes in viral prevalence^{7,11}, as the intercept represents a 361 fixed number U for a theoretical x=0, (*i.e.*, a number of individuals with very severe COVID-19, while no 362 cases are detected), or (b) changes in testing accuracy^{7,11}, with intercept values closer to zero reflecting 363 364 optimal accuracy. Rolling 3-weeks regression windows could be employed to detect dynamic changes of 365 the pandemic. The study of all the available confounding factors (e.g., prevalence of new virus variant, 366 changes in sampling strategies, changes in testing parameters, characteristics of areas infected, 367 lockdown and other measures' status, ages of tested and infected individuals, etc.) is required to discern 368 which exact change is responsible for the observed new disease dynamics, and the SDR derived 369 regression analysis can provide significant hints as to the direction of the change. In any of the above 370 cases, an important shift of the SDR would signify an important change in the pandemic parameters, 371 which in turn would dictate a specific course of action for the authorities, appropriate for each case.

372 In Table 2 we contrast the regression parameters (*i.e.*, slope, intercept and R^2) against important 373 factors of the ongoing pandemic, such as, Delta variant prevalence, vaccination levels, and lockdown 374 periods. What is most notable is the stable slope decrease of the regression equations, over all 6 periods 375 examined, which is compatible with a decrease in population-level severity/virulence. This is to be 376 expected, given the long periods of the applied lockdown measures and the ongoing mass vaccination 377 program in the country (reaching 50% population coverage of fully vaccinated individuals on 8/8/2021). 378 As presented in the previous paragraph, another factor that can possibly lower the SDR slope is a 379 significant change in sampling parameters, in a way where the group of asymptomatic individuals that 380 are being tested becomes considerably increased, a situation that results inherently to fewer detected

It is made available under a CC-BY 4.0 International license .

cases than the group of symptomatic individuals. While it is hard to discern the potential contribution of each factor with just the publicly available data, it is, nonetheless, possible to calculate a 9.5-fold total drop in the observed severity between the beginning and ending of the six periods (17/10/2020 -> 8/8/2021), after adjusting for the obvious contribution of the change in the average number of cases and tests (Table 2):

- 386 Δ (slope, 17/10/2020 -> 8/8/2021): (2*10⁻⁶ / 7*10⁻⁸) = 28.6 Unadjusted fold change in severity

387

 Δ (average SDR denominator, 17/10/2020 -> 8/8/2021): (1,602*55,190) / (1,142*25,585) = 3 -

389
$$\Delta$$
(observed severity, 17/10/2020 -> 8/8/2021): 28.6 / 3 = 9.5 - Adjusted fold change in severity

390

391 On the contrary, the intercept oscillates considerably between periods, ranging from +28.6 to -392 3.9. As explained previously, increases of the intercept may be attributed to greater viral spread in the 393 community, as was the case in the second period (11/2/2021 - 21/4/2021), when Athens, the capital, saw a great increase in infected cases, which signaled the beginning of the 3rd wave of the pandemic in 394 395 Greece. Besides viral spread, the other factor that influences the intercept is the accuracy of the tests 396 performed, i.e., potential false positives and false negatives, due to poor test specificity, test sensitivity, 397 or yet undetectable levels of the virus in asymptomatic infected individuals who simply got tested too 398 early in the course of the disease. Regarding Delta variant prevalence (B.1.617.2), representing 90% of 399 cases in Greece on 8/8/2021, it doesn't appear to be affecting the severity of the disease (*i.e.*, a slope 400 increase), however it is possibly contributing to the intercept increase from 16/6/2021 onwards, with its greater transmissibility potential, as reported by other studies¹⁸. Overall, the slope and intercept of SDR-401 402 based regression equations offer an additional layer of information, which, in conjunction with other 403 metrics and parameters, may create a better understanding of the pandemic's dynamics.

We called this new metric Severity Detection Rate, as its representation of the percentage of very severe COVID-19 outcomes is modulated by the number of tests performed. It is essentially a standardization of the very severe cases ratio over the infected individuals, with the rate of daily testing. In other words, the Severity Detection Rate becomes representative of the proportion of people who

It is made available under a CC-BY 4.0 International license .

408 need ICUs out of the total cases once a sufficient threshold of daily testing rate (hence 'detection rate')409 is achieved.

410 As presented in the Methods section, for a more complete examination, we also defined the 411 percentage of patients who require ICU admission, per new cases detected (t-i) days ago, as ICU Rate 412 (IR). If, in theory, the total number of tests became equal to the entire population of a country (or the 413 area of interest), then the SDR metric would be the same as the IR metric, as the 'number of tests' 414 parameter would be removed from the denominator (as redundant), and both would practically 415 represent the true percentage of critical patients per infected individual. In order to assess the 416 predictive potential of the IR metric, we have repeated for IR every piece of analysis that was performed 417 on the SDR metric throughout this study.

418 Regarding forecasting, the conclusion drawn by this parallel analysis is that the IR metric 419 performed as well as the SDR metric, in the analyzed example (Figures 3-S, 4-S, Table 2-S). On top of 420 this, the IR metric would probably have the advantage of simplicity when communicated in the general 421 public, as it represents a more comprehensible concept: the number of very severe cases per infected 422 individuals. We therefore believe that the IR metric may be used in cases where the population-level 423 COVID-19 testing surveillance of the pandemic is well established, by efficient and sufficient testing. 424 Nonetheless, we support that by including the number of daily tests performed, the SDR metric is 425 inherently more suitable for a wider range of surveillance scenarios, e.g., when the testing strategies 426 and pandemic parameters (e.g., number & type of tests, geographical/ occupational/ age targeting, 427 contact tracing efficacy, transmissibility of the virus, etc.) are more volatile in time. In different 428 countries, or in specific areas of interest, it is still possible for the IR-based monitoring to fail to return 429 regression coefficients as strong as in our studied example. In those cases, it would be necessary to 430 switch to SDR-based monitoring to ensure that a threshold of sufficient testing has been reached (i.e., 431 plateau formation). In any case, although more studied examples are required to better understand the 432 potential practical differences between the two metrics, since they both showed equal forecasting 433 performances, we believe that SDR is the more well-rounded metric, which can be efficiently used in 434 potentially very diverse situations of pandemic surveillance.

435

436

It is made available under a CC-BY 4.0 International license .

439 Table 2. Regression equations per distinct period of stable Severity Detection Rate, with respective average numbers of observed daily needs for 440 new ICU admissions, cases detected and tests performed, vaccination coverage at the beginning of each period, percent of Delta variant prevalence 441 in cases detected, along with important dates and comments that potentially influenced the course of the pandemic in Greece (each period is color-442 coded with reference to the respective 'distinct periods of testing levels', in Table 1).

443

Distinct periods of stable Severity Detection Rate						
17/10/2020 - 10/2/2021 11/2/2021 - 21/4/2021 22/4/2021 - 22/5/2021 23/5/2021 - 15/6/2021 16/6/2021 - 6/7/2021 7/7/2021 - 8/8/2021						
SDR ₁₂ Regression	2*10^6x + 5.7511	5*10^7x + 28.6	4*10^7x + 25.948	4*10^-7x - 3.8899	2*10^7x + 6.7379	7*10∿8x + 5.2217
SDR ₁₂ R ²	0.981	0.833	0.84	0.962	0.954	0.933
average U (∆U)	56 [-]	69 [+ 2 3%]	73 [+6%]	34 [-53%]	13 [-62%]	13 [0%]
average cases (∆cases)	1,142 [-]	1847 [+62%]	2,453 [+33%]	1887 [-23%]	763 [-60%]	1602 [+110%]
average tests (∆tests)	25,585 [-]	39847 [+56%]	49,580 [+24%]	48976 [-1%]	37651 [-23%]	55190 [+47%]
Vaccination levels @period start	-	1/2/2021	10/4/2021	11/5/2021	4/6/2021	25/6/2021
at east 1-dose [%pop]	0	244,652 [2.28%]	1,435,513 [13.39%]	2,561,871 [23.90%]	3,854,962 [35.96%]	4,678,468 [43.64%]
complete vaccination	0	46,295 [0.43%]	733,840 [6.85%]	1,335,798 [12.46%]	2,253,347 [21.02%]	3,461,165 [32.29%]
Delta variant prevalence (B.1.617.2)	-	-	~0.2% (22/5/21)	~1% (29/5/21)	~5% (12/6/21), ~61% (3/7/21)	~76% (10/7/21) ¹⁷
Important dates & comments	"2nd wave", national ockdown (7/11/20) ¹²	Athens (hosting half of Greece's population) majorly affected, full capital lockdown (9/2/21), "3rd wave" ¹³	service workers must perform and report weekly self tests (19/4/21) ¹⁴	lockdown lifted progressively (14/5/21) ¹⁵ , all employees must perform and report weekly self-tests (24/5/21) ¹⁶	Indoor dining, amusement parks and other entertainment services reopen (7/6/21) ¹⁵	possible"4th wave"

444

445

446

447

448

It is made available under a CC-BY 4.0 International license

It is made available under a CC-BY 4.0 International license .

449

450 **Conclusions**

451 Taken together, the monitoring of the Severity Detection Rate and the forecasting of number U 452 (i.e., daily needs for ICU) should be viewed as integral parts of the currently employed epidemiological 453 toolbox, i.e., the positivity rate, efficient contact tracing for determination of the basic reproduction number R₀^{19,20}, and wastewater-based surveillance^{21,22}. The metric introduces the goal for authorities to 454 455 minimize its variation by means of a sufficient number of daily tests and an adequate sampling strategy. 456 Once this goal is achieved, accurate forecasting of daily needs for new ICU admissions becomes possible. 457 With accurate forecasting, number U becomes in essence a quantitative metric for the severity of the 458 pandemic.

459 In Figure 6 we detail all the proposed steps for population-level surveillance of COVID-19 460 pandemic using the Severity Detection Rate metric. For monitoring SDR Standard Deviation, a minimum of 3-weeks rolling window interval is suggested empirically, as this interval includes the roughly 2-week 461 lag period between case detection and ICU intubation. The recommended surveillance model provides 462 463 three distinct advantages: (1) a measurable threshold for adequacy of tests performed, (2) important 464 qualitative information regarding the current dynamics of the pandemic (virulence, prevalence, testing 465 accuracy, etc.) that are reflected by changes in the slope/intercept of the regression analysis, and (3) the 466 ability to accurately predict the ICU needs, t+I days ahead.

We strongly believe that the explicit tracking of this novel metric enhances the visibility of viral spread and dynamics and may procure an accurate outlook of the upcoming needs for ICU admissions well in advance, which should serve as an early warning system for COVID-19 health establishments and resources. We therefore suggest further study of Severity Detection Rate with data from more countries, as well as at a local level wherever possible, to confirm the proposed functionality and utility of this metric.

473





476 Figure 6. Stepwise schema detailing the logic and requirements for population-level surveillance of
477 COVID-19 pandemic with the use of the Severity Detection Rate metric.

It is made available under a CC-BY 4.0 International license .

478 **References**

Dowdy D, D'Souza G. COVID-19 Testing: Understanding the "Percent Positive". Johns
 Hopkins Bloomberg School of Public Health. Accessed 31 August 2021. Available from:
 https://www.jhsph.edu/covid-19/articles/covid-19-testing-understanding-the-percent-positive.html

- Siddarth D, Katz R, Graeden E, Analytics T, Allen D, Tsai T. 2020. Evidence Roundup: Why
 positive test rates need to fall below 3%. Harvard Global Health Institute. Accessed 31 August 2021.
 Available from: <u>https://globalhealth.harvard.edu/evidence-roundup-why-positive-test-rates-need-to-</u>
- 485 <u>fall-below-3</u>

486 3. Hellenic National Public Health Organization. Daily Reports COVID-19. Accessed 31
487 August 2021. Available from: <u>https://eody.gov.gr/epidimiologika-statistika-dedomena/ektheseis-covid-</u>
488 19/

489 4. Greek Government's official community on Viber network. Official COVID-19 update.
490 Accessed 31 August 2021. Available from:

491 <u>https://invite.viber.com/?g2=AQAVNiDVlfjrlEtaYLf1s2sUzRrpfLVlfLVg4J8wkdNKMUSnUcQWJxnXH0Os1h</u>
 492 <u>eH&lang=el</u>

Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DKW, Bleicker T, Brünink S,
 Schneider J, Schmidt ML, Mulders DGJC, Haagmans BL, van der Veer B, van den Brink S, Wijsman L,
 Goderski G, Romette J-L, Ellis J, Zambon M, Peiris M, Goossens H, Reusken C, Koopmans MPG, Drosten
 C. 2020. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. Euro Surveill.
 25(3):pii=2000045. doi.org: 10.2807/1560-7917.ES.2020.25.3.2000045.

Lambert-Niclot S, Cuffel A, Le Pape S, Vauloup-Fellous C, Morand-Joubert L, Roque Afonso AM, Le Goff J, Delaugerre C. 2020. Evaluation of a Rapid Diagnostic Assay for Detection of SARS CoV-2 Antigen in Nasopharyngeal Swabs. J Clin Microbiol. 23;58(8):e00977-20. doi: 10.1128/JCM.00977 20.

 502
 7.
 Mercer TR, Salit M. 2021. Testing at scale during the COVID-19 pandemic. Nat Rev Genet

 503
 22, 415–426. doi: 10.1038/s41576-021-00360-w

5048.Elbe S, Buckland-Merrett G. 2017. Data, disease and diplomacy: GISAID's innovative505contribution to global health. Glob Chall. 10;1(1):33-46. doi: 10.1002/gch2.1018.

It is made available under a CC-BY 4.0 International license .

9. Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, Sagulenko P, Bedford T,
 Neher RA. 2018. Nextstrain: real-time tracking of pathogen evolution. Bioinformatics 1;34(23):4121 4123. doi: 10.1093/bioinformatics/bty407.

50910.Forster P, Forster L, Renfrew C, Forster M. Phylogenetic network analysis of SARS-CoV-2510genomes. 2020. Proc Natl Acad Sci U S A. 28;117(17):9241-9243. doi: 10.1073/pnas.2004999117.

11. Nichols JD, Bogich TL, Howerton E, Bjørnstad ON, Borchering RK, Ferrari M, Haran M,
Jewell J, Pepin KM, Probert WJM, Pulliam JRC, Runge MC, Tildesley M, Viboud C, Shea K. 2021. Strategic
testing approaches for targeted disease monitoring can be used to inform pandemic decision-making.
PLoS Biol 19(6): e3001307. doi: 10.1371/journal.pbio.3001307

515 12. Krinis N. 2020. Greece to Enter Lockdown to Fight Second Covid-19 Wave. Greek Travel
516 Pages. Accessed 31 August 2021. Available from: <u>https://news.gtp.gr/2020/11/05/greece-enter-</u>
517 <u>lockdown-fight-second-covid-19-wave</u>

51813.Koutantou A. 2021. Greek premier orders full lockdown in Athens after surge in519coronaviruscases.Reuters.Accessed31August2021.Availablefrom:520https://www.reuters.com/world/greek-premier-orders-full-lockdown-athens-after-surge-coronavirus-

521 <u>cases-2021-02-09</u>

14. Reuters Staff. 2021. Greece orders COVID self-testing for service workers. Reuters.
 Accessed 31 August 2021. Available from: <u>https://www.reuters.com/article/health-coronavirus-greece-</u>
 <u>tests-idUSL8N2M740B</u>

525 15. ESN COVID-19 Official Announcements & News. Timeline for the loosening of the 526 lockdown measures in Greece. 2021. Erasmus Student Network Greece. Accessed 31 August 2021. 527 Available from: https://esngreece.gr/covid-19-official-announcements-news

528 16. GTP editing team. 2021. All Employees in Greece Must Self-test for Covid-19. Greek
529 Travel Pages. Accessed 31 August 2021. Available from: <u>https://news.gtp.gr/2021/05/24/all-employees-</u>
530 greece-must-self-test-covid-19

531 17. European Centre for Disease Prevention and Control. SARS-CoV-2 variants of concern as
532 of 26 August 2021. Accessed 31 August 2021. Available from: <u>https://www.ecdc.europa.eu/en/covid-</u>
533 19/variants-concern

It is made available under a CC-BY 4.0 International license .

- <u>18. Centers for disease control and prevention. Delta Variant: What We Know About the</u>
 <u>Science.</u> Accessed 31 August 2021. Available from: https://www.cdc.gov/coronavirus/2019 ncov/variants/delta-variant.html
- 537 19. Macdonald G (1952). The analysis of equilibrium in malaria. Tropical Diseases Bulletin.
 538 49 (9): 813–829. ISSN 0041-3240. PMID 12995455.
- 539 20. Delamater PL, Street EJ, Leslie TF, Yang YT, Jacobsen KH. 2019. Complexity of the basic
 540 reproduction number (R0). Emerg. Infect. Dis. 25, 1–4.
- 21. Wu Y, Guo C, Tang L, Hong Z, Zhou J, Dong X, Yin H, Xiao Q, Tang Y, Qu X, Kuang L, Fang
 X, Mishra N, Lu J, Shan H, Jiang G, Huang X. 2020. Prolonged presence of SARS-CoV-2 viral RNA in faecal
 samples. Lancet Gastroenterol Hepatol. 5(5):434-435. doi: 10.1016/S2468-1253(20)30083-2. Epub 2020
 Mar 20.
- 545 22. Polo D, Quintela-Baluja M, Corbishley A, Davey LJ, Andrew CS, David WG, Jesús LR. 2020. 546 Making waves: Wastewater-based epidemiology for COVID-19 – approaches and challenges for 547 surveillance and prediction. Water Research, 186, 116404. doi: 10.1016/j.watres.2020.116404.
- 548
- 549

550

551