1 Detection and genome sequencing of SARS-CoV-2 belonging to the B.1.1.7 lineage in the Philippines

2

3	Francis A. Tablizo ¹ , Cynthia P. Saloma ^{2,7} *, Marc Jerrone R. Castro ¹ , Kenneth M. Kim ¹ , Maria Sofia L.
4	Yangzon ¹ , Carlo M. Lapid ¹ , Benedict A. Maralit ³ , Marc Edsel C. Ayes ⁴ , Jan Michael C. Yap ¹ , Jo-Hannah S.
5	Llames³, Shiela Mae M. Araiza³, Kris P. Punayan³, Irish Coleen A. Asin³, Candice Francheska B.
6	Tambaoan ³ , Asia Louisa U. Chong ³ , Karol Sophia Agape R. Padilla ³ , Rianna Patricia S. Cruz ¹ , El King D.
7	Morado ¹ , Joshua Gregor A. Dizon ¹ , Eva Maria Cutiongco-de la Paz ^{2,7} , Alethea R. de Guzman ⁵ , Razel Nikka
8	M. Hao ⁶ , Arianne A. Zamora ⁵ , Devon Ray Pacial ⁵ , Juan Antonio R. Magalang ⁵ , Marissa Alejandria ⁷ , Celia
9	Carlos ⁷ , Anna Ong-Lim ⁷ , Edsel Maurice Salvaña ⁷ , John Q. Wong ⁷ , Jaime C. Montoya ⁷ , and Maria Rosario
10	Singh-Vergeire ⁷
11	
12	* To whom correspondence should be addressed. Email address: <u>cpsaloma@up.edu.ph</u>
13	
14	¹ Core Facility for Bioinformatics, Philippine Genome Center, University of the Philippines System
15	² Philippine Genome Center, University of the Philippines System
16	³ DNA Sequencing Core Facility, Philippine Genome Center, University of the Philippines System
17	4 Clinical Genomics Laboratory, Philippine Genome Center, University of the Philippines System
18	⁵ Epidemiology Bureau, Department of Health, Philippines
19	⁶ Disease Prevention and Control Bureau, Department of Health, Philippines
20	⁷ Inter-Agency Task Force on Emerging Infectious Diseases (IATF) Task Force on COVID-19 Variants,
21	Department of Health, Philippines

22 ABSTRACT

We report the sequencing and detection of 39 Severe Acute Respiratory Syndrome Coronavirus 2 (SARS CoV-2) samples containing lineage-defining mutations specific to viruses belonging to the B.1.1.7 lineage
 (UK variant) in the Philippines.

26

27 ANNOUNCEMENT

The Coronavirus Disease 2019 (COVID-19) is an infectious disease that has gained pandemic status from the World Health Organization, with millions of cases and deaths recorded worldwide. This global health crisis is caused by viruses referred to as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), a member of the genus *Betacoronavirus* (Coronaviridae) together with the causative agents of the first SARS outbreak in 2003 and the Middle East Respiratory Syndrome (MERS) in 2012.

33

In this study, we present the genome sequences of 39 cases of COVID-19 in the Philippines caused by viruses belonging to SARS-CoV-2 lineage B.1.1.7, more commonly known as the United Kingdom (UK) variant. This particular SARS-CoV-2 variant was initially identified in the United Kingdom and has been reported to cause a surge of COVID-19 infections in the said country (Kirby, 2021). Initial studies also suggest that the B.1.1.7 viruses appear to have replicative advantage (Grabowski et al., 2021) and are more transmissible (Leung et al., 2021).

40

A total of 39 Philippine SARS-CoV-2 samples were classified under the B.1.1.7 lineage. **Table 1** shows the primary consensus assembly metrics for these samples. The average depth of coverage across all the sequences is 1,128x, with 33 of the samples carrying all 17 hallmark mutations of the B.1.1.7 lineage as listed in the PANGO Lineages report for the B.1.1.7 variant of concern (<u>https://cov-</u> <u>lineages.org/global report B.1.1.7.html</u>).

4	5
---	---

47 The detection of B.1.1.7 from returning overseas Filipino workers and in the community highlights the 48 need for genomic surveillance at the country's ports of entry and the general population to monitor the 49 importation and local transmission of emerging variants of concern that may impact the public health 50 response to the SARS-CoV-2 pandemic in the Philippines. 51 52 53 **METHODS** 54 In order to detect the entry of B.1.1.7 infection in the Philippines, nasopharyngeal swabs were collected 55 between December 10, 2020 and January 31, 2021 from COVID-19 cases detected in returning overseas 56 Filipinos, as well as from local case clusters mainly from the Cordillera Administrative Region of the 57 country, among others. The collected swab samples were subjected to RNA extraction using the QIAamp 58 Viral RNA Mini Kit, the product of which were then used as templates for the amplicon-based Illumina 59 COVIDSeq Test (https://www.illumina.com/products/by-type/ivd-products/covidseq.html) sequencing 60 workflow. Afterwards, the resulting sequence reads were mapped to the reference SARS-CoV-2 genome 61 (NCBI Accession Number NC 045512.2) using Minimap2 (Li, 2018) version 2.17-r941, with the '-x sr' 62 parameter for short accurate genomic reads alignment.

63

Primer clipping and quality trimming, intrahost variant calling, removal of reads associated with mismatched primer indices, and consensus sequence assembly were all done following the suggested workflow of iVar (Grubaugh et al., 2019) version 1.2.2, using default parameters. The consensus variants were identified by comparing the resulting assemblies with the reference sequence using MUMmer (Kurtz et al., 2004) as implemented in the RATT software (Otto et al., 2011). Lastly, SARS-CoV-2 lineage

69	classifications were assigned using Pangolin version 2.3.2 (github.com/cov-lineages/pangolin; Rambaut						
70	et al., 2020).						
71							
72							
73	PUBLIC AVAILABILITY AND ACCESSION NUMBERS						
74	The consensus sequence assemblies reported in this study are deposited in the $\underline{EpiCoV^{TM}}$ database of						
75	the GISAID. The accession codes are shown in Table 1 .						
76							
77							
78	ACKNOWLEDGEMENT						
79	This project was supported by a Genomics Biosurveillance grant from the Philippine Department of						
80	Health, and a Department of Science and Technology – Philippine Council for Health Research and						
81	Development grant to BM and the University of the Philippines. The authors would also like to thank the						
82	contributing institutions to the Philippine Genomic Biosurveillance Network.						
83							
84	The protocols used in this study were reviewed and approved by the Single Joint Research Ethics Board						
85	of the Department of Health, with approval code SJREB-2021-11, as part of a larger research program						
86	entitled "A retrospective study on the national genomic surveillance of COVID-19 transmission in the						
87	Philippines by SARS-CoV-2 genome sequencing and bioinformatics analysis".						

88 **REFERENCES**

- Grabowski F., Preibisch G., Kochanczyk M., and Lipniacki T. (2021). SARS-CoV-2 Variant Under
 Investigation 202012/01 has more than twofold replicative advantage. medRXiV.
 DOI: https://doi.org/10.1101/2020.12.28.20248906.
- 92 Grubaugh N., Gangavarapu K., Quick J., et al. (2019). An amplicon-based sequencing framework for
- accurately measuring intrahost virus diversity using PrimalSeq and iVar. Genome Biology. 20(8).
 DOI:https://doi.org/10.1186/s13059-018-1618-7.
- 95 Kirby T. (2021). New variant of SARS-COV-2 in UK causes surge of COVID-19. Lancet Respiratory
 96 Medicine. DOI:https://doi.org/10.1016/S2213-2600(21)00005-9.
- 97 Kurtz S., Phillippy A., Delcher A., Smoot M., Shumway M., Antonescu C., and Salzberg S. (2004). Versatile
- 98 and open software for comparing large genomes. Genome Biology. 5:R12.
- 99 Leung K., Shum M., Leung G., Lam T., and Wu J. (2021). Early transmissibility assessment of the N501Y
- 100 mutant strains of SARS-CoV-2 in the United Kingdom, October to November 2020. EuroSurveillance.

101 26(1):pii=2002106. DOI:https://doi.org/10.2807/1560-7917.ES.2020.26.1.2002106.

- Li H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 34:3094-3100.
- 103 doi:10.1093/bioinformatics/bty191.
- Otto T., Dillon G., Degrave W., and Berriman M. (2011). RATT: rapid annotation transfer tool. Nucleic
 Acids Research. 39(9):e57. DOI:https://doi.org/10.1093/nar/gkq1268.
- 106 Rambaut, A., Holmes, E.C., O'Toole, Á. et al. (2020). A dynamic nomenclature proposal for SARS-CoV-2
- 107 lineages to assist genomic epidemiology. Nature Microbiology. 5:1403–140.
 108 DOI:https://doi.org/10.1038/s41564-020-0770-5.
- Shu, Y., and McCauley, J. (2017). GISAID: Global initiative on sharing all influenza data from vision to
 reality. EuroSurveillance. 22(13). DOI:10.2807/1560-7917.ES.2017.22.13.30494.

Sample Code	GISAID Accession Code	Collection Date	% GC Content	Sample Source	Mean Coverage Depth	# Consensus SNPs	# B.1.1.7 SNPs	% N	Length
PH-PGC-00315	EPI ISL 1081793	29-Dec-20	37.32	ROF	1,201.27	52	17	1.76	29,884
PH-PGC-00317	<u>EPI ISL 1081794</u>	29-Dec-20	37.48	ROF	1,194.12	50	17	1.30	29,884
PH-PGC-00401	<u>EPI ISL 1081795</u>	10-Dec-20	37.22	ROF	996.07	51	17	2.12	29,884
PH-PGC-00986	<u>EPI ISL 1081796</u>	7-Jan-21	37.14	ROF	1,164.12	50	17	2.26	29,884
PH-PGC-02005	<u>EPI ISL 1081797</u>	4-Jan-21	31.86	CAR	508.19	44	14	16.82	29,884
PH-PGC-02008	<u>EPI ISL 1081798</u>	3-Jan-21	35.45	CAR	740.89	48	17	6.87	29,884
PH-PGC-02009	<u>EPI ISL 1081799</u>	3-Jan-21	34.31	CAR	689.44	45	14	10.19	29,884
PH-PGC-02033	<u>EPI ISL 1081800</u>	5-Jan-21	37.38	CAR	1,362.04	49	17	1.52	29,884
PH-PGC-02127	<u>EPI ISL 1081801</u>	7-Jan-21	33.83	CAR	973.77	45	14	11.36	29,884
PH-PGC-02131	<u>EPI ISL 1081802</u>	7-Jan-21	36.53	CAR	1,326.01	49	17	3.77	29,885
PH-PGC-02133	EPI_ISL_1081803	7-Jan-21	34.19	CAR	892.30	49	16	10.20	29,884
PH-PGC-02152	EPI_ISL_1081804	9-Jan-21	36.16	CAR	1,086.44	38	16	4.80	29,884
PH-PGC-02181	<u>EPI ISL 1081805</u>	8-Jan-21	36.40	CAR	882.19	50	17	4.18	29,884
PH-PGC-02183	EPI_ISL_1081806	8-Jan-21	37.06	CAR	1,210.23	51	17	2.42	29,884
PH-PGC-02184	EPI_ISL_1081807	8-Jan-21	35.20	CAR	891.29	48	16	7.54	29,884
PH-PGC-02185	<u>EPI ISL 1081808</u>	8-Jan-21	37.33	CAR	1,131.33	49	17	1.69	29,884
PH-PGC-02225	EPI_ISL_1081809	8-Jan-21	37.13	CAR	1,303.29	51	17	2.14	29,884
PH-PGC-02408	EPI_ISL_1081810	7-Jan-21	37.92	ROF	1,652.89	51	17	0.10	29,884
PH-PGC-02434	<u>EPI ISL 1081811</u>	12-Jan-21	37.70	ROF	1,168.26	53	17	0.78	29,884
PH-PGC-02630	EPI_ISL_1081812	16-Jan-21	37.83	ROF	1,307.23	48	17	0.41	29,884
PH-PGC-02725	EPI_ISL_1081813	14-Jan-21	37.16	ROF	1,140.03	52	17	2.04	29,884
PH-PGC-02730	EPI ISL 1081814	16-Jan-21	37.87	ROF	1,620.34	52	17	0.13	29,884

Table 1. Primary consensus sequence assembly metrics.

ROF – Returning Overseas Filipino

CAR – Cordillera Administrative Region

NCR – National Capital Region

NM – Northern Mindanao

% N – Percentage of ambiguous base calls ("N" content) in the consensus sequence assembly. High % N values lead to lower % GC content.

Sample Code	NCBI Accession Code	Collection Date	% GC Content	Sample Source	Mean Coverage Depth	# Consensus SNPs	# B.1.1.7 SNPs	% N	Length
PH-PGC-02732	<u>EPI ISL 1081815</u>	17-Jan-21	37.89	ROF	724.58	52	17	0.25	29,884
PH-PGC-02733	<u>EPI ISL 1081816</u>	17-Jan-21	37.88	ROF	1,499.29	51	17	0.15	29,884
PH-PGC-02740	<u>EPI ISL 1081817</u>	19-Jan-21	32.63	ROF	769.66	30	12	14.07	29,894
PH-PGC-02745	<u>EPI ISL 1081818</u>	19-Jan-21	30.64	ROF	679.99	32	12	19.71	29,894
PH-PGC-02756	<u>EPI ISL 1081819</u>	15-Jan-21	37.01	CAR	1,037.94	50	17	2.23	29,884
PH-PGC-02770	<u>EPI ISL 1081820</u>	15-Jan-21	35.34	ROF	971.34	44	14	6.34	29,885
PH-PGC-02793	<u>EPI ISL 1081821</u>	19-Jan-21	37.65	ROF	1,290.90	50	17	0.90	29,884
PH-PGC-02812	<u>EPI ISL 1081822</u>	24-Jan-21	37.91	CAR	1,654.09	50	17	0.18	29,884
PH-PGC-02826	<u>EPI ISL 1081823</u>	21-Jan-21	37.85	CAR	1,346.54	52	17	0.31	29,884
PH-PGC-02845	<u>EPI ISL 1081824</u>	13-Jan-21	37.74	CAR	1,454.34	51	17	0.58	29,884
PH-PGC-02851	<u>EPI ISL 1081825</u>	11-Jan-21	37.87	CAR	1,563.14	49	17	0.28	29,884
PH-PGC-02886	<u>EPI ISL 1081826</u>	16-Jan-21	37.67	CAR	1,411.43	51	17	0.86	29,884
PH-PGC-03690	<u>EPI ISL 1081829</u>	22-Dec-20	19.13	Calabarzon	421.38	20	5	50.35	29,900
PH-PGC-03846	<u>EPI ISL 1081830</u>	24-Jan-21	37.87	ROF	1,458.88	49	17	0.16	29,884
PH-PGC-03939	<u>EPI ISL 1081831</u>	31-Jan-21	37.89	ROF	2,082.45	50	16	0.27	29,885
PH-PGC-03978	<u>EPI ISL 1081832</u>	25-Jan-21	37.25	NCR	990.68	53	16	2.01	29,884
PH-PGC-03995	EPI_ISL_1081841	25-Jan-21	23.29	NM	206.76	22	7	37.77	29,896

ROF – Returning Overseas Filipino

CAR – Cordillera Administrative Region

NCR – National Capital Region

NM – Northern Mindanao

% N – Percentage of ambiguous base calls ("N" content) in the consensus sequence assembly. High % N values lead to lower % GC content.