

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

COVID-19 and Cognitive Impairment: Severity, Evolution, and Functional Impact during Inpatient Rehabilitation

Ruchi Patel MA OTR/L¹, Irene Savrides MA OTR/L¹, Christine Cahalan MS OTR/L¹, Gargi Doulatani MA², Michael W. O'Dell MD^{1,2}, Joan Toglia PhD OTR/L^{1,2,4}, & Abhishek Jaywant, PhD^{1,2,3}

¹Department of Rehabilitation Medicine, NewYork-Presbyterian Hospital/Weill Cornell Medical Center

²Department of Rehabilitation Medicine, Weill Cornell Medicine

³Department of Psychiatry, Weill Cornell Medicine

⁴School of Health and Natural Sciences, Mercy College

Manuscript Word Count: 1196 words

Corresponding Author:

Abhishek Jaywant, PhD

525 E 68th St, F-1232, New York NY 10065

Email: abj2006@med.cornell.edu

Tel: 857-225-5783

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

Abstract

Objective: To determine the frequency, magnitude, and change in cognitive impairment in patients with COVID-19 undergoing acute inpatient rehabilitation. We secondarily evaluated correlates of cognitive impairment and the relationship between cognition and functional gain.

Design: Cross-sectional observational study with assessments at admission and discharge

Setting: Acute inpatient rehabilitation unit within a large, urban academic medical center

Participants: 77 patients hospitalized for COVID-19 and subsequently admitted to an inpatient rehabilitation unit between March-August 2020, 45 of whom were re-assessed at discharge.

Interventions: N/A

Main Outcome Measures: Montreal Cognitive Assessment (MoCA) scores on admission and discharge (when available) and Quality Indicator for Self-Care (QI-SC) scores on admission and discharge.

Results: 62/77 (80.5%) of patients demonstrated cognitive deficits on the MoCA at admission: 39/77 (50.6%) were mildly impaired, 20/77 (26%) moderately impaired, and 3/77 (3.9%) severely impaired. Cognitive impairment was associated with a prior history of delirium, but not age or length of acute care hospitalization. 32/45 (71.1%) patients with discharge scores improved and met the MoCA minimally clinically important difference (MCID); however, 35/45 (77.8%) continued to score in the impaired range. Patients who met the MoCA MCID demonstrated significantly greater QI-SC score gains than those that did not meet the MCID ($p=.02$).

Conclusion: Cognitive impairment is common among hospitalized COVID-19 patients requiring acute inpatient rehabilitation. Cognitive impairment improves over the course of inpatient

42 rehabilitation, and is associated with functional gain. Nonetheless, cognitive deficits frequently
43 remain present at discharge, indicating the need for systematic assessment and follow-up,
44 especially given the association with functional outcome.

45 **Key Words:** Coronavirus, SARS-CoV-2, Cognition, Activities of Daily Living, Rehabilitation

46 **1.Introduction**

47 Acute and critical illness frequently result in cognitive impairment, affecting function and quality
48 of life¹. COVID-19 can result in respiratory distress, hypoxia, ICU admission,
49 intubation/ventilation, delirium, and vascular complications, all associated with cognitive
50 impairment². Cognitive deficits lead to dependence in activities of daily living³ (ADLs),
51 prompting the need for assessing and treating cognitive dysfunction to optimize function.

52 Studies on COVID-19 show a high frequency of cognitive impairment in mild to severe disease
53 expressions. Among community-dwelling patients with mild cases of COVID-19, 21% of
54 patients experienced a decline in cognitive functioning⁴. 81% of hospitalized but post-acute
55 COVID-19 patients score abnormally on neuropsychological measures of attention, processing
56 speed, and executive function⁵. There is still limited knowledge on how cognitive deficits evolve
57 and impact functional outcome. One study found 80% of COVID-19 patients on a rehabilitation
58 unit had cognitive deficits, with continued impairment in over half of the patients at 1 month
59 follow-up⁶; however, the relationship to functional gain was not evaluated. This study aims to
60 determine the magnitude and change in cognitive impairment, correlates of cognitive
61 impairment, and the relation between cognition and function, in a relatively large sample of
62 COVID-19 patients admitted to inpatient rehabilitation.

63

64 **2.Methods**

65 94 patients hospitalized for COVID-19 were admitted to an inpatient rehabilitation unit (IRU)
66 from March-August 2020. Seventeen patients were not administered our cognitive measure as
67 they were either deemed as not exhibiting cognitive impairment during instrumental ADLs or

68 underwent other cognitive testing, resulting in a final sample of 77. Patients were medically
69 stable, able to tolerate 3 hours/day of therapy, and had potential to gain function. Infection
70 control guidelines were followed, allowing for admission of positive COVID-19 patients and in-
71 person administration of our cognitive measure. All data were extracted from retrospective chart
72 reviews under approval of the Weill Cornell Medicine Institutional Review Board. 19/77 patients
73 underwent a separate neuropsychological evaluation during their admission, the results of which
74 have been published⁵; the results described here are novel.

75 Demographic/clinical data extracted included age, gender, prior level of function, intubation
76 history and length, delirium history, and IRU and acute care lengths of stay (LOS) (Table 1).

77 The Montreal Cognitive Assessment (MoCA) was administered by occupational therapists (OT)
78 72 hours before/after admission to the IRU. The minimal clinical important difference (MCID)
79 of 3 points was used⁷. Scores range from 0-30 and are stratified as follows: ≥ 26 =normal, 18-
80 25=mild impairment, 11-17=moderate impairment, and ≤ 10 =severe impairment⁸. Admission
81 MoCA assessments were administered on all 77 subjects and discharge MoCA assessments on
82 45/77. Three patients received the Blind MoCA, which was converted to 30-point scale per
83 established guidelines⁸. 32 patients did not receive discharge assessment as they either scored
84 normal on the initial assessment, were unavailable, and/or declined.

85 The Quality Indicator for Self-Care (QI-SC) is a standard of care functional assessment for
86 inpatient rehabilitation facilities. Items scored on a scale of 1 to 6 include eating, grooming,
87 toileting, bathing, upper body dressing, lower body dressing, and donning/doffing footwear.
88 Total score ranges from 7, indicating total assistance, to 42, indicating independence⁹. The QI-
89 SC is reliable and discriminates between the quality of care among rehabilitation facilities⁹. OTs
90 assigned admission and discharge QI-SC scores to all participants.

91 Statistical analyses used non-parametric tests to evaluate group differences (Mann-Whitney U),
92 within-group change from admission to discharge (Wilcoxon Signed-Ranks Test), association
93 between continuous variables (Spearman rank-order correlations), and chi-square tests for
94 categorical variables. Effect sizes are reported as Cohen's d.

95

96 **3.Results**

97 80.5% of patients demonstrated cognitive deficits on the MoCA at admission, of whom 15/77
98 were in the normal range, 39/77 had mild deficits, 20/77 had moderate deficits, and 3/77 had
99 severe deficits (Figure 1). Admission MoCA scores were higher for patients who did not
100 previously have delirium ($M=22.03$, $SD=4.64$) than patients who previously had delirium
101 ($M=18.95$, $SD=5.8$), $U=466.5$, $p=.02$, Cohen's $d=.58$. Admission MoCA did not differ based on
102 whether a patient was intubated ($U=454.5$, $p=.82$), age ($r_s=-.15$, $p=.19$) or acute hospitalization
103 length of stay ($r_s=-.08$, $p=.50$).

104 Patients with discharge MoCA scores ($n=45$) did not differ from those who were not
105 administered the MoCA at discharge ($n=32$) in age, gender, prior neurologic history, prior
106 delirium, prior intubation, or prior hypoxia (all p 's $>.05$). These 45 patients with admission and
107 discharge MoCA scores improved on the MoCA ($M=4.02$, $SD=3.67$, $Z=5.08$, $p<.001$, Cohen's
108 $d=1.10$) and QI-SC ($M=12.2$, $SD=6.18$, $Z=5.47$, $p<.001$, Cohen's $d=1.97$). 32/45 patients
109 improved and met the MoCA MCID at discharge. This group of patients had a greater increase in
110 QI-SC scores ($M=13.88$, $SD=5.01$) versus patients who did not meet the MCID or declined
111 ($M=9.46$, $SD=7.79$), Mann-Whitney $U=113.0$, $p=.02$, Cohen's $d=0.75$, a relationship that
112 persisted in a linear regression model accounting for age and delirium. Despite overall

113 improvement, 28/45 patients still had mild deficits on the MoCA at discharge, 7 had moderate
114 deficits, none had severe deficits, and 10 scored in the normal range.

115

116 **4.Discussion**

117 Among 77 COVID-19 patients undergoing acute inpatient rehabilitation, 80.5% of patients
118 demonstrated cognitive impairment, the majority of whom had mild deficits. This number is
119 consistent with studies using neuropsychological assessments to evaluate cognition in
120 hospitalized COVID-19 patients⁵⁻⁶ and expands on these findings with a larger sample than
121 previous studies. Consistent with the literature, delirium during acute hospitalization is
122 associated with greater cognitive impairment at admission¹⁰, emphasizing the importance of
123 early detection and management of delirium in COVID-19. The lack of association between
124 admission cognitive impairment and intubation is also consistent with prior work.⁵

125 COVID-19 patients exhibited significant improvement in cognition with a large effect size.
126 32/45 (71.1%) percent exhibited clinically meaningful improvement, possibly reflecting natural
127 recovery, improvement in residual acute mental status changes, and/or rehabilitation intervention
128 effects. Mean QI-SC scores for this subgroup improved with a large effect size, signifying that
129 patients required less assistance for basic ADLs. Notably, significantly greater functional
130 improvement was seen in patients who exhibited a clinically meaningful improvement in
131 cognition than those that did not. Cognitive ability is associated with short- and long-term
132 independence with self-care tasks in older adults³ and our results reinforce the importance of
133 targeting cognition to improve functional gains in COVID-19. COVID-19 patients warrant
134 further follow-up as a majority continued to demonstrate mild deficits at discharge.

135 *4.1. Study Limitations*

136 Discharge cognitive data was not available for 32/77 patients, limiting the ability to determine
137 the true prevalence of cognitive deficits in our sample, although there were no
138 demographic/clinical differences between the 45 with and 32 without discharge cognitive data.
139 Further research is necessary regarding the underlying etiology of the cognitive deficits as
140 imaging results and the effects of psychiatric illness⁵ were not addressed in this sample. Finally,
141 this sample included a relatively sick group of patients from the first wave of COVID-19;
142 findings may not apply to milder disease presentations or from subsequent waves of illness.

143

144 **5. Conclusion**

145 This study confirms the presence of cognitive impairment in COVID-19 patients with prolonged
146 hospitalization requiring inpatient rehabilitation and provides new insights into the evolution of
147 cognitive status. As most patients demonstrated mild/moderate cognitive deficits on admission,
148 routine cognitive screening should be conducted for COVID-19 patients. Despite improvement
149 during inpatient rehabilitation, a majority demonstrated mild cognitive deficits at discharge,
150 indicating ongoing cognitive needs. As cognitive improvement is associated with functional
151 gain, interventions targeting cognitive dysfunction in the acute inpatient rehabilitation setting
152 may be important in optimizing independence.

153

154

155

156 **Acknowledgements:** The authors received no funding to complete this study. All authors have
157 no conflicts or competing interests to declare. AJ is funded through a Career Development
158 Award from the National Center of Medical Rehabilitation Research/ Eunice Kennedy Shriver
159 National Institute of Child Health and Human Development (1K12HD093427-04), via a
160 subaward from Georgetown University. We thank the staff of the IRU for assisting in cognitive
161 and functional assessments, and for caring for, our COVID-19 patients.

162

163 **References**

164 1. Ehlenbach WJ, Hough CL, Crane PK, et al. Association between acute care and critical illness
165 hospitalization and cognitive function in older adults. *JAMA*. 2010;303(8):763–770.

166 doi:10.1001/jama.2010.167

167 2. Wilcox ME, Brummel NE, Archer K, Ely EW, Jackson JC, Hopkins RO. Cognitive
168 dysfunction in ICU patients: risk factors, predictors, and rehabilitation interventions. *Crit Care*
169 *Med*. 2013 Sep;41(9 Suppl 1): S81-98. doi: 10.1097/CCM.0b013e3182a16946. PMID:
170 23989098.

171 3. Millán-Calenti JC, Tubío, J., Pita-Fernández, S., Rochette, S., Lorenzo, T., Maseda, A.
172 Cognitive impairment as predictor of functional dependence in an elderly sample. *Archives of*
173 *Gerontology and Geriatrics*. 2012;54(1):197-201. <https://doi.org/10.1016/j.archger.2011.02.010>.

174 4. Del Brutto, O.H., Wu, S., Mera, R.M., Costa, A.F., Recalde, B.Y. and Issa, N.P. Cognitive
175 decline among individuals with history of mild symptomatic SARS-CoV-2 infection: A
176 longitudinal prospective study nested to a population cohort. *Eur J Neurol*. 2021

177 <https://doi.org/10.1111/ene.14775> Online ahead of print.

- 178 5. Jaywant A, Vanderlind WM, Alexopoulos GS, Fridman CB, Perlis RH, Gunning FM.
179 Frequency and profile of objective cognitive deficits in hospitalized patients recovering from
180 COVID-19. *Neuropsychopharmacology*. 2021 Feb 15:1–6. doi: 10.1038/s41386-021-00978-8.
181 Epub ahead of print. PMID: 33589778; PMCID: PMC7884062.
- 182 6. Alemanno F, Houdayer E, Parma A, Spina A, Del Forno A, Scatolini A, Angelone S,
183 Brugliera L, Tettamanti A, Beretta L, Iannaccone S. COVID-19 cognitive deficits after
184 respiratory assistance in the subacute phase: A COVID-rehabilitation unit experience. *PLoS One*.
185 2021 Feb 8;16(2):e0246590. doi: 10.1371/journal.pone.0246590. PMID: 33556127; PMCID:
186 PMC7870071. eCollection 2021.
- 187 7. Wu CY, Hung SJ, Lin KC, Chen KH, Chen P, Tsay PK. Responsiveness, minimal clinically
188 important difference, and validity of the MoCA in stroke rehabilitation. *Occup Ther Int*. 2019
189 Apr 14:2517658. doi: 10.1155/2019/2517658. PMID: 31097928; PMCID: PMC6487084.
- 190 8. www.mocatest.org Accessed March 1st, 2021.
- 191 9. Pardasaney PK, Deutsch A, Iriundo-Perez J, Ingber MJ, McMullen T. Measuring inpatient
192 rehabilitation facility quality of care: Discharge self-care functional status quality measure. *Arch*
193 *Phys Med Rehabil*. 2018 Jun;99(6):1035-1041. doi: 10.1016/j.apmr.2017.02.023. Epub 2017
194 Mar 28. PMID: 28363701.
- 195 10. Hopkins RO, Jackson JC. Short- and long-term cognitive outcomes in intensive care unit
196 survivors. *Clin Chest Med*. 2009 Mar;30(1):143-53, ix. doi: 10.1016/j.ccm.2008.11.001. PMID:
197 19186286.
- 198

Table 1. Demographics and clinical characteristics of main sample and subsample with discharge MoCA. Patients without discharge MoCA are included for comparison.			
	N=77 (Full Sample)	N= 45 (Patients with discharge MoCA)	N=32 (Patients without a discharge MoCA)
Gender			
Male	49 (63.6%)	30 (66.7%)	19 (59.4%)
Female	28 (36.4%)	15 (33.3%)	13 (40.6%)
Age	61.03 (15.67)	61.98 (14.11)	59.69 (17.77)
Acute hospitalization length of stay	37.03 (31.8)	39.4 (34.11)	33.69 (28.4)
Inpatient rehabilitation unit length of stay	13.43 (6.47)	14.44 (6.06)	12 (6.84)
Documented delirium	44 (57.1%)	28 (62.2%)	16 (50%)
Intubation	59 (76.6%)	35 (77.8%)	24 (75%)
Intubation length	13.31 (10.53)	13.75 (10.03)	12.67 (11.37)
Documented hypoxia	69 (89.6%)	39 (86.7%)	30 (93.8%)
History of neurological diagnosis	2 (2.6%)	1 (2.2%)	1 (3.1%)
Initial MoCA scores	20.29 (5.47)	17.96 (4.71)	23.47 (5.06)
Discharge MoCA scores		22.02 (4.23)	
Initial QI-SC score	25.75 (6.21)	24.96 (4.94)	26.88 (7.59)
Discharge QI-SC score	36.57 (8.37)	37.11 (6.97)	35.81 (10.09)


199 Note. MoCA = Montreal Cognitive Assessment; QI-SC = Quality Indicator for Self-Care; All
200 values are N (%) or Mean (SD).

201

202 **Figure 1.** Classification of cognitive impairment by score on the Montreal Cognitive Assessment
203 (MoCA) at admission in the full sample, admission in the subsample with discharge scores, and
204 discharge in the latter sample.

205

206



207