Essential Emergency and Critical Care – a consensus among global clinical experts

Carl Otto Schell,^{1,2,3*} Karima Khalid,^{4,5} Alexandra Wharton-Smith,⁶ Jacquie Narotso Oliwa,^{7,8} Hendry Robert Sawe,⁹ Nobhojit Roy,^{1,10,11} Alex Sanga,¹² John C. Marshall,¹³ Jamie Rylance,¹⁴ Claudia Hanson,^{1,15} Raphael Kazidule Kayambankadzanja,^{16,17} Lee A. Wallis,¹⁸ Maria Jirwe,¹⁹ Tim Baker,^{1,5,20} and the EECC Collaborators**

- 1. Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden
- 2. Centre for Clinical Research Sörmland, Uppsala University, Eskilstuna, Sweden
- 3. Department of Medicine Nyköping Hospital, Nyköping, Sweden
- 4. Muhimbili University of Allied Sciences, Dar es Salaam, Tanzania
- 5. Ifakara Health Institute, Dar es Salaam, Tanzania
- 6. Department of Global Health and Development, Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, UK
- 7. Health Services Unit, KEMRI-Wellcome Trust Research Programme, Nairobi, Kenya
- 8. Department of Paediatrics & Child Health, University of Nairobi, Nairobi, Kenya
- 9. Emergency Medicine Department, Muhimbili University of Allied Sciences, Dar es Salaam, Tanzania
- 10. The George Institute of Global Health, New Delhi, India
- 11. WHO Collaborating Centre for Research in Surgical Care Delivery in LMICs, BARC hospital, Mumbai, India
- 12. Ministry of Health, Community Development, Gender, Elderly and Children, Dodoma, Tanzania
- 13. Departments of Surgery and Critical Care Medicine, University of Toronto, Toronto, Canada
- 14. Malawi Liverpool Wellcome Trust, Blantyre, Malawi
- 15. Department of Disease control, Faculty of Infectious and Tropical disease London School of Hygiene and Tropical Medicine, UK
- 16. Department of Anaesthesia and Intensive Care, Queen Elizabeth Central Hospital, Blantyre, Malawi
- 17. School of Public Health and Family Medicine, College of Medicine, Blantyre, Malawi
- 18. Division of Emergency Medicine, University of Cape Town, South Africa
- 19. Department of Health Sciences, Red Cross University College, Huddinge, Sweden
- 20. Department of Clinical Research, London School of Hygiene and Tropical Medicine, UK

* Corresponding Author

Carl Otto Schell, Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden <u>Carl.Schell@ki.se</u>

** The EECC Collaborators

Adam Asghar	Ben Morton
Adam Laytin	Bhakti Sarang,
Adrian J. Holloway	Bharath Kumar Tirupakuzhi Vijayaraghavan,
Ahmed Rhassane El Adib	Bobby King
Alexia Michaelides	Brian Rice
Alvaro Coronado Munoz	C Louise Thwaites
Amos Muzuka	Chian Wern Tai
Analía Fernández	Christian Owoo
Andrea B Pembe	Cornelius Sendagire
Andreas Wellhagen	Dan Brun Petersen
Andrew G. Smith	Daniel Tatay
Anita Gadgil	David Lee Skinner
Anna Hvarfner	Denis Kinyua
Anuja Abayadeera	Dhruva Ghosh
Asya Agulnik	Diptesh Aryal
Aurélie Godard	Donald Mlombwa
Boot E enhanced int reports new research that has not been certified	by been review and should not be used to guide clinical practice.
Bargo Mahamat Yousif	Edwin R. Lugazia

Elisabeth Riviello Elizabeth M Molyneux Ellena Heyns **Emmanuel Fru Nsutebu** Erika Montalvo Ernesto Gerardo Moreno Esther Banda Kanyangira **Fiona Muttalib** Francis Mupeta Franco Diaz Fred Bulamba Furaha Nzanzu Blaise Pascal Gavin Wooldridge Gibonce Mwakisambwe Guy A. Richards Hala Ammar Halinder S. Mangat Hasanein H. Ghali Hiral A. Shah Hoi Ping Shum Ibrahim Salim Abdullahi Ignacio Martin-Loeches Ingrid T. Von Der Osten Jacob Mcknight James S. Lee Jane Kasozi Namagga Jaran Eriksen Jasmine Armour-Marshall John Kellett John Z Metcalfe Jolene Moore Jonas Blixt Josephine Langton Juan Gutierrez Mejia Juan Ignacio Silesky-Jiménez Kapil Dev Soni Karl Martin Kohne Kathryn Rowan Kazuhiro Yokobatake Kent Doi Kristina Rudd Kwame Asante Akuamoah-Boateng Lars Irestedt Lia I. Losonczy Lina Zhang Lisa Kurland Lorna Guinness Lovenish Bains Margaret Nyaika Markus Castegren Matthew Loftus Matti Reinikainen Megan Cox Mervyn Mer Michael Jaung Michael S. Lipnick Mike English

Miklos Lipcey Monty Khajanchi Mpoki Ulisubisya Mulinda Nyirenda Märit Amanda Halmin Naman Shah Natalie L. Cobb Nathan D. Nielsen Neill KJ Adhikari Neville Vlok Nicholas Risko Niranjan Kissoon Ntogwiachu Daniel Kobuh Oscar Fernández Rostello Patricia Duque Paul D Sonenthal Paul Patrick Mwasapi Peter Baker Petronella Bjurling-Sjöberg **Piedad Sarmiento** Pryanka Relan **Raymond Towey Rebecca Silvers** Rehema Mlay **Rich Branson Richard J Wang Richard Kojan Richard Peter Von Rahden Richard Venn Rob Mac Sweeney** Rodrigo Genaro Arduini **Rodwell Gundo Ruyumbu Sixtus** Samson Kwazizira Mndolo Samuel Akech Shada Rouhani Siriel Nanzia Massawe Stefan Swartling Peterson Steven A Webb Sunkaru Touray Susana Guido Teresa Kortz Theodoros Aslanidis **Thomas Weiser** Tom Bashford Traci A. Wolbrink Usha Lalla V. Theodore Barnett Vijay Christopher Kannan Vincent loos Waleed S. Eldebsy Wangari Waweru-Siika Wezzie Kumwenda Mwafulirwa William Obeng Wim Van Damme Yasein Omer Zione Banda

Abstract

Background

Globally, critical illness results in millions of deaths every year. Although many of these deaths are potentially preventable, the basic, life-saving care of critically ill patients are often overlooked in health systems. Essential Emergency and Critical Care (EECC) has been devised as the care that should be provided to all critically ill patients in all hospitals in the world. EECC includes the effective care of low cost and low complexity for the identification and timely treatment of critically ill patients across all medical specialities. This study aimed to specify the content of EECC and additionally, given the surge of critical illness in the ongoing pandemic, the essential diagnosis-specific care for critically ill patients with COVID-19.

Methods

A Delphi process was conducted to seek consensus (>90% agreement) in a diverse panel of global clinical experts. The panel was asked to iteratively rate proposed treatments and actions based on previous guidelines and the WHO/ICRC's Basic Emergency Care. The output from the Delphi was adapted iteratively with specialist reviewers into a coherent and feasible EECC package of clinical processes plus a list of hospital resource requirements.

Results

The 269 experts in the Delphi panel had clinical experience in different acute medical specialties from 59 countries and from all resource settings. The agreed EECC package contains 40 clinical processes and 67 hospital readiness requirements. The essential diagnosis-specific care of critically ill COVID-19 patients has an additional 7 clinical processes and 9 hospital readiness requirements.

Conclusion

The study has specified the content of the essential emergency and critical care that should be provided to all critically ill patients. Implementation of EECC could be an effective strategy to reduce preventable deaths worldwide. As critically ill patients have high mortality rates, especially where trained staff or resources are limited, even small improvements would have a large impact on survival. EECC has a vital role in the effective scale-up of oxygen and other care for critically ill patients in the COVID-19 pandemic. Policy makers should prioritise EECC, increase its coverage in hospitals, and include EECC as a component of universal health coverage.

Key words

Critical illness, Health Priorities, Health systems, Critical Care, Emergency care, Quality of Health Care, Triage, Oxygen, Global Health, COVID-19

Introduction

Critical illness, when defined as a state of ill health with vital organ dysfunction and a high risk of imminent death, is common in hospitals throughout the world [1-6]. It is the most severe form of acute illness due to any underlying condition and results in millions of deaths globally every year [1, 5]. The COVID-19 pandemic has led to increased morbidity and mortality with a surge in critical illness worldwide [7-9].

Many of the deaths due to critical illness are potentially preventable [10-12]. In critical illness, the patient's airway, breathing, or circulation may become compromised, and early identification of the problem and timely care can be lifesaving. Unfortunately, this care is frequently a neglected part of healthcare. The basic, life-saving clinical processes may be overlooked in specialised care [12] and in settings of both high [13-15] and low resources [16-18]. In hospitals all over the world, guidelines, equipment, and routines focusing on the care of critically ill patients, are often missing for adult[19] and paediatric patients[11], in emergency units [20], in wards [21] and in intensive care units [22]. Improving the way healthcare manages critical illness could save many lives [11, 23, 24].

To improve outcomes for critically ill patients by means that are feasible to deliver in all hospital wards and settings, the Essential Emergency and Critical Care (EECC) concept was devised [25]. EECC is defined as the care that should be provided to all critically ill patients of all ages in all hospitals in the world. It is distinguished by three principles. First, priority to those with the most urgent clinical need, including both early identification and timely care. Second, provision of the life-saving treatments that support and stabilise failing vital organ functions. And third, a focus on effective care of low cost and low complexity.

The clinical processes that comprise the essential care of critically ill patients, and the resources required for those processes have not previously been specified. As critically ill patients can be suffering from any underlying condition, EECC is conceptualised to be integrated into all acute clinical specialties. We therefore sought consensus among a diverse group of global clinical experts with the aim of specifying the content of EECC. An additional aim, given the ongoing pandemic, was consensus around the essential diagnosis-specific care for critically ill patients with COVID-19.

Methods

The study used three phases. (Figure 1) Firstly, a consensus was sought about the treatments and actions (T&A) in EECC using a modified Delphi technique [26]. Secondly, the output from the Delphi was adjusted into a coherent, user-friendly, and feasible package of clinical processes. And thirdly, a list of requirements for hospitals to be ready to provide the care was developed.



Figure 1. Summary of the process

EECC Essential Emergency and critical care T&A Treatment and actions WHO World Health Organisation

Phase One

An online, three-round modified Delphi process was conducted in November and December 2020. The Delphi method uses anonymous responses from an expert panel to iteratively posed questions and controlled feedback to reach consensus on the topic of interest [26]. A Delphi process was chosen for this study as EECC is new, its content has not been previously specified and a large group of diverse experts was required.

To be part of the panel, experts needed to have clinical experience of caring for critically ill patients. To ensure the involvement of a diverse range of experts, it was decided that at least 50% of the invitations to participate in the panel should be sent to experts with substantial experience of working in low- and middle-income countries, and there should be a balance between clinical experience (work in general wards, emergency units, intensive care units);

specialty (paediatrics, obstetrics, medicine, surgery, intensive care, anaesthesia and emergency care); profession (doctors, nurses, other health professionals); location, and gender. A list of potential participants was made from a mapping of stakeholders, the literature across all acute medical specialties, the researchers' networks and additional purposive and snowball sampling for under-represented groups. Additionally, a link to a screening survey was sent to global professional networks, specialist societies and on social media to identify further potential participants. A total of 895 experts were invited to participate, and those who accepted provided written informed consent.

EECC consists of clinical processes of care. To enable rating by the Delphi panel, clinical processes were disassembled into individual treatments & actions (T&A). The T&A concern the identification of critical illness; care of critical illness, and the diagnosis-specific care of critically ill COVID-19 patients. To be included, all T&A were required to meet two a-priori defined criteria: *effectiveness** and *feasibility**. Additionally, *universality** was required for the identification and care of critical illness and *relevance** was required for the diagnosis-specific care of critically ill patients with COVID-19. A draft list of potential T&A was developed based on clinical guidelines and tools from related specialties [27-38] and aligned with the WHO/ICRC's Basic Emergency Care [39]. The draft list was revised by specialist reviewers – a group of senior clinicians, researchers, and policy makers, with expertise in paediatrics, medicine, emergency medicine, anaesthesia and intensive care, critical care nursing, obstetrics and gynaecology, and surgery.

Three Delphi rounds were deemed sufficient to address the aim while avoiding attrition and poor response rates. A four-point Likert scale (strongly disagree, disagree, agree, strongly agree) with a 'do-not know' option was used for the panel to rate their opinion about the inclusion of each T&A in EECC [40-42]. Consensus was achieved when more than 90% of respondents selected "agree" or "strongly agree", excluding "don't know" responses. The experts were able to provide free-text comments, which were analysed to identify appropriate, relevant changes to the wording of T&A for clarity of understanding, and to identify newly proposed T&A. After the first round, newly proposed T&A that fulfilled the EECC criteria for potential inclusion were revised after input from the specialist reviewers and included for assessment by the panel. T&A that did not reach consensus in the previous round were presented for re-assessment in rounds two and three, together with a visual representation of the spread of previous responses.

As the Delphi panel was diverse, it was considered that there may be different opinions about the inclusion of T&As between experts with particular a-priori defined characteristics. These subgroups of experts were those with work experience in a low-income country or not; those who are doctors or not; those with clinical experience in emergency care and those without;

^{*} *Effectiveness:* Established or proven to be safe and to reduce mortality. (compression to stop bleeding is effective; treating with leech therapy is not)

Feasibility: Low-cost and low complexity. Possible to provide in a low-staffed, low-resourced setting without the immediate presence of a doctor (placing a comatose patient in the recovery position (lateral position) is feasible; continuous haemodialysis is not).

Universality: Supports vital organ function rather than being the definitive care of a diagnosis. (IV fluids for shock are universal; thrombolytic therapy is not).

Relevance: Established or proven to be a treatment for COVID-19.

and those with clinical experience in intensive care and those without. The levels of agreement in each subgroup were assessed and presented for all the T&As that reached consensus.

Phase Two

After the Delphi, slight adjustments were made to the wording of the T&A that had reached consensus to ensure language consistency. The T&A were reassembled back into clinical processes to increase overall coherence and feasibility of the EECC package, with the goal of user-friendliness for health system implementation and quality improvement work. The adjustments were done in an iterative process with the same specialist reviewers as in Phase One to ensure relevance for all acute medical specialties. The final package of clinical processes was organised into those relevant for identification, for care, and general processes.

Phase Three

A provisional list of hospital readiness requirements for the provision of the clinical processes were developed using existing WHO tools, guidelines for related specialties, facility preparedness lists [29, 32, 34, 35, 37-39, 43, 44] and the experience and knowledge of the study team. The specialist reviewers provided iterative input into the provisional list, approving suggested items, adding relevant items from their clinical specialties, and suggesting modifications. Based on previous work and following consultation with health economists and procurement experts, the final list of requirements was agreed and arranged into eight categories: equipment, consumables, drugs, human resources, training, routines, guidelines, and infrastructure.

Results

Phase One

Of the 895 invited experts, 269 participated in the first round of the Delphi when the majority of the decisions were made (30% response rate). In Round Two, 228 experts participated (85% of those in Round One) and Round Three included 194 experts (85% of those in Round Two). The panel comprised experts from diverse resource settings, clinical settings, specialties, and professions **(Table 1)**. The panel included experts from 59 countries (**Figure 2**) and 38% were female.

	Number of	Proportion of
	experts	experts
	(N=269)	(%)
Resource setting*		
High-income country	139	52
Middle-income country	115	43
Low-income country	177	66
Do not know	2	1
Clinical setting*		
General ward	153	57
Emergency unit	179	67
High dependency unit	153	57
Intensive care unit	232	86
Operating theatre	102	38
Other	15	6
Specialty*		
Emergency Care	93	35
Intensive Care	190	71
Anaesthesia	59	22
Medicine	39	15
Surgery	20	7
Paediatrics	47	17
Obstetrics/Gynaecology	13	5
Other	25	9
Profession*		
Doctor	212	79
Nurse	40	15
Midwife	6	2
Clinical Officer	9	3
Other	17	6
Gender		
Female	102	38
Male	165	62

Table 1. The characteristics of the expert panel in the Delphi (first round)

*As the experts were asked to select all that apply, the sum of the percentages may exceed 100



Figure 2. Expert panel locations. The locations of the expert panel in the Delphi Round One

Of the 57 T&A for EECC in Round One, consensus was reached for 49. In Round Two, 29 newly proposed T&A were added to the eight remaining from Round One, of which two had been re-worded for clarity. Out of these 37, consensus was reached for 17. The remaining 20, of which another two had been re-worded for clarity, were included in Round Three. Consensus was reached for nine of the final 20 T&A. In total, consensus was reached for 75 out of 86 proposed T&A, including 54 of the original 57. (Supplementary table 1)

Of the seven T&A for the essential diagnosis-specific care of critically ill COVID-19 in Round One, all reached consensus for inclusion. In Round Two, two newly proposed T&A were added. Neither of these reached consensus in Round Two or Round Three.

Analyses of participant sub-groups did not reveal substantial divergence from the overall results. For the T&A that reached 90% agreement in the panel, agreement was not below 80% in any subgroup. (**Supplementary tables 2-4**)

Phase Two and Three

After the Delphi, the T&A that had reached consensus were reassembled into a final userfriendly and feasible package of EECC containing 40 clinical processes – 30 identification and care processes and 10 general processes. (**Panel 1**) All T&A for the care of critical illness were included, with some rewording and reordering. Eleven T&A for the identification of critical illness were not included, so that the package could be feasible for triage in all hospitals, and were added as an addendum (outside the remit of EECC), in order to underscore their importance in settings where staff have sufficient time and expertise.

The list of hospital readiness requirements for EECC contained 67 items, (fourteen for identification and 53 for essential care). (Panel 2)

The essential diagnosis-specific care of critically ill COVID-19 patients consisted of an additional seven clinical processes and nine hospital readiness requirements. (**Panel 3**)

PANEL 1. The clinical processes of Essential Emergency and Critical Care

IDENTIFICATION OF CRITICAL ILLNESS Critical illness is identified as soon as possible so timely care can be provided.
1. The hospital uses vital signs-based triage to identify critical illness
1.1 Triage/identification of critical illness includes the use of these vital signs
1.1.1 Pulse rate
1.1.2 Blood pressure
1.1.3 Respiratory rate
1.1.4 Oxygen saturation (SpO2)
1.1.5 Temperature
1.1.6 Level of consciousness (eg. "AVPU", "ACVPU" or Glasgow Coma Scale)
1.1.7 Presence of abnormal airway sounds heard from the bedside (eg. snoring, gurgling, stridor)
1.1.8 The overall condition of the patient (health worker's concern that the patient is critically ill)
1.2 Triage/identification of critical illness is conducted at these times
1.2.1 When a patient arrives at hospital seeking acute care
1.2.2 For hospital in-patients, at least every 24 hours, unless otherwise prescribed, with increased
frequency for patients who are at risk of becoming critically ill or who are critically ill, and then less frequently again when patients are stabilising
1.2.3 When a health worker, or the patient or guardian, is concerned that a patient may be critically ill
1.2.4 During and after surgery or anaesthesia
1.2.5 During and after transport/transfer of a patient who is critically ill or at risk of becoming critically ill
1.2.6 Following a treatment or action (re-evaluation)

PANEL 1 continued...

CARE OF CRITICAL ILLNESS

Essential care of processes when	f critical illness is initiated as soon as critical illness is identified and involves these clinical appropriate:
AIRWAY	 Placing the patient in the recovery position (lateral position) Age-appropriate airway positioning (eg. chin lift or jaw thrust in adults, neutral position in young children)
Care for a blocked or	4. Removal of any visible foreign body from the mouth or use of age-appropriate chest
threatened	5. Suction for secretions that are obstructing the airway
airway	6. Insertion of an oro-pharyngeal (Guedel) airway
BREATHING	7 Ontimizing the nation (so sitting up or prope)
Care for	 Oxygen therapy using nasal prongs, facemask, or mask with a reservoir bag (non re-
respiratory	breathing mask) 9 Bag-valve-mask ventilation in threatened or manifest respiratory arrest
distress	
CIRCULATION	 Optimizing the patient position (eg. lying flat, head-down, raised-legs, lateral tilt in pregnancy)
Care for a	11. Compression and elevation to stop bleeding
threatened	12. Appropriate bolus of intravenous fluid 13. Oral rehydration solution or other appropriate oral fluids for dehydration without shock
shock	14. Intramuscular adrenaline for anaphylaxis
	15. Uterine massage and/or oxytocin when indicated
REDUCED	16. Treating an unconscious patient as having a threatened airway (eg. recovery position
LEVEL	17. Dextrose (iv or buccal) in unconsciousness or seizures unless bedside blood glucose
Cana fam a	testing rules out hypoglycaemia or there is a clear alternative cause
reduced level	19. Quick-acting anti-seizure medication (eg. intravenous/rectal diazepam, or magnesium
of	sulphate in pregnancy/post-partum)
consciousness	
	21. Insertion of an intravenous cannula when critical illness is identified
OTHER CARE	22. Insertion of an intraosseous cannula, if indicated, if an intravenous cannula is not
IN EECC	23. Stabilizing the cervical spine in possible cervical spine injury
	24. Appropriate antibiotics for sepsis
Other	26. Keeping the patient warm using blankets and other means (including skin-to-skin care
immediate or	for babies)
ongoing care	27. Feeding (including breastreeding for bables), naso-gastric feeding and dextrose for nutrition and to avoid hypoglycaemia
illness	 Prevention of delirium (eg. sleep hygiene, provision of the patient's glasses or hearing aid)
	29. Regular turning of immobilised patients
	30. Mobilising the patient as early as possible

PANEL 1 continued...

GENERAL PROCESSES

Care is provided according to these general processes:

- 1. Assistance from additional or senior staff is sought when a critically ill patient is identified
- 2. Essential Emergency and Critical Care (EECC) is respectful and patient-centred
- 3. EECC is provided without considering the patient's ability to pay
- 4. Critically ill patients are cared-for in locations that facilitate observation and care (eg. designated beds, a bay or a unit for critically ill patients)
- 5. Infection, Prevention and Control (IPC) measures are used including hand hygiene and separation of patients with a suspected or confirmed contagious disease from those without
- 6. Communication is clear, including:
 - Within the care team when a patient is identified as critically ill (eg. verbal communication, at staff handovers, visible colour-coding)
 - Within the care team about the planned EECC (eg. continue oxygen therapy, give intravenous fluids)
 - Documentation in the patient notes about the vital signs, when critical illness has been identified and the treatments and actions conducted
 - Effective and respectful communication with the patient and family
- 7. If there is poor response to treatment, or if the patient deteriorates, other indicated EECC clinical processes are used
- 8. Clinical processes are discontinued that are no longer indicated (eg. if a patient improves or if they are deemed to no longer be in the patient's best interest)
- 9. It is recognised when EECC alone is not sufficient to manage the critical illness
- 10. EECC is integrated with care that is outside the scope of EECC (eg. the need for prompt investigations, definitive treatment of underlying conditions including following disease-specific best-practice guidelines, end-of-life care, referral)

Addendum: Extended identification of critical illness

To maintain feasibility of the EECC package, only a limited number of signs for the identification of critical illness are included. However, if time and expertise allow, there are additional signs that are not part of EECC that aid the identification of critical illness:

- Presence of respiratory distress (eg. unable to complete sentences; accessory muscle use; chest recessions; grunting or head nodding)
- Cyanosis
- Capillary refill time
- Cold or warm extremities
- Presence of severe dehydration (eg. decreased skin turgor; dry mucous membranes; sunken fontanelle)
- Confused, agitated or disoriented mental state
- Presence of prostration or lethargy
- Presence of a generalized seizure
- Inability to stand or walk without help
- Inability to breastfeed or feed in a young child
- Presence of severe acute malnutrition

PANEL 2. The Hospital Readiness Requirements for Essential Emergency and Critical Care

IDENTIFICATION OF CRITICAL ILLNESS

The following items are required for a hospital to be ready for the identification of critically ill patients:

CATEGORY	ITEM
1.1. EQUIPMENT	1.1.1 Clock with second hand
	1.1.2 Pulse oximeter & probe
	1.1.3 Blood pressure measuring equipment (eg. sphygmomanometer with a stethoscope)
	1.1.4 Blood pressure cuffs of different paediatric and adult sizes
	1.1.5 Light source (lamp or flashlight)
	1.1.6 Thermometer
1.2 CONSUMABLES	1.2.1 Soap or hand disinfectant
	1.2.2 Examination gloves
1.3 DRUGS	None
1.4 HUMAN	1.4.1 Health workers with the ability to identify critical illness 24h/day
RESOURCES	
1.5 TRAINING	1.5.1 The health workers are trained in the identification of critical illness
1.6 ROUTINES	1.6.1 Routines for the identification of critical illness
1.7 GUIDELINES	1.7.1 Guidelines for the identification of critical illness
1.8	1.8.1 Designated triage area (area for the identification of critical illness) in the Out-Patient
INFRASTRUCTURE	Department or Emergency Unit (area of the hospital where patients arrive)
	1.8.2 Running water

PANEL 2 continued ...

CARE OF CRITICAL ILLNESS

The following items are required for a hospital to be ready to provide the care of critically ill patients:

CATEGORY	ITEM
2.1 EQUIPMENT	2.1.1 Suction machine (electric or manual)
	2.1.3 Oxygen supply 24h/day (cylinder, concentrator (with electricity supply) or piped
	oxygen)
	2.1.4 Flow meter (if using cylinder or piped oxygen)
	2.1.5 Leak-free connectors from oxygen source to tubing
	2.1.6 Bag Valve Mask (resuscitator) – neonatal, paediatric and adult sizes
	2.1.7 Sharps disposal container
	2.1.8 External heat source
2.2 CONSUMABLES	2.2.1 Suction catheters of paediatric and adult sizes
	2.2.2 Guedel airways of paediatric and adult sizes
	2.2.3 Pillows
	2.2.4 Oxygen tubing
	2.2.5 Oxygen nasal prongs
	2.2.6 Oxygen face masks of paediatric and adult sizes
	2.2.7 Oxygen face masks with reservoir bags of paediatric and adult sizes
	2.2.8 Masks for Bag Valve Mask (resuscitator) – neonatal, paediatric and adult sizes
	2.2.9 Compression bandages
	2.2.10 Plasters or tape
	2.2.11 Gauze
	2.2.12 Intravenous cannulas of paediatric and adult sizes
	2.2.13 Intravenous giving sets
	2.2.14 Skin disinfectant for cannulation
	2.2.15 Syringes
	2.2.16 Nutrition
	2.2.17 Nasogastric tubes
	2.2.18 Lubricant for nasogastric tube insertion
	2.2.19 Intramuscular needles
	2.2.20 Intraosseous cannulas of different sizes
	2.2.21 Blankets
	2.2.22 Facemasks for Infection Prevention and Control
	2.2.23 Aprons or gowns
	2.2.24 Charts/notes for documentation
	2.2.25 Pens
2.3 DRUGS	2.3.1 Oral rehydration solution
	2.3.2 Intravenous crystalloid fluids (eg. normal saline or Ringer's Lactate)
	2.3.3 Intravenous dextrose fluid (eg. 5%, 10% or 50%)
	2.3.5 Adrenaline
	2.3.6 Appropriate antibiotics
	2.3.7 Diazepam
	2.3.6 Widghesium Sulphate
	2.3.9 Paralelelation
2.4.1.1.1.4.4.1.1	2.5.10 Local andesthetic (eg. 2% lighticalite) (eg. for intraosseous cannulation)
	2.4.1 meaith workers with the ability to care for critically ill patients 24hrs/day
RESUURCES	2.4.2 Senior health worker who can be called to assist with the care of critically ill patients
	241115/Udy

PANEL 2 continued ...

2.5 TRAINING	2.5.1 The health workers are trained in the care of critically ill patients
2.6 ROUTINES	2.6.1 Routines for managing critically ill patients
	2.6.2 Routine for the provision of EECC without taking into account patients' ability to pay
	2.6.3 Routines for who and how to call to seek senior help 24hrs/day, 7 days/week
	2.6.4 Routines for integrating EECC with other care including the definitive care of the
	underlying condition (eg. use of condition-specific guidelines)
2.7 GUIDELINES	2.7.1 Guidelines for the essential care of critically ill patients
2.8	2.8.1 Designated space for the care of critically ill patients (eg. a bay, ward, high
INFRASTRUCTURE	dependency unit)
	2.8.2 Areas for separating and managing patients with a suspected or confirmed
	contagious disease from those without

PANEL 3. The essential diagnosis-specific care for critically ill patients with COVID-19

CLINICAL PROCESSES

- 1. The Essential Emergency and Critical Care (EECC) clinical processes as specified for all critical illnesses
- 2. Personal Protective Equipment (PPE) that is appropriate for COVID-19 as part of Infection, Prevention and Control (IPC)
- 3. Monitoring oxygen saturation using pulse oximetry at least every 6 hours, unless otherwise prescribed
- 4. Intermittent prone positioning
- 5. Low molecular weight heparin or other anticoagulant
- 6. Corticosteroid
- 7. Antibiotics in patients with suspected bacterial superinfection

HOSPITAL READINESS REQUIREMENTS

Critically ill patients with COVID-19 require the same hospital readiness for EECC as other critically ill patients. For the provision of the essential diagnosis specific care of critically ill patients with COVID-19, the following additional items are required:

CATEGORY	ITEM
3.1 EQUIPMENT	None
3.2 CONSUMMABLES	3.2.1 Facemasks appropriate for COVID-19 (eg. N95)
	3.2.2 Eye protection or face shields
3.3 DRUGS	3.3.1 Low-molecular weight heparin (eg. enoxaparin or dalteparin) or other
	anticoagulant
	3.3.2 Corticosteroid (eg. dexamethasone)
3.4 HUMAN	3.4.1 Health workers with the ability to care for critically ill COVID-19 patients
RSOURCE	24hrs/day
3.5 TRAINING	3.5.1 The health workers are trained in essential care of critically ill COVID-19
	patients
3.6 ROUTINES	3.6.1 Routines for care of critically ill COVID-19 patients
3.7 GUIDELINES	3.7.1 Guidelines for essential care of critically ill COVID-19 patients
3.8	3.8.1 Areas for separating and managing patients with suspected or confirmed
INFRASTRUCTURE	COVID-19 from those without

Discussion

We have specified the content of essential emergency and critical care (EECC) based on consensus among global clinical experts. While the EECC approach is new, the included clinical processes are commonly used in the care of sick patients and can be seen in WHO publications and specialist society standards and guidelines [29, 32-35, 38, 39, 45]. The contribution of this study is the specification of a baseline bundle of care interventions that should be provided

when needed to all critically ill patients in all hospitals in the world. This marks a break from previous guidelines that tend to be specialty-specific, condition-specific or location-specific, or that specify care that may be too complex and costly to provide in all hospital settings.

The EECC approach

EECC is an approach that supports priority-setting in health systems. In this regard, it has parallels to the approaches used in the WHO's Essential Medicines List [37], Interagency Integrated Triage Tool [29], Emergency Triage and Treatment for Children [32] and Universal Health Coverage [36]. EECC emphasises the identification and care of the critically ill, and the provision of the life-saving supportive care that is of low cost and of low complexity [25]. EECC can be seen as a unifying concept for such aspects of patient management found in WHO and specialist guidelines, triage, early warning systems and rapid response teams [28, 29, 39, 46, 47]. To maintain focus on life-saving supportive care and to be useful across all specialties, EECC does not include the definitive care of the underlying diagnoses. Instead, EECC is intended to complement speciality-based care and existing guidelines and does not aim to include all the care a patient needs - as well as EECC, patients should receive diagnostics, definitive and symptomatic care of their condition, additional nursing care, and if available, higher levels of emergency and critical care. It seeks to bridge the quality gap that is commonly found between the current care of critical illness and best-practice guidelines [12, 48, 49]. To ensure feasibility in settings with restricted human resources, EECC is designed to enable tasksharing between health professionals [50]. It should be noted that not all the EECC clinical processes will be needed in the care of every critically ill patient – they should be seen as essential "tools in the tool-box" for health workers to use when required. To operationalise the EECC approach, it is intended that the content specified here is used to develop tools for quality monitoring, teaching and integration into other guidelines and recommendations.

EECC complements the current healthcare organisation

The basic clinical processes specified in EECC have been overlooked in healthcare [11, 13-15, 18, 20, 51, 52] In UK hospitals, half of patients received substandard basic vital organ support prior to intensive care and 31% of preventable deaths were associated with absent clinical monitoring [13, 14]. In Malawi, 89% of adult hypoxic patients, and 75% of children dying from pneumonia in hospital did not receive oxygen [17, 18]. The usual organisational set-up of health services may be one underlying reason for this. Specialist units with a primary function of delivering the definitive management for one disease group may under-estimate the effort needed to maintain core processes and competences in the supportive management of critically ill patients. Innovative and specialised treatments and technologies may become preferred to those that are basic and long-standing [53]. By targeting a feasible, lowest baseline quality for critically ill patients throughout hospital settings, EECC provides a complimentary approach to the current organisation that safeguards the provision of basic life-saving actions, enhancing the impact of hospital care for all acute conditions.

EECC in the COVID-19 pandemic

EECC has added importance in a situation causing a substantial amount of severe disease and the Delphi panel agreed that EECC should be part of the care of critically ill patients with COVID-19. In addition, the agreed essential diagnostic-specific care for COVID-19 can assist in decisions about the priorities of care when the pandemic threatens to overwhelm available resources. All of the COVID-19 specific processes are well established and are included in the

WHO COVID-19 clinical management guidance [30]. The WHO guidance, and others [54], additionally include recommendations for advanced critical care (such as mechanical ventilation, vasopressors and extracorporeal oxygenation), which may be difficult to rapidly scale-up in settings of low resources. Advanced critical care can be necessary to save the lives of some patients, but has a high cost per recovery and risks diverting scarce resources to a few individuals [55-60]. Fortunately the focus has shifted in the global pandemic response from advanced critical care towards securing basic oxygen delivery systems [61, 62] underscored by statements from the WHO and other partners [63, 64]. The impact of this shift, in and beyond the pandemic, could be even greater if the necessary processes for the effective use of oxygen and other care specified in EECC were included in the scale-up.

Strengths and limitations

Our use of a consensus method with a large expert panel from diverse clinical and resource settings, specialties, and geographical locations gives the specified content legitimacy. The high response rate for this type of study during an ongoing pandemic illustrates the interest that experts had in the project's aims. The high level of consensus (>90%) for the included clinical processes promotes confidence in the final package. However, the Delphi method does have limitations. It is expert-opinion based and is limited by the make-up of the panel. Only English language speakers were included, experts were not included from all countries, and the expedited timeline of the project due to the need for results that could impact the global response to the COVID-19 pandemic may have excluded experts who could have provided additional input. The initial content presented to the panel was aligned with WHO initiatives [39], and developed by a diverse specialist team, but the possibility remains that alternative methods would have led to a different output. The study did not address the underlying evidence-base for the included clinical processes, the impact, or the potential opportunity costs of increasing the coverage of EECC in hospitals - such system-wide effects warrant careful evaluation during EECC implementation. It should be noted that, while policy makers were involved throughout the process, the EECC content has not been ratified by the WHO or governmental ministries of health – the method has been primarily scientific. The findings should be seen as the first version of the EECC content, as recommended by global clinicians and researchers, one that could be incorporated into WHO and other global and national programmes and that should subsequently be improved and updated as new knowledge arises.

Implications

Implementation of EECC could be an effective strategy as part of the current calls to save lives through improved quality of care in health systems [65] – a "low-hanging fruit". Critically ill patients have high mortality rates in all hospital settings, especially where trained staff or resources are limited, and even small improvements in outcomes would have a large impact. EECC has a vital role in the ongoing COVID-19 pandemic, for the care of the surge of critically ill patients and for optimising the impact of the efforts to scale-up oxygen. Policy makers at global, national and regional levels aiming to reduce preventable deaths should focus on improved coverage of EECC and inclusion of EECC as part of universal health coverage[36].

Conclusion

The content of essential emergency and critical care – and the essential care of critically ill patients with COVID-19 – has been specified using an inclusive global consensus. The content consists of effective, low-cost, and low-complexity life-saving care that is still frequently overlooked. The time has come to ensure that all patients in the world receive this care.

Declarations

Ethics approval and consent to participate

The study was approved by the London School of Hygiene and Tropical Medicine Research Ethics Committee (Ref: 22575). All participants in the Delphi panel provided written consent.

Competing interests

JM declares personal fees from Gilead Pharmaceuticals, support for meetings from Sphingotec, participation in advisory boards for AKPA Pharma and AM Pharma, and the position of Associate Editor at Critical Care Medicine. JR declares grants from Wellcome Trust, NIHR, and the position of Vice Chair of the Adult and Child Lung Health Section of the Union (unpaid). TB declares personal fees from UNICEF, the World Bank, USAID and the Wellcome Trust, all outside the submitted work. The other co-authors declare no conflicts of interest.

Funding

This work was supported by the Wellcome Trust [221571/Z/20/Z], as part of the 'Innovation in low-and middle-income countries' Flagship. COS received grants from the Centre for Clinical Research Sörmland, Uppsala University [DLL-941999].

Author contributions

COS conceptualised and designed the study, acquired and analysed the data, and developed the first draft of the manuscript. TB and KK conceptualised and designed the study, and acquired and analysed the data. AWS and MJ contributed to the design, acquisition and analysis of data. LAW provided input into study design and JNO, HRS, NR, JCM, JR, CH and RKK contributed to the design and analysis. All the authors interpreted the findings, critically revised the manuscript, and approved the final version.

Acknowledgements

We thank the World Federation of Intensive and Critical Care (WFICC) for supporting the study. We also thank all experts who participated in the Delphi panel, those who took part in piloting the questions, and Anders Wennerstrand for communications advice.

Abbreviations

ACVPU	Alert, confusion, voice, pain, unresponsive
AVPU	Alert, voice, pain, unresponsive
EECC	Essential Emergency and Critical Care
ICRC	The International Committee of the Red Cross
IPC	Infection prevention and control
SpO2	Saturation of haemoglobin with oxygen as measured by pulse oximetry
T&A	Treatments and Actions

WHO The World Health Organization

References

- 1. Adhikari NK, Fowler RA, Bhagwanjee S, Rubenfeld GD: Critical care and the global burden of critical illness in adults. *Lancet* 2010, **376**(9749):1339-1346.
- 2. Dart PJ, Kinnear J, Bould MD, Mwansa S, Rakhda Z, Snell D: An evaluation of inpatient morbidity and critical care provision in Zambia. *Anaesthesia* 2017, **72**(2):172-180.
- 3. Bell MB, Konrad D, Granath F, Ekbom A, Martling CR: **Prevalence and sensitivity of MET-criteria in a Scandinavian University Hospital**. *Resuscitation* 2006, **70**(1):66-73.
- 4. Kruisselbrink R, Kwizera A, Crowther M, Fox-Robichaud A, O'Shea T, Nakibuuka J, Ssinabulya I, Nalyazi J, Bonner A, Devji T *et al*: **Modified Early Warning Score (MEWS) Identifies Critical Illness among Ward Patients in a Resource Restricted Setting in Kampala, Uganda: A Prospective Observational Study.** *PLoS ONE* 2016, **11**(3):e0151408.
- Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, Colombara DV, Ikuta KS, Kissoon N, Finfer S: Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study. The Lancet 2020, 395(10219):200-211.
- 6. Razzak J, Usmani MF, Bhutta ZA: Global, regional and national burden of emergency medical diseases using specific emergency disease indicators: analysis of the 2015 Global Burden of Disease Study. *BMJ Global Health* 2019, 4(2):e000733.
- 7. Karlinsky A, Kobak D: **The World Mortality Dataset: Tracking excess mortality across countries during the COVID-19 pandemic.** *Preprint at medRxiv* 2021.
- Endris BS, Saje SM, Metaferia ZT, Sisay BG, Afework T, Mengistu YG, Fenta EH, Gebreyesus SH, Petros A, Worku A: Excess Mortality in the Face of COVID-19: Evidence from Addis Ababa Mortality Surveillance Program. Preprint with the Lancet 2021.
- 9. Tyrrell CS, Mytton OT, Gentry SV, Thomas-Meyer M, Allen JLY, Narula AA, McGrath B, Lupton M, Broadbent J,
 Ahmed A: Managing intensive care admissions when there are not enough beds during the COVID-19 pandemic:
 a systematic review. Thorax 2021, 76(3):302-312.
- 10. Jacob ST, Banura P, Baeten JM, Moore CC, Meya D, Nakiyingi L, Burke R, Horton CL, Iga B, Wald A *et al*: **The impact of early monitored management on survival in hospitalized adult Ugandan patients with severe sepsis: a prospective intervention study**. *Crit Care Med* 2012, **40**(7):2050-2058.
- 11. Molyneux E, Ahmad S, Robertson A: **Improved triage and emergency care for children reduces inpatient mortality in a resource-constrained setting**. *Bulletin of the World Health Organization* 2006, **84**(4):314-319.
- 12. Roy N, Veetil DK, Khajanchi MU, Kumar V, Solomon H, Kamble J, Basak D, Tomson G, von Schreeb J: Learning from 2523 trauma deaths in India-opportunities to prevent in-hospital deaths. *BMC health services research* 2017, 17(1):1-8.
- 13. Hogan H, Healey F, Neale G, Thomson R, Vincent C, Black N: **Preventable deaths due to problems in care in English acute hospitals: a retrospective case record review study**. *BMJ quality & safety* 2012, **21**(9):737–745.
- 14. McQuillan P, Pilkington S, Allan A, Taylor B, Short A, Morgan G, Nielsen M, Barrett D, Smith G: **Confidential** inquiry into quality of care before admission to intensive care. *Bmj* 1998, **316**(7148):1853-1858.
- 15. Goulet H, Guerand V, Bloom B, Martel P, Aegerter P, Casalino E, Riou B, Freund Y: **Unexpected death within 72** hours of emergency department visit: were those deaths preventable? *Critical Care* 2015, **19**(1):1-7.
- 16. Floyd J, Wu L, Burgess DH, Izadnegahdar R, Mukanga D, Ghani AC: **Evaluating the impact of pulse oximetry on** childhood pneumonia mortality in resource-poor settings. *Nature* 2015, **528**(7580):S53.
- King C, Banda M, Bar-Zeev N, Beard J, French N, Makwenda C, McCollum E, Mdala M, Nisar YB, Phiri T: Careseeking patterns amongst suspected paediatric pneumonia deaths in rural Malawi. Gates Open Research 2020, 4(178):178.
- 18. Kayambankadzanja RK, Schell CO, Mbingwani I, Mndolo SK, Castegren M, Baker T: **Unmet need of essential treatments for critical illness in Malawi**. *Priprint at medRxiv* 2021.
- 19. Baker T, Lugazia E, Eriksen J, Mwafongo V, Irestedt L, Konrad D: **Emergency and critical care services in Tanzania:** a survey of ten hospitals. *BMC Health Serv Res* 2013, **13**:140.

- 20. Reynolds TA, Sawe H, Rubiano AM, Shin SD, Wallis L, Mock CN: **Strengthening Health Systems to Provide Emergency Care**. In: *Disease Control Priorities: Improving Health and Reducing Poverty*. 3rd edn. Washington (DC): The International Bank for Reconstruction and Development / The World Bank; 2017.
- 21. Abdu M, Wilson A, Mhango C, Taki F, Coomarasamy A, Lissauer D: Resource availability for the management of maternal sepsis in Malawi, other low-income countries, and lower-middle-income countries. International Journal of Gynecology & Obstetrics 2018, 140(2):175-183.
- 22. Baker T, Schell CO, Lugazia E, Blixt J, Mulungu M, Castegren M, Eriksen J, Konrad D: Vital Signs Directed Therapy: Improving Care in an Intensive Care Unit in a Low-Income Country. *PLoS One* 2015, **10**(12):e0144801.
- 23. Diaz JV, Riviello ED, Papali A, Adhikari NK, Ferreira JC: Global critical care: moving forward in resource-limited settings. *Annals of global health* 2019, **85**(1):3.
- 24. Schell CO, Beane A, Kayambankadzanja RK, Khalid K, Haniffa R, Baker T: **Global Critical Care: Add Essentials to the Roadmap**. *Annals of global health* 2019, **85**(1):97.
- 25. Schell CO, Gerdin Warnberg M, Hvarfner A, Hoog A, Baker U, Castegren M, Baker T: **The global need for essential** emergency and critical care. Crit Care 2018, **22**(1):284.
- 26. Dalkey NC: An experimental study of group opinion: The Delphi method. *Futures* 1969, **1**(5):408-426.
- 27. Ward A: Advanced Medical Life Support Assessment for the Medical Patient. Burlington: Jones and Bartlett; 2018.
- 28. Henry S: **ATLS 10th edition offers new insights into managing trauma patients**. *Bulletin of the American College of Surgeons* 2018.
- 29. World Health Organization: Clinical care for severe acute respiratory infection: toolkit: COVID-19 adaptation. Geneva: World Health Organization; 2020.
- 30. COVID-19 clinical management: living guidance, 25 January 2021 [https://www.who.int/publications/i/item/WHO-2019-nCoV-clinical-2021-1]
- 31. Helping Mothers Survive Essential Care for Labor & Birth (African Graphics) [http://reprolineplus.org/resources/helping-mothers-survive-essential-care-labor-birth-african-graphics]
- 32. World Health Organization: Emergency Triage Assessment and Treatment (ETAT) course. Genveva, Switzerland; 2005.
- 33. APLS: The Pediatric Emergency Medicine Resource [https://www.aap.org/en-us/continuing-medicaleducation/life-support/APLS-The-Pediatric-Emergency-Medicine-Resource/Pages/APLS-The-Pediatric-Emergency-Medicine-Resource.aspx]
- 34. Monitoring emergency obstetric care : a handbook [https://apps.who.int/iris/handle/10665/44121]
- 35. World Health Organization: Guidelines for essential trauma care: World Health Organization; 2004.
- 36. UHC Compendium. Health interventions for Universal Health Coverage [https://www.who.int/universal-healthcoverage/compendium]
- 37. WHO Model Lists of Essential Medicines [https://www.who.int/groups/expert-committee-on-selection-and-useof-essential-medicines/essential-medicines-lists]
- 38. World Health Organization: WHO recommendations on newborn health: guidelines approved by the WHO Guidelines Review Committee. In.: World Health Organization; 2017.
- 39. World Health Organization: **BASIC EMERGENCY CARE: approach to the acutely ill and injured**: World Health Organization; 2018.
- 40. Vogel C, Zwolinsky S, Griffiths C, Hobbs M, Henderson E, Wilkins E: **A Delphi study to build consensus on the** definition and use of big data in obesity research. *International Journal of Obesity* 2019, **43**(12):2573-2586.
- 41. Lozano LM, García-Cueto E, Muñiz J: Effect of the number of response categories on the reliability and validity of rating scales. *Methodology* 2008, **4**(2):73-79.
- 42. Akins RB, Tolson H, Cole BR: **Stability of response characteristics of a Delphi panel: application of bootstrap data expansion**. *BMC medical research methodology* 2005, **5**(1):1-12.
- 43. Marshall JC, Bosco L, Adhikari NK, Connolly B, Diaz JV, Dorman T, Fowler RA, Meyfroidt G, Nakagawa S, Pelosi P: What is an intensive care unit? A report of the task force of the World Federation of Societies of Intensive and Critical Care Medicine. Journal of critical care 2017, **37**:270-276.
- 44. Priority medical devices list for the COVID-19 response and associated technical specifications: interim guidance,19 November 2020 [https://apps.who.int/iris/handle/10665/336745]
- 45. Cecconi M, Hernandez G, Dunser M, Antonelli M, Baker T, Bakker J, Duranteau J, Einav S, Groeneveld AJ, Harris T: Fluid administration for acute circulatory dysfunction using basic monitoring: narrative review and expert panel recommendations from an ESICM task force. Intensive care medicine 2019, **45**(1):21-32.
- 46. Smith GB, Prytherch DR, Meredith P, Schmidt PE, Featherstone PI: **The ability of the National Early Warning** Score (NEWS) to discriminate patients at risk of early cardiac arrest, unanticipated intensive care unit admission, and death. *Resuscitation* 2013, **84**(4):465-470.
- 47. Winters BD, Weaver SJ, Pfoh ER, Yang T, Pham JC, Dy SM: Rapid-response systems as a patient safety strategy: a systematic review. Annals of internal medicine 2013, **158**(5 Pt 2):417-425.
- 48. Baelani I, Jochberger S, Laimer T, Otieno D, Kabutu J, Wilson I, Baker T, Dunser MW: Availability of critical care resources to treat patients with severe sepsis or septic shock in Africa: a self-reported, continent-wide survey of anaesthesia providers. *Crit Care* 2011, **15**(1):R10.

- 49. Maaløe N, Ørtved AMR, Sørensen JB, Dmello BS, van den Akker T, Kujabi ML, Kidanto H, Meguid T, Bygbjerg IC, van Roosmalen J: The injustice of unfit clinical practice guidelines in low-resource realities. *The Lancet Global Health* 2021.
- 50. Callaghan M, Ford N, Schneider H: A systematic review of task- shifting for HIV treatment and care in Africa. *Hum Resour Health* 2010, **8**:8.
- 51. Duke T, Wandi F, Jonathan M, Matai S, Kaupa M, Saavu M, Subhi R, Peel D: Improved oxygen systems for childhood pneumonia: a multihospital effectiveness study in Papua New Guinea. *Lancet* 2008, 372(9646):1328-1333.
- 52. Luettel D, Beaumont K, Healey F: **Recognising and responding appropriately to early signs of deterioration in hospitalised patients**. In. London: The National Patient Safety Agency - NHS; 2007.
- 53. Dessap AM: Frugal innovation for critical care. Intensive care medicine 2019, **45**(2):252-254.
- Alhazzani W, Møller MH, Arabi YM, Loeb M, Gong MN, Fan E, Oczkowski S, Levy MM, Derde L, Dzierba A:
 Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). Intensive care medicine 2020, 46(5):854-887.
- 55. Cleary SM, Wilkinson T, Tchuem CT, Docrat S, Solanki G: **Cost-effectiveness of intensive care for hospitalized COVID-19 patients: experience from South Africa**. *BMC Health Serv Res* 2021, **21**(1):1-10.
- 56. Barasa E, Kairu A, Maritim M, Were V, Akech S, Mwangangi M: **Examining unit costs for COVID-19 case** management in Kenya. *BMJ global health* 2021, **6**(4):e004159.
- 57. Chang DW, Shapiro MF: Association between intensive care unit utilization during hospitalization and costs, use of invasive procedures, and mortality. *JAMA internal medicine* 2016, **176**(10):1492-1499.
- Baker T, Schell CO, Petersen DB, Sawe H, Khalid K, Mndolo S, Rylance J, McAuley DF, Roy N, Marshall J: Essential care of critical illness must not be forgotten in the COVID-19 pandemic. *The Lancet* 2020, 395(10232):1253-1254.
- 59. Malawi Biomedical Equipment for COVID-19 Case Management [https://path.azureedge.net/media/documents/Malawi_Biomedical_Equipment_Assessment_Report_PATH_2021 .01.25_final.pdf]
- 60. Mantena S, Rogo K, Burke TF: **Re-Examining the Race to Send Ventilators to Low-Resource Settings**. *Respiratory Care* 2020, **65**(9):1378-1381.
- 61. Stein F, Perry M, Banda G, Woolhouse M, Mutapi F: **Oxygen provision to fight COVID-19 in sub-Saharan Africa**. *BMJ Global Health* 2020, **5**(6):e002786.
- 62. COVID-19 Oxygen Needs Tracker [https://www.path.org/programs/market-dynamics/covid-19-oxygen-needstracker/]
- 63. Usher AD: Medical oxygen crisis: a belated COVID-19 response. The Lancet 2021, 397(10277):868-869.
- Fore HH, Ghebreyesus TA, Watkins K, Greenslade L, Berkley S, Bassat Q, Duneton P, Klugman K, Golden A: Leveraging the COVID-19 response to end preventable child deaths from pneumonia. *The Lancet* 2020, 396(10264):1709-1711.
- 65. Kruk M, Gage A, Arsenault C: **High-quality health systems in the Sustainable Development Goals era: time for a revolution**. *Lancet Global Health* 2018, **6**(11):E1162-E1162.