

Multi-tract multi-symptom relationships in pediatric concussions

Guido I. Guberman¹, Sonja Stojanovski^{2,3}, Eman Nishat^{2,3}, Alain Ptito¹, Danilo Bzdok^{4,5,6}, Anne Wheeler^{2,3}, Maxime Descoteaux⁷

1. Department of Neurology and Neurosurgery, Faculty of medicine, McGill University, Montreal, Quebec, Canada

2. Department of Physiology, Faculty of Medicine, University of Toronto, Toronto, ON, Canada

3. Neuroscience and Mental Health, The Hospital for Sick Children, Toronto, ON, Canada

4. McConnell Brain Imaging Centre (BIC), Montreal Neurological Institute (MNI), Faculty of Medicine, McGill University, Montreal, Quebec, Canada

5. Department of Biomedical Engineering, Faculty of Medicine, School of Computer Science, McGill University, Montreal, Quebec, Canada

6. Mila - Quebec Artificial Intelligence Institute, Montreal, Quebec, Canada

7. Department of Computer Science, Sherbrooke University, Sherbrooke, QC, Canada

Abstract:

The heterogeneity of white matter damage and symptoms in concussions has been identified as a major obstacle to therapeutic innovation. In contrast, the vast majority of diffusion MRI studies on concussion have traditionally employed group-comparison approaches. Such studies do not consider heterogeneity of damage and symptoms in concussion. To parse concussion heterogeneity, the present study combines diffusion MRI (dMRI) and multivariate statistics to investigate multi-tract multi-symptom relationships. Using dMRI data from a sample of 306 children ages 9 and 10 with a history of concussion from the Adolescent Brain Cognitive Development Study (ABCD study), we built connectomes weighted by classical and emerging diffusion measures. These measures were combined into two informative indices, the first capturing a mixture of patterns suggestive of microstructural complexity, the second representing almost exclusively axonal density. We deployed pattern-learning algorithms to jointly decompose these connectivity features and 19 behavioural measures that capture well-known symptoms of concussions. We found idiosyncratic symptom-specific multi-tract connectivity features, which would not be captured in traditional univariate analyses. Multivariable connectome-symptom correspondences were stronger than all single-tract/single-symptom associations. Multi-tract connectivity features were also expressed equally across different sociodemographic strata and their expression was not accounted for by injury-related variables. In a replication dataset, the expression of multi-tract connectivity features predicted adverse psychiatric outcomes after accounting for other psychopathology-related variables. By defining cross-demographic multi-tract multi-symptom relationships to parse concussion heterogeneity, the present study can pave the way for the development of improved stratification strategies that may contribute to the success of future clinical trials and the improvement of concussion management.

Introduction

Concussion afflicts approximately 600 per 100,000 individuals every year.¹ Its incidence rate is rising in children and adolescents,² and compared to adult populations, the impact of concussions on pediatric brains is understudied.³ Despite considerable funding devoted to clinical and basic research, no major advances in therapeutics have been achieved to date.⁴ A root cause of this stagnation appears to be a contradiction: while all concussions are treated equally in clinical trials and research studies, they are characterized by extensive heterogeneity in their pathophysiology, clinical presentation, symptom severity and duration.^{4,5} Concussion heterogeneity across patients has been identified as a major hurdle in advancing concussion care.^{4,5}

Due to shearing forces transmitted during injury, the brain's white matter is especially vulnerable to concussion.^{6,7} Decades of research have studied white matter structure in individuals who sustain concussions. However, most studies continue to assume consistent, one-to-one structure/symptom relationships and employ traditional group comparisons,^{8,9} averaging out the diffuse and likely more idiosyncratic patterns of brain structure abnormalities in favour of shared ones. Hence, the extant literature suggests that a large proportion of the clinical and research studies have not adequately accounted for clinical and neuropathological concussion heterogeneity.

To remedy this shortcoming, a growing number of studies aimed to parse the clinical heterogeneity in concussions by algorithmically partitioning patients into discrete subgroups based on symptoms.¹⁰⁻¹² Other studies aim instead to account for heterogeneity in white matter structure alterations.¹³⁻¹⁵ Ware et al.¹⁵ built individualized maps of white matter abnormalities which revealed substantial inter-subject variability in traumatic axonal injury and minimal consistency of subject-level effects. Taylor et al.¹⁴ computed a multivariate summary measure of white matter structure across 22 major white matter bundles which achieved better classification accuracy of concussed patients from healthy controls compared to single tract measures. Hence, studies have attempted to address concussion heterogeneity in symptoms and in white matter structure. However, no prior studies have considered both sources of heterogeneity simultaneously.

White matter alterations due to concussion are diffuse and can elicit several symptoms that may interact with each other in complex ways.^{4,5,16} For instance, two individuals may suffer a concussion and develop sleep problems. The first may have damaged white matter tracts related to sleep/wakefulness control, whereas the second may have damaged tracts related to mood, causing depression-like symptoms, which include sleep problems. These two individuals will thus display a common symptom but will have overall different symptom profiles and different white matter damage profiles. Parsing concussion heterogeneity requires accounting for these dynamic, multi-tract multi-symptom relationships.

In the present study, we leveraged advanced diffusion MRI (dMRI) methods as well as a double-multivariate approach to parse concussion heterogeneity in white matter structure and symptoms simultaneously in a large sample of previously-concussed children. Multi-tract multi-

symptom relationships captured more information than traditional univariate approaches. Expression of multi-tract connectivity features was not driven by sociodemographic strata and injury-related variables. Finally, after accounting for univariate variables found to be related to adverse psychiatric outcome in the discovery dataset (n=214), we found that expression of one multi-tract connectivity feature predicted adverse psychiatric outcome in a replication dataset (n=92).

Methods

Participants

Participants in this study were obtained from the world's largest child development study of its kind – the ongoing longitudinal Adolescent Brain Cognitive Development Study (ABCD study; <https://abcdstudy.org/>), data release 2.0 (<https://data-archive.nimh.nih.gov/abcd>). The ABCD Study acquired data from 11,874 children aged 9 to 10 years (mean age = 9.49 years) from across the United States (48% girls; 57% Caucasian, 15% African American, 20% Hispanic, 8% other).¹⁷ Additional information about the ABCD study can be found in Garavan et al.¹⁸

History of concussion

Parents completed a modified version of the Ohio State University TBI Identification Method (OSU-TBI-ID)¹⁹. We included participants who reported a head injury without loss of consciousness but with memory loss and/or a head injury with loss of consciousness for less than 30 minutes (n=434). Due to missing or incomplete data, corrupted files, data conversion errors, and images rated by the ABCD study team as being of poor quality, the final sample of participants with usable data was 345. After processing, images were visually inspected by two independent raters (G.I.G., S.S.). Images that were deemed of low quality after processing by both raters were removed (n=39), leading to a final sample of 306 participants. We randomly divided the sample into a discovery dataset (70%, n=214) and a replication dataset (20%, n=92). Figure 1 summarizes the subject selection procedure.

Symptom-oriented measures

To probe various aspects of concussion symptomatology, we used items collected from assessments available in the ABCD dataset. These items, as well as the concussion symptom they are meant to probe are outlined in Table 1.

MRI Acquisition

MRI scans were acquired across 21 sites, with data coming from 28 different scanners. Details about the acquisition protocols and image specifications are outlined in Casey et al 2018.²⁰ Multi-shell dMRI scans had 96 diffusion-weighted directions, with 6 directions of b=500 s/mm², 15 directions of b=1000s/mm², 15 directions of b=2000s/mm², and 60 directions of b=3000s/mm². The b=2000 shell was excluded from the data processing. In addition, scans had 6 or 7 b=0s/mm² images, depending on scanner type. Lastly, a reverse b0 image was included for each participant.

Processing

We used Tractoflow²¹ to process dMRI and T1-weighted scans. Tractoflow is a novel diffusion MRI processing pipeline, incorporating state-of-the-art functions from FSL, Dipy, and MRtrix into NextFlow. The processing steps are summarized in Theaud et al.²¹ Important deviations from the default parameters utilized by Tractoflow are as follows: 1. We used gray-white matter interface seeding, as this method accounts for the length bias introduced by white-matter seeding;²² 2. We used 24 seeds-per-voxel with the objective of obtaining approximately 2 million streamlines across the entire brain. We used the b=0, 500, and 1000 shells to perform tensor fitting, and the b=0 and 3000 shells to perform Constrained Spherical Deconvolution (CSD) (Descoteaux et al. 2009; Tournier et al. 2007).^{23,24} We obtained group-average fiber-response functions from voxels with high (>0.70) fractional anisotropy (FA). Lastly, we created tractograms using a probabilistic particle-filtering tractography algorithm (Girard et al. 2014).²²

Connectivity matrices

The post-processing workflow is illustrated in Figure 2. To construct connectivity matrices, we used Freesurfer on McGill's CBrain platform²⁵ to fit the Desikan-Killiani Tourville (DKT)²⁶ and *aseg* atlases onto the processed T1-images that had been transformed to DWI space during processing (Figure 2A). We removed redundant and irrelevant labels from the fitted atlas (a list of retained labels is supplied in supplementary material), yielding a final atlas with 76 labels. We then thresholded matrices such that a connection was only retained if it was found to be successfully reconstructed (defined as the presence of at least one streamline) across 90% of participants.²⁷ Results using different thresholds are presented in supplementary material. We then weighted thresholded connectomes by FA, mean, radial, and axial diffusivities (MD, RD, AD respectively), number of fiber orientations (NuFO), and apparent fiber density along fixels (AFDf) (Figure 2B). The first four measures are derived from the tensor model, whereas the latter two are based on fiber orientation distribution functions (fODFs) obtained from CSD.^{28,29} Simulation studies have shown that AFD is more specifically related to axonal density, and by computing it along "fixels" (fiber elements), axonal density specific to particular fiber populations can be studied independently of crossing fibers.²⁸ Although individual diffusion measures are related to different aspects of neuropathology, together they provide more information than when considered separately.²⁷ A recent framework based on principal component analysis (PCA) has been proposed to recombine diffusion measures into biologically-interpretable measures of white matter structure.³⁰ We therefore performed PCA on the concatenated set of standardized measures across subjects and connections, generating connectivity matrices weighted by principal component (PC) scores (Figure 2C).

Additional data transformations

We imputed missing connectivity (prior to the PCA), symptom, and nuisance data (sex, pubertal stage, handedness, scanner) by randomly selecting non-missing data from other participants in the same dataset. We reverse-coded cognitive scores, such that increasing scores in all symptom data reflected more problems. From connectivity and symptom data, we regressed out the following nuisance variables: sex, pubertal stage, scanner (only for connectivity data), and handedness.

Pattern-learning pipeline

Feature selection. To reduce the number of connectivity features included in the partial least squares correlation (PLSc) analysis, we selected the 200 connectivity features most correlated (based on Pearson correlations) with any symptom score. This solution is becoming increasingly adopted for high-dimensional variable sets (Figure 2D).^{31,32}

PLSc. We performed PLSc analyses using the *tepPLS* function from the *texposition* package.³³ PLSc involves singular value decomposition on the covariance matrix between connectivity and symptom features, creating pairs of multi-tract and multi-symptom features called multi-tract multi-symptom relationships. Each multi-tract multi-symptom relationship encapsulates a linear combination of connectivity features (“multi-tract features”), a linear combination of symptom scores (“multi-symptom features”), and an eigenvalue (reflective of the amount of explained covariance between connectivity and symptom features). Each multi-tract multi-symptom relationship is constructed so as to explain a successively smaller portion of the covariance between symptoms and connectivity features. We constructed the largest number of possible multi-tract multi-symptom relationships, given the dimensionality of the behavioral variable set ($k=19$) (Figure 2D).

Selection and interpretation of multi-tract multi-symptom pairs. To select the multi-tract multi-symptom pairs to retain for interpretation, we performed permutation testing (2000 iterations). This procedure randomly shuffles row labels for the connectivity features, without replacement, repeats the PLSc and computes eigenvalues at every permutation. We calculated p -values as the proportion of permutations that yielded eigenvalues that exceeded the original amount.

To interpret symptom and connectivity weights of significant ($p<0.05$) multi-tract multi-symptom pairs, we performed bootstrap analyses (2000 iterations), using the *BOOT4PLSC* command from the *texposition* package. At each iteration, labels for data were drawn with replacement, the entire PLSc was repeated and the weights for all pairs were obtained. This process yields a sampling distribution of weights for each connectivity and symptom feature.³⁴ The ratio of the original weights to the standard error of each feature’s bootstrap distribution can be interpreted as a z-score, which yielded so-called ‘bootstrap ratios’. We used a value of 1.96 to determine which variables significantly contributed to each particular significant pair.

Comparison of multivariate against univariate approaches

To compare information captured by the PLSc and univariate approaches, we first divided participants based on whether they had T-scores above 70, a well-recognized clinical threshold,³⁵ in the Child Behavior Checklist (CBCL) Depression, Attention Problems, Anxiety Disorder, or Aggression scales.³⁵ Using a threshold of $p<0.05$, we computed univariate comparisons of connectivity (PC scores) between individuals with and those without clinical-level psychopathology, thus identifying psychopathology-related univariate connectivity features.

We were interested in comparing how many of these features were also found to significantly contribute to each multi-tract connectivity feature. To do so, we computed a measure of percent overlap as follows:

$$\%Overlap = \frac{C_{sig}}{(S_u + S_m - C_{sig})} \times 100,$$

where C_{sig} refers to the number of connections flagged as significant in both approaches, S_u to the number of connections flagged as significant in the univariate approach, and S_m to the number of connections flagged as significant in the multivariate approach. This measure can account for the apparent high overlap that can arise when S_u and S_m are not equivalent in size.

Relation to TBI-related and sociodemographic factors

We addressed whether expression of multi-tract connectivity features was related to injury-specific and sociodemographic factors. Injury-related variables included: the time between the last-documented injury and testing, the cause of injury, and the total number of documented mTBIs. Sociodemographic variables included: sex, total combined family income in the last 12 months, and race/ethnicity. We used the following categories for race/ethnicity: “Asian” (Asian Indian, Chinese, Filipino, Japanese, Korean, Vietnamese, Other Asian), AIAN (“American Indian”/Native American, Alaska Native), NHPI (Native Hawaiian, Guamanian, Samoan, Other Pacific Islander), Non-Hispanic White, Non-Hispanic Black, Hispanic, Other, and Multiple.³⁶ To illustrate the influence of these sociodemographic factors we created scatter plots illustrating expression of connectivity latent factors color-coded by sociodemographic factors (Supplementary Figure 1).

Prediction of clinical outcome

We used the Kiddie-Schedule for Affective and Psychiatric Disorders in School Age Children (KSADS), a gold-standard tool, to assess the presence of pediatric psychiatric disorders.³⁷ We used the presence of any current psychiatric diagnosis as indicating an adverse psychiatric outcome (55% of discovery set, 55% of replication set). We created separate multivariable logistic regressions to predict adverse psychiatric outcomes using 1) expression of multi-tract connectivity features, 2) psychopathology measures from the CBCL, or 3) psychopathology-related univariate connectivity features. From each set of models, we retained the variables that were significantly related to adverse psychiatric outcome. We then tested whether expression of multi-tract connectivity features could predict adverse psychiatric outcome in the replication dataset.

To do so, we first projected connectivity data from the replication dataset onto the same principal components retained in the discovery dataset. Next, we selected the same 200 connections that had been retained based on univariate feature selection, and projected these connections onto the multi-tract connectivity spaces obtained from the discovery dataset. Finally, we performed a logistic regression model incorporating all the predictors found to be significant in the discovery dataset, applied on data from the replication dataset.

The objective of these models was to assess whether expression of multi-tract connectivity features could predict adverse psychiatric outcomes in the replication dataset after accounting for other psychopathology-related factors (CBCL behavioural measures, univariate connectivity measures).

RESULTS

Sample

Out of 434 participants with a history of mTBI, 306 (127F/179M) had usable data (Figure 1). Table 2 outlines sociodemographic and injury-related factors, as well as handedness and sex. The majority had sustained an injury over 1 year prior to the study. Nuisance variables were well-balanced between participants in the discovery and the replication set.

Recombined measures of white matter tract microstructure

The PCA yielded two biologically-interpretable components that together explained 97% of the variance in dMRI measures (Figure S2). The first (PC1) represented all diffusion measures except AFDf, and likely reflects microstructural complexity. The second (PC2) represented AFDf exclusively, and likely reflects axonal density. Because we retained two PCs, we performed two PLSc analyses.

Multi-tract multi-symptom relationships

Each PLSc analysis yielded 19 latent modes of covariance (termed here “multi-tract multi-symptom relationships”), each consisting of a pair of multi-tract connectivity and multi-symptom features. Based on permutation testing, 18 multi-tract multi-symptom pairs were retained from PC1 and 14 from PC2. For brevity, only a few informative pairs are presented herein, but an illustration of all significant pairs for all thresholds can be found in supplementary material (Figures S3-S6).

For PC1, similarly to PC2, the first multi-tract multi-symptom pair broadly represented all symptoms (Figure 3A), capturing general problems. Interestingly, this pair implicated several callosal tracts (Figure 5). The third multi-tract multi-symptom pair obtained from PC2 represented mostly sleep and cognitive problems, whereas the fourth pair represented more strongly mood, sleep, and somatic problems (Figure 3B and C).

Multivariate vs univariate approaches

Correlations between multi-tract and multi-symptom features were higher than all univariate correlations between the PC1 and PC2 scores of every retained tract and every symptom (“single-tract single-symptom relationships”) (Figure 4).

We identified 28 individuals with scores above 70 in CBCL Depression, Attention Problems, Anxiety Disorder, or Aggression scales, considered to be the clinical range.⁹ Figure 3 illustrates the expression of multi-symptom and multi-tract connectivity features from three pairs. In clinical trials and research studies, these 28 individuals would be grouped together. However, when considering all symptoms, these individuals appear to be markedly different. Accordingly, these individuals were differentiable by their expression of multi-symptom and multi-tract connectivity features (Figure 3).

We compared PC1 and PC2 scores across all 200 connections between individuals with and without clinical-level CBCL scores, and calculated the percent overlap between each multi-tract connectivity feature and the set of tracts found to be significant in univariate comparisons. The percent overlap scores are presented in Figure 5. Notably, the highest overlap occurred with multi-tract connectivity feature 1 (22%) from both PLSc analyses (Figure 5). Univariate psychopathology-related features (Figure 5 red and blue chord plots) and multi-tract connectivity feature 1 (Figure 5 violet chord plot) implicated several callosal tracks, whereas

fewer callosal tracks were implicated in multi-tract connectivity features 3 and 4 (Figure 5 turquoise and green chord plots).

Relationship with sociodemographic and injury-related factors

The expression of multi-tract connectivity features was equivalent across sociodemographic strata defined by sex, total combined household income, and race/ethnicity (Supplementary Figure 1). Out of 32 retained multi-tract multi-symptom pairs, time-since-injury was only significantly correlated to the expression of two multi-tract connectivity features (Table S1) and no multi-symptom features. Only one multi-tract connectivity feature (feature 2 from the first PLSc) was significantly different between groups defined by injury cause (Figure S7). Only two multi-tract connectivity features, feature 17 from the first PLSc (Figure S8), and feature 8 from the second PLSc (Figure S9).

Prediction of clinical outcome

Using separate multivariable logistic regressions, we found expression of four multi-tract connectivity features from PC1, one behavioural measure (CBCL Attention Problems), and no single tract features that were significantly related to adverse psychiatric outcome. No other variables, including no multi-tract connectivity features obtained from PC2 scores, were related to adverse psychiatric outcome. We projected the data from the replication set onto the PCs obtained from the discovery set, selected the same 200 connections, and computed the expression of the multi-tract connectivity features significantly related to adverse psychiatric outcome in the discovery set. We incorporated these multi-tract connectivity features with other significant variables, using the replication data, in a single multivariable logistic regression. We found that a one-unit increase in the expression of multi-tract connectivity feature 4 significantly increased the odds of an adverse psychiatric outcome in the replication set 2.31 times ($p=0.02$) after controlling for univariate, psychopathology-related behavioural features. This multi-tract connectivity feature was paired with a multi-symptom feature that implicated sleep and somatic problems (Figure 6).

Sensitivity analyses

Results using different thresholds ($t=85\%$, 95% , 100%) for the connectomes are outlined in supplementary material. All results were consistent, including PCs, the weights of the multi-symptom features, the strength of multivariate vs univariate relationships, the trends for the % overlap scores, and the significance of adverse psychiatric outcome predictions using multi-tract connectivity feature expression. Results were less consistent with the $t=100\%$ threshold, which demonstrated instead little to no significant multi-tract/multi-symptom pairs based on permutation testing. Correlations between expression of multi-tract connectivity features at different thresholds are illustrated in supplementary material. Notably, for $t=85\%$, 90% , and 95% , correlations between the expression of the corresponding multi-tract connectivity features (e.g.: multi-tract feature 1 from $t=85\%$, multi-tract feature 1 from $t=95\%$) all have high correlations, indicating how similar these features are, whereas these correlations were lower when comparing against $t=100\%$ (Figure S10).

Discussion

In the present study we leveraged novel dMRI methods and a double-multivariate approach to parse heterogeneity in white matter structure and symptoms in a large sample of previously-concussed children. Applying PLSc on biologically-interpretable measures of dMRI obtained from PCA, we found cross-demographic latent multi-tract multi-symptom relationships that captured more information than traditional approaches and predicted meaningful clinical outcomes in unseen data. These results both recapitulated well-known findings from the concussion literature and revealed new insights about white matter structure/symptom relationships.

In response to decades of failed attempts to translate basic science findings into successful clinical trials and novel therapies, concussion heterogeneity has been identified as a major obstacle.^{4,5} Heterogeneity in symptoms, impact of injury on brain structure and function, and pre-injury factors pose a particular problem for most concussion neuroimaging studies which have traditionally employed univariate comparisons between concussed and healthy or orthopedic injury control groups, or between patients with and without persistent symptoms.^{8,9} These sources of variability are believed to be problematic because they decrease the statistical power needed for group comparisons and multivariable models to detect the often-subtle effects of concussions.³⁸ To overcome this challenge, landmark initiatives such as the IMPACT,³⁸ InTBIR,³⁹ CENTER TBI,⁴⁰ and TRACK TBI⁴¹ aim to standardize and pool multi-center data collected across sociodemographic strata, to identify and statistically correct for pre-injury factors known to impact brain structure, and develop diagnostic and prognostic tools leveraging multimodal data and increasingly sophisticated machine-learning approaches.

In this study, we posited that concussion heterogeneity is also problematic because by pooling across patients, idiosyncratic patterns of connectivity that may be more symptom-specific are sacrificed in favour of shared ones. By assuming that symptoms map cleanly and consistently onto shared connectivity abnormalities in a one-to-one fashion, erroneous inferences could be made about relationships between group-level patterns of connectivity differences and specific symptoms. Our results are consistent with this idea: univariate comparisons between a group displaying clinical-level psychopathology and the rest of the sample identified connectivity features that mostly overlapped with the first multi-tract multi-symptom pair obtained from both PLSc analyses. These pairs, which accounted for the most covariance, reflected general problems, not specifically psychopathology. Both these pairs and the univariate “psychopathology-related” connections implicated several callosal tracts. These results suggest that univariate comparisons, even when performed in such a way as to identify a symptom-specific set of connectivity features, identified only the most consistent group-level connectivity differences at the expense of more symptom-specific idiosyncratic ones. These findings are consistent with most prior concussion literature, which strongly implicate the corpus callosum,⁹ but extend this literature by suggesting that that the central importance of the corpus callosum may have resulted from pooling across concussed subjects.

Rather than a clean and consistent set of single-tract single-symptom relationships, this study suggests concussions may be best conceptualized as a multiplicity of multi-tract multi-symptom combinations. Multi-tract relationships may be driven by the metabolic demands

imposed by the network structure of the brain, which is known to predict the course of several brain diseases,⁴² by biomechanical constraints imposed by the skull and other structures exposing certain areas to more shearing strain,⁴³ or by both factors simultaneously.⁴⁴ These possibilities need to be tested further.

Few recent studies have parsed heterogeneity in concussion symptoms using clustering analyses.¹⁰⁻¹² The reported subgroups differed from those found in the present study. Differences between symptom profiles arose because our multi-symptom features are associated with brain structure and not driven by variability in symptoms alone. Further, we did not group patients into discrete subgroups. Although whether patients cleanly fit into discrete subtypes has not been explicitly tested in our study, expression of multi-tract and multi-symptom features do not suggest any obvious clusters. The existence of concussion subtypes needs to be explored further.

Prior studies have attempted to address heterogeneity in white matter structure in concussions.¹³⁻¹⁵ Using different approaches, these studies generated point summaries that accounted for the high-dimensional variability of white matter structure to better distinguish patients from controls. Our approach offers an important advantage. Rather than collapsing the highly variable information provided by white matter structure, our approach attempts to find symptom-specific patterns of white matter structure, which has potential to lead to new treatment targets.

After accounting for behavioural and single-tract predictors of adverse psychiatric outcome, the expression of one multi-tract connectivity feature significantly predicted adverse psychiatric outcome in a holdout dataset. The multi-symptom feature that corresponded to this significant multi-tract feature implicated mostly sleep and somatic problems. Sleep problems are known to be linked to psychiatric disorders, both as symptoms, and risk factors.⁴⁵ Longitudinal studies are needed to elucidate whether these multi-tract features reflect abnormalities that increase the risk of later adverse psychiatric outcomes, or instead reflect abnormalities associated with existing psychiatric diagnoses. Nonetheless, the prediction of a meaningful clinical outcome on unseen data is particularly promising.

Limitations and Strengths

The present results should be considered in light of methodological limitations. Data on mTBI occurrence was collected retrospectively. Participants did not have baseline data, and additionally had highly variable times since injury. Most individuals with concussions recover from their injury⁴⁶ which should have led to a concussed group where most participants were similar to healthy controls. Interestingly, our PCA yielded combinations of diffusion measures that differed from those of two prior studies that have used this approach.^{30,47} These prior studies used samples of typically-developing children without neurological insults and found similar PCs. To the extent that the PCs reported in these prior studies reflect healthy neurotypical brains, our PCs suggest that our concussed sample was not as similar in white matter structure to healthy controls as expected. However, variable time since injury made the interpretation of patterns of microstructure difficult. Further, due to the cross-sectional nature

of this data, the difference between symptoms and pre-existing characteristics are difficult to discern, especially since some behavioural measures often believed to be symptoms of concussion, such as attention problems, can also be risk factors for injury.⁴⁸ Heterogeneity has several forms, including in symptoms, duration, severity, neuropathology,⁷ lesion location,¹⁵ sociodemographics,³⁸ genetics,⁴⁹ behaviour,⁴⁸ pre-injury comorbidities,⁵⁰ and environmental differences, including access to and quality of care.⁵¹ These factors have been theorized to interact in complex ways.⁴ This study only addressed a minority of these complex relationships, further studies integrating even more sets of data are needed to address these other drivers of heterogeneity.

Conversely, this study leveraged some of the most recent and important advances in dMRI to address the major limitations of conventional approaches. We used high quality multi-shell dMRI data,⁵² as well as modelling approaches, tractography techniques, and microstructural measures robust to crossing fibers, partial volume effects, and connectivity biases.^{22-24,28-30} We used PCA to combine dMRI measures into meaningful indices of white matter structure. Lastly, we used gold-standard measures of psychiatric illness to predict clinical outcomes. Future iterations of this work will need better control of time since injury, longitudinal follow-up, broader assessment of clinical outcomes, and further development of predictive models.

Conclusion

In conclusion, leveraging advanced dMRI and a pattern-learning algorithm to parse concussion heterogeneity, we have found clinically-meaningful cross-demographic latent representations of multi-tract multi-symptom relationships. As the field moves towards large-scale studies aiming to statistically control for sociodemographic sources of heterogeneity to detect a putative consistent white matter signature of concussion across patients, the insights gained from this study should be taken into consideration: informative, clinically-meaningful, symptom-specific patterns of connectivity differences may be lost when pooling across concussed patients. This insight is an important step in improving stratification strategies for clinical trials and the identification of treatment targets.

References

- 1 Cassidy, J. D. *et al.* Incidence, risk factors and prevention of mild traumatic brain injury: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil. Med.*, 28-60 (2004).
- 2 Zhang, A. L., Sing, D. C., Rugg, C. M., Feeley, B. T. & Senter, C. The Rise of Concussions in the Adolescent Population. *Orthop J Sports Med* **4**, 2325967116662458, doi:10.1177/2325967116662458 (2016).
- 3 Mayer, A. R. *et al.* Advanced biomarkers of pediatric mild traumatic brain injury: Progress and perils. *Neurosci Biobehav Rev* **94**, 149-165, doi:10.1016/j.neubiorev.2018.08.002 (2018).
- 4 Kenzie, E. S. *et al.* Concussion As a Multi-Scale Complex System: An Interdisciplinary Synthesis of Current Knowledge. *Front Neurol* **8**, 513, doi:10.3389/fneur.2017.00513 (2017).
- 5 Hawryluk, G. W. & Bullock, M. R. Past, Present, and Future of Traumatic Brain Injury Research. *Neurosurg Clin N Am* **27**, 375-396, doi:10.1016/j.nec.2016.05.002 (2016).
- 6 Armstrong, R. C., Mierzwa, A. J., Marion, C. M. & Sullivan, G. M. White matter involvement after TBI: Clues to axon and myelin repair capacity. *Exp Neurol* **275 Pt 3**, 328-333, doi:10.1016/j.expneurol.2015.02.011 (2016).
- 7 Bigler, E. D. & Maxwell, W. L. Neuropathology of mild traumatic brain injury: relationship to neuroimaging findings. *Brain Imaging Behav* **6**, 108-136, doi:10.1007/s11682-011-9145-0 (2012).
- 8 Dodd, A. B., Epstein, K., Ling, J. M. & Mayer, A. R. Diffusion tensor imaging findings in semi-acute mild traumatic brain injury. *J Neurotrauma* **31**, 1235-1248, doi:10.1089/neu.2014.3337 (2014).
- 9 Hulkower, M. B., Poliak, D. B., Rosenbaum, S. B., Zimmerman, M. E. & Lipton, M. L. A decade of DTI in traumatic brain injury: 10 years and 100 articles later. *AJNR Am J Neuroradiol* **34**, 2064-2074, doi:10.3174/ajnr.A3395 (2013).
- 10 Langdon, S., Konigs, M., Adang, E., Goedhart, E. & Oosterlaan, J. Subtypes of Sport-Related Concussion: a Systematic Review and Meta-cluster Analysis. *Sports Med* **50**, 1829-1842, doi:10.1007/s40279-020-01321-9 (2020).
- 11 Si, B. *et al.* Sub-classifying patients with mild traumatic brain injury: A clustering approach based on baseline clinical characteristics and 90-day and 180-day outcomes. *PLoS One* **13**, e0198741, doi:10.1371/journal.pone.0198741 (2018).
- 12 Yeates, K. O. *et al.* Derivation and Initial Validation of Clinical Phenotypes of Children Presenting with Concussion Acutely in the Emergency Department: Latent Class Analysis of a Multi-Center, Prospective Cohort, Observational Study. *J Neurotrauma* **36**, 1758-1767, doi:10.1089/neu.2018.6009 (2019).
- 13 Stojanovski, S. *et al.* Microstructural abnormalities in deep and superficial white matter in youths with mild traumatic brain injury. *Neuroimage Clin* **24**, 102102, doi:10.1016/j.nicl.2019.102102 (2019).
- 14 Taylor, P. N., Moreira da Silva, N., Blamire, A., Wang, Y. & Forsyth, R. Early deviation from normal structural connectivity: A novel intrinsic severity score for mild TBI. *Neurology* **94**, e1021-e1026, doi:10.1212/WNL.0000000000008902 (2020).

- 15 Ware, J. B. *et al.* Inter-Subject Variability of Axonal Injury in Diffuse Traumatic Brain Injury. *J Neurotrauma* **34**, 2243-2253, doi:10.1089/neu.2016.4817 (2017).
- 16 Iverson, G. L. Network Analysis and Precision Rehabilitation for the Post-concussion Syndrome. *Front Neurol* **10**, 489, doi:10.3389/fneur.2019.00489 (2019).
- 17 Volkow, N. D. *et al.* The conception of the ABCD study: From substance use to a broad NIH collaboration. *Dev Cogn Neurosci* **32**, 4-7, doi:10.1016/j.dcn.2017.10.002 (2018).
- 18 Garavan, H. *et al.* Recruiting the ABCD sample: Design considerations and procedures. *Dev Cogn Neurosci* **32**, 16-22, doi:10.1016/j.dcn.2018.04.004 (2018).
- 19 Corrigan, J. D. & Bogner, J. Initial reliability and validity of the Ohio State University TBI Identification Method. *J. Head Trauma Rehabil.* **22**, 318-329, doi:10.1097/01.HTR.0000300227.67748.77 (2007).
- 20 Casey, B. J. *et al.* The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. *Dev Cogn Neurosci* **32**, 43-54, doi:10.1016/j.dcn.2018.03.001 (2018).
- 21 Theaud, G. *et al.* TractoFlow: A robust, efficient and reproducible diffusion MRI pipeline leveraging Nextflow & Singularity. *Neuroimage* **218**, 116889, doi:10.1016/j.neuroimage.2020.116889 (2020).
- 22 Girard, G., Whittingstall, K., Deriche, R. & Descoteaux, M. Towards quantitative connectivity analysis: reducing tractography biases. *Neuroimage* **98**, 266-278, doi:10.1016/j.neuroimage.2014.04.074 (2014).
- 23 Descoteaux, M., Deriche, R., Knosche, T. R. & Anwander, A. Deterministic and probabilistic tractography based on complex fibre orientation distributions. *IEEE Trans Med Imaging* **28**, 269-286, doi:10.1109/TMI.2008.2004424 (2009).
- 24 Tournier, J. D., Calamante, F. & Connelly, A. Robust determination of the fibre orientation distribution in diffusion MRI: non-negativity constrained super-resolved spherical deconvolution. *Neuroimage* **35**, 1459-1472, doi:10.1016/j.neuroimage.2007.02.016 (2007).
- 25 Sherif, T. *et al.* CBRAIN: a web-based, distributed computing platform for collaborative neuroimaging research. *Front Neuroinform* **8**, 54, doi:10.3389/fninf.2014.00054 (2014).
- 26 Klein, A. & Tourville, J. 101 labeled brain images and a consistent human cortical labeling protocol. *Front Neurosci* **6**, 171, doi:10.3389/fnins.2012.00171 (2012).
- 27 Guberman, G. I., Houde, J. C., Ptito, A., Gagnon, I. & Descoteaux, M. Structural abnormalities in thalamo-prefrontal tracks revealed by high angular resolution diffusion imaging predict working memory scores in concussed children. *Brain Struct Funct* **225**, 441-459, doi:10.1007/s00429-019-02002-8 (2020).
- 28 Raffelt, D. *et al.* Apparent Fibre Density: a novel measure for the analysis of diffusion-weighted magnetic resonance images. *Neuroimage* **59**, 3976-3994, doi:10.1016/j.neuroimage.2011.10.045 (2012).
- 29 Dell'Acqua, F., Simmons, A., Williams, S. C. & Catani, M. Can spherical deconvolution provide more information than fiber orientations? Hindrance modulated orientational anisotropy, a true-tract specific index to characterize white matter diffusion. *Hum Brain Mapp* **34**, 2464-2483, doi:10.1002/hbm.22080 (2013).

- 30 Chamberland, M. *et al.* Dimensionality reduction of diffusion MRI measures for improved tractometry of the human brain. *Neuroimage* **200**, 89-100, doi:10.1016/j.neuroimage.2019.06.020 (2019).
- 31 Boulesteix, A. L. PLS dimension reduction for classification with microarray data. *Stat Appl Genet Mol Biol* **3**, Article33, doi:10.2202/1544-6115.1075 (2004).
- 32 Wang, H. T. *et al.* Finding the needle in a high-dimensional haystack: Canonical correlation analysis for neuroscientists. *Neuroimage* **216**, 116745, doi:10.1016/j.neuroimage.2020.116745 (2020).
- 33 Beaton, D., Chin Fatt, C. R. & Abdi, H. An ExPosition of multivariate analysis with the singular value decomposition in R. *Computational Statistics & Data Analysis* **72**, 176-189, doi:10.1016/j.csda.2013.11.006 (2014).
- 34 McIntosh, A. R. & Lobaugh, N. J. Partial least squares analysis of neuroimaging data: applications and advances. *Neuroimage* **23 Suppl 1**, S250-263, doi:10.1016/j.neuroimage.2004.07.020 (2004).
- 35 Petty, C. R. *et al.* The child behavior checklist broad-band scales predict subsequent psychopathology: A 5-year follow-up. *J Anxiety Disord* **22**, 532-539, doi:10.1016/j.janxdis.2007.04.003 (2008).
- 36 Heeringa, S. G. & Berglund, P. A. *bioRxiv*, doi:10.1101/2020.02.10.942011 (2020).
- 37 Kaufman, J. *et al.* Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (K-SADS-PL): initial reliability and validity data. *J Am Acad Child Adolesc Psychiatry* **36**, 980-988, doi:10.1097/00004583-199707000-00021 (1997).
- 38 Maas, A. I. *et al.* Advancing care for traumatic brain injury: findings from the IMPACT studies and perspectives on future research. *Lancet Neurol* **12**, 1200-1210, doi:10.1016/S1474-4422(13)70234-5 (2013).
- 39 Tosetti, P. *et al.* Toward an international initiative for traumatic brain injury research. *J Neurotrauma* **30**, 1211-1222, doi:10.1089/neu.2013.2896 (2013).
- 40 Maas, A. I. *et al.* Collaborative European NeuroTrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI): a prospective longitudinal observational study. *Neurosurgery* **76**, 67-80, doi:10.1227/NEU.0000000000000575 (2015).
- 41 Bodien, Y. G. *et al.* Optimizing Outcome Assessment in Multicenter TBI Trials: Perspectives From TRACK-TBI and the TBI Endpoints Development Initiative. *J Head Trauma Rehabil* **33**, 147-157, doi:10.1097/HTR.0000000000000367 (2018).
- 42 Crossley, N. A. *et al.* The hubs of the human connectome are generally implicated in the anatomy of brain disorders. *Brain* **137**, 2382-2395, doi:10.1093/brain/awu132 (2014).
- 43 Hernandez, F. *et al.* Lateral impacts correlate with falx cerebri displacement and corpus callosum trauma in sports-related concussions. *Biomech Model Mechanobiol* **18**, 631-649, doi:10.1007/s10237-018-01106-0 (2019).
- 44 Anderson, E. D., Giudice, J. S., Wu, T., Panzer, M. B. & Meaney, D. F. Predicting Concussion Outcome by Integrating Finite Element Modeling and Network Analysis. *Front Bioeng Biotechnol* **8**, 309, doi:10.3389/fbioe.2020.00309 (2020).
- 45 Krystal, A. D. Psychiatric disorders and sleep. *Neurol Clin* **30**, 1389-1413, doi:10.1016/j.ncl.2012.08.018 (2012).

- 46 Leddy, J. J., Sandhu, H., Sodhi, V., Baker, J. G. & Willer, B. Rehabilitation of Concussion and Post-concussion Syndrome. *Sports Health* **4**, 147-154, doi:10.1177/1941738111433673 (2012).
- 47 Geeraert, B. L., Chamberland, M., Lebel, R. M. & Lebel, C. Multimodal principal component analysis to identify major features of white matter structure and links to reading. *PLoS One* **15**, e0233244, doi:10.1371/journal.pone.0233244 (2020).
- 48 Guberman, G. I. *et al.* A Prospective Study of Childhood Predictors of Traumatic Brain Injuries Sustained in Adolescence and Adulthood. *Can J Psychiatry* **65**, 36-45, doi:10.1177/0706743719882171 (2020).
- 49 Stojanovski, S. *et al.* Polygenic Risk and Neural Substrates of Attention-Deficit/Hyperactivity Disorder Symptoms in Youths With a History of Mild Traumatic Brain Injury. *Biol Psychiatry* **85**, 408-416, doi:10.1016/j.biopsych.2018.06.024 (2019).
- 50 Yue, J. K. *et al.* Pre-injury Comorbidities Are Associated With Functional Impairment and Post-concussive Symptoms at 3- and 6-Months After Mild Traumatic Brain Injury: A TRACK-TBI Study. *Front Neurol* **10**, 343, doi:10.3389/fneur.2019.00343 (2019).
- 51 Yue, J. K. *et al.* Concussion and Mild-Traumatic Brain Injury in Rural Settings: Epidemiology and Specific Health Care Considerations. *J Neurosci Rural Pract* **11**, 23-33, doi:10.1055/s-0039-3402581 (2020).
- 52 Jones, D. K., Knosche, T. R. & Turner, R. White matter integrity, fiber count, and other fallacies: the do's and don'ts of diffusion MRI. *Neuroimage* **73**, 239-254, doi:10.1016/j.neuroimage.2012.06.081 (2013).

Figure 1. Flowchart describing the participant selection procedure.

Figure 2. Schematic illustration of the study’s workflow. A. After processing DWI images using tractoflow, we applied the DKT parcellation onto each participant’s T1-weighted image that had been registered to DWI space. From each tractogram, we extracted streamlines connecting each pair of labels in the DKT parcellation. Using this approach, we built a binary connectivity matrix that displayed for all 306 subjects in the full dataset (rows), whether (black) or not (white) a streamline existed between each pair of labels (columns). B. We split the dataset into a discovery set (n=214) and a replication set (n=92). On the discovery set, we thresholded connectomes, only keeping connections that existed across 90% of participants (a threshold of 100% is illustrated here for simplicity). We then constructed connectomes of 6 scalar diffusion measures (Fractional Anisotropy (FA), Axial Diffusivity (AD), Mean Diffusivity (MD), Radial Diffusivity (RD), Apparent Fiber Density along fixels (AFDf), and Number of Fiber Orientations (NuFO)), by computing the average measures across each bundle (weighted by the streamline count across each voxel to reduce the impact of spurious streamlines). C. We stacked all columns from each connectivity matrix, creating vectors every pair of subject/connection features. We then joined together these vectors into one matrix, showing pairs of subject/connection features (rows) for every diffusion measure (columns). We then performed principal component analysis (PCA) on these matrices. Given that each pair of subject/connection feature was treated as an observation, principal component (PC) scores could be calculated for each subject/connection feature, allowing us to reconstruct connectomes, weighted by PC scores rather than the original diffusion measures. D. From these connectomes of PC scores, we selected 200 features on the basis of univariate Pearson correlations with symptom-oriented measures. We then performed partial least squares correlation on these PC-weighted features and symptom measures, which allowed us to obtain multi-tract/multi-symptom relationships, pairs of multi-tract connectivity features (“MCF”) and multi-symptom features (“MSF”). Each multi-variate feature is composed of linear combinations (weighted sums, illustrated by the black arrows called “weights”) of variables from its corresponding feature set.

Figure 3. Illustration of 3 multi-tract multi-symptom relationships obtained from PLSc on selected PC1 features (A), or selected PC2 features (B and C). Left column: Polar plots displaying the weights of all 19 symptom-oriented measures on multi-symptom features. Each circle is color-coded to match with the corresponding scatter plots. Bars pointing away from the circle center illustrate positive weights, bars pointing towards the circle center represent negative weights. White stars illustrate symptoms that were found to significantly contribute to the multi-tract multi-symptom pair based on bootstrapping analyses. The bar graph underneath the polar plots illustrates the percent covariance explained by each multi-tract multi-symptom pair, with the currently-shown pair highlighted. Right column: Scatter plots showing the expression of multi-tract connectivity features (x-axis) and multi-symptom features (y-axis). In each plot, the same 6 representative participants are labelled (1 through 6), with each plot showing the scaled symptom-oriented *measures* (i.e.: not the expression of multi-symptom features) for two participants, one expressing low levels of a multi-tract multi-symptom pair, the other expressing high levels of the pair. For each illustrated participant, positive bars illustrate symptoms that are higher than the sample average, negative bars represent symptoms that are lower than the sample average. Participants with clinical levels of CBCL Depression, Attention Problems, Anxiety Disorder, or Aggression scores are illustrated in black. A. Illustration of multi-tract multi-symptom pair 1 obtained from the PLSc performed on selected PC1 features. B. Illustration of multi-tract multi-symptom pair 3 obtained from the PLSc performed on selected PC2 features. C. Illustration of multi-tract multi-symptom pair 4 obtained from the PLSc performed on selected PC2 features.

Figure 4. Histogram illustrating correlation coefficients for all possible single-tract single-symptom relationships (blue), and between corresponding connectivity and symptom latent factors (pink) (referred to as “multi-tractmulti-symptom relationships”).

Figure 5. Line plot showing the percent overlap between univariate analyses and each connectivity latent factor. Highest overlap occurred for the LF1 from both PLSc analyses. Chord plots shown above graph illustrate which connections were found to be significant for univariate comparisons of PC1 features (red), univariate comparisons of PC2 features (blue), multi-tract connectivity feature 1 from the PLSc performed on PC1 features (violet), multi-tract connectivity feature 3 from the PLSc performed on PC2 features (turquoise), and multi-tract connectivity feature 4 from the PLSc performed on PC2 features (green). Inter-hemispheric connections are identified in each chord plot by a black outline, illustrating how many interhemispheric connections are present for univariate comparisons and for multi-tract connectivity feature 1, but less so for multi-tract connectivity features 3 and 4. The percent overlap score for each of the three illustrated multi-tract connectivity features are identified in the line plot with a circle of the corresponding color.

Figure 6. A. Bar plot illustrating the average expression of multi-tract connectivity feature 4 for participants with and without adverse psychiatric outcomes. B. Polar plot illustrating weights of individual symptoms contributing to multi-symptom feature 4.

Table 1.

Questionnaire - Description	Symptom Measured	Respondent
CBCL – Headaches	Headaches	Parent
CBCL – Nausea, feels sick	Nausea	Parent
CBCL – Vomiting, throwing up	Vomiting	Parent
CBCL – Feels dizzy or lightheaded	Dizziness	Parent
CBCL – Overtired without good reason	Fatigue	Parent
SDS – The child experiences daytime sleepiness	Drowsiness	Parent
SDS – The child has difficulty getting to sleep at night	Trouble falling asleep	Parent
CBCL – Sleep more than most kids during day and/or night	Sleep more than usual	Parent
CBCL – Sleeps less than most kids	Sleep less than usual	Parent
CBCL – Depression (DSM) T score	Sadness	Parent
CBCL – Anxiety Disorder (DSM) T score	Nervousness	Parent
CBCL – Attention Problems T score	Difficulty concentrating	Parent
NIH Toolbox Picture Sequence Memory Test – Fully-Corrected T-score	Sequence Memory (difficulty remembering)	Child
NIH Toolbox List Sorting Working Memory Test – Fully-Corrected T-score	Working memory (difficulty remembering)	Child
RAVLT Short Delay Trial VI – Total Correct	Short recall (difficulty remembering)	Child
RAVLT Long Delay Trial VII – Total Correct	Long recall (difficulty remembering)	Child
CBCL Aggression T score	Irritability	Child


Table outlining all behavioural measures used in analyses, along with the corresponding symptom they reflect. CBCL: Child Behavior Checklist. SDS: Sleep Disturbance Scale. NIH: National Institutes of Health. DSM: Diagnostics and Statistics Manual. RAVLT: Ray Auditory Verbal Learning Test.

Table 2.


Demographic and injury data	Discovery set (n=214)	Replication set (n=92)
Interview Age		
Mean (SD)	9.57 (0.496)	9.54 (0.501)
Median [Min, Max]	10.0 [9.00, 10.00]	10.0 [9.00, 10.0]
Sex		
F	88 (41.1%)	39 (42.4%)
M	126 (58.9%)	53 (57.6%)
Pubertal Stage		
Early	41 (19.2%)	18 (19.6%)
Mid	58 (27.1%)	19 (20.7%)
Prepubertal	115 (53.7%)	52 (56.5%)
Late	0 (0%)	3 (3.3%)
Race/Ethnicity		
Asian	2 (0.9%)	2 (2.2%)
Hispanic	27 (12.6%)	18 (19.6%)
Multiple	18 (8.4%)	8 (8.7%)
Non-Hispanic Black	14 (6.5%)	11 (12.0%)
Non-Hispanic White	151 (70.6%)	52 (56.5%)
Other	2 (0.9%)	1 (1.1%)
Combined Family Income		
<5K	5 (2.3%)	5 (5.4%)
\$5,000 - \$11,999	5 (2.3%)	1 (1.1%)
\$12,000-\$15,999	3 (1.4%)	2 (2.2%)
\$16,000-\$24,999	5 (2.3%)	3 (3.3%)
\$25,000-\$34,999	12 (5.6%)	4 (4.3%)
\$35,000-\$49,999	12 (5.6%)	5 (5.4%)
\$50,000-\$74,999	34 (15.9%)	16 (17.4%)
\$75,000-\$99,999	31 (14.5%)	13 (14.1%)
\$100,000-\$199,000	76 (35.5%)	27 (29.3%)
>\$200,000	31 (14.5%)	16 (17.4%)
Handedness		
LH	10 (4.7%)	10 (10.9%)
RH	175 (81.8%)	69 (75%)
Mixed	29 (13.6%)	13 (14.1%)
Injury Mechanism		
Fall/hit by object	135 (63.1%)	48 (52.2%)
Fight/shaken	2 (0.9%)	3 (3.3%)
Multiple	10 (4.7%)	5 (5.4%)
Motor vehicle collision	14 (6.5%)	3 (3.3%)
Unknown	53 (24.8%)	33 (35.9%)
Time Since Injury		
Mean (SD)	3.22 (2.79)	3.23 (2.60)

Median [Min, Max]	2.00 [0.00, 11.0]	2.50 [0.00, 9.00]
Total TBIs		
Unknown	53 (24.8%)	33 (35.9%)
1	151 (70.6%)	54 (58.7%)
2	9 (4.2%)	5 (5.4%)
3	1 (0.5%)	0 (0%)

Note: Participants with “Unknown” Injury Mechanism and Total TBIs reported sustaining a TBI but no mechanism of injury was endorsed.




medRxiv preprint doi: <https://doi.org/10.1101/2021.04.01.21254814>; this version posted April 7, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

A


+



=

**C**

medRxiv preprint doi: <https://doi.org/10.1101/2021.04.01.21254814>; this version posted April 7, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

FA

Weighting


B

Thresholding


→ 92 (replication set)









+MD
+RD

Connectivity Features**Symptoms****D**



Weights







medRxiv preprint doi: <https://doi.org/10.1101/2021.04.01.21254814>; this version posted April 7, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Single-Tract Single-Symptom Relationships




Multi-Tract Multi-Symptom Relationships





A



B

