

Survey of Behaviour Attitudes Towards Preventive Measures Following COVID-19

Vaccination

Authors:

Daniella Rahamim-Cohen MB BS, MSc¹; Sivan Gazit, MD MA²; Galit Perez, MN MA²; Barak Nada, BSc²; Shay Ben Moshe, MSc^{3,2}; Miri Mizrahi-Reuveni MD MHA¹; Joseph Azuri MD MHA^{1,4}; Tal Patalon, MD²

¹ Health Division, Maccabi Healthcare Services, Tel Aviv, Israel.

² Kahn Sagol Maccabi (KSM) Research & Innovation Center, Maccabi Healthcare Services, Israel.

³ Ben-Gurion University, Beer Sheva, Israel.

⁴ Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel.

The authors declare they have no conflict of interest.

Corresponding author: Daniella Rahamim-Cohen

cohen_dani@mac.org.il, HaMared 27, Tel Aviv, 68125, Israel

Abstract:

Following the widespread vaccination program for COVID-19 carried out in Israel, a survey was conducted to preliminarily assess behavior changes in the vaccinated population, prior to the expected upcoming policy change as to mask wearing and social distancing regulation in Israel. 200 people answered at least one question pertaining to preventive behaviour. Among the respondents, 21.1% reported a decrease in mask wearing compared to 47.3% who reported a decrease in social distancing. There was no difference in these measures between the sexes. However, people under the age of 50 were more likely to decrease mask wearing (28.1%) and decrease social distancing (56.1%), as compared with people over the age of 50 (17.2% and 41.8%, respectively). Among health care workers, there was a minimal decrease in mask wearing (1/23 people) compared to a more widespread decrease in social distancing (10/23). These data suggest that preventive attitudes change following COVID-19 vaccination, with less adherence to social distancing as compared to mask wearing, and should be taken

into account when planning public policy in the future.

medRxiv preprint doi: <https://doi.org/10.1101/2021.04.12.21255304>; this version posted April 13, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. All rights reserved. No reuse allowed without permission.

Introduction:

Since the onset of the COVID-19 global pandemic in late 2019, there have been over 130,000 million reported cases with over 3 million deaths worldwide. Most countries set various guidelines and procedures in an attempt to curb the spread of the virus, including the use of personal protective equipment (PPE) and maintaining social distancing. These measures were found to be very effective in reducing viral transmission rate¹ and were the cornerstones for health policy in tackling virus spread in most countries around the world, manifesting in lockdowns and various enforcement measures to ensure compliance.

With the introduction of vaccines against the virus, the question arises about a possible change in personal attitudes and behaviours to the recommendations for mask wearing and keeping social distancing.

Evidence from previous vaccine rollouts suggests that there may be a decrease in adherence to preventive behaviours. Following the Lyme disease vaccine rollout, there was a decrease in the use of light color clothing and tick repellent². An evaluation of behaviour following the influenza vaccine's rollout showed that people interacted with more people and in larger groups³.

Israel began vaccinating its population at the end of December 2020, using the Pfizer/BioNtech BNT162b2 COVID-19 vaccine. The vaccination program was implemented through HMO's and hospitals, with clear prioritization – beginning with the elderly and

medical staff, until the current situation where any citizen over the age of 16 is eligible to receive a vaccine.

As of April 8th, 52.8% of the Israeli population has completed the course of two vaccinations⁴ with over 5 million people vaccinated. Since the beginning of the program, Israel has seen a steady decrease in confirmed COVID-19 cases and mortality. As a result of the improvement in morbidity and mortality and the high vaccination uptake, the government has lifted many of the restrictions imposed on the population over the past year, opening shops, restaurants, and schools. Some are accessible only to fully vaccinated individuals. However, the policy to wear masks indoors and outdoors and maintain social distancing is still in effect and has not officially changed.

The possible effect of the COVID-19 vaccination program on adherence to the rules and regulations surrounding personal protective behaviour is central to formulating health policy regarding the virus in the vaccination and post-vaccination era. The UK has prepared an executive summary regarding the possible impact of widespread vaccination on adherence outcomes and how they can be mitigated⁵. To assess Israel's situation and explore if there are behavioural changes in the population following vaccination, we conducted a survey to identify behaviour attitudes in the general vaccinated population within Maccabi Healthcare Services (MHS).

Method:

A survey was sent by a text message to 2500 subjects, randomly selected, above the age of 18, vaccinated or not vaccinated, from MHS, the second largest HMO in Israel, serving

approximately 2.5 million people (about a quarter of the Israeli population) with nationwide distribution. The survey was conducted between 25/3/21- 7/4/21. Participants were asked to compare mask-wearing and social distancing before and after the vaccination. Demographics included age, sex, and being a healthcare professional.

The following analysis is preliminary, as we are currently conducting further research exploring the influence of the upcoming healthcare policy changes.

The study was approved by the MHS institutional review board.

Results:

Of 2500 surveys sent, 200 people aged 18-80 years responded to at least one question. The average time for answering was 3:39 minutes. Thirteen respondents were not vaccinated with even one vaccination. The main results are described in Table 1.

Following the second vaccination, 39 respondents reported “lower mask wearing” than before receiving the vaccination (21.1%) compared to 140 who did not change their mask wearing behaviour (75.7%) [Figure 1]. There was no substantial difference in terms of sex or being a healthcare professional in terms of decreased mask wearing; however, more people aged less than 50 decreased their mask wearing, 28.1% (16/57) compared to the group aged 50 and above, 17.2% (21/122).

When comparing social distancing behavior before and after vaccination, 89 (48.4%) reported no change, whereas 87 (47.3%) respondents reported a decrease in social distancing [Figure

2]. There was only a slight difference between males and females (38/80, 47.5%; 49/104, 47.1%, respectively).

Of the respondents aged 50 and above, 41.8% (51/122) reported keeping “less social distancing”, while 53.3% (65/122) reported no change, whereas respondents below the age of 50 reported 56.1% (32/57) and 40.4% (23/57) respectively.

Only one healthcare professional reported decreased “mask wearing” following the vaccine (1/23), in comparison to almost half (10/23) of the healthcare professionals who reported a decrease in “social distancing” following the vaccine.

Discussion

Consistent with previous findings^{2,3}, the survey showed decreased social distancing and mask wearing in specific population groups following COVID-19 vaccination. The reported decrease in mask wearing was less prevalent than the reported decrease in social distancing, which occurred in approximately half the surveyed population. In the subgroup of health professionals, it is clear that maintaining mask-wearing is still the prevailing practice, as opposed to maintaining social distancing.

Moreover, the survey suggests that people under the age of 50 are less likely to maintain the preventive behaviours of mask wearing and social distancing when compared to the older population.

Although the data above is not surprising, it should be taken into account when devising health policy regarding COVID-19 preventive behaviour in populations that have been widely vaccinated.

Limitations

The current survey is a preliminary effort to assess health behaviour in vaccinated population, prior to the upcoming policy change as to mask wearing and social distancing regulation in Israel. The cohort is relatively small, without statistical analysis. Availability of cell phone use among different populations (e.g. ultraorthodox, elderly) is of consideration, as further studies will address these issues.

References:

1. Chu, D. K. *et al.* Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet* **395**, 1973–1987 (2020).
2. Brewer, N. T., Cuite, C. L., Herrington, J. E. & Weinstein, N. D. Risk compensation and vaccination: Can getting vaccinated cause people to engage in risky behaviors? *Ann. Behav. Med.* **34**, 95–99 (2007).
3. Reiber, C. *et al.* Change in human social behavior in response to a common vaccine. *Ann. Epidemiol.* **20**, 729–733 (2010).
4. COVID-19 in Israel dashboard. (2021).
5. *SPI-B: Possible impact of the COVID-19 vaccination programme on adherence to rules and guidance about personal protective behaviours aimed at preventing spread of the virus.*

Figures and Tables

Table 1. Main results.

		All	Male	Female	Age <50	Age ≥50	Health workers
Mask usage post vaccination	Responded	185	80	105	57	122	23
	Lower	39 (21.1%)	16 (20.5%)	23 (21.9%)	16 (28.1%)	21 (17.2%)	1 (4.3%)
	No change	140 (75.7%)	62 (77.5%)	78 (74.3%)	40 (70.2%)	97 (79.5%)	22 (95.7%)
	Higher	6 (3.2%)	2 (2.5%)	4 (3.8%)	1 (1.8%)	4 (3.3%)	0 (0%)
Social distancing post vaccination	Responded	184	80	104	57	122	23
	Lower	87 (47.3%)	38 (47.5%)	49 (47.1%)	32 (56.1%)	51 (41.8%)	10 (43.5%)
	No change	89 (48.4%)	39 (48.8%)	50 (48.1%)	23 (40.4%)	65 (53.3%)	12 (52.2%)
	Higher	8 (4.3%)	3 (3.8%)	5 (4.8%)	2 (3.5%)	6 (4.9%)	1 (4.3%)

Figure 1. Mask wearing behaviour post vaccination




Figure 2. Social distancing practices post vaccination

