

1 **Estimating local outbreak risks and the effects of non-**
2 **pharmaceutical interventions in age-structured populations:**
3 **SARS-CoV-2 as a case study**

4
5 **Authors**

6 Francesca A. Lovell-Read^{1*}, Silvia Shen^{1,2}, Robin N. Thompson^{3,4}

7
8 **Affiliations**

9 ¹Mathematical Institute, University of Oxford, Oxford, UK

10 ²Pembroke College, University of Oxford, Oxford, UK

11 ³Mathematics Institute, University of Warwick, Coventry, UK

12 ⁴The Zeeman Institute for Systems Biology and Infectious Disease Epidemiology Research,
13 University of Warwick, Coventry, UK

14
15 *** Corresponding author**

16 Address: Merton College, Merton Street, Oxford, OX1 4JD

17 E-mail: francesca.lovell-read@merton.ox.ac.uk. Tel: +44 (0)1865 273525.

18
19 **Abstract**

20 During the COVID-19 pandemic, non-pharmaceutical interventions (NPIs) including school
21 closures, workplace closures and social distancing policies have been employed worldwide to
22 reduce transmission and prevent local outbreaks. However, transmission and the effectiveness of

23 NPIs depend strongly on age-related factors including heterogeneities in contact patterns and
24 pathophysiology. Here, using SARS-CoV-2 as a case study, we develop a branching process
25 model for assessing the risk that an infectious case arriving in a new location will initiate a local
26 outbreak, accounting for the age distribution of the host population. We show that the risk of a
27 local outbreak depends on the age of the index case, and we explore the effects of NPIs targeting
28 individuals of different ages. Social distancing policies that reduce contacts outside of schools
29 and workplaces and target individuals of all ages are predicted to reduce local outbreak risks
30 substantially, whereas school closures have a more limited impact. In the scenarios considered
31 here, when different NPIs are used in combination the risk of local outbreaks can be eliminated.
32 We also show that heightened surveillance of infectious individuals reduces the level of NPIs
33 required to prevent local outbreaks, particularly if enhanced surveillance of symptomatic cases is
34 combined with efforts to find and isolate nonsymptomatic infected individuals. Our results
35 reflect real-world experience of the COVID-19 pandemic, during which combinations of intense
36 NPIs have reduced transmission and the risk of local outbreaks. The general modelling
37 framework that we present can be used to estimate local outbreak risks during future epidemics
38 of a range of pathogens, accounting fully for age-related factors.

39

40 **Keywords**

41 mathematical modelling; infectious disease epidemiology; outbreak probability; SARS-CoV-2;
42 COVID-19; age-structured models; non-pharmaceutical interventions.

43

44 **1. Introduction**

45 Throughout the COVID-19 pandemic, policy makers worldwide have relied on non-
46 pharmaceutical interventions (NPIs) to limit the spread of SARS-CoV-2. Commonly introduced
47 NPIs have included school closures, workplace closures and population-wide social distancing
48 policies, all of which aim to reduce the numbers of contacts between individuals and disrupt
49 potential chains of transmission [1-4]. Similar measures have previously been adopted for
50 countering other infectious diseases such as Ebola and pandemic influenza [5-7], and are likely
51 to remain a key line of defence against emerging pathogens that are directly transmitted between
52 hosts. NPIs are particularly important when no effective treatment or vaccine is available, and
53 they are also beneficial when vaccination programmes are being rolled out [8-10]. If vaccines do
54 not prevent transmission completely, then NPIs may be important even when vaccination is
55 widespread [11]. However, the negative economic, social and non-disease health consequences
56 of NPIs have been widely discussed, with the impact of school closures on the academic progress
57 and wellbeing of school-aged individuals a particular concern [7, 12-16]. Therefore, assessing
58 the effectiveness of different NPIs at reducing transmission is critical for determining whether or
59 not they should be used.

60
61 Since NPIs such as school and workplace closures affect distinct age groups within the
62 population, when evaluating their effectiveness it is important to account for age-dependent
63 factors that influence transmission. Multiple studies have documented marked heterogeneities in
64 the patterns of contacts between individuals in different age groups, with school-aged individuals
65 tending to have more contacts each day than older individuals [17-23]. Since close contact
66 between individuals is a key driver of transmission for respiratory pathogens such as influenza

67 viruses and SARS-CoV-2, these contact patterns influence transmission dynamics and
68 consequently the effects of interventions that target different age groups [18, 19, 24-27].
69 Additionally, many diseases are characterised by significant age-related variations in
70 pathophysiology. For example, for SARS-CoV-2, children may be less susceptible to infection
71 than adults [27-31], and more likely to experience asymptomatic or subclinical courses of
72 infection [28, 31-36]. Since the secondary attack rate (the proportion of close contacts that lead
73 to new infections) from asymptomatic or subclinical hosts is lower than from hosts with clinical
74 symptoms [37-42], children are likely to be less infectious on average than older individuals who
75 are at increased risk of developing symptoms [43-45].

76
77 Previous studies have used age-stratified deterministic transmission models to investigate the
78 effects of NPIs on COVID-19 epidemic peak incidence and timing. Prem *et al.* [46] projected the
79 outbreak in Wuhan, China, over a one year period under different control scenarios, and
80 demonstrated that a period of intense control measures including school closures, a 90%
81 reduction in the workforce and a significant reduction in other social mixing could delay the
82 epidemic peak by several months. Zhang *et al.* [27] predicted that eliminating all school contacts
83 during the outbreak period would lead to a noticeable decrease in the peak incidence and a later
84 peak; however, they did not take differences between symptomatic and asymptomatic cases into
85 account explicitly. In contrast, Davies *et al.* [31] used estimates of age-dependent susceptibility
86 and clinical fraction fitted to the observed age distribution of cases in six countries to
87 demonstrate that school closures alone were unlikely to reduce SARS-CoV-2 transmission
88 substantially. Davies *et al.* [47] subsequently concluded that a combination of several strongly

89 enforced NPIs would be necessary to avoid COVID-19 cases exceeding available healthcare
90 capacity in the UK.

91
92 Rather than considering the entire epidemic curve, here we focus on estimating the probability
93 that cases introduced to a new location trigger a local outbreak as opposed to fading out with few
94 cases. Localised clusters of transmission have been a feature of the COVID-19 pandemic [48-
95 50], and assessing the risk that such local outbreaks occur requires a stochastic model in which
96 the pathogen can either invade or fade out. Stochastic branching process models have been
97 applied previously to assess outbreak risks for many pathogens without considering different age
98 groups explicitly [51-55], and extended to consider adults and children as two distinct groups
99 [56]. However, the significant heterogeneities in contact patterns and pathophysiology between
100 individuals across the full range of ages have never previously been considered in estimates of
101 local outbreak risks. Here, we develop an age-structured branching process model that can be
102 used to estimate the probability of a local outbreak occurring for index cases of different ages,
103 and demonstrate how the age-dependent risk profile changes when susceptibility to infection and
104 clinical fraction vary with age.

105
106 We use the model to investigate the effects on the local outbreak probability of NPIs that reduce
107 the numbers of contacts between individuals. Specifically, we use location-specific contact data
108 for the UK detailing the average numbers of daily contacts occurring in school, in the workplace
109 and elsewhere [17] to model the impacts of school closures, workplace closures and broader
110 social distancing policies. We demonstrate that, for SARS-CoV-2, contacts occurring outside
111 schools and the workplace are a key driver of sustained transmission. Thus, population-wide

112 social distancing policies that affect individuals of all ages lead to a substantial reduction in the
113 risk of local outbreaks. In contrast, since school-aged individuals only make up around one
114 quarter of the UK population and tend to have large numbers of contacts outside school, school
115 closures are predicted to have only a limited effect when applied as the sole NPI.

116
117 We then go on to consider the impacts of mixed strategies made up of multiple NPIs, as well as
118 additional NPIs that do not only reduce numbers of contacts. Specifically, we show that rigorous
119 surveillance and effective isolation of infected hosts can reduce the level of contact-reducing
120 NPIs required to achieve substantial reductions in the risk of local outbreaks. Although we use
121 SARS-CoV-2 as a case study, our approach can be applied more generally to explore the effects
122 of NPIs on the risk of outbreaks of any pathogen for which age-related heterogeneities play a
123 significant role in transmission dynamics.

124

125 **2. Methods**

126 **2.1 Mathematical model**

127 We considered a branching process model in which the population was divided into 16 age
128 groups, denoted G_1, G_2, \dots, G_{16} . The first 15 groups represent individuals aged 0-74, divided into
129 five-year intervals (0-4, 5-9, 10-14 etc.). The final group represents individuals aged 75 and over.
130 The total number of individuals in age group G_k is denoted N_k . Infected individuals in each age
131 group G_k are classified into compartments representing asymptomatic (A_k), presymptomatic (P_k)
132 or symptomatic (S_k) hosts, where an individual in the A_k compartment does not develop
133 symptoms at any time during their course of infection.

134

135 An infected individual of any type in group G_k may generate new infections in any age group. In
136 our model, the rate at which a single infected symptomatic individual in group G_k generates
137 infections in group G_j is given by

$$138 \quad \beta_{kj} = B\tau_k\omega_jC_{kj}.$$

139 Here, τ_k represents the infectivity of individuals in group G_k , ω_j represents the susceptibility to
140 infection of individuals in group G_j , C_{kj} represents the daily number of unique contacts a single
141 individual in group G_k has with individuals in group G_j , and B is a scaling factor that can be used
142 to set the reproduction number of the pathogen being considered (see Section 2.2). Since the
143 initial phase of potential local outbreaks are the focus of this study, we did not account for
144 depletion of susceptible hosts explicitly. The relative transmission rates from presymptomatic
145 and asymptomatic individuals compared to symptomatic individuals are given by the scaled
146 quantities $\eta\beta_{kj}$ and $\theta\beta_{kj}$, respectively, where η and θ were chosen so that the proportions of
147 transmissions generated by presymptomatic and asymptomatic hosts were in line with literature
148 estimates [57]. The parameter ξ_k represents the proportion of asymptomatic infections in group
149 G_k , so that a new infection in group G_k either increases A_k by one (with probability ξ_k) or
150 increases P_k by one (with probability $1 - \xi_k$).

151
152 A presymptomatic individual in group G_k may go on to develop symptoms (transition from P_k
153 to S_k) or be detected and isolated (so that P_k decreases by one). A symptomatic individual in
154 group G_k may be detected and isolated as a result of successful surveillance, or may be removed
155 due to self-isolation, recovery or death (so that S_k decreases by one in either case). Similarly, an
156 asymptomatic individual in group G_k may be detected and isolated or recover (so that A_k
157 decreases by one). A schematic of the different possible events in the model is shown in Fig 1.


158

159 The parameter λ represents the rate at which presymptomatic individuals develop symptoms, so
160 that the expected duration of the presymptomatic infectious period is $1/\lambda$ days in the absence of
161 surveillance of nonsymptomatic infected individuals. Similarly, the expected duration of the
162 asymptomatic infectious period in the absence of surveillance is $1/\nu$ days. The parameter μ
163 represents the rate at which symptomatic individuals are removed as a result of self-isolation,
164 recovery or death, so that the duration of time for which they are able to infect others is $1/\mu$
165 days.

166

167 For each group G_k , the rate at which symptomatic individuals are detected and isolated as a result
168 of enhanced surveillance is determined by the parameter ρ_k . Analogously, the parameter σ_k
169 governs the rate at which presymptomatic and asymptomatic individuals in G_k are detected and
170 isolated. We assumed that surveillance measures targeting nonsymptomatic hosts are equally
171 effective for those who are presymptomatic and those who are asymptomatic, and therefore used
172 the same rate of isolation due to surveillance for both of these groups.

173



174

175 **Fig 1. The branching process model used in our analyses.** Schematic showing the different possible
 176 events in the branching process model and the rates at which they occur. The parameters of the model
 177 are described in the text and in Tables 1, 2 and 3.

178

179 **2.2 Reproduction number**

180 The effective reproduction number, R , represents the expected number of secondary infections
 181 generated by a single infected individual during their entire course of infection, accounting for
 182 interventions that are in place. Here, we take a heuristic approach to derive the following
 183 expression for R :

184
$$R = \sum_{k=1}^{16} \frac{N_k}{N} \sum_{l=1}^{16} \left[\frac{\xi_k \theta \beta_{kl}}{\nu + \sigma_k} + (1 - \xi_k) \left(\frac{\eta \beta_{kl}}{\lambda + \sigma_k} + \frac{\lambda}{\lambda + \sigma_k} \frac{\beta_{kl}}{\mu + \rho_k} \right) \right], \quad (1)$$

185 where $N = N_1 + \dots + N_{16}$ is the total population size. To obtain this expression, we first consider
 186 the expected number of secondary infections an infected individual in age group G_k will generate
 187 in age group G_l . If an individual in age group G_k experiences a fully asymptomatic course of
 188 infection, which occurs with probability ξ_k , they will generate new infections in age group G_l at

189 rate $\theta\beta_{kl}$ and recover or be isolated at rate $\nu + \sigma_k$. Therefore, the total number of infections they
190 are expected to cause in age group G_l is $\theta\beta_{kl}/(\nu + \sigma_k)$. If instead the individual in age group G_k
191 experiences a symptomatic course of infection, which occurs with probability $1 - \xi_k$, whilst
192 presymptomatic they will generate new infections in age group G_l at rate $\eta\beta_{kl}$ and be isolated or
193 develop symptoms at rate $\lambda + \sigma_k$. Thus, the total number of infections they are expected to cause
194 in age group G_l whilst presymptomatic is $\eta\beta_{kl}/(\lambda + \sigma_k)$. If they go on to develop symptoms
195 before being isolated, which occurs with probability $\lambda/(\lambda + \sigma_k)$, applying similar reasoning they
196 are expected to cause $\beta_{kl}/(\mu + \rho_k)$ new infections in age group G_l whilst symptomatic.
197 Combining these possibilities leads to the term in square brackets in expression (1), which is then
198 summed over all possible age groups G_l of the infectee. Finally, to obtain the full expression (1)
199 we take a weighted average across all possible age groups G_k of the infector, where the weights
200 N_k/N represent the proportions of the population belonging to each age group. This corresponds
201 to the assumption that the initial infected host is more likely to belong to an age group containing
202 more individuals than an age group with fewer individuals.

203

204 In the absence of interventions, i.e. when $\sigma_k = \rho_k = 0$ (representing no enhanced isolation as a
205 result of surveillance) and β_{kl} is calculated using contact patterns that are characteristic of
206 normal behaviour, the effective reproduction number, R , is equal to the basic reproduction
207 number, R_0 .

208

209 **2.3 Model parameterisation**

210 The numbers of individuals in each age group (values of N_k) were chosen according to United
211 Nations age demographic data for the UK [58] (Fig 2A). The daily numbers of contacts between

212 individuals in each age group (values of C_{kj}) were set according to the 16x16 ‘contact matrix’
213 for the UK, in which the (k, j) th entry represents the expected daily number of unique contacts
214 an individual in age group G_k has with individuals in age group G_j [17]. In addition to matrices
215 representing ‘all’ contacts (Figure 2B), we also considered matrices detailing a breakdown into
216 ‘school’, ‘work’, ‘home’ and ‘other’ contacts (Figures 2C-F), allowing us to investigate the
217 effects of control interventions that reduce contacts in each of these settings.

218
219 Since we considered SARS-CoV-2 as a case study, we used studies conducted during the
220 COVID-19 pandemic to inform the epidemiological parameters of our model. Despite previous
221 research assessing the relationships between age and factors such as susceptibility to SARS-
222 CoV-2 infection or the propensity to develop symptoms, there is some variation in estimated
223 parameters between different studies. To test the robustness of our results to this uncertainty, we
224 conducted our analyses under three different scenarios (A, B and C). In scenario A, we assumed
225 that susceptibility to infection (values of ω_k) and the proportion of hosts who experience a fully
226 asymptomatic course of infection (values of ξ_k) are independent of age. In scenario B,
227 susceptibility was assumed to vary with age but the proportion of asymptomatic infections is
228 independent of age. In scenario C, we allowed both susceptibility and the asymptomatic
229 proportion to vary with age. The values used for the parameters ω_k and ξ_k in each of these three
230 scenarios are shown in Table 1 (see also [31]).

231
232 In all scenarios considered, the inherent infectivity was not assumed to be age-dependent (i.e.
233 $\tau_k = 1$ for all values of k). In other words, the expected infectiousness of infected hosts in
234 different age groups was governed solely by the proportion of asymptomatic infections in that

235 age group. We chose the scaling factors η and θ for the relative transmission rates from
 236 presymptomatic and asymptomatic individuals compared to symptomatic individuals so that the
 237 proportions of infections arising from each of these groups were in line with literature estimates
 238 (see Table 3 and [57]).

239
 240 In the absence of enhanced isolation, we set the expected duration of the presymptomatic
 241 infectious period and the time for which symptomatic individuals are able to infect others to be
 242 $1/\lambda = 2$ days and $1/\mu = 8$ days, respectively [59-62]. The asymptomatic infectious period was
 243 then chosen so that all infected individuals are expected to be infectious for the same period (i.e.
 244 $1/\nu = 10$ days). In our initial analysis, we set the isolation rates ρ_k and σ_k equal to 0 for all
 245 k ; later, we considered the effects of increasing these rates.

246
 247 Initially, we fixed $R_0 = 3$ (in line with initial estimates of SARS-CoV-2 transmissibility [63-66],
 248 before the emergence of more transmissible variants) and used expression (1) to determine the
 249 appropriate corresponding value of the scaling factor B . Later, when considering the impact of
 250 NPIs on the probability of a local outbreak, we retained this value of B and used expression (1)
 251 to determine how the reproduction number changes as a result of the control implemented.

252
 253 **Table 1. Baseline values of age-dependent parameters.** Values used for the age-dependent relative
 254 susceptibility to infection (ω_k) and the proportion of infections that are asymptomatic (ξ_k) for each of the
 255 scenarios A, B and C [31].

Age group (G_k)	Relative susceptibility (ω_k)		Asymptomatic proportion (ξ_k)	
	Scenario A	Scenarios B & C	Scenarios A & B	Scenario C
G_1, G_2 (0-9)	1.0	0.4	0.584	0.71
G_3, G_4 (10-19)		0.38	(weighted average of age-dependent	0.79
G_5, G_6 (20-29)		0.79		0.73

G_7, G_8 (30-39)		0.86	values of ξ_k in scenario C)	0.67
G_9, G_{10} (40-49)		0.8		0.6
G_{11}, G_{12} (50-59)		0.82		0.51
G_{13}, G_{14} (60-69)		0.88		0.37
G_{15}, G_{16} (70+)		0.74		0.31

256

257 **Table 2. Baseline values of scenario-independent parameters.** Values used for the parameters that were
 258 assumed not to vary between scenarios A, B and C. We also considered strategies involving enhanced
 259 surveillance ($\rho_k, \sigma_k > 0$) – see Fig 6.

Parameter	Meaning	Baseline value	Justification
R_0	Expected number of secondary infections generated by a single infectious host in the absence of interventions	$R_0 = 3$	Within estimated range for original SARS-CoV-2 virus [63-66]
λ	Rate at which presymptomatic hosts develop symptoms	$\lambda = 1/2 \text{ days}^{-1}$	[59]
μ	Rate at which symptomatic hosts are removed due to self-isolation, recovery or death	$\mu = 1/8 \text{ days}^{-1}$	[60-62]
ν	Rate at which asymptomatic hosts are removed due to recovery or death	$\nu = 1/10 \text{ days}^{-1}$	Chosen so that, in the absence of interventions, the expected duration of infection is identical for all infected hosts ($\frac{1}{\nu} = \frac{1}{\lambda} + \frac{1}{\mu}$)
τ_k	(Relative) infectivity parameter	$\tau_k = 1$ for $k = 1, \dots, 16$	Assumed
ρ_k	Isolation rate due to surveillance of symptomatic individuals	$\rho_k = 0$ for $k = 1, \dots, 16$	N/A
σ_k	Isolation rate due to surveillance of nonsymptomatic individuals	$\sigma_k = 0$ for $k = 1, \dots, 16$	N/A

260

261 **Table 3. Baseline values of scenario-dependent scaling parameters.** Values used for the scaling parameters
 262 B, η and θ for each of the scenarios A, B and C.

Parameter	Meaning	Value			Justification
		Scenario A	Scenario B	Scenario C	
B	Transmission rate scaling factor	0.0386	0.0559	0.0607	Chosen so that $R_0 = 3$ [63-66]
η	Relative transmission rate of presymptomatic hosts compared to symptomatic hosts	4.83	4.83	4.83	Chosen so that the proportion of all infections arising from presymptomatic hosts is 0.489 [57]

θ	Relative transmission rate of asymptomatic hosts compared to symptomatic hosts	0.149	0.149	0.130	Chosen so that the proportion of all infections arising from entirely asymptomatic hosts is 0.106 [57]
----------	--	-------	-------	-------	--

263

264 **2.4 Probability of a local outbreak**

265 The probability that an infected individual in a particular age group initiates a local outbreak

266 when they are introduced into the population was calculated using the branching process model.

267 One possible approach for approximating the age-dependent local outbreak probability using a

268 branching process model is to run a large number of stochastic simulations of the model starting

269 from a single infected individual in a particular age group, and record the proportion of

270 simulations in which the pathogen does not fade out after only a small number of infections [67].

271 This would then need to be repeated for index cases of different ages. Here, we instead take an

272 analytic approach, and derive a nonlinear system of simultaneous equations that determine the

273 age-dependent outbreak probabilities, as described below. The local outbreak probabilities are

274 then obtained by solving these equations numerically, and are analogous to the probabilities that

275 would be derived from the simulation approach in the limit of infinitely many simulations. The

276 benefit of our analytic approach is that it does not require a large number of stochastic

277 simulations to be run.

278

279 The probability of a local outbreak not occurring (i.e. pathogen fadeout occurs), starting from a

280 single symptomatic (or presymptomatic, asymptomatic respectively) infectious individual in age

281 group G_k , was denoted by x_k (y_k, z_k). Beginning with a single symptomatic individual in G_k ,

282 the possibilities for the next event are as follows:

283 1. The infected individual in G_k infects a susceptible individual in G_j , so that either A_j
284 increases by one (with probability ξ_j) or P_j increases by one (with probability $(1 - \xi_j)$).

285 This occurs with probability

$$286 \alpha_{kj} = \frac{\beta_{kj}}{\mu + \rho_k + \sum_{l=1}^{16} \beta_{kl}}.$$

287 2. The infected individual in G_k recovers, dies or is isolated before infecting anyone else, so
288 that S_k decreases to zero (and there are no infected individuals left in the population).

289 This occurs with probability

$$290 \gamma_k = \frac{\mu + \rho_k}{\mu + \rho_k + \sum_{l=1}^{16} \beta_{kl}}.$$

291 If there are no infectious hosts present in the population, then a local outbreak will not occur.

292 Therefore, assuming that chains of transmission arising from infectious individuals are
293 independent, the probability that no local outbreak occurs beginning from a single symptomatic
294 individual in G_k is

$$295 x_k = x_k \left[\sum_{j=1}^{16} \alpha_{kj} (\xi_j z_j + (1 - \xi_j) y_j) \right] + \gamma_k. \quad (2)$$

296 Similarly, beginning instead with a single presymptomatic individual in G_k , the possibilities for
297 the next event are:

298 1. The presymptomatic infected individual in G_k infects a susceptible individual in G_j , so
299 that (as before) either A_j increases by one (with probability ξ_j) or P_j increases by one
300 (with probability $(1 - \xi_j)$). This occurs with probability

$$301 \tilde{\alpha}_{kj} = \frac{\eta \beta_{kj}}{\lambda + \sigma_k + \eta \sum_{l=1}^{16} \beta_{kl}}.$$

302 2. The infected individual in G_k develops symptoms (transitions from P_k to S_k). This occurs
 303 with probability

$$304 \quad \delta_k = \frac{\lambda}{\lambda + \sigma_k + \eta \sum_{l=1}^{16} \beta_{kl}}.$$

305 3. The infected individual in G_k is isolated before infecting anyone else, so that S_k decreases
 306 by one. This occurs with probability

$$307 \quad \phi_k = \frac{\sigma_k}{\lambda + \sigma_k + \eta \sum_{l=1}^{16} \beta_{kl}}.$$

308 Therefore, the probability that no local outbreak occurs beginning from a single presymptomatic
 309 individual in G_k is

$$310 \quad y_k = y_k \left[\sum_{j=1}^{16} \tilde{\alpha}_{kj} (\xi_j z_j + (1 - \xi_j) y_j) \right] + \delta_k x_k + \phi_k. \quad (3)$$

311 Similarly, the probability z_k that a local outbreak does not occur starting from a single
 312 asymptomatic individual in G_k satisfies the equation

$$313 \quad z_k = z_k \left[\sum_{j=1}^{16} \hat{\alpha}_{kj} (\xi_j z_j + (1 - \xi_j) y_j) \right] + \epsilon_k, \quad (4)$$

314 where

$$315 \quad \hat{\alpha}_{kj} = \frac{\theta \beta_{kj}}{\nu + \sigma_k + \theta \sum_{l=1}^{16} \beta_{kl}} \quad \text{and} \quad \epsilon_k = \frac{\nu + \sigma_k}{\nu + \sigma_k + \theta \sum_{l=1}^{16} \beta_{kl}}.$$

316
 317 The system of simultaneous equations (2) – (4) can be solved numerically to obtain x_k, y_k and
 318 z_k (here, we did this using the MATLAB nonlinear system solver *fsolve*). Specifically, we take
 319 the minimal non-negative solution, as is standard when calculating extinction probabilities using
 320 branching process models [55, 68]. Then, for each k , the probability of a local outbreak

321 occurring beginning from a single symptomatic (or presymptomatic, asymptomatic respectively)
322 individual in group G_k is given by $1 - x_k(1 - y_k, 1 - z_k)$.

323
324 Throughout, we consider the probability p_k of a local outbreak occurring beginning from a single
325 nonsymptomatic individual in group G_k arriving in the population at the beginning of their
326 infection:

$$327 \quad p_k = \xi_k(1 - z_k) + (1 - \xi_k)(1 - y_k).$$


328
329 The average local outbreak probability, P , which is defined as the probability of a local outbreak
330 when the index case is chosen randomly from the population, is also considered. The value of P
331 is therefore a weighted average of the p_k values, where the weights correspond to the proportion
332 of the population represented by each group:

$$333 \quad P = \frac{1}{N} \sum_{k=1}^{16} N_k p_k.$$

334 This reflects an assumption that the index case is more likely to come from an age group with
335 more individuals than an age group with fewer individuals.

336
337 All computing code used to implement the above methods was written in MATLAB version
338 R2019a, and is available at <https://github.com/francescalovellread/age-dependent-outbreak-risks>.

339



340

341 **Fig 2. Age demographic and age-structured contact patterns for the United Kingdom. A.**

342 United Nations age demographic data for the UK in 2020, split into five-year age groups (where the
343 final age group contains all ages 75+) [58]. B. Heat map of UK ‘all contacts’ matrix, representing the
344 expected daily number of unique contacts that an individual in each age group G_k has with
345 individuals in each other age group G_j [17]. C. The analogous figure to B, but showing only the
346 subset of ‘all’ contacts that occur in schools (‘school’ contacts). D. The analogous figure to C, but

347 showing only ‘workplace’ contacts. E. The analogous figure to C, but showing only ‘home’ contacts.
348 F. The analogous figure to C, but showing only ‘other’ contacts (i.e. all contacts outside schools,
349 workplaces or homes).

350

351 **3. Results**

352 **3.1 Effect of the age of the index case on the risk of a local outbreak**

353 We first considered the probability that a single infected individual in a particular age group G_k
354 initiates a local outbreak when introduced into a new host population. This quantity was
355 calculated for each of the three scenarios A, B and C (Fig 3).


356

357 In scenario A, the variation in the local outbreak risk for introduced cases of different ages is
358 driven solely by the numbers of contacts between individuals. As a result, due to their higher
359 numbers of daily contacts, school- and working-age individuals are more likely to trigger a local
360 outbreak than children under five or adults over 60, with index cases aged 15-19 posing the
361 highest risk (Fig 3A). These findings do not change significantly when susceptibility is allowed
362 to vary with age in scenario B (Fig 3B). However, in scenario C, assuming that the clinical
363 fraction also varies between age groups alters the age-dependent risk profile substantially. This is
364 because asymptomatic individuals are assumed to be less infectious than symptomatic
365 individuals, and therefore an index case in an age group with a high proportion of asymptomatic
366 infections is less likely to initiate a local outbreak. In this scenario, index cases aged 40 or over
367 had a disproportionately high probability of generating a local outbreak, with individuals aged
368 70-74 presenting the highest risk (Fig 3C). These individuals are more likely to develop
369 symptoms than younger individuals (Table 1), leading to a higher expected infectiousness. In
370 contrast, individuals under the age of 40 had a below average probability of generating a local

371 outbreak, with individuals aged 10-14 presenting the lowest risk. Noticeably, individuals aged 5-
372 19 presented relatively low risks, despite the high numbers of contacts occurring among these
373 age groups (Fig 2B). In this scenario, the large number of contacts was offset by the fact that
374 individuals in these age groups are more likely to be asymptomatic and consequently less
375 infectious than older individuals (Table 1). Therefore, an index case in one of these age groups is
376 likely to lead to fewer secondary transmissions. Furthermore, the contact patterns between
377 individuals in these age groups are highly assortative with respect to age (Fig 2B). Therefore, in
378 addition to the index case being less infectious, a high proportion of the contacts they make are
379 with individuals who are also likely to be less infectious, as well as being less susceptible to
380 infection in the first place.

381
382 We performed our subsequent analyses for each of the three scenarios A, B and C, with
383 qualitatively similar results. The figures shown in the main text are for scenario C, since we
384 deem this scenario to be the most realistic for SARS-CoV-2 transmission, but the analogous
385 results for scenarios A and B are presented in Supplementary Figs S1-6.

386



387

388

389

390

391

392

Fig 3. The probability of a local outbreak depends on the age of the index case. A. The probability that a single infected individual in any given age group triggers a local outbreak (grey bars) for scenario A, in which susceptibility and clinical fraction are assumed constant across all age groups. The weighted average local outbreak probability P is shown by the black horizontal line. B. The analogous figure to A but for scenario B, in which clinical fraction is assumed constant across

393 all age groups but susceptibility varies with age (Table 1). C. The analogous figure to A but for
394 scenario C, in which both susceptibility and clinical fraction vary with age (Table 1).

395

396 **3.2 Effect of the target age group on NPI effectiveness**

397 We next considered the effects of NPIs that reduce the number of contacts between individuals
398 on the probability that an introduced case will lead to a local outbreak. To approximate the
399 relative effects of school closures, workplace closures and population-wide social distancing
400 policies, we calculated the age-dependent risk profiles when each of these types of contact were
401 excluded from the overall contact matrix.


402

403 First, we removed all ‘school’ contacts from the total contact matrix (Fig 4A). For scenario C,
404 removing ‘school’ contacts led to a 4.2% reduction in the average probability of a local outbreak
405 (from 0.449 to 0.430). This small reduction is unsurprising for scenario C, since in that scenario
406 school-aged infected individuals are assumed to be more likely to be asymptomatic than other
407 infected individuals, and therefore their expected infectiousness is lower. However, even for
408 scenarios A and B, in which school-aged individuals present the greatest risk of triggering a local
409 outbreak, the effectiveness of removing ‘school’ contacts alone at reducing the local outbreak
410 probability was limited (reductions of 7.2% and 4.75% respectively; see Supplementary Figs
411 S1A, S4A). In each scenario, the reduction in risk was predominantly for school-aged index
412 cases, with the risk from index cases of other ages only slightly reduced. Second, we considered
413 the effects of removing ‘work’ contacts from the total contact matrix (Fig 4B). This led to a more
414 substantial 25.4% reduction in the average probability of a local outbreak for scenario C (with
415 corresponding reductions of 19.0% and 24.0% for scenarios A and B respectively; see
416 Supplementary Figs S1B, S4B). As well as reducing the risk of a local outbreak from an index

417 case of working age, removing ‘work’ contacts also reduced the probability of a local outbreak
418 occurring starting from a school-aged individual. This is because closing workplaces helps to
419 block chains of transmission that begin with an infected child. For example, a transmission chain
420 involving a child transmitting to an adult at home, followed by subsequent spread around the
421 adult’s workplace, will be less likely to occur. Third, we investigated the effect of removing all
422 ‘other’ contacts, reflecting perfect social distancing being observed outside of the home, school
423 or workplace (Fig 4C). This had the most significant effect of the three types of contact-reducing
424 intervention considered, reducing the probability of a local outbreak by 41.7% for scenario C
425 (and 30.7% or 33.2% for scenarios A and B, respectively).

426
427 In the three cases described above, we considered complete reductions in ‘school’, ‘work’ and
428 ‘other’ contacts, respectively. In practice, such complete elimination of contacts is unlikely. We
429 therefore also considered partial reductions in ‘school’, ‘work’ and ‘other’ contacts, and
430 compared the resulting reductions in the local outbreak probability (Fig 4D). For any given
431 percentage reduction in contacts, reducing ‘other’ contacts always led to the largest reduction in
432 the local outbreak probability (see also Supplementary Figs S1D, S4D). This suggests that
433 reducing social contacts outside schools and workplaces can be an important component of
434 strategies to reduce the risk of local outbreaks of SARS-CoV-2. However, this alone is not
435 enough to eliminate the risk of local outbreaks entirely. For greater risk reductions using contact-
436 reducing NPIs, a mixed approach involving combinations of reductions in ‘school’, ‘workplace’
437 and ‘other’ contacts is needed.

438



439

440

441

442

443

444

445

446

447

448

449 3.3 Mixed strategies for reducing the local outbreak risk

450 Next, we considered the effects of combining reductions in 'school', 'work' and 'other' contacts

451 on the local outbreak probability (Fig 5; analogous results for scenarios A and B are shown in

452 Supplementary Figs S2 and S5). We allowed reductions in 'school' and 'work' contacts to vary

453 between 0% and 100% whilst ‘other’ contacts were reduced by 25%, 50% or 75% (Fig 5A,B,C,
454 respectively).


455

456 Since NPIs have negative economic, social and non-disease health consequences, policy makers
457 may choose to implement public health measures in which the risk of local outbreaks is not
458 eliminated completely. These results provide contact reduction targets for mixed strategies in
459 which the local outbreak probability is reduced to a pre-specified ‘acceptable’ level. For
460 example, to reduce the local outbreak probability to 0.25, ‘other’ contacts could be reduced by
461 25% from the baseline level, and ‘school’ and ‘home’ contacts reduced as indicated by the red
462 dotted contour marked ‘0.25’ in Fig 5A. Alternatively, to achieve the same local outbreak risk,
463 ‘other’ contacts can instead be reduced by 50% or by 75% with the degree of ‘school’ and
464 ‘work’ reductions chosen according to the contours marked ‘0.25’ in Figs 5B,C respectively.

465

466 If a policy maker wishes to eliminate the local outbreak risk entirely using contact-reducing
467 NPIs, for the model parameterisation considered very significant reductions in multiple types of
468 contacts are needed in combination. For example, even if all ‘school’ and ‘work’ contacts are
469 removed, ‘other’ contacts must be reduced by 66% for the overall average local outbreak
470 probability to fall below 0.01 (Fig 5D). Since such substantial reductions in multiple types of
471 contacts are unlikely to be possible, this suggests that contact-reducing NPIs must be combined
472 with other interventions, such as effective surveillance and isolation strategies, to eliminate local
473 outbreaks.

474



475

476

477

478

479

480

481

482

483

484

485 3.4 Effect of surveillance on the contact-reducing NPIs required for local outbreak control

486 We considered whether or not low local outbreak probabilities can be achieved using limited

487 contact-reducing NPIs in combination with other interventions. Specifically, the effects of

488 surveillance and isolation of infected individuals (through e.g. contact tracing) as well as
489 reducing contacts in schools, workplaces and other locations, were assessed. While results are
490 shown for scenario C in Fig 6, analogous results for scenarios A and B are presented in
491 Supplementary Figs S3 and S6.


492

493 Initially, we considered the effect of increasing the rate at which symptomatic and/or
494 nonsymptomatic infected individuals are detected and isolated as a result of surveillance, in the
495 absence of contact-reducing NPIs (i.e. with no reduction in the number of contacts between
496 individuals compared to the baseline case in Fig 2B) (Fig 6A,B). For symptomatic hosts, this
497 represents an enhanced rate of isolation compared to the baseline rate of self-isolation already
498 present in the model. Isolation of nonsymptomatic hosts was more effective at reducing the local
499 outbreak probability than isolation of symptomatic hosts (Figs 6A,B), although of course this is
500 more challenging to achieve [55]. However, if fast isolation of nonsymptomatic hosts could be
501 achieved through efficient large-scale testing (potentially in combination with contact tracing
502 [69]), the probability of local outbreaks could be reduced substantially through this measure
503 alone.

504

505 We then demonstrated the effects of combining contact-reducing NPIs with enhanced isolation
506 of infected hosts due to infection surveillance. First, we increased the enhanced isolation rate of
507 symptomatic individuals to $\rho_k = 1/2 \text{ days}^{-1}$. In the absence of other interventions, this reduced
508 the local outbreak probability by 22.0% (Figure 6C). With this level of surveillance, the local
509 outbreak risk could be reduced below 0.01 with a reduction in ‘work’ and ‘other’ contacts of
510 around 73% each, for example.

511
512 Finally, keeping the enhanced isolation rate of symptomatic individuals equal to $\rho_k =$
513 $1/2 \text{ days}^{-1}$, we increased the isolation rate of nonsymptomatic individuals to $\sigma_k = 1/7 \text{ days}^{-1}$.
514 In this case, the local outbreak probability without contact-reducing NPIs fell by 59.4%
515 compared to a situation without enhanced surveillance (Fig 6E), and the reductions in ‘work’ and
516 ‘other’ contacts needed to bring the local outbreak probability below 0.01 were significantly
517 smaller (Figure 6F). For example, if ‘work’ contacts can be reduced by 50%, then ‘other’
518 contacts only need to be reduced by 43%. This indicates that effective surveillance of both
519 symptomatic and nonsymptomatic individuals can substantially lower the extent of contact-
520 reducing NPIs that are required to achieve substantial reductions in local outbreak risks.



521
 522 **Fig 6. Surveillance as part of a mixed strategy to reduce the local outbreak probability.** A. The
 523 effect of increasing the isolation rate of symptomatic (red line) or nonsymptomatic infected hosts (blue
 524 line) on the average probability of a local outbreak (P), in the absence of contact-reducing NPIs. The
 525 isolation rates ρ_k and σ_k are varied in turn between 0 days⁻¹ and 1 days⁻¹. B. The effect of
 526 simultaneously varying the isolation rate of symptomatic and nonsymptomatic hosts on the average
 527 probability of a local outbreak (P), again without contact-reducing NPIs. C. The age-dependent

528 probability of a local outbreak when the isolation rate of symptomatic individuals is $\rho_k = 1/2 \text{ days}^{-1}$,
529 without contact-reducing NPIs or surveillance of nonsymptomatic infected individuals (purple bars and
530 solid line). Pale grey bars and black dash-dotted line represent the original local outbreak probabilities
531 without any contact-reducing NPIs or enhanced surveillance (as in Fig 3C). D. The effect of reducing
532 ‘work’ and ‘other’ contacts when the isolation rate of symptomatic infected individuals is $\rho_k =$
533 $1/2 \text{ days}^{-1}$, as in C, without surveillance of nonsymptomatic infected individuals. E,F. The analogous
534 figures to C,D, with enhanced surveillance of both symptomatic and nonsymptomatic infected hosts
535 ($\rho_k = 1/2 \text{ days}^{-1}$ and $\sigma_k = 1/7 \text{ days}^{-1}$).

536

537 **4. Discussion**

538 During the COVID-19 pandemic, public health measures that reduce the numbers of contacts
539 between individuals have been implemented in countries globally. These measures include
540 school closures, workplace closures and population-wide social distancing policies. Contact-
541 reducing NPIs have been shown to be effective at reducing SARS-CoV-2 transmission, and have
542 also been used previously during influenza pandemics [5, 70-72]. However, long-term
543 implementation of these measures has negative social, psychological and economic
544 consequences [7, 12-16]. It is therefore important to assess the effectiveness of different contact-
545 reducing NPIs at lowering transmission and preventing local outbreaks, in order to design
546 effective targeted control strategies that avoid unnecessarily strict measures.

547

548 Here, we constructed a branching process model to estimate the risk of local outbreaks under
549 different contact-reducing NPIs and different levels of surveillance for symptomatic and
550 nonsymptomatic infected individuals. Unlike previous approaches for estimating outbreak risks
551 using branching processes [51-56], we considered the effects of age-related heterogeneities

552 affecting transmission for infected individuals of a wide range of ages, including age-dependent
553 variations in social mixing patterns, susceptibility to infection and clinical fraction. Using SARS-
554 CoV-2 as a case study, we demonstrated that the risk that an introduced case initiates a local
555 outbreak depends on these age-related factors and on the age of the introduced case (Fig 3), as
556 well as the age-structure of the local population.

557

558 We used our model to assess the effects of reducing the numbers of contacts that occur in school,
559 in the workplace and elsewhere. Of the three contact-reducing NPIs considered, removing
560 ‘school’ contacts had the smallest effect on the probability of observing a local outbreak, even
561 when age-dependent variations in susceptibility and clinical fraction were ignored (Figs 4A,D,
562 Supplementary Figs S1A,D, and S4,A,D). This can be attributed to the fact that school closures
563 predominantly reduce contacts between individuals aged 5-19, who only account for
564 approximately 23% of the total population [58]. Additionally, these individuals tend to have
565 large numbers of contacts outside of the school environment (Figs 2E,F). Therefore, compared to
566 other measures, interrupting within-school transmission may have only a limited effect on
567 transmission in the wider population, particularly when school-aged individuals are less
568 susceptible to infection and more likely to experience subclinical courses of infection. In
569 contrast, reducing contacts that occur outside schools or workplaces was the most effective
570 intervention, significantly lowering the local outbreak risk across all age groups, and for those
571 aged over 60 in particular (Figs 4C,D, and Supplementary Figs S1C,D and S4C,D). This could
572 explain the success of social distancing strategies worldwide for reducing observed COVID-19
573 cases and deaths.

574

575 Mixed strategies combining reductions in ‘school’, ‘work’ and ‘other’ contacts led to greater
576 reductions in the local outbreak probability than individual interventions (Figs 5A-C), but very
577 large reductions in all three types of contact were required to eliminate the risk of local outbreaks
578 entirely (Fig 5D). However, implementing effective surveillance to identify infected hosts
579 (followed by isolation) led to substantial reductions in the risk of local outbreaks even in the
580 absence of other control measures (Figs 6A,B). In the scenarios considered here, with an
581 efficient surveillance strategy in place, significantly smaller reductions in ‘work’ and ‘other’
582 contacts were needed to render the local outbreak probability negligible, even when ‘school’
583 contacts were not reduced at all (Figs 6C-F). This supports the use of surveillance that targets
584 both symptomatic and nonsymptomatic individuals, such as contact tracing and isolation
585 strategies or population-wide diagnostic testing, to help prevent local outbreaks [55].

586
587 Although here we used SARS-CoV-2 as a case study, our model provides a framework for
588 estimating the risk of local outbreaks in age-structured populations that can be adapted for other
589 pathogens, provided sufficient data are available to parametrise the model appropriately. The
590 effects of age-structure on local outbreak risks may vary for pathogens with different
591 epidemiological characteristics. For influenza-A viruses, for example, susceptibility to infection
592 tends to decrease with age, whilst the risk of an infection leading to severe symptoms is greater
593 both for the elderly and for the very young [31, 73, 74]. This is in contrast to SARS-CoV-2, for
594 which children are more likely to experience subclinical courses of infection. In this study, we
595 used age demographic and contact data for the UK, but equivalent data for other countries are
596 available and can easily be substituted into our model to estimate outbreak risks elsewhere [17,
597 58].

598

599 One caveat of the results for SARS-CoV-2 presented here is that, although the epidemiological
600 parameters of our model were chosen to be consistent with reported literature estimates, there is
601 considerable variation between studies. In particular, the precise age-dependent variation in
602 susceptibility and clinical fraction remains unclear, and the relative infectiousness of
603 asymptomatic, presymptomatic and symptomatic hosts has not been determined exactly.
604 Furthermore, the inherent transmissibility of SARS-CoV-2 is now higher than in the initial stage
605 of the pandemic, due to the appearance of novel variants. To explore ongoing local outbreak
606 risks due to SARS-CoV-2, it would be necessary to update the model to reflect the increased
607 transmissibility of the Delta variant [75]. Due to the uncertainty in model parameter values, we
608 conducted sensitivity analyses to explore the effects of varying the parameters of the model on
609 our results (Supplementary Figs S1-12). In each case that we considered, our main conclusions
610 were unchanged: the probability that an introduced case initiates a local outbreak depends on
611 age-dependent factors affecting pathogen transmission and control, with widespread
612 interventions and combinations of NPIs reducing the risk of local outbreaks most significantly.
613

614 An important limitation of our approach to modelling contact-reducing NPIs is that we made a
615 standard assumption in our main analyses that ‘school’, ‘work’ and ‘other’ contacts are
616 independent [27, 31, 47]. In other words, reducing the numbers of contacts in one location did
617 not affect the numbers of contacts occurring in another. In reality, this is unlikely to be the case.
618 For example, closing schools also affects workplace contacts, as adults may then work from
619 home in order to fulfil childcare requirements. Additionally, the contact data that we used
620 represent the number of unique contacts per day and do not reflect the numbers of repeated

621 contacts with the same person, which affect the risk of transmission between individuals. These
622 assumptions could in principle be removed, if relevant data become available – for example, data
623 describing the effects of school closures on numbers of contacts in other locations. To
624 demonstrate how changes in multiple types of contact due to NPIs could be implemented in the
625 model, we conducted a supplementary analysis in which we investigated the effects of removing
626 all ‘school’ contacts and allowing for concurrent changes in ‘work’, ‘home’ and ‘other’ contacts
627 (Supplementary Figs S13-14). These results support our conclusion that school closures are
628 unlikely to have a substantial impact on SARS-CoV-2 transmission when applied as the sole
629 NPI. The benefit of school closures could even potentially be outweighed by the possible
630 secondary effects on other types of contacts. An improved understanding of how NPIs affect
631 different types of contact is important for more accurate assessments of interventions in future.

632
633 Despite the simplifications made, our model provides a useful framework for estimating the risk
634 of local outbreaks and the effects of NPIs. Different measures can be considered in combination
635 in the model to develop strategies for lowering local outbreak risks. Our results emphasise the
636 importance of quantifying age-dependent factors that affect transmission dynamics, such as
637 susceptibility to infection and the proportion of hosts who develop clinical symptoms, for
638 individuals of different ages. As we have shown, it is crucial to take age-dependent factors into
639 account when assessing local outbreak risks and designing public health measures.

640

641 **Declaration of interests**

642 We declare no competing interests.

643

644 **Authors' contributions**

645 Conceptualisation: All authors. Methodology: FALR, RNT. Investigation: FALR, SS.

646 Writing - original draft: FALR, RNT. Writing - review and editing: All authors. Supervision:

647 RNT.

648

649 **Data availability**

650 All computer code used in this paper was written in MATLAB version R2019a, and is available

651 at the following GitHub repository: <https://github.com/francescalovellread/age-dependent->

652 [outbreak-risks](#).

653

654 **Funding**

655 FALR acknowledges funding from the Biotechnology and Biological Sciences Research Council

656 (BBSRC), grant number BB/M011224/1. SS was funded by a BBSRC Research Experience

657 Placement and a Rokos award for undergraduate research from Pembroke College Oxford. RNT

658 was supported by UK Research and Innovation (UKRI) via grant EP/V053507/1. The funders

659 had no role in the study design; the collection, analysis and interpretation of data; the writing of

660 the report; or the decision to submit the article for publication.

661

662 **References**

663 [1] Perra, N. 2021 Non-pharmaceutical interventions during the COVID-19 pandemic: A review.

664 *Phys. Rep.* **913**, 1-52. (doi:10.1016/j.physrep.2021.02.001).

- 665 [2] Regmi, K. & Lwin, C.M. 2020 Impact of non-pharmaceutical interventions for reducing
666 transmission of COVID-19: a systematic review and meta-analysis protocol. *BMJ Open* **10**,
667 e041383. (doi:10.1136/bmjopen-2020-041383).
- 668 [3] Imai, N., Gaythorpe, K.A.M., Abbott, S., Bhatia, S., van Elsland, S., Prem, K., Liu, Y. &
669 Ferguson, N.M. 2020 Adoption and impact of non-pharmaceutical interventions for COVID-19.
670 *Wellcome Open Res* **5**, 59-59. (doi:10.12688/wellcomeopenres.15808.1).
- 671 [4] Thompson, R.N., Hollingsworth, T.D., Isham, V., Arribas-Bel, D., Ashby, B., Britton, T.,
672 Challenor, P., Chappell, L.H.K., Clapham, H., Cunniffe, N.J., et al. 2020 Key questions for
673 modelling COVID-19 exit strategies. *Proc. R. Soc. Lond. B. Biol. Sci.* **287**, 20201405.
674 (doi:10.1098/rspb.2020.1405).
- 675 [5] Hatchett, R.J., Mecher, C.E. & Lipsitch, M. 2007 Public health interventions and epidemic
676 intensity during the 1918 influenza pandemic. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 7582.
677 (doi:10.1073/pnas.0610941104).
- 678 [6] Kirsch, T.D., Moseson, H., Massaquoi, M., Nyenswah, T.G., Goodermote, R., Rodriguez-
679 Barraquer, I., Lessler, J., Cumings, D.A.T. & Peters, D.H. 2017 Impact of interventions and the
680 incidence of ebola virus disease in Liberia—implications for future epidemics. *Health Policy*
681 *Plan.* **32**, 205-214. (doi:10.1093/heapol/czw113).
- 682 [7] Berkman, B.E. 2008 Mitigating Pandemic Influenza: The Ethics of Implementing a School
683 Closure Policy. *J. Public Health Manag. Pract.* **14**, 372-378.
684 (doi:10.1097/01.PHH.0000324566.72533.0b).
- 685 [8] Moore, S., Hill, E.M., Tildesley, M.J., Dyson, L. & Keeling, M.J. 2021 Vaccination and non-
686 pharmaceutical interventions for COVID-19: a mathematical modelling study. *Lancet Infect Dis*
687 **21**, 793-802. (doi:10.1016/S1473-3099(21)00143-2).

- 688 [9] Spinelli, M.A., Glidden, D.V., Gennatas, E.D., Bielecki, M., Beyrer, C., Rutherford, G.,
689 Chambers, H., Goosby, E. & Gandhi, M. 2021 Importance of non-pharmaceutical interventions
690 in lowering the viral inoculum to reduce susceptibility to infection by SARS-CoV-2 and
691 potentially disease severity. *Lancet Infect Dis* **21**, E296-E301. (doi:10.1016/S1473-
692 3099(20)30982-8).
- 693 [10] Thompson, R.N., Hill, E.M. & Gog, J.R. 2021 SARS-CoV-2 incidence and vaccine escape.
694 *Lancet Infect Dis* **21**, 913-914. (doi:10.1016/S1473-3099(21)00202-4).
- 695 [11] Sachak-Patwa, R., Byrne, H.M., Dyson, L. & Thompson, R.N. 2021 The risk of SARS-
696 CoV-2 outbreaks in low prevalence settings following the removal of travel restrictions.
697 *Communications Medicine* **1**, 39. (doi:10.1038/s43856-021-00038-8).
- 698 [12] Donohue, J.M. & Miller, E. 2020 COVID-19 and School Closures. *JAMA* **324**, 845-847.
699 (doi:10.1001/jama.2020.13092).
- 700 [13] Silverman, M., Sibbald, R. & Stranges, S. 2020 Ethics of COVID-19-related school
701 closures. *Can. J. Public Health* **111**, 462-465. (doi:10.17269/s41997-020-00396-1).
- 702 [14] Sadique, M.Z., Adams, E.J. & Edmunds, W.J. 2008 Estimating the costs of school closure
703 for mitigating an influenza pandemic. *BMC Public Health* **8**, 135. (doi:10.1186/1471-2458-8-
704 135).
- 705 [15] Brooks, S.K., Webster, R.K., Smith, L.E., Woodland, L., Wessely, S., Greenberg, N. &
706 Rubin, G.J. 2020 The psychological impact of quarantine and how to reduce it: rapid review of
707 the evidence. *The Lancet* **395**, 912-920. (doi:10.1016/S0140-6736(20)30460-8).
- 708 [16] Loades, M.E., Chatburn, E., Higson-Sweeney, N., Reynolds, S., Shafran, R., Brigden, A.,
709 Linney, C., McManus, M.N., Borwick, C. & Crawley, E. 2020 Rapid Systematic Review: The
710 Impact of Social Isolation and Loneliness on the Mental Health of Children and Adolescents in

- 711 the Context of COVID-19. *J. Am. Acad. Child Adolesc. Psychiatry* **59**, 1218-1239.e1213.
712 (doi:10.1016/j.jaac.2020.05.009).
- 713 [17] Prem, K., Cook, A.R. & Jit, M. 2017 Projecting social contact matrices in 152 countries
714 using contact surveys and demographic data. *PLoS Comp. Biol.* **13**, e1005697.
715 (doi:10.1371/journal.pcbi.1005697).
- 716 [18] Mossong, J., Hens, N., Jit, M., Beutels, P., Auranen, K., Mikolajczyk, R., Massari, M.,
717 Salmaso, S., Tomba, G.S., Wallinga, J., et al. 2008 Social Contacts and Mixing Patterns Relevant
718 to the Spread of Infectious Diseases. *PLoS Med.* **5**, e74. (doi:10.1371/journal.pmed.0050074).
- 719 [19] Danon, L., Read, J.M., House, T.A., Vernon, M.C. & Keeling, M.J. 2013 Social encounter
720 networks: characterizing Great Britain. *Proc. R. Soc. Lond. B. Biol. Sci.* **280**, 20131037.
721 (doi:10.1098/rspb.2013.1037).
- 722 [20] Leung, K., Jit, M., Lau, E.H.Y. & Wu, J.T. 2017 Social contact patterns relevant to the
723 spread of respiratory infectious diseases in Hong Kong. *Sci. Rep.* **7**. (doi:10.1038/s41598-017-
724 08241-1).
- 725 [21] Béraud, G., Kazmerczak, S., Beutels, P., Levy-Bruhl, D., Lenne, X., Mielcarek, N.,
726 Yazdanpanah, Y., Boëlle, P.-Y., Hens, N. & Dervaux, B. 2015 The French Connection: The First
727 Large Population-Based Contact Survey in France Relevant for the Spread of Infectious
728 Diseases. *PLoS One* **10**, e0133203. (doi:10.1371/journal.pone.0133203).
- 729 [22] Ibuka, Y., Ohkusa, Y., Sugawara, T., Chapman, G.B., Yamin, D., Atkins, K.E., Taniguchi,
730 K., Okabe, N. & Galvani, A.P. 2016 Social contacts, vaccination decisions and influenza in
731 Japan. *J. Epidemiol. Community Health* **70**, 162-167. (doi:10.1136/jech-2015-205777).

- 732 [23] Read, J.M., Lessler, J., Riley, S., Wang, S., Tan, L.J., Kwok, K.O., Guan, Y., Jiang, C.Q. &
733 Cummings, D.A.T. 2014 Social mixing patterns in rural and urban areas of southern China. *Proc.*
734 *R. Soc. Lond. B. Biol. Sci.* **281**, 20140268. (doi:10.1098/rspb.2014.0268).
- 735 [24] Read, J.M., Edmunds, W.J., Riley, S., Lessler, J. & Cummings, D.A.T. 2012 Close
736 encounters of the infectious kind: methods to measure social mixing behaviour. *Epidemiol.*
737 *Infect.* **140**, 2117-2130. (doi:10.1017/S0950268812000842).
- 738 [25] Wallinga, J., Teunis, P. & Kretzschmar, M. 2006 Using Data on Social Contacts to Estimate
739 Age-specific Transmission Parameters for Respiratory-spread Infectious Agents. *Am. J.*
740 *Epidemiol.* **164**, 936-944. (doi:10.1093/aje/kwj317).
- 741 [26] Kucharski, A.J., Kwok, K.O., Wei, V.W.I., Cowling, B.J., Read, J.M., Lessler, J.,
742 Cummings, D.A. & Riley, S. 2014 The Contribution of Social Behaviour to the Transmission of
743 Influenza A in a Human Population. *PLoS Pathog.* **10**, e1004206.
744 (doi:10.1371/journal.ppat.1004206).
- 745 [27] Zhang, J., Litvinova, M., Liang, Y., Wang, Y., Wang, W., Zhao, S., Wu, Q., Merler, S.,
746 Viboud, C., Vespignani, A., et al. 2020 Changes in contact patterns shape the dynamics of the
747 COVID-19 outbreak in China. *Science* **368**, 1481-1486. (doi:10.1126/science.abb8001).
- 748 [28] Munro, A.P. & Faust, S.N. 2020 COVID-19 in children: current evidence and key
749 questions. *Curr. Opin. Infect. Dis.* **33**, 540-547. (doi:10.1097/QCO.0000000000000690).
- 750 [29] Viner, R.M., Mytton, O.T., Bonell, C., Melendez-Torres, G.J., Ward, J., Hudson, L.,
751 Waddington, C., Thomas, J., Russell, S., van der Klis, F., et al. 2021 Susceptibility to SARS-
752 CoV-2 Infection Among Children and Adolescents Compared With Adults: A Systematic
753 Review and Meta-analysis. *JAMA Pediatrics* **175**, 143-156.
754 (doi:10.1001/jamapediatrics.2020.4573).

- 755 [30] Goldstein, E., Lipsitch, M. & Cevik, M. 2020 On the effect of age on the transmission of
756 SARS-CoV-2 in households, schools and the community. *J. Infect. Dis.* **223**, 362-369.
757 (doi:10.1093/infdis/jiaa691).
- 758 [31] Davies, N.G., Klepac, P., Liu, Y., Prem, K., Jit, M., Pearson, C.A.B., Quilty, B.J.,
759 Kucharski, A.J., Gibbs, H., Clifford, S., et al. 2020 Age-dependent effects in the transmission
760 and control of COVID-19 epidemics. *Nat. Med.* **26**, 1205-1211. (doi:10.1038/s41591-020-0962-
761 9).
- 762 [32] Ludvigsson, J.F. 2020 Systematic review of COVID-19 in children shows milder cases and
763 a better prognosis than adults. *Acta paediatrica (Oslo, Norway : 1992)* **109**, 1088-1095.
764 (doi:10.1111/apa.15270).
- 765 [33] Dong, Y., Mo, X., Hu, Y., Qi, X., Jiang, F., Jiang, Z. & Tong, S. 2020 Epidemiology of
766 COVID-19 Among Children in China. *Pediatrics* **145**, e20200702. (doi:10.1542/peds.2020-
767 0702).
- 768 [34] Siebach, M.K., Piedimonte, G. & Ley, S.H. 2021 COVID-19 in childhood: Transmission,
769 clinical presentation, complications and risk factors. *Pediatr. Pulmonol.* **56**, 1342-1356.
770 (doi:10.1002/ppul.25344).
- 771 [35] Mehta, N.S., Mytton, O.T., Mullins, E.W.S., Fowler, T.A., Falconer, C.L., Murphy, O.B.,
772 Langenberg, C., Jayatunga, W.J.P., Eddy, D.H. & Nguyen-Van-Tam, J.S. 2020 SARS-CoV-2
773 (COVID-19): What Do We Know About Children? A Systematic Review. *Clin. Infect. Dis.* **71**,
774 2469-2479. (doi:10.1093/cid/ciaa556).
- 775 [36] Han, M.S., Choi, E.H., Chang, S.H., Jin, B.-L., Lee, E.J., Kim, B.N., Kim, M.K., Doo, K.,
776 Seo, J.-H., Kim, Y.-J., et al. 2021 Clinical Characteristics and Viral RNA Detection in Children

- 777 With Coronavirus Disease 2019 in the Republic of Korea. *JAMA Pediatrics* **175**, 73-80.
778 (doi:10.1001/jamapediatrics.2020.3988).
- 779 [37] Pollock, A.M. & Lancaster, J. 2020 Asymptomatic transmission of covid-19. *BMJ* **371**,
780 m4851. (doi:10.1136/bmj.m4851).
- 781 [38] Buitrago-Garcia, D., Egli-Gany, D., Counotte, M.J., Hossmann, S., Imeri, H., Ipekci, A.M.,
782 Salanti, G. & Low, N. 2020 Occurrence and transmission potential of asymptomatic and
783 presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis. *PLoS*
784 *Med.* **17**, e1003346. (doi:10.1371/journal.pmed.1003346).
- 785 [39] Byambasuren, O., Cardona, M., Bell, K., Clark, J., McLaws, M.-L. & Glasziou, P. 2020
786 Estimating the extent of asymptomatic COVID-19 and its potential for community transmission:
787 Systematic review and meta-analysis. *JAMMI* **5**, 223-234. (doi:10.3138/jammi-2020-0030).
- 788 [40] Koh, W.C., Naing, L., Chaw, L., Rosledzana, M.A., Alikhan, M.F., Jamaludin, S.A., Amin,
789 F., Omar, A., Shazli, A., Griffith, M., et al. 2020 What do we know about SARS-CoV-2
790 transmission? A systematic review and meta-analysis of the secondary attack rate and associated
791 risk factors. *PLoS One* **15**, e0240205. (doi:10.1371/journal.pone.0240205).
- 792 [41] Madewell, Z.J., Yang, Y., Longini, I.M., Jr., Halloran, M.E. & Dean, N.E. 2020 Household
793 Transmission of SARS-CoV-2: A Systematic Review and Meta-analysis. *JAMA Network Open*
794 **3**, e2031756-e2031756. (doi:10.1001/jamanetworkopen.2020.31756).
- 795 [42] Qiu, X., Nergiz, A.I., Maraolo, A.E., Bogoch, I.I., Low, N. & Cevik, M. 2021 Defining the
796 role of asymptomatic and pre-symptomatic SARS-CoV-2 transmission - a living systematic
797 review. *Clin. Microbiol. Infect.* **27**, 511-519. (doi:10.1016/j.cmi.2021.01.011).
- 798 [43] Pijls, B.G., Jolani, S., Atherley, A., Derckx, R.T., Dijkstra, J.I.R., Franssen, G.H.L.,
799 Hendriks, S., Richters, A., Venemans-Jellema, A., Zalpuri, S., et al. 2021 Demographic risk

800 factors for COVID-19 infection, severity, ICU admission and death: a meta-analysis of 59
801 studies. *BMJ Open* **11**, e044640. (doi:10.1136/bmjopen-2020-044640).

802 [44] Verity, R., Okell, L.C., Dorigatti, I., Winskill, P., Whittaker, C., Imai, N., Cuomo-
803 Dannenburg, G., Thompson, H., Walker, P.G.T., Fu, H., et al. 2020 Estimates of the severity of
804 coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis* **20**, 669-677.
805 (doi:10.1016/S1473-3099(20)30243-7).

806 [45] Chen, T., Dai, Z., Mo, P., Li, X., Ma, Z., Song, S., Chen, X., Luo, M., Liang, K., Gao, S., et
807 al. 2020 Clinical Characteristics and Outcomes of Older Patients with Coronavirus Disease 2019
808 (COVID-19) in Wuhan, China: A Single-Centered, Retrospective Study. *J. Gerontol. A Biolo.*
809 *Sci. Med. Sci.* **75**, 1788-1795. (doi:10.1093/gerona/glaa089).

810 [46] Prem, K., Liu, Y., Russell, T.W., Kucharski, A.J., Eggo, R.M., Davies, N., Flasche, S.,
811 Clifford, S., Pearson, C.A.B., Munday, J.D., et al. 2020 The effect of control strategies to reduce
812 social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a modelling study.
813 *Lancet Public Health* **5**, e261-e270. (doi:10.1016/S2468-2667(20)30073-6).

814 [47] Davies, N.G., Kucharski, A.J., Eggo, R.M., Gimma, A., Edmunds, W.J. & Centre for the
815 Mathematical Modelling of Infectious Diseases COVID-19 working group. 2020 Effects of non-
816 pharmaceutical interventions on COVID-19 cases, deaths, and demand for hospital services in
817 the UK: a modelling study. *Lancet Public Health* **5**, e375-e385. (doi:10.1016/S2468-
818 2667(20)30133-X).

819 [48] Liu, T., Gong, D., Xiao, J., Hu, J., He, G., Rong, Z. & Ma, W. 2020 Cluster infections play
820 important roles in the rapid evolution of COVID-19 transmission: A systematic review. *Int. J.*
821 *Infect. Dis.* **99**, 374-380. (doi:10.1016/j.ijid.2020.07.073).

- 822 [49] Leclerc, Q., Fuller, N., Knight, L., CMMID COVID-19 Working Group, Funk, S. & Knight,
823 G. 2020 What settings have been linked to SARS-CoV-2 transmission clusters? [version 2; peer
824 review: 2 approved]. *Wellcome Open Res* **5**. (doi:10.12688/wellcomeopenres.15889.2).
- 825 [50] Rossman, H., Keshet, A., Shilo, S., Gavrieli, A., Bauman, T., Cohen, O., Shelly, E., Balicer,
826 R., Geiger, B., Dor, Y., et al. 2020 A framework for identifying regional outbreak and spread of
827 COVID-19 from one-minute population-wide surveys. *Nat. Med.* **26**, 634-638.
828 (doi:10.1038/s41591-020-0857-9).
- 829 [51] Althaus, C.L., Low, N., Musa, E.O., Shuaib, F. & Gsteiger, S. 2015 Ebola virus disease
830 outbreak in Nigeria: Transmission dynamics and rapid control. *Epidemics* **11**, 80-84.
831 (doi:10.1016/j.epidem.2015.03.001).
- 832 [52] Thompson, R.N., Jalava, K. & Obolski, U. 2019 Sustained transmission of Ebola in new
833 locations: more likely than previously thought. *Lancet Infect Dis* **19**, 1058-1059.
834 (doi:10.1016/S1473-3099(19)30483-9).
- 835 [53] Thompson, R.N. 2020 Novel Coronavirus Outbreak in Wuhan, China, 2020: Intense
836 Surveillance Is Vital for Preventing Sustained Transmission in New Locations. *J. Clin. Med.* **9**,
837 498. (doi:10.3390/jcm9020498).
- 838 [54] Hellewell, J., Abbott, S., Gimma, A., Bosse, N.I., Jarvis, C.I., Russell, T.W., Munday, J.D.,
839 Kucharski, A.J., Edmunds, W.J., Sun, F., et al. 2020 Feasibility of controlling COVID-19
840 outbreaks by isolation of cases and contacts. *Lancet Glob. Health* **8**, e488-e496.
841 (doi:10.1016/S2214-109X(20)30074-7).
- 842 [55] Lovell-Read, F.A., Funk, S., Obolski, U., Donnelly, C.A. & Thompson, R.N. 2021
843 Interventions targeting nonsymptomatic cases can be important to prevent local outbreaks:
844 SARS-CoV-2 as a case-study. *J. R. Soc. Interface* **18**, 20201014. (doi:10.1098/rsif.2020.1014).

- 845 [56] Nishiura, H., Cook, A.R. & Cowling, B.J. 2011 Assortativity and the Probability of
846 Epidemic Extinction: A Case Study of Pandemic Influenza A (H1N1-2009). *Interdiscip.*
847 *Perspect. Infect. Dis.* **2011**, 194507. (doi:10.1155/2011/194507).
- 848 [57] Ferretti, L., Wymant, C., Kendall, M., Zhao, L., Nurtay, A., Abeler-Dorner, L., Parker, M.,
849 Bonsall, D. & Fraser, C. 2020 Quantifying SARS-CoV-2 transmission suggests epidemic control
850 with digital contact tracing. *Science* **368**, eabb6936. (doi:10.1126/science.abb6936).
- 851 [58] United Nations, Department of Economic and Social Affairs, Population Division. 2019
852 World Population Prospects 2019, Online Edition, Rev. 1. Available at:
853 <https://population.un.org/wpp/Download/Standard/Population/>.
- 854 [59] Wei, W.E., Li, Z., Chiew, C.J., Yong, S.E., Toh, M.P. & Lee, V.J. 2020 Presymptomatic
855 Transmission of SARS-CoV-2 — Singapore, January 23–March 16, 2020. *Morb. Mortal. Weekly*
856 *Rep.* **69**, 411-415. (doi:10.15585/mmwr.mm6914e1external icon).
- 857 [60] Arons, M.M., Hatfield, K.M., Reddy, S.C., Kimball, A., James, A., Jacobs, J.R., Taylor, J.,
858 Spicer, K., Bardossy, A.C., Oakley, L.P., et al. 2020 Presymptomatic SARS-CoV-2 Infections
859 and Transmission in a Skilled Nursing Facility. *N. Engl. J. Med.* **382**, 2081-2090.
860 (doi:10.1056/NEJMoa2008457).
- 861 [61] Bullard, J., Dust, K., Funk, D., Strong, J.E., Alexander, D., Garnett, L., Boodman, C., Bello,
862 A., Hedley, A., Schiffman, Z., et al. 2020 Predicting Infectious Severe Acute Respiratory
863 Syndrome Coronavirus 2 From Diagnostic Samples. *Clin. Infect. Dis.* **71**, 2663-2666.
864 (doi:10.1093/cid/ciaa638).
- 865 [62] Wolfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Muller, M.A.,
866 Niemeyer, D., Jones, T.C., Vollmar, P., Rothe, C., et al. 2020 Virological assessment of

- 867 hospitalized patients with COVID-2019. *Nature* **581**, 465-469. (doi:10.1038/s41586-020-2196-
868 x).
- 869 [63] Lai, C.-C., Shih, T.-P., Ko, W.-C., Tang, H.-J. & Hsueh, P.-R. 2020 Severe acute respiratory
870 syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The
871 epidemic and the challenges. *Int. J. Antimicrob. Agents* **55**, 105924.
872 (doi:10.1016/j.ijantimicag.2020.105924).
- 873 [64] Liu, Y., Gayle, A.A., Wilder-Smith, A. & Rocklöv, J. 2020 The reproductive number of
874 COVID-19 is higher compared to SARS coronavirus. *J. Travel Med.* **27**, taaa021.
875 (doi:10.1093/jtm/taaa021).
- 876 [65] Zhai, P., Ding, Y., Wu, X., Long, J., Zhong, Y. & Li, Y. 2020 The epidemiology, diagnosis
877 and treatment of COVID-19. *Int. J. Antimicrob. Agents* **55**, 105955-105955.
878 (doi:10.1016/j.ijantimicag.2020.105955).
- 879 [66] Zhao, S., Lin, Q., Ran, J., Musa, S.S., Yang, G., Wang, W., Lou, Y., Gao, D., Yang, L., He,
880 D., et al. 2020 Preliminary estimation of the basic reproduction number of novel coronavirus
881 (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the
882 outbreak. *Int. J. Infect. Dis.* **92**, 214-217. (doi:10.1016/j.ijid.2020.01.050).
- 883 [67] Thompson, R.N., Gilligan, C.A. & Cunniffe, N.J. 2020 Will an outbreak exceed available
884 resources for control? Estimating the risk from invading pathogens using practical definitions of
885 a severe epidemic. *J. R. Soc. Interface* **17**, 20200690. (doi:10.1098/rsif.2020.0690).
- 886 [68] Norris, J.R. 1997 *Markov Chains (Cambridge Series in Statistical and Probabilistic*
887 *Mathematics)*. Cambridge, Cambridge University Press.

- 888 [69] Hart, W.S., Maini, P.K. & Thompson, R.N. 2021 High infectiousness immediately before
889 COVID-19 symptom onset highlights the importance of continued contact tracing. *Elife* **10**.
890 (doi:10.7554/eLife.65534).
- 891 [70] Huang, Q.S., Wood, T., Jelley, L., Jennings, T., Jefferies, S., Daniells, K., Nesdale, A.,
892 Dowell, T., Turner, N., Campbell-Stokes, P., et al. 2021 Impact of the COVID-19
893 nonpharmaceutical interventions on influenza and other respiratory viral infections in New
894 Zealand. *Nat. Commun.* **12**, 1001. (doi:10.1038/s41467-021-21157-9).
- 895 [71] Flaxman, S., Mishra, S., Gandy, A., Unwin, H.J.T., Mellan, T.A., Coupland, H., Whittaker,
896 C., Zhu, H., Berah, T., Eaton, J.W., et al. 2020 Estimating the effects of non-pharmaceutical
897 interventions on COVID-19 in Europe. *Nature* **584**, 257-261. (doi:10.1038/s41586-020-2405-7).
- 898 [72] Borse, R.H., Behravesh, C.B., Dumanovsky, T., Zucker, J.R., Swerdlow, D., Edelson, P.,
899 Choe-Castillo, J. & Meltzer, M.I. 2011 Closing Schools in Response to the 2009 Pandemic
900 Influenza A H1N1 Virus in New York City: Economic Impact on Households. *Clin. Infect. Dis.*
901 **52**, S168-S172. (doi:10.1093/cid/ciq033).
- 902 [73] Greer, A.L., Tuite, A. & Fisman, D.N. 2010 Age, influenza pandemics and disease
903 dynamics. *Epidemiol. Infect.* **138**, 1542-1549. (doi:10.1017/S0950268810000579).
- 904 [74] Clohisey, S. & Baillie, J.K. 2019 Host susceptibility to severe influenza A virus infection.
905 *Crit. Care* **23**, 303. (doi:10.1186/s13054-019-2566-7).
- 906 [75] Campbell, F., Archer, B., Laurenson-Schafer, H., Jinnai, Y., Konings, F., Batra, N., Pavlin,
907 B., Vandemaele, K., Van Kerkhove, M.D., Jombart, T., et al. 2021 Increased transmissibility and
908 global spread of SARS-CoV-2 variants of concern as at June 2021. *Eurosurveillance* **26**,
909 2100509. (doi:10.2807/1560-7917.ES.2021.26.24.2100509).

