

Quantifying COVID-19 vaccination hesitancy during early vaccination rollout in Canada

Authors: Xuyang Tang (1), Hellen Gelband (1), Nico Nagelkerke (1), Isaac I. Bogoch (2), Patrick Brown (1), Ed Morawski (2), Theresa Tam (1), Angus Reid (1), Prabhat Jha (1) for the Action to beat coronavirus/Action pour battre le coronavirus (Ab-C) Study Investigators*

Affiliations:

- (1) Centre for Global Health Research, Unity Health Toronto and Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada
- (2) University Health Network, Ontario Canada
- (3) Angus Reid Forum, Toronto, Ontario, Canada

Corresponding author and address:

Prabhat Jha, MD, DPhil,
30 Bond Street
Toronto, ON, M5B 1W8, Canada
Email: Prabhat.jha@utoronto.ca

Role of the funding source

The research was funded by an unrestricted grant from Pfizer Global Medical Grants, Unity Health Foundation and the Canadian COVID-19 Immunity Task Force. The funders had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Declaration of interests: We declare no competing interests.

Abstract

Background: Understanding vaccination hesitancy during early vaccination rollout in Canada can help the government's vaccination efforts in education and outreach, which may help eventually achieving herd immunity. This study uses an online survey to assess vaccination hesitancy in population subgroups in Canada.

Method: Panel members from the nationally representative Angus Reid Forum were randomly invited to complete an online survey on their experiencing with COVID-19 symptoms and testing, as well as intention to vaccination against COVID-19. Respondents were asked "when a vaccine against the coronavirus becomes available to you, will you get vaccinated or not?" Vaccination hesitancy was defined as choosing "No – I will not get a coronavirus vaccination" as a response.

Results: 14,621 panel members (46% male and 53% female) completed the survey. Although the respondents overrepresent age 60+ and higher levels of education, other demographics, the prevalences of smoking, obesity, diabetes and hypertension were comparable to the Canadian national census and health surveys. COVID-19 vaccination hesitancy is relatively low overall (9%). Being a resident of Alberta (predicted probability = 15%), aged 40-59 (OR = 0.87, 0.78 – 0.97, predicted probability = 12%), identifying as a visible minority (OR = 0.56, 0.37 – 0.84, predicted probability = 15%), having some college level education or lower (predicted probability = 14%), or living in households of at least 5 are related to greater vaccination hesitancy (OR = 0.82, 0.76 – 0.88, predicted probability = 13%).

Conclusion: Our study enhances the understanding of COVID-19 vaccination hesitancy and identifies key population groups with higher vaccination hesitancy. As the Canadian COVID-19 vaccination effort continues, policymakers may focus outreach, education, and other efforts on these groups, which also represent groups with higher risks for contracting and dying from COVID-19. Furthermore, Canada would need to vaccinate virtually the entire population to reach herd immunity due to its relatively low infection level, and a high vaccination hesitancy would be a major hurdle to achieving that.

Introduction

Although the Canadian COVID-19 mortality rate is low compared to that in the United States and some European countries, many parts of Canada experienced a significant third wave, driven by more infectious variants and premature lifting of public health measures.¹⁻³

Canada started vaccinating healthcare workers and nursing home residents in December 2020, and has prioritized the rest of the population by mortality risk (e.g. age, medical comorbidity) and some provinces further prioritize based on risk of disease acquisition (e.g., essential workers; living in highly impacted regions).^{4,5} Progress in recent weeks has been strong with about 1% of Canadians being vaccinated with at least one dose per day in recent weeks.⁶ Vaccination hesitancy may thwart the effort to reach high vaccine coverage. As a part of the ongoing Action to Beat Coronavirus (Ab-C) study⁷, we report on levels and predictors of self-reported vaccination hesitancy in a reasonably representative sample of Canadians.

Methods

Study design and survey

The original study design and participants' description can be found in earlier reports that examined participants' symptoms related to COVID-19 and the seroprevalence of SARS-CoV-2 antibodies.⁸⁻¹⁰ Most demographic characteristics of the respondents in the Ab-C study were similar to those from the 2016 Census.^{9,11,12} The older age distribution of the respondents Ab-C versus the Census is a result of intentionally oversampling individuals age 60 and older. Also, the Ab-C cohort has a somewhat higher education level than the national profile (23% with a Bachelor's degree or higher in Canada and 43% in our sample), hence all analyses are adjusted for education levels.¹³ Nevertheless, the overall demographic distribution of the surveyed panel was similar to the Canadian Census distribution, and the prevalences of smoking, diabetes, hypertension was also similar to national surveys.¹⁴⁻¹⁷ To learn about vaccine hesitancy, we re-surveyed the original cohort¹⁰ about their experience with symptoms related to COVID-19 since the last survey, their experience with COVID-19 testing, and their intention to be vaccinated.

Survey question

Participants were asked "when a vaccine against the coronavirus becomes available to you, will you get vaccinated or not?" The five response choices were: 1. "No – I will not get a coronavirus vaccination," 2. "Not sure," 3. "Yes – I will eventually get a vaccination, but I will wait awhile first," 4. "Yes – I will get a vaccination as soon as one becomes available to me," 5. "I have already been vaccinated."

Because the action and opportunity to obtain a vaccine is largely determined by the government's policy on vaccination eligibility rather than respondents' own decision making, we assume that those who responded "Yes – I will get a vaccination as soon as one becomes available to me" and those who chose "I have already been vaccinated" have similar levels of intention to vaccinate. Hence, we combined these two groups into one category and created a 4-category variable: 1 = "No – I will not get a coronavirus vaccination," 2 = "Not sure," 3 = "Yes – I will eventually get a vaccination, but I will wait awhile first," 4 = "Yes – I will get a vaccination as soon as one becomes available to me or I have already been vaccinated."

Analytical method

Based on previous Ab-C reports, we know that the survey sample is largely representative of the Canadian non-nursing home Census population, but with a smaller proportion with some college education or below and a higher proportion with a bachelor degree or higher. To adjust for this, we applied the same survey weights as in prior reports in the ordered logit models that examine the odds of having stronger intention to vaccinate.

Prior to calculating odds ratios, parallel line assumptions (also known as the proportional odds assumption, i.e. the relationship between demographics and different thresholds of the outcome is proportional) were tested for each demographic factor at the 0.05 level within the survey setting, and all variables did meet the parallel line assumption. The overall model also meets the parallel line assumption with an adjusted Wald test. Sensitivity analysis that a used weighted generalized ordered logit testing the parallel line assumption and relaxing the assumption if variables fail to meet it had shown that the results were the same as the weighted ordered logit where the parallel line assumption held (data not shown). Marginal effects of the weighted ordered logit were also calculated to show the predicted probabilities of each vaccination intention level by demographic group. All analyses were done in Stata 16.¹⁸

Ethics

The Ab-C study was approved by the IRB of St. Michael's Hospital, Unity Health Toronto.

Results

Between January and March 2021, 14,621 participants out of 20,359 polled responded to the survey. The respondents are demographically comparable to the Canadian population in terms of age, gender, and household size. (Table 1), with the exception of the proportion of the Indigenous group (8.9% in the Ab-C and

5% in Canada) and as mentioned above, educational attainment (43% bachelor's degree or higher in Ab-C and 23% in Canada). In contrast, Ab-C participants also have similar prevalences of obesity, diabetes and hypertension as the national population.

Among Ab-C participants, 9% have no intention to vaccinate. (Table 1) Alberta (16%) and other Prairie provinces (14%) have the highest proportion who do not intend to vaccinate compared to less than 10% in other provinces. Participants aged 40-59 have the highest vaccination hesitancy, with 11.6% reporting no intention to vaccinate, compared to less than 10% in other age groups. Men have higher vaccination hesitancy (11%) than women (8%), and those with some college level education or less have stronger vaccination hesitancy (14%) than participants with other levels of education. Other groups with stronger than average vaccination hesitancy include those identifying themselves as visible minority (12%), Indigenous (15%), and those living in households of five or more people (16%).

Table 2 presents participants' weighted odds ratios of vaccination intention and the predicted probabilities to not vaccinate for each demographic group. Compared to the reference groups, the demographics with lower odds of having strong intention to vaccinate (and thus higher vaccination hesitancy) include age 40-59 years (OR = 0.87, 0.78 – 0.97, predicted probability = 12%), being a visible minority (OR = 0.56, 0.37 – 0.84, predicted probability = 15%), and belonging to a household of five or more people (OR = 0.82, 0.76 – 0.88, predicted probability = 13%). Being from Alberta and having some college level education or lower also have high predicted probability of vaccination hesitancy at 15% and 14%, respectively. The predicted probability of unwillingness for key demographic groups are also displayed in Figure 1.

Furthermore, anyone who identifies with 2 of the 3 demographics with the highest predicted probability for vaccination hesitancy (i.e. being from Alberta, having some college education or lower, being a visible minority) has a predicted probability of 2% for having no intention to vaccinate (0.21 and weighted n = 238 for from Alberta and being a visible minority, 2% and weighted n = 631 for being a visible minority and having some college level education or lower, 2% and weighted n = 600 for from Alberta and having some college level education or lower). For anyone who is from Alberta with some college level education or lower and identifies as a visible minority, the predicted probability of having no intention to vaccinate is 3%, although the sample size is very small (weighted n = 89).

Discussion and next steps

More than 9% of 14,621 participants from the Ab-C study reported having no intention to vaccinate against COVID-19. Being a resident of Alberta, aged 40-59, identifying as a visible minority, having some college level education or lower, or living in households of at least 5 are related to greater vaccination hesitancy. The key differences between the Canadian population and the Ab-C cohort are the oversampling of Forum panel members at least 60 years of age and the high proportion of respondents with a Bachelor's degree or higher. Both groups have lower vaccination hesitancy compared to their counter parts in the Ab-C survey, making the overall 9% vaccination hesitancy a conservative estimate.

Applying the 9% vaccination hesitancy to Canada as a whole, the approximately 3 million adult Canadians hesitant to vaccinate may play an important role in continuing viral transmission. This population may interact with the approximate 7.2 million Canadians below age 18,¹⁹ whom are not likely to be vaccinated for the foreseeable future. Moreover, vaccine effectiveness is about 90% for most of the vaccines available in Canada,²⁰ which implies that about 3 million vaccinated people don't have the full immunity. Collectively, these represent over a third of the total population. Within particular sub-groups, such as visible minorities, greater proportions of vaccine hesitancy paired with exposure to children might play a role in fueling sub-waves of transmission within local network.^{21,22} Thus, reducing vaccine hesitancy overall (and eventually vaccinating children) are important.

As the Canadian COVID-19 vaccination effort continues, policymakers may focus outreach, education, and other efforts on these groups, which also represent groups with higher risks for contracting and dying from COVID-19.²³ As Canada's vaccination campaign continues, with a goal of vaccinating the majority of the population by September 2021,⁴ we will periodically re-survey the Ab-C cohort and report on vaccination hesitancy and status.

References

1. Liu S. "Things are going to get worse": Canada's 7-day average reaches more than 8,200 COVID-19 cases. CTV News. Published 2021. Accessed April 27, 2021. <https://www.ctvnews.ca/health/coronavirus/things-are-going-to-get-worse-canada-s-7-day-average-reaches-more-than-8-200-covid-19-cases-1.5396978>
2. Miller A. Canada is losing the race between vaccines and variants as the 3rd wave worsens. CBC News. Published 2021. Accessed April 27, 2021. <https://www.cbc.ca/news/health/coronavirus-variants-canada-covid-19-vaccine-third-wave-1.5978394>
3. Aziz S. Have COVID-19 variants pushed Canada into a third wave of the pandemic? Global News. Published 2021. Accessed April 27, 2021. <https://globalnews.ca/news/7700045/ontario-third-wave-covid-19-new-variants/>
4. Government of Canada. Vaccines for COVID-19: Shipments and deliveries. Canada.ca. Published 2021. Accessed April 27, 2021. <https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection/prevention-risks/covid-19-vaccine-treatment/vaccine-rollout.html>
5. Ministry of Health. COVID-19 vaccines for Ontario. COVID-19 (coronavirus) in Ontario. Published 2021. Accessed April 27, 2021. <https://covid-19.ontario.ca/covid-19-vaccines-ontario>
6. Government of Canada. COVID-19 vaccination coverage in Canada. Canada.ca. Published 2021. Accessed April 27, 2021. <https://health-infobase.canada.ca/covid-19/vaccination-coverage/>
7. Centre for Global Health Research, Angus Reid Forum Inc., Unity Health Toronto, University of Toronto. Ab-C | Action to Beat Coronavirus. Abcstudy.ca. Published 2020. Accessed April 27, 2021. <https://www.abcstudy.ca/>
8. Action to beat coronavirus/Action pour battre le coronavirus (Ab-C) Study Investigators, Jha P. COVID Seroprevalence, Symptoms and Mortality During the First Wave of SARS-CoV-2 in Canada. *medRxiv*. Published online March 5, 2021:2021.03.04.21252540. doi:10.1101/2021.03.04.21252540
9. Wu DC, Jha P, Lam T, et al. Predictors of self-reported symptoms and testing for COVID-19 in Canada using a nationally representative survey. Picone GA, ed. *PLOS ONE*. 2020;15(10):e0240778. doi:10.1371/journal.pone.0240778
10. Tang X, Gelband H, Lam T, Nagelkerke N, Reid A, Jha P. A national study of self-reported COVID symptoms during the first viral wave in Canada. *medRxiv*. Published online October 5, 2020:2020.10.02.20205930. doi:10.1101/2020.10.02.20205930
11. Statistics Canada. Estimates of population (2016 Census and administrative data), by age group and sex for July 1st, Canada, provinces, territories, health regions (2018 boundaries) and peer groups. Statistics Canada. Published 2021. Accessed April 27, 2021. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710013401>
12. Statistics Canada. Census Profile, 2016 Census. Statistics Canada. Published 2020. Accessed April 27, 2021. <https://www12.statcan.gc.ca/census-recensement/2016/dp-pd/prof/index.cfm?Lang=E>
13. Statistics Canada. Education Highlight Tables, 2016 Census - Highest level of educational attainment (general) by selected age groups 15 years and over, both sexes, % distribution 2016, Canada, provinces and territories, 2016 Census – 25% Sample data. Canada.ca. Published 2017. Accessed April 27, 2021. <https://www12.statcan.gc.ca/census-recensement/2016/dp-pd/hlt-fst/edu-sco/Table.cfm?Lang=E&T=11&Geo=00&SP=1&view=2&age=1&sex=1>
14. Statistics Canada. Smoking, 2016. Health Fact Sheets. Published 2017. Accessed April 27, 2021. <https://www150.statcan.gc.ca/n1/pub/82-625-x/2017001/article/54864-eng.htm>
15. Statistics Canada. *Overweight and Obese Adults, 2018.*; 2019. Accessed April 27, 2021. <https://www150.statcan.gc.ca/n1/en/catalogue/82-625-x201900100005>
16. DeGuire J, Clarke J, Rouleau K, Roy J, Bushnik T. Blood pressure and hypertension. *Health Reports*. 2019;30(2):14-21. doi:10.25318/82-003-x201900200002
17. Public Health Agency of Canada. Chapter 1: Diabetes in Canada: Facts and figures from a public health perspective – Burden - Canada.ca. Canada.ca. Published 2011. Accessed April 27, 2021. <https://www.canada.ca/en/public-health/services/chronic-diseases/reports-publications/diabetes/diabetes-canada-facts-figures-a-public-health-perspective/chapter-1.html>
18. StataCorp. Stata Statistical Software: Release 16. Published online 2019.

19. Statistics Canada. Population estimates on July 1st, by age and sex. Statistics Canada. Published 2021. Accessed April 27, 2021. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000501>
20. Olliaro P, Torreele E, Vaillant M. COVID-19 vaccine efficacy and effectiveness—the elephant (not) in the room. *The Lancet Microbe*. 2021;0(0). doi:10.1016/S2666-5247(21)00069-0
21. Bilal U, Tabb LP, Barber S, Diez Roux A v. Spatial Inequities in COVID-19 Testing, Positivity, Confirmed Cases, and Mortality in 3 U.S. Cities. *Annals of Internal Medicine*. Published online March 30, 2021:M20-3936. doi:10.7326/M20-3936
22. Hong B, Bonczak BJ, Gupta A, Thorpe LE, Kontokosta CE. Exposure density and neighborhood disparities in COVID-19 infection risk. *Proceedings of the National Academy of Sciences of the United States of America*. 2021;118(13). doi:10.1073/pnas.2021258118
23. Statistics Canada. *Delivering Insights through Data for a Better Canada COVID-19 in Canada: A One-Year Update on Social and Economic Impacts*. Catalogue No. 11-631-X2021001.; 2021. Accessed April 27, 2021. www.statcan.gc.ca

Table 1. Unweighted survey sample characteristics (n = 14,621)

	Canada		Ab-C	Ab-C respondents who "Will not vaccinate" by demographics	
	%	n	%	n	row %
Total		14,621	100%	1355	9.3%
Province					
Ontario	38%	5,766	39.4%	448	7.8%
British Columbia	13%	2,761	18.9%	199	7.2%
Quebec	23%	2,299	15.7%	191	8.3%
Alberta	12%	1,839	12.6%	301	16.4%
Manitoba & Saskatchewan	7%	1,027	7%	142	13.8%
Atlantic provinces	7%	929	6.4%	74	8%
Age groups					
18 to 39 years	49%	4,270	29.2%	396	9.3%
40 to 59 years	28%	5,106	34.9%	590	11.6%
60 to 69 years	12%	3,472	23.7%	257	7.4%
70 years or older	11%	1,773	12.1%	112	6.3%
Sex					
Male	49%	6,742	46.1%	736	10.9%
Female	51%	7,777	53.2%	602	7.7%
Prefer to self-describe		102	0.7%	17	16.7%
Education					
Some college or lower	45%	3,402	23.3%	468	13.8%
Some university	32%	4,935	33.8%	564	11.4%
Bachelor or higher	23%	6,284	43%	323	5.1%
Visible minority					
No		12,526	85.7%	1,098	8.8%
Yes	22%	2,095	14.3%	257	12.3%
Ethnicity					
Not Indigenous		13,317	91.1%	1,164	8.7%
Indigenous	5%	1,304	8.9%	191	14.6%
Household size					
One person	28%	2,678	18.3%	262	9.8%
Two people	34%	6,315	43.2%	501	7.9%
Three people	15%	2,415	16.5%	219	9.1%
Four people	14%	2,163	14.8%	204	9.4%
Five or more people	8%	1,050	7.2%	169	16.1%
Weight status					
Under or normal weight	37%	4,318	29.5%	396	9.2%

Overweight	37%	4,716	32.3%	447	9.5%
Obese	27%	3,914	26.8%	337	8.6%
Unknown		1,673	11.4%	175	10.5%
Smoking status					
Never	46%	7,310	50%	632	8.6%
Former or current	54%	7,063	48.3%	693	9.8%
Unknown		248	1.7%	30	12.1%
Diabetic					
No		13,064	89.4%	1,222	9.4%
Yes	9%	1,424	9.7%	116	8.1%
Unknown		133	0.9%	17	12.8%
Hypertensive					
No		10,419	71.3%	1,026	9.8%
Yes	23%	3,980	27.2%	299	7.5%
Unknown		222	1.5%	30	13.5%
Note					
British Columbia includes Yukon. Manitoba & Saskatchewan include Northwest Territories.					


Sources:

- a. Census Canada 2016^{11-13, 19}
- b. Health Fact Sheets^{14, 15}
- c. Health Reports¹⁶
- d. Public Health Agency of Canada¹⁷

Table 2. Results of the weighted ordered logit for intention to vaccinate and predicted probability for vaccination hesitancy

	n	Weighted OR	Weighted 95% CI	Prob	SE
Province					
Ontario	5,766	Ref		0.092**	(0.012)
British Columbia	2,761	1.09	(0.67, 1.77)	0.085**	(0.017)
Quebec	2,299	1.01	(0.72, 1.43)	0.091**	(0.014)
Alberta	1,839	0.58	(0.17, 2.01)	0.146*	(0.047)
Manitoba & Saskatchewan	1,027	0.70	(0.21, 2.28)	0.126*	(0.031)
Atlantic provinces	929	0.95	(0.52, 1.74)	0.096**	(0.014)
Age groups					
18 to 39 years	4,270	Ref		0.107**	(0.022)
40 to 59 years	5,106	0.87**	(0.78, 0.97)	0.120**	(0.022)
60 to 69 years	3,472	1.41**	(1.03, 1.92)	0.079**	(0.012)
70 years or older	1,773	1.75***	(1.50, 2.04)	0.065**	(0.012)
Sex					
Male	6,742	Ref		0.110**	(0.020)
Female	7,777	1.23*	(0.97, 1.56)	0.091**	(0.017)
Prefer to self-describe	102	0.76	(0.38, 1.53)	0.138*	(0.037)
Education					
Some college or lower	3,402	Ref		0.140**	(0.027)
Some university	4,935	1.33**	(1.10, 1.62)	0.110**	(0.020)
Bachelor or higher	6,284	2.83***	(1.96, 4.09)	0.056**	(0.008)
Visible minority					
No	12,526	Ref		0.092**	(0.018)
Yes	2,095	0.56**	(0.37, 0.84)	0.150**	(0.016)
Ethnicity					
Not Indigenous	13,317	Ref		0.100**	(0.018)
Indigenous	1,304	0.95	(0.86, 1.05)	0.104**	(0.017)
Household size					
One person	2,678	Ref		0.113**	(0.024)
Two people	6,315	1.28**	(1.09, 1.50)	0.091**	(0.017)
Three people	2,415	1.19	(0.87, 1.62)	0.097**	(0.015)
Four people	2,163	1.20	(0.82, 1.75)	0.097**	(0.013)
Five or more people	1,050	0.82***	(0.76, 0.88)	0.134**	(0.028)
Weight status					
Under or normal weight	4,318	Ref		0.107**	(0.016)
Overweight	4,716	1.11	(0.75, 1.63)	0.098**	(0.021)
Obese	3,914	1.18**	(1.05, 1.34)	0.092**	(0.016)
Unknown	1,673	0.93	(0.59, 1.48)	0.113**	(0.023)
Smoking status					
Never	7,310	Ref		0.100**	(0.019)

Former or current	7,063	0.99	(0.85, 1.15)	0.101**	(0.017)
Unknown	248	0.83	(0.57, 1.22)	0.117**	(0.025)
Diabetic					
No	13,064	Ref		0.101**	(0.018)
Yes	1,424	1.11	(0.85, 1.44)	0.092**	(0.021)
Unknown	133	0.77	(0.32, 1.85)	0.126*	(0.032)
Hypertensive					
No	10,419	Ref		0.104**	(0.018)
Yes	3,980	1.21**	(1.09, 1.33)	0.088**	(0.018)
Unknown	222	0.72*	(0.47, 1.10)	0.137**	(0.016)
Notes:					
British Columbia includes Yukon. Manitoba & Saskatchewan include Northwest Territories.					
*** p<0.01, ** p<0.05, * p<0.1					



Notes:

1. British Columbia includes Yukon. Manitoba & Saskatchewan include Northwest Territories.
2. Upper 95% CI in dash line.

Figure 1. Probabilities of unwillingness to vaccinate for key demographic groups among participants in the Ab-C study