1	SARS-CoV-2 Heterogeneity by Ethnicity in Los Angeles		
2	Lao-Tzu Allan-Blitz, MD ¹ Fred Hertlein, ME, MBA, ² Jeffrey D. Klausner, MD, MPH ³		
3			
4	1. Brigham and Women's Hospital, Boston, MA		
5	2. Curative Inc. San Dimas, CA		
6	3. Department of Preventative Medicine, Keck School of Medicine, University of		
7	Southern California		
8			
9	Compliance with Ethical Standards:		
10	- Disclosures: Dr. Allan-Blitz has served as a consultant for Curative Inc. Dr.		
11	Klausner has served as the medical director of Curative Inc. during the		
12	observation period. Fred Hertlein has served as an employee of Curative Inc.		
13	during the observation period.		
14	- Ethical Approval: This article does not contain any studies with human		
15	participants performed by any of the authors. The Mass General Brigham		
16	institutional review board deemed the analysis of de-identified data did not		
17	constitute human subjects' research (2020P003530).		
18	- No informed consent was necessary.		
19			
20	Acknowledgements and Funding: The authors would like to acknowledge the city of Los		
21	Angeles. There was no funding for this study.		
22			

23 Word Count: 1,219

- 24 Abstract Word Count: 116
- 25 References: 19
- 26
- 27 Corresponding Author Contact Information:
- 28 Lao-Tzu Allan-Blitz
- 29 Department of Medicine, Brigham and Women's Hospital
- 30 75 Francis Street
- 31 Boston, MA 02115, USA
- 32 Email: <u>lallan-blitz@partners.org</u>, phone: (805) 896-5313
- 33
- 34 Alternative Author Information:
- 35 Dr. Jeffrey D. Klausner
- 36 Department of Preventative Medicine
- 37 Keck School of Medicine
- 38 University of Southern California
- 39 Los Angeles, CA 90033, USA
- 40 Email: jdklausner@med.usc.edu
- 41
- 42

43 Abstract

44	Recent studies have identified notable disparities in SARS-CoV-2 infection risk
45	among ethnic minorities. We evaluated SARS-CoV-2 test results from individuals
46	presenting for testing in Los Angeles between June-December, 2020. We calculated
47	prevalence ratios for various employment categories. Among 518,914 test results, of
48	which 295,295 (56.9%) were from individuals reporting Hispanic ethnicity, SARS-CoV-2
49	positivity was 16.5% among Hispanic individuals compared to 5.0% among non-
50	
50	Hispanic individuals (<i>p</i> -value<0.01). The prevalence ratios comparing Hispanic and non-
50	Hispanic individuals (<i>p</i> -value<0.01). The prevalence ratios comparing Hispanic and non- Hispanic individuals was highest for members of the media (PR=6.7; 95% CI 4.3-10.4),
51	Hispanic individuals was highest for members of the media (PR=6.7; 95% CI 4.3-10.4),
51 52	Hispanic individuals was highest for members of the media (PR=6.7; 95% CI 4.3-10.4), government employees (PR=4.0; 95% CI 3.3-4.9), and agricultural workers (PR=4.0;

57 Background

58 The impact of the SARS-CoV-2 pandemic has been disproportionately felt among Hispanic communities (1-3). The complex and interconnected structural and 59 60 systemic factors that likely contribute to such disparities (4) are also likely driving the heterogeneity with which SARS-CoV-2 spreads (5, 6). Ironically, however, the broad 61 62 business closures enforced in order to curb the pandemic may be less efficacious 63 among minority and poorer communities, as such communities are less able to limit 64 their mobility (e.g. working from home or limit within household contact due to dense 65 housing and household occupancy) and are less likely to work in settings with regular 66 testing or adequate ventilation, all of which may increase risk of SARS-CoV-2 infection 67 (7). Even worse, such communities who have less financial security at the outset may 68 feel the economic impacts of public health interventions most acutely (8, 9). 69 A new strategy, therefore, is desperately needed. In the wake of previous 70 epidemics, targeted public health interventions have been made possible via a thorough 71 understanding of factors that facilitate the transmission of disease. In response to the 72 Human Immunodeficiency Virus (HIV) epidemic, for example, the HIV Prevention Trials 73 Network has developed innumerable, novel, and cost-effective interventions that are 74 tailored directly towards factors that drive infection (10). Similarly, targeted interventions 75 may be possible for the SARS-CoV-2 pandemic if we are able to develop a more 76 thorough understanding of the specific drives among sub-populations at highest risk. 77 We therefore aimed to evaluate the risk of SARS-CoV-2 infection associated with

different employment categories among Hispanic compared to White individuals, in
order to unwind the details of factors that contribute to the disparity in burden of

infection. We performed that analysis among a community-based sample in Los Angles
as Los Angeles county is among the most populous counties in the United States and
one of the most diverse, with 48.6% of those surveyed in the national census identifying
as Hispanic (11).

84

85 Methods

86 We collected data from individuals presenting for SARS-CoV-2 polymerase chain 87 reaction testing to one of 250 drive-through centers in Los Angeles county. Individuals 88 presented between June-December, 2020. Specimens were collected via oral swabs 89 and processed as has been previously reported (12). Data collected at the time of 90 testing included age, gender, race and ethnicity, reported exposure to an individual 91 known to be infected with SARS-CoV-2, report of symptoms (muscle aches, cough, 92 fevers, diarrhea, nausea, headache, congestion, sore throat, anosmia, and shortness of 93 breath), and type of employment from a pre-specified list. Laboratory data collected 94 included human and viral cycle threshold values, date and time of testing. 95 We included data in our analysis for each individual test if there were no missing

values for ethnicity, test result, and employment category. We then calculated the

97 prevalence of SARS-CoV-2 test positivity by ethnicity and employment category as well

98 as prevalence ratios and 95% confidence intervals (CI) for each employment category.

99 The Mass General Brigham institutional review board deemed the analysis of de-

identified data did not constitute human subjects' research (2020P003530).

101

102 Results

Overall, we analyzed 518,914 individual tests results, of which 296,052 (57.1%) 103 104 were collected from women and 295,295 (56.9%) reported Hispanic ethnicity. The mean 105 age of the population was 35 years (standard deviation 15.2). In all, 247,974 (47.8%) 106 individuals reported any symptoms at the time of testing. SARS-CoV-2 positivity was 107 16.5% among Hispanic individuals compared to 5.0% among non-Hispanic individuals 108 (p-value<0.01). The percent of individuals who reported symptoms at the time of testing 109 among those who tested positive was comparable among Hispanic (70.9%) and non-110 Hispanic individuals (69.8%). The table shows SARS-CoV-2 positivity by employment 111 category as well as prevalence ratios and 95% confidence intervals.

112

113 Discussion

From a community-based sample of individuals presenting for testing in Los Angeles county, we identified a significant disparity in SARS-CoV-2 test positivity among Hispanic compared to non-Hispanic individuals. And though descriptive, we also identified notable heterogeneity in risk based on reported employment among Hispanic communities.

119 Many recent studies have demonstrated the disproportionate burden of SARS-120 CoV-2 infection suffered by minority races and ethnicities (1-4, 13-16). There are 121 numerous factors that likely contribute to those findings, including the higher proportion 122 of immigrant families without health insurance limiting the ability to seek care, language 123 barriers impacting the quality of care received, financial insecurity resulting in need for 124 potentially higher-risk employment or limiting the ability to work from home (17), and 125 housing insecurity leading to denser household occupancy and thus increased risk for

126 transmission. The tangled web of those factors as well as innumerable others creates 127 significant challenges in developing public health strategies that optimally address the 128 needs of such communities. In order to untangle such a web, we must first be able to 129 precisely describe the specific scenarios that perpetuate those disparities. Testing 130 centers can be used to gather detailed exposure data (e.g. exposure to gyms, places of 131 worship, parks, museums, hospitals, nursing homes, overnight camps, hotels, beaches, 132 movie theaters, bars, restaurants, and airports) as well as dates of those exposures in 133 order to understand when and how the transmission chains occur. Furthermore, such 134 data should be collected among individuals who test positive as well as negative, as the 135 comparison of risk will be essential in evaluating which types of exposures truly drive 136 transmission.

Our results demonstrated notably higher prevalence ratios for Hispanic 137 138 individuals employed as members of the media, agriculture workers, and government 139 personnel compared to non-Hispanics with similar employment. A recent study similarly 140 found increased risk for SARS-CoV-2 positivity among kitchen workers of minority races 141 (18). Beyond the known structural inequities that most likely contribute to such 142 differences, there may be ongoing disparities in division of labor within the workforce of 143 essential workers in specific roles that additionally compounds the heighted risk for 144 transmission. For example, Hispanic members of the media may be more likely than 145 non-Hispanic members to work in settings that place them at higher risk for SARS-CoV-146 2 transmission and acquisition. One survey found that Hispanic individuals were more 147 likely than White individuals to lose their job during the pandemic (19), which may 148 suggest that the jobs held by Hispanic individuals were different in substance than their

149 White counterparts, and thus reflect ongoing discrepant positions held by different 150 ethnicities. We did not have sufficient data, however, to evaluate other confounding 151 factors, thus further research is needed to more deeply evaluate such findings. 152 Our study was also limited in that all participants were tested in one geographic 153 region, thus potentially limiting the generalizability of our results. However, given the 154 diversity within Los Angeles (mirrored partially among our sample), and the fact that Los 155 Angeles in particular is an important case study of the ongoing socioeconomic 156 disparities surrounding the SARS-CoV-2 pandemic, we do not feel that limitation 157 reduces the importance of our results. Our study is additionally limited in that we were 158 unable to adjust our analysis based on socioeconomic status, and thus our results may 159 be confounded, thus further research is warranted.

160

161 Conclusion

162 Our results add further support to the existing body of research identifying 163 notable disparities in SARS-CoV-2 test positivity among ethnic minorities. Furthermore, 164 in an attempt to explore the heterogeneity in SARS-CoV-2 transmission within Hispanic compared to non-Hispanic communities we reported prevalence ratios for various 165 166 employment categories, noting that several demonstrated significantly higher risk for 167 infection. Further research is needed in order to precisely identify factors driving 168 transmission within high-risk populations and thereby develop tailored public health 169 interventions.

171 References

172 Ogedegbe G, Ravenell J, Adhikari S, et al. Assessment of Racial/Ethnic Disparities in 1. Hospitalization and Mortality in Patients With COVID-19 in New York City. JAMA Netw Open. 173 174 2020;3(12):e2026881. 175 2. Anand S, Montez-Rath M, Han J, et al. Prevalence of SARS-CoV-2 antibodies in a large 176 nationwide sample of patients on dialysis in the USA: a cross-sectional study. Lancet. 2020. 177 3. Macias Gil R, Marcelin JR, Zuniga-Blanco B, Marquez C, Mathew T, Piggott DA. COVID-19 178 Pandemic: Disparate Health Impact on the Hispanic/Latinx Population in the United States. J 179 Infect Dis. 2020. 180 4. Page KR, Flores-Miller A. Lessons We've Learned - Covid-19 and the Undocumented Latinx Community. N Engl J Med. 2021;384(1):5-7. 181 182 5. Britton T, Ball F, Trapman P. A mathematical model reveals the influence of population 183 heterogeneity on herd immunity to SARS-CoV-2. Science. 2020;369(6505):846-9. 184 6. Centers for Disease Control and Prevention. COVID Data Tracker: Integrated County 185 View. Available at: https://covid.cdc.gov/covid-data-186 tracker/?CDC AA refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-187 ncov%2Fcases-updates%2Fcases-in-us.html#county-view Accessed 11/30/20. 2020. 188 7. Chang S PE, Wei Koh P, Geradin J, Redbird B, Grusky D, Leskovec J. Mobility network 189 models of COVID-19 explain inequities and inform reopening. Nature 190 https://doiorg/101038/s41586-020-2923-3. 2020. 191 8. Usman M, Ali Y, Riaz A, Riaz A, Zubair A. Economic perspective of coronavirus (COVID-192 19). J Public Aff. 2020:e2252.

- 193 9. Nicola M, Alsafi Z, Sohrabi C, et al. The socio-economic implications of the coronavirus
- 194 pandemic (COVID-19): A review. Int J Surg. 2020;78:185-93.
- 195 10. HIV Prevention Trials Network. Available at: <u>https://www.hptn.org/</u> Accessed February
- 196 3, 2021.
- 197 11. United States Census Bureau. Table B03002. Hispanic or Latino Origin by Race. Available198 at:
- 199 https://data.census.gov/cedsci/table?q=B03002&g=0500000US06037&tid=ACSDT1Y2019.B030
- 200 <u>02&hidePreview=true</u> Accessed February 2, 2021.
- 201 12. Kojima N, Turner F, Slepnev V, et al. Self-Collected Oral Fluid and Nasal Swab Specimens
- 202 Demonstrate Comparable Sensitivity to Clinician-Collected Nasopharyngeal Swab Specimens for
- the Detection of SARS-CoV-2. Clin Infect Dis. 2020.
- 204 13. Wadhera RK, Wadhera P, Gaba P, et al. Variation in COVID-19 Hospitalizations and
- 205 Deaths Across New York City Boroughs. JAMA. 2020;323(21):2192-5.
- 206 14. Killerby ME L-GR, Haight SC, Schrodt CA, England L, Gomes DJ, Shamout M, Pettrone K,
- 207 O'Laughlin K, Kimball A, Blau EF, Burnett E, Ladva CN, Szablewski CM, Tobin-D'Angelo M,
- 208 Oosmanally N, Drenzek DVM, Murphy DJ, Blum JM, Hollberg J, Lefkove B, Brown FW,
- 209 Shimabukuro R, Midgley CM, Tate JE. Characteristics Associated with Hospitalization Among
- 210 Patients with COVID-19 Metropolitan Atlanta, Georgia, March-April 2020. MMWR Morb
- 211 MOrtal Wkly Rep.2020;69:790-794.
- 212 15. Price-Haywood EG, Burton J, Fort D, Seoane L. Hospitalization and Mortality among
- 213 Black Patients and White Patients with Covid-19. N Engl J Med. 2020;382(26):2534-43.

- 214 16. Millett GA, Jones AT, Benkeser D, et al. Assessing differential impacts of COVID-19 on
- black communities. Ann Epidemiol. 2020;47:37-44.
- 216 17. News Release: Bureau of Labor Statistics. US Department of Labor. Job Flexibilities and
- 217 Work Schedules 2017-2018. Published September 2019. Available at:
- 218 <u>https://www.bls.gov/news.release/pdf/flex2.pdf</u> Accessed February 5th, 2021.
- 219 18. Chen YH GM, Riley A, Balmes J, Duchowny Kate, Harrison R, Matthay E, Bibbins-Domingo
- 220 K. Excess mortality associated with the COVID-19 pandemic among Californians 18–65 years of
- 221 age, by
- 222 occupational sector and occupation: March through October 2020. medRxiv. Available at:
- 223 <u>https://www.medrxiv.org/content/10.1101/2021.01.21.21250266v1.full.pdf</u> 2021.
- 19. Guskin E. Hispanics are Almost Twice as Likely as Whites to Have Lost Their Jobs Amid
- the Pandemic, Poll Finds. The Washington Post. 2020.
- 226

227 228

229

231 Table: SARS-CoV-2 Positivity Among Hispanic and Non-Hispanic Individuals, Los

232 Angeles June-December, 2020

233

	SARS-CoV-2 Positivity			
	Hispanic (n=295,295)	Non-Hispanic (n=223,619)	Prevalence Ratio (95% CI)	
Overall	16.5%	5.0%	3.3 (3.2 – 3.4)	
Employment* (n=518,914)				
Agriculture (n=4,415)	25.9%	6.5%	4.0 (3.2 – 5.0)	
Construction (n=23,584)	16.4%	5.0%	3.2 (2.9 – 3.6)	
Correctional Facility (n=2,038)	18.3%	7.1%	2.6 (1.9 – 3.4)	
Delivery or Ride-Share (n=16,746)	12.4%	3.2%	3.9 (3.3 – 4.3)	
Disability Care (n=18,829)	18.1%	8.7%	2.1 (1.9 – 2.3)	
Education (n=59,595)	13.1%	3.6%	3.6 (3.4 – 3.9)	
Elderly Care (n=10,294)	9.0%	2.7%	3.3 (2.8 – 4.1)	
Emergency Services (n=13,173)	14.4%	5.7%	2.5 (2.2 – 2.9)	
Frist Responder (n=15,150)	11.6%	4.2%	2.8 (2.4 – 3.1)	
Food Services (n=85,497)	21.2%	6.5%	3.3 (3.1 – 3.4)	
Government Personnel (n=13,646)	7.3%	1.8%	4.0 (3.3 – 4.9)	
Grocery Store (n=30,818)	12.7%	3.5%	3.6 (3.2 – 4.0)	
Health Care (n=157,684)	14.0%	4.7%	3.0 (2.9 – 3.1)	
Member of Media (n=4,715)	5.3%	0.8%	6.7 (4.3 – 10.4)	
Public Transportation (n=9,039)	20.1%	9.1%	2.2 (1.9 – 2.5)	
Retail or Manufacturing (n=92,958)	20.8%	6.9%	3.0 (2.9 – 3.1)	

* Data were collected beginning September 2020

235

236 The table shows SARS-CoV-2 test positivity among Hispanic and non-Hispanic

individuals by type of employment as well as the prevalence ratio and 95% confidence

238 intervals for each employment category.