

# Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines

Adam V Wisnewski<sup>1\*</sup>, Carrie A Redlich<sup>1</sup>, Kathy Kamath<sup>2</sup>, Queenie-Ann Abad<sup>1</sup>, Richard F Smith<sup>1</sup>, Louis Fazen<sup>1</sup>, Romero Santiago<sup>1</sup>, Jian Liu<sup>1</sup>, Julian Campillo Luna<sup>1</sup>, Brian Martinez<sup>2</sup>, Elizabeth Baum-Jones<sup>2</sup>, Rebecca Waitz<sup>2</sup>, Winston A Haynes<sup>2</sup>, John C Shon<sup>2</sup>

<sup>1</sup>Department of Medicine, Yale University School of Medicine, New Haven, CT 06520

<sup>2</sup>Serimmune, Inc., Goleta, CA, 93117, USA

\*Corresponding Author: Adam Wisnewski, PO Box 208057, Yale University School of Medicine, New Haven, CT 06520, email: [adam.wisnewski@yale.edu](mailto:adam.wisnewski@yale.edu), phone: (203) 737-4054, fax: (203) 785-3826

## ABSTRACT

Reverse vaccinology is an evolving approach for improving vaccine effectiveness and minimizing adverse responses by limiting immunizations to critical epitopes. Towards this goal, we sought to identify immunogenic amino acid motifs and linear epitopes of the SARS-CoV-2 spike protein that elicit IgG in COVID-19 mRNA vaccine recipients. Paired pre/post vaccination samples from N=20 healthy adults, and post-vaccine samples from an additional N=13 individuals were used to immunoprecipitate IgG targets expressed by a bacterial display random peptide library, and preferentially recognized peptides were mapped to the spike primary sequence. The data identify several distinct amino acid motifs recognized by vaccine-induced IgG, a subset of those targeted by IgG from natural infection, which may mimic 3-dimensional conformation (mimotopes). Dominant linear epitopes were identified in the C-terminal domains of the S1 and S2 subunits (aa 558-569, 627-638, and 1148-1159) which have been previously associated with SARS-CoV-2 neutralization in vitro and demonstrate identity to bat coronavirus and SARS-CoV, but limited homology to non-pathogenic human coronavirus. The identified COVID-19 mRNA vaccine epitopes should be considered in the context of variants, immune escape and vaccine and therapy design moving forward.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

## INTRODUCTION

The humoral immune response to vaccination is thought to be an important component of protective immunity (1, 2). The first mRNA vaccines for COVID-19 have been shown to elicit antibodies with significant neutralizing capacity in vitro and protect against severe disease in vivo (3-6). The SARS-CoV-2 spike protein epitopes that induce specific IgG in COVID-19 mRNA vaccine recipients remain incompletely characterized and could form the basis of future “epitope vaccines” for SARS-CoV-2 (7-9).

Despite limited knowledge of immunogenic SARS-CoV-2 spike epitopes in vaccinated individuals, immunodominant epitopes in naturally infected COVID-19 patients have been identified based on recognition by serum IgG (10-18). The receptor binding domain (RBD) is an important region that contains conformational epitopes encoded by non-contiguous regions of the spike protein (19). Dominant linear epitopes are located in the CTDs of S1 and S2, the S1/S2 cleavage site, and the fusion peptide region of S2 (10-16). Importantly, dominant linear epitopes and specific conformational epitopes have been shown to mediate viral neutralization in vitro (10, 13, 16, 20).

The present study mapped COVID-19 mRNA vaccine epitopes using Serum Epitope Repertoire Analysis (SERA), a technique based on a high throughput random bacterial peptide display technology (12). SERA’s unbiased, whole proteome approach to epitope discovery is ideal for mapping the targets of humoral responses to vaccination. The primary focus herein is on de novo responses of naïve individuals, whose immune system’s specificity was not influenced by prior SARS-CoV-2 exposure. Pre vs. post vaccine serum samples were available for most participants to help ensure immune specificity was a result of vaccination, not pre-existing cross-reactivity. Similarities and differences in epitopes induced by mRNA vaccine vs. natural COVID-19, and their association with viral neutralization are discussed, along with preliminary findings from vaccinated individuals that previously had COVID-19.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

RESULTS AND DISCUSSION

*Identification of amino acid motifs recognized by vaccine-induced serum IgG:* Amino acid motifs preferentially recognized by serum IgG from individuals after COVID-19 mRNA vaccination were readily identified by SERA analysis as previously described (12). The relative increase in amino acid motif-specific IgG vs. pre-pandemic controls is shown in Figure 1 for (A) N=20 subjects without prior COVID-19 before Pfizer-BioNTech immunization, (B) N=20 subjects without prior COVID-19 after immunization with the Pfizer-BioNTech COVID-19 vaccine, (C) N=8 subjects without prior COVID-19 after immunization with Moderna vaccine, and (D) 5 subjects with prior COVID-19 immunized with either Pfizer-BioNTech (N=2) or Moderna (N=3) mRNA vaccines. Additional information on the study subjects and amino acid motifs are provided in supplemental materials (Tables S1 and S2).

Some preferentially recognized motifs share overlapping sequence and their IgG recognition show excellent correlation ( $r_s > 0.9$ ) with each other (see supplemental materials Figure S1). Preferentially recognized motifs without overlapping amino acids also showed significant ( $p < 0.001$ ), but more limited correlation with one another; the most substantial between K[FY][LM]XFXQ and P[EDT]WRXY had an  $r_s$  of 0.66 (supplemental materials Figure S2). There was significant ( $p < 0.044$ ) but weak to moderate ( $r_s = 0.35-0.46$ ) correlation of ELISA OD against spike antigen with several amino acid motifs, but no significant correlation with gender (see Supplemental Material Correlation Matrix 1). A moderate inverse relationship between age and recognition of several amino acid motifs were observed, the most significant ( $p < 0.012$ ) for overlapping amino acid motifs KWXXFQQ and K[FY][LM]XFXQ ( $r_s = -0.44$  see supplemental materials Figure S2). The preferentially recognized amino acid motifs do not map to the primary sequence of SARS-CoV-2 proteome but may mimic 3-dimensional epitopes (mimotopes) formed by non-continuous regions of the spike antigen and are a subset of those preferentially recognized by IgG from hospitalized COVID-19 patients as previously reported (12, 21).

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

*Identification of linear SARS-CoV-2 spike epitopes.* Linear epitopes of COVID-19 mRNA vaccines were identified by analyzing the specificity of IgG from available pre/post vaccine serum samples, using a protein-based immunome wide association study (PIWAS) approach as previously described (12, 22). The linear epitope (LE)-1 with the highest PIWAS values was also the most commonly recognized by IgG from >2/3 of the vaccinated subjects (Figure 2). LE-1 (aa 558-569) is located in the C-terminal domain (CTD) of the spike S1 subunit and contains the core amino acids present in dominant linear epitopes of COVID-19 patients (S14P5, S1-93, S1-55, S556-570), independently described by several other research teams, shown in Figure 3 (10, 11, 14, 16, 18). Previously described epitopes containing the LE-1 core sequence FLPFQQ mediate neutralization in vitro based on studies with epitope-affinity purified/depleted COVID-19 patient serum, and through competitive inhibition assays with clinical isolates (10, 16). We have previously reported a correlation of LE-1 specific IgG in COVID-19 patients with serum neutralizing titers in vitro (13). In the present vaccinated population, we observed a significant ( $p < 0.0004$ ) association of LE-1 recognition with total spike IgG (ELISA OD) and several different amino acid motifs, with strongest  $r_s$  values (>0.6) against overlapping motifs FXXELXX[WY]F and [FY]XX[QE]LXXWF, which contain portions of the FLPFQQ core of LE-1 (see supplemental materials Figure S3).

The linear epitope (LE-2) with the second highest average PIWAS score was recognized by IgG from 2/3 of the vaccinated subjects (Fig. 2). LE-2 (aa 627-638) is also located in the CTD of S1 and largely overlaps (10/12 amino acids) epitope S1-105, previously defined by Yi et al. (see Figure 3), based on recognition by serum from naturally infected subjects (13). S1-105 mediates a significant portion of COVID-19 patients' serum neutralizing capacity against a SARS-CoV-2 clinical isolate in limited testing (13). Recognition of LE-2 (PIWAS score) was significantly correlated with that of each of the 10 amino acid motifs defined in Figure 1 and previously published (12), but not age, gender or total spike IgG.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

A third linear epitope (LE-3) in the S2 CTD, near the 2<sup>nd</sup> heptad repeat region (HD2), is shared by a majority of vaccinated individuals. LE-3 (aa 1148-1159) contains the core amino acids present in dominant linear epitopes S2-78, S2-47, and 1147-1158 independently described by three other research teams in COVID-19 patients and shown in Figure 3 (10, 11, 13, 14). These epitopes mediate neutralization in vitro, have been experimentally observed in SARS-CoV (20), and predicted bioinformatically for SARS-CoV-2 (23-25). Recognition of LE-3 (PIWAS) was significantly ( $p < 0.0003$ ) associated with that of overlapping motifs P[EDT]WRXY and PXWR[IV]Y ( $r > 0.58$ ), but not age, sex or total IgG. An adjacent linear epitope, LE-4, spanning amino acids 1164-1173 is also recognized by IgG from more than ½ of the vaccinated subjects (Figures 2 and 4).

Additional linear spike epitopes recognized by IgG from multiple post mRNA vaccinated subjects include 12-mers centered around spike amino acids 580, 610, 665, and surrounding the S1/S2 cleavage loci ~ 685-700 (Figure 4). A complete list of the PIWAS scores for all 12-mer linear epitopes of the spike protein for each study subjects is provided as supplementary material (Table S3).

*Uniqueness of linear epitopes.* Severe outcomes of SARS-CoV-2 infections are thought in part to arise from the novelty of the virus and the naïve immune status of most humans. The dominant vaccine linear epitopes LE-1, LE-2, and LE-3 show identity or near identity to bat coronavirus and SARS-CoV, but not non-pathogenic human coronaviruses (Figure 3). Further homology searches of the linear spike epitopes identify limited sequence identity outside of organisms to which humans are rarely exposed, such as thermophilic bacteria (supplemental materials List S1). The present data support the hypothesis that humoral immune naivete contributes to SARS-CoV-2's pathogenicity and identifies specific deficiencies in the normal adult immune repertoire readily corrected by mRNA vaccination.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

*IgG responses in mRNA vaccine recipients vs. natural infection.* We compared the linear epitope recognition by IgG from the different groups of vaccinated subjects in the present study and our prior studies of COVID-19 patients (12). Similar specificity for dominant linear epitopes were observed in individuals that received mRNA vaccine from Pfizer-BioNTech or Moderna (Figure 4), regardless of prior COVID-19 status. However, vaccinated individuals with prior COVID-19 exhibited significantly ( $p < 0.001$ ) higher PIWAS values for dominant epitopes LE-1, LE-2, and LE-3 than those without prior infection. Vaccine-induced IgG specificity overlaps substantially with that of COVID-19 patients (12) and can occur de novo, or augment naturally occurring responses (see pre/post vaccine examples in supplemental material Figure S4). The most notable difference in specificity between vaccine vs. infection induced IgG, is the lack of specificity for linear epitopes in the fusion peptide region of the spike-S2 domain by vaccine-elicited IgG. Despite the induction of specific IgG to the S2 fusion peptide domain in natural SARS-CoV-2 infection, vaccination does little to boost this response compared to the dominant linear epitopes (Figure 4, and supplemental materials Figure S4).

*RBD epitopes.* This study did not identify vaccine epitopes of the spike RBD domain, which is a major target for neutralizing antibodies in natural infection. The findings are consistent with prior reports that did not detect spike RBD linear epitopes and may be due to multiple reasons including limitations in sensitivity and frequency/sample size, as well as the RBD's complex secondary structure, which is thought to form immunogenic 3-D conformations, or non-continuous epitopes (11, 13, 17, 26). RBD specific antibodies are a minor proportion of total anti-spike IgG and linear RBD epitopes are not prominent IgG epitopes in natural SARS-CoV-2 infection (12, 13, 27). Nonetheless, the RBD is an important target of neutralizing antibodies that develop in vivo during natural infection (28), and possibly a driver for the evolution of mutations given its critical role in the virus life cycle (29, 30). Many of the SARS-CoV-2 variants that have evolved to date contain mutations that affect the RBD and result in decreased

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

neutralization by RBD-specific antibodies (31), but not mAbs that bind outside the receptor-binding motif (32, 33). In this context dominant linear neutralizing epitopes may provide an increased role in immunity and compensate for loss of neutralization activity due to decreased IgG recognition of variant RBDs, and perhaps contribute to current mRNA vaccine efficacy in populations with high prevalence of B.1.1.7 and B.1.351 variants (34). Mutations that result in escape from dominant linear epitopes outside the RBD might also adversely affect host immunity and vaccine efficacy, particularly if compounded by RBD escape variants. It is thus important to be vigilant in surveillance for immune escape from dominant linear epitopes as well as conformational RBD affected epitopes in new variants.

*Summary:* Spike protein epitopes of COVID-19 mRNA vaccines were identified through immunomics / reverse vaccinology. Dominant epitopes included specific amino acid motifs that may reflect 3-D conformations and linear epitopes of the spike protein, particularly in the C-terminal domain of S1. COVID-19 mRNA vaccine epitopes share identity with bat coronavirus, SARS-CoV (but not the spike protein of non-pathogenic human coronaviruses), and epitopes that trigger IgG during natural SARS-CoV-2 infection, which have been previously associated with SARS-CoV-2 neutralization (10, 13, 16, 20). The finding of increased IgG response in vaccinated individuals with prior infection (vs. naïve individuals) is consistent with previous reports and highlights the effectiveness of “booster shots” towards robust humoral immunity (35). In summary, we identified epitopes of COVID-19 mRNA vaccines that trigger specific IgG responses previously associated with SARS-CoV-2 neutralizing activity and which may form the basis of future diagnostics, therapeutics, and focused vaccine development (e.g., multivalent peptide).

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

MATERIALS AND METHODS

Blood samples from vaccine recipients: Subjects were working aged adults employed in the health care industry. Blood was obtained by venipuncture in a serum separator tube (BD Vacutainer®, Franklin Lakes, NJ) and the serum fraction was separated following centrifugation and stored at -80°C. Four groups of samples were studied; (A) N=20 subjects without prior COVID-19 before immunization, (B) N=20 subjects without prior COVID-19 after immunization with the Pfizer-BioNTech COVID-19 vaccine, (C) N=8 subjects without prior COVID-19 after immunization with Moderna vaccine, and (D) 5 subjects with prior COVID-19 immunized with either Pfizer-BioNTech (N=2) or Moderna (N=3). The investigation was approved by the Yale University Institutional Review Board (IRB) and all study subjects provided written informed consent before participating in the study.

ELISAs: ELISAs were performed as previously described (36, 37). Triton X-100 and RNase A were added to serum samples at final concentrations of 0.5% and 0.5mg/mL respectively and incubated at room temperature (RT) for 30 minutes before use to reduce risk from any potential virus in serum. 96-well MaxiSorp plates (ThermoFisher Scientific, Waltham, MA) were coated with 50 µL/well of recombinant SARS Cov-2 spike ectodomain or nucleocapsid protein (Sino Biological, Wayne, PA) at a concentration of 1 µg/mL in NaCO<sub>3</sub> buffer pH 9.6 and incubated overnight at 4°C. ELISA plate blocked for 1h at RT with 200 µL of PBS with 3% milk powder. Serum was diluted 1:100 in PBS with 0.05% Tween20, 1% milk powder and 100 µL was added for 1 hr at RT. Plates were washed three times with PBS-T (PBS with 0.1% Tween-20) and incubated with 50 µL of HRP anti-Human IgG Antibody (Pharmingen, 1:2,000) for 1 h at RT. Plates were developed with 100 µL of TMB Substrate Reagent Set (ThermoFisher) and the reaction was stopped when an internal pooled serum positive control sample reaches an OD of 1.0 at 650 nm, by the addition of 2 N sulfuric acid. Plates were then read at a wavelength of 450 nm with 650 nm reference calibration.



*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

SERA: A description of the SERA assay has been previously published (9). Briefly, serum was incubated with a fully random 12-mer bacterial display peptide library ( $1 \times 10^{10}$  diversity, 10-fold oversampled) at a 1:25 dilution in a 96-well, deep well plate format. Antibody-bound bacterial clones were selected with 50  $\mu$ L Protein A/G Sera-Mag SpeedBeads (GE Life Sciences, cat#17152104010350) (IgG) final assay dilution 1:100 (ThermoFisher). The selected bacterial pools were resuspended in growth media and incubated at 37°C shaking overnight at 300 RPM to propagate the bacteria. Plasmid purification, PCR amplification of peptide encoding DNA, barcoding with well-specific indices was performed as described. Samples were normalized to a final concentration of 4nM for each pool and run on the Illumina NextSeq500. After SERA screening, we applied two complementary discovery tools, IMUNE and PIWAS, to identify antigens and epitopes involved in the SARS-CoV-2 immune response.

PIWAS analysis: We applied the previously published PIWAS method (22) to identify antigen and epitope signals against the Uniprot reference SARS-CoV-2 proteome (UP000464024) (38). The PIWAS analysis was run on the IgG SERA data with a single sample per COVID-19 patient versus 497 discovery pre-pandemic controls, and 1500 validation controls used for normalization. Additional parameters include: a smoothing window size of 5 5mers and 5 6mers; z-score normalization of kmer enrichments; maximum peak value; and generation of epitope level tiling data.

IMUNE-based motif discovery: Peptide motifs representing epitopes or mimotopes of SARS CoV-2 specific antibodies were discovered using the IMUNE algorithm as previously described (39). Peptide patterns identified by IMUNE were clustered using a PAM30 matrix and combined into motifs. A motif was classified positive in a given sample if the enrichment was  $\geq 4$  times the standard deviation above the mean of a training control set. Thirty eight motifs were enriched in 406 unique confirmed COVID-19 cases from four separate cohorts of COVID patients compared to 1500 pre-pandemic controls.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

Statistical analysis: GraphPad Prism (v8) and Microsoft Excel for Windows 10 (v16.0.13001.20254) were used for statistical analyses. Statistical differences were analyzed with Mann-Whitney U-test.

Nonparametric Kruskal-Wallis test was used to perform multiple comparisons between groups analyzed.

Correction for multiple comparison was performed with Dunn's test. Logistic regression or nonparametric

Spearman Spearman's rank correlation were used to compute associations between pairs of data.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

AUTHOR CONTRIBUTIONS: AW, CR, QA, RFS, RS, LF, JL, JCL obtained samples and performed ELISAs, WAH, KK, JCS performed bacteriophage peptide immunoprecipitation, SERA analysis and bioinformatics. AW wrote and all authors edited, reviewed and approved the final version of the manuscript.

ACKNOWLEDGEMENTS: We are grateful to the study participants, without which the research could not have been done. The work was supported by funds from the National Institute of Occupational Safety and Health, 5T03OH008607-15.

COMPETING INTERESTS: All authors have completed the ICMJE uniform disclosure form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) and declare support from the National Institute of Occupational Safety and Health (5T03OH008607-15); no financial relationships with any organizations that might have an interest in the submitted work in the previous three years; and no other relationships or activities that could appear to have influenced the submitted work.

SUPPLEMENTAL MATERIALS: Supplemental material with available data on correlation between IgG specificities, PIWAS values, and examples of post-COVID/Post-COVID-vaccine samples.

Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines




Figure 1. Heat map identifying amino acid motifs preferentially recognized by IgG from vaccinated subjects. Amino acid motifs were identified among the peptides immunoprecipitated by IgG from vaccine recipients using IMUNE algorithm and the level of enrichment (fold increase) in serum of individual subjects relative to pre-pandemic control subjects is depicted in the heat map. Samples 1-20 are pre-vaccine and samples 21-40 are the same subjects after Pfizer-BioNTech COVID-19 mRNA vaccination. Subjects 41-48 are from subjects that received Moderna vaccine. Samples 49-53 are vaccinated subjects that previously had COVID-19.




Figure 2. Linear epitope mapping of spike protein in mRNA vaccine recipients. The relative specificity for linear epitopes or PIWAS score (Y-axes, color coded key to right) are graphed across the spike primary sequence (amino acid numbers on X-axis) for paired samples from subjects# 1-20 in panel A (pre-vaccine) and panel B (post-vaccine) that received Pfizer-BioNTech vaccine. Panel C shows data from subjects # 21-28, post Moderna vaccine and panel D shows vaccinated subjects# 29-33 that previously had COVID-19

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

A.

AA Sequence	Epitope Name	Ref
<b>FLPFQQFGRDIA</b>	<b>LE-1</b>	(12)
NKK <b>FLPFQQFGRDIA</b>	S556-570	(15)
TESNKK <b>FLPFQQFGRDIA</b>	S14P5	(10, 16)
TESNKK <b>FLPFQQ</b>	S1-93	(10, 14)
ESNKK <b>FLPFQQFGRDIADTT</b>	S1-55	(13)
<b>DQLTPTWRVYST</b>	<b>LE-2</b>	-
HAD <b>QLTPTWRVY</b>	S1-105	(10, 14)
<b>FKEELDKYFKNH</b>	<b>LE-3</b>	-
SF <b>FKEELDKYFKN</b>	1147-1158	(11)
DS <b>FKEELDKYFKNHTSPDVD</b>	S2-47	(13)
<b>FKEELDKYFKNH</b>	S2-78	(10, 14)

B.

SARS-CoV-2	<b>FL*PFQQFGRDIA</b>	<b>DQLT**PTWRVYST</b>	<b>FKEELDKYFKNH</b>
BtCoV-RaTG13	--*-----	----**-----	-----
SARS-CoV	-Q*-----VS	----**-----	-----
MERS-CoV	GV*R---VYDAQN	YSRSTRSMLKRRD*	-QD---EF---V
HCoV-OC43	YYVSW-NLLYDSN	R--Q**--INYFD**	-----QW---Q
HCoV-HKU1	YYVSW-NLLYDSN	**--ATQ*****	--S--SHW---Q
HCoV-NL63	LAGGITYVSNMGN	YG-QNLLQLP****	VNKT-QEFAQ-L
HCoV-2229E	--NGITYTSTSGN	YGFSNBVVEMP***	VNKT-QELSYKL

Figure 3. Alignment of dominant linear vaccine epitopes with those induced by infection and in other coronaviruses. Alignment of vaccine epitopes with those previously described in natural SARS-CoV-2 infection (A). Alignment of vaccine epitopes with homologous regions in other coronaviruses (B). \* are spaces included to maximize alignment.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*




Figure 4. Dominant linear epitopes in COVID-19 mRNA vaccine recipients. The relative IgG binding (average PIWAS, Y-axes) to different linear epitopes (spike amino acid number, X-axis) is highlighted for the C-terminal domains of S1 (A) and S2 (B). Data are from N=20 Pfizer-BioNTech, N=8 Moderna vaccine recipients, and N=5 prior COVID-19 patients that received either Pfizer or Moderna vaccine as labeled in the key in upper right. The red line corresponds to the 95% quantile PIWAS score for a prior cohort of COVID+ subjects (12). Note the highest PIWAS scores for LE-1 and LE-2 as labeled and limited vaccine induced IgG towards fusion peptide region, aa 788-806 (40).

## REFERENCES

1. Amanna IJ, Carlson NE, and Slifka MK. Duration of Humoral Immunity to Common Viral and Vaccine Antigens. *New England Journal of Medicine*. 2007;357(19):1903-15.
2. Poland GA, Ovsyannikova IG, and Kennedy RB. SARS-CoV-2 immunity: review and applications to phase 3 vaccine candidates. *Lancet*. 2020;396(10262):1595-606.
3. Anderson EJ, Roupheal NG, Widge AT, Jackson LA, Roberts PC, Makhene M, et al. Safety and Immunogenicity of SARS-CoV-2 mRNA-1273 Vaccine in Older Adults. *N Engl J Med*. 2020;383(25):2427-38.
4. Jackson LA, Anderson EJ, Roupheal NG, Roberts PC, Makhene M, Coler RN, et al. An mRNA Vaccine against SARS-CoV-2 - Preliminary Report. *N Engl J Med*. 2020;383(20):1920-31.
5. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. 2020;383(27):2603-15.
6. Walsh EE, Frenck RW, Jr., Falsey AR, Kitchin N, Absalon J, Gurtman A, et al. Safety and Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. *N Engl J Med*. 2020;383(25):2439-50.
7. Palatnik-de-Sousa CB, Soares IS, and Rosa DS. Editorial: Epitope Discovery and Synthetic Vaccine Design. *Front Immunol*. 2018;9:826.
8. Rappuoli R, and Covacci A. Reverse vaccinology and genomics. *Science*. 2003;302(5645):602.
9. Naz A, Shahid F, Butt TT, Awan FM, Ali A, and Malik A. Designing Multi-Epitope Vaccines to Combat Emerging Coronavirus Disease 2019 (COVID-19) by Employing Immuno-Informatics Approach. *Front Immunol*. 2020;11:1663.



*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

10. Amrun SN, Lee CY, Lee B, Fong SW, Young BE, Chee RS, et al. Linear B-cell epitopes in the spike and nucleocapsid proteins as markers of SARS-CoV-2 exposure and disease severity. *EBioMedicine*. 2020;58:102911.
11. Farrera-Soler L, Daguer JP, Barluenga S, Vadas O, Cohen P, Pagano S, et al. Identification of immunodominant linear epitopes from SARS-CoV-2 patient plasma. *PLoS One*. 2020;15(9):e0238089.
12. Haynes WA, Kamath K, Bozekowski J, Baum-Jones E, Campbell M, Casanovas-Massana A, et al. High-resolution mapping and characterization of epitopes in COVID-19 patients. *medRxiv*. 2020:2020.11.23.20235002.
13. Li Y, Lai DY, Zhang HN, Jiang HW, Tian X, Ma ML, et al. Linear epitopes of SARS-CoV-2 spike protein elicit neutralizing antibodies in COVID-19 patients. *Cell Mol Immunol*. 2020;17(10):1095-7.
14. Li Y, Ma M-l, Lei Q, Wang F, Lai D-y, Hou H, et al. Linear epitope landscape of SARS-CoV-2 Spike protein constructed from 1,051 COVID-19 patients. *medRxiv*. 2020:2020.07.13.20152587.
15. Lu S, Xie XX, Zhao L, Wang B, Zhu J, Yang TR, et al. The immunodominant and neutralization linear epitopes for SARS-CoV-2. *Cell Rep*. 2021;34(4):108666.
16. Poh CM, Carissimo G, Wang B, Amrun SN, Lee CY-P, Chee RS-L, et al. Two linear epitopes on the SARS-CoV-2 spike protein that elicit neutralising antibodies in COVID-19 patients. *Nature Communications*. 2020;11(1):2806.
17. Yi Z, Ling Y, Zhang X, Chen J, Hu K, Wang Y, et al. Functional mapping of B-cell linear epitopes of SARS-CoV-2 in COVID-19 convalescent population. *Emerg Microbes Infect*. 2020;9(1):1988-96.
18. Shrock E, Fujimura E, Kula T, Timms RT, Lee IH, Leng Y, et al. Viral epitope profiling of COVID-19 patients reveals cross-reactivity and correlates of severity. *Science*. 2020;370(6520):eabd4250.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

19. He Y, Lu H, Siddiqui P, Zhou Y, and Jiang S. Receptor-binding domain of severe acute respiratory syndrome coronavirus spike protein contains multiple conformation-dependent epitopes that induce highly potent neutralizing antibodies. *J Immunol.* 2005;174(8):4908-15.
20. He Y, Zhou Y, Wu H, Luo B, Chen J, Li W, et al. Identification of immunodominant sites on the spike protein of severe acute respiratory syndrome (SARS) coronavirus: implication for developing SARS diagnostics and vaccines. *J Immunol.* 2004;173(6):4050-7.
21. Gulati S, Pennington MW, Czerwinski A, Carter D, Zheng B, Nowak NA, et al. Preclinical Efficacy of a Lipooligosaccharide Peptide Mimic Candidate Gonococcal Vaccine. *mBio.* 2019;10(6).
22. Haynes WA, Kamath K, Daugherty PS, and Shon JC. Protein-based Immunome Wide Association Studies (PIWAS) for the discovery of significant disease-associated antigens. *bioRxiv.* 2020:2020.03.18.997759.
23. Bhattacharya M, Sharma AR, Patra P, Ghosh P, Sharma G, Patra BC, et al. Development of epitope-based peptide vaccine against novel coronavirus 2019 (SARS-CoV-2): Immunoinformatics approach. *J Med Virol.* 2020;92(6):618-31.
24. Parvez S, and Preeti S. *Prediction of T and B Cell Epitopes in the Proteome of SARS-CoV-2 for Potential Use in Diagnostics and Vaccine Design.* 2020.
25. Ahmed SF, Quadeer AA, and McKay MR. Preliminary Identification of Potential Vaccine Targets for the COVID-19 Coronavirus (SARS-CoV-2) Based on SARS-CoV Immunological Studies. *Viruses.* 2020;12(3).
26. Piccoli L, Park Y-J, Tortorici MA, Czudnochowski N, Walls AC, Beltramello M, et al. Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike Receptor-Binding Domain by Structure-Guided High-Resolution Serology. *Cell.* 2020;183(4):1024-42.e21.
27. Voss WN, Hou YJ, Johnson NV, Delidakis G, Kim JE, Javanmardi K, et al. Prevalent, protective, and convergent IgG recognition of SARS-CoV-2 non-RBD spike epitopes. *Science.* 2021:eabg5268.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

28. Greaney AJ, Loes AN, Crawford KHD, Starr TN, Malone KD, Chu HY, et al. Comprehensive mapping of mutations in the SARS-CoV-2 receptor-binding domain that affect recognition by polyclonal human plasma antibodies. *Cell Host Microbe*. 2021;29(3):463-76.e6.
29. Zahradník J, Marciano S, Shemesh M, Zoler E, Chiaravalli J, Meyer B, et al. SARS-CoV-2 RBD in vitro evolution follows contagious mutation spread, yet generates an able infection inhibitor. *bioRxiv*. 2021:2021.01.06.425392.
30. Van Egeren D, Novokhodko A, Stoddard M, Tran U, Zetter B, Rogers M, et al. Risk of evolutionary escape from neutralizing antibodies targeting SARS-CoV-2 spike protein. *medRxiv*. 2020:2020.11.17.20233726.
31. Shen X, Tang H, Pajon R, Smith G, Glenn GM, Shi W, et al. Neutralization of SARS-CoV-2 Variants B.1.429 and B.1.351. *New England Journal of Medicine*. 2021.
32. Collier DA, De Marco A, Ferreira IATM, Meng B, Datir RP, Walls AC, et al. Sensitivity of SARS-CoV-2 B.1.1.7 to mRNA vaccine-elicited antibodies. *Nature*. 2021;593(7857):136-41.
33. Garcia-Beltran WF, Lam EC, St. Denis K, Nitido AD, Garcia ZH, Hauser BM, et al. Multiple SARS-CoV-2 variants escape neutralization by vaccine-induced humoral immunity. *Cell*. 2021;184(9):2372-83.e9.
34. Abu-Raddad LJ, Chemaitelly H, and Butt AA. Effectiveness of the BNT162b2 Covid-19 Vaccine against the B.1.1.7 and B.1.351 Variants. *New England Journal of Medicine*. 2021.
35. Assis R, Jain A, Nakajima R, Jasinskas A, Kahn S, Palma A, et al. Distinct SARS-CoV-2 antibody reactivity patterns elicited by natural infection and mRNA Vaccination. *bioRxiv*. 2021:2021.04.15.440089.
36. Mahajan S, Redlich CA, Wisnewski AV, Fazen LE, Rao LV, Kuppusamy K, et al. Performance of Abbott Architect, Ortho Vitros, and Euroimmun Assays in Detecting Prior SARS-CoV-2 Infection. *medRxiv*. 2020:2020.07.29.20164343.

*Immunogenic Amino Acid Motifs and Linear Epitopes of COVID-19 mRNA Vaccines*

37. Amanat F, Stadlbauer D, Strohmeier S, Nguyen THO, Chromikova V, McMahon M, et al. A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nature Medicine*. 2020;26(7):1033-6.
38. UniProt: the universal protein knowledgebase. *Nucleic Acids Res*. 2017;45(D1):D158-d69.
39. Pantazes RJ, Reifert J, Bozekowski J, Ibsen KN, Murray JA, and Daugherty PS. Identification of disease-specific motifs in the antibody specificity repertoire via next-generation sequencing. *Sci Rep*. 2016;6:30312.
40. Xia S, Zhu Y, Liu M, Lan Q, Xu W, Wu Y, et al. Fusion mechanism of 2019-nCoV and fusion inhibitors targeting HR1 domain in spike protein. *Cellular & Molecular Immunology*. 2020;17(7):765-7.