

1 **A clinical observational analysis of aerosol emissions from dental procedures.**

2 **Authors**

3 T. Dudding^{&1,2,3}, S. Sheikh^{&4}, F. Gregson⁴, J. Haworth^{3,5}, S. Haworth^{1,2,3}, B.G. Main^{2,3,6}, A.J. Shrimpton⁷,
4 F.W. Hamilton^{1,2,8}, AERATOR group, A.J. Ireland^{3,5}, N.A. Maskell⁹, J.P. Reid⁴, B.R. Bzdek^{%4}, M.
5 Gormley^{%1,2,3}

6

7 ¹ MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University
8 of Bristol, Bristol, UK.

9 ² Department of Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK.

10 ³ Bristol Dental Hospital and School, University of Bristol, Bristol, UK.

11 ⁴ School of Chemistry, University of Bristol, Bristol, UK.

12 ⁵ Royal United Hospital Bath, Combe Park, Bath, UK.

13 ⁶ Bristol Centre for Surgical Research, Population Health Sciences, Bristol Medical School, Bristol, UK.

14 ⁷ School of Physiology, Pharmacology & Neuroscience, University of Bristol, Bristol, UK.

15 ⁸ Infection Sciences, Southmead Hospital, North Bristol NHS Trust, Bristol, UK.

16 ⁹ Academic Respiratory Unit, University of Bristol, UK.

17 [&] Joint first authors

18 [%] Joint last authors

19

20

21

22

23 Corresponding author: Mark Gormley

24 University of Bristol Dental Hospital and School

25 Lower Maudlin Street

26 Bristol, BS1 2LY

27 Email : mark.gormley@bristol.ac.uk

28 Tel: +44(0) 117 33 10097

29

30 **Word count:** 3,143 (excluding abstracts, acknowledgments, figure legends and references)

31 **Reference count:** 25

32 **Figure/table count:** 5

33

34 **Keywords:** COVID-19; Coronavirus; Dental public health; Epidemiology; Ultrasonics; Restorative

35 dentistry

36

37

38

39

40

41

42

43

44 **Abstract**

45 Aerosol generating procedures (AGPs) are defined as any procedure releasing airborne particles <5
46 μm in size from the respiratory tract. There remains uncertainty about which dental procedures
47 constitute AGPs. We quantified the aerosol number concentration generated during a range of
48 periodontal, oral surgery and orthodontic procedures using an aerodynamic particle sizer, which
49 measures aerosol number concentrations and size distribution across the 0.5–20 μm diameter size
50 range. Measurements were conducted in an environment with a sufficiently low background to
51 detect a patient’s cough, enabling confident identification of aerosol. Phantom head control
52 experiments for each procedure were performed under the same conditions as a comparison.
53 Where aerosol was detected during a patient procedure, we assessed whether the size distribution
54 could be explained by the non-salivary contaminated instrument source in the respective phantom
55 head control procedure using a two-sided unpaired t-test (comparing the mode widths ($\log(\sigma)$) and
56 peak positions ($D_{p,c}$)). The aerosol size distribution provided a robust fingerprint of aerosol emission
57 from a source. 41 patients underwent fifteen different dental procedures. For nine procedures, no
58 aerosol was detected above background. Where aerosol was detected, the percentage of procedure
59 time that aerosol was observed above background ranged from 12.7% for ultrasonic scaling, to
60 42.9% for 3-in-1 air + water syringe. For ultrasonic scaling, 3-in-1 syringe use and surgical drilling, the
61 aerosol size distribution matched the non-salivary contaminated instrument source, with no
62 unexplained aerosol. High and slow speed drilling produced aerosol from patient procedures with
63 different size distributions to those measured from the phantom head controls (mode widths $\log(\sigma)$)
64 and peaks ($D_{p,c}$), $p < 0.002$) and, therefore, may pose a greater risk of salivary contamination. This
65 study provides evidence for sources of aerosol generation during common dental procedures,
66 enabling more informed evaluation of risk and appropriate mitigation strategies.

67

68

69 Introduction

70 Transmission of respiratory diseases, such as severe acute respiratory syndrome coronavirus 2
71 (SARS-CoV-2), the causative virus for the coronavirus disease 2019 (COVID-19), can occur through
72 direct or indirect physical contact, droplet inhalation or airborne transmission [1]. Aerosols and
73 droplets are created when the surface tension of a fluid is overcome by force, for example from air
74 turbines within dental drills [2]. High viral loads present in the course of COVID-19 infection make
75 dental aerosols a plausible source of infective particles [2-4]. A recent study has demonstrated that
76 asymptomatic patients attending dental care settings can be positive for SARS-CoV-2 [5].

77 Understanding and managing the disease risk posed by dental aerosols is important to protect
78 patients and dental teams.

79

80 Aerosol generating procedures (AGPs) may result in respiratory disease transmission, and are
81 defined as any procedure that can result in the release of airborne particles $<5 \mu\text{m}$ in size from the
82 respiratory tract of an individual [6]. Recent work has shown that dental AGPs generate particles
83 with a size distribution sufficiently wide to potentially incorporate SARS-CoV-2 virions ($0.05\text{--}0.15$
84 μm) [7]. Aerosol generated during dental procedures is of respirable size, can remain suspended in
85 air around the dental team [8], and is removed primarily by ventilation parameters of the room. By
86 contrast, larger splatter particles ($>50 \mu\text{m}$), often follow a ballistic trajectory, and are rarely detected
87 more than 2 m from the source during dental procedures [9].

88

89 Potential AGPs have attracted additional mandatory infection control practices, including personal
90 protective equipment (PPE), ensuring adequate ventilation and allowing additional 'fallow' time
91 between patients to enhance aerosol dispersion [10]. In dentistry there remains uncertainty about
92 which procedures constitute an AGP, with a recent systematic review highlighting this evidence gap
93 [11]. Previous studies suffer from inherent drawbacks, using instruments with limited sensitivity to

94 accurately detect aerosol, such as culture or settling plates, which do not account for the suspension
95 of particles or those removed through ventilation [12-14]. Others have employed simulation on
96 phantom heads [15, 16], which may not accurately capture the real clinical scenario. Some recent
97 studies have used high-resolution electrical low-pressure impactor particle sizers [7] and portable
98 scanning mobility particle sizers [17], to capture the smaller aerosol particles (<10 µm) likely
99 produced during dental AGPs. However, this work has also been performed in phantom heads rather
100 than on dental patients.

101

102 For any dental procedure on a patient, there are three aerosol sources to consider. First, the host
103 (patient) aerosol generated during breathing, speaking, or coughing may be infectious to the dental
104 team in close proximity [18]. The second source of aerosol is the instrument generated aerosol,
105 which is not considered infectious as there is no physical interaction with the host. Finally, there is
106 salivary-contaminated aerosol generated by the action of the instrument in a potentially infectious
107 host, which might be infectious. One challenge is separating this salivary-contaminated aerosol from
108 the non-salivary contaminated instrument source. In this study we quantified the aerosol number
109 concentration, in the 0.5–20 µm size range, produced during a wide range of dental procedures in a
110 real-world clinical setting. We also aimed to determine whether aerosol detected was intrinsically
111 generated from the non-salivary contaminated dental instrument, or was likely to be contaminated
112 using aerosol size distribution analysis and modelling with phantom head controls. These
113 measurements were conducted in an environment with an aerosol background concentration low
114 enough to reliably detect a patient’s cough, allowing robust detection of any aerosol generated
115 during dental procedures [19].

116

117 **Methods**

118 *Ethical approval and patient recruitment*

119 This study obtained ethical approval as part of the AERosolisation And Transmission Of SARS-CoV-2
120 in Healthcare Settings (AERATOR) study via the Northwest Research Ethics Committee (Ref:
121 20/NW/0393) and was conducted in accordance with the STROBE guidelines (Strengthening the
122 Reporting of Observational Studies in Epidemiology). Adult patients >18 years old on waiting lists
123 requiring either periodontal, oral surgery or orthodontic treatment were recruited consecutively.
124 Each patient was contacted via telephone, received an information leaflet via post, and provided
125 written consent on the day of treatment.

126

127 *Environment and equipment*

128 An Aerodynamic Particle Sizer (APS) (TSI Incorporated, model 3321, Shoreview, NM, USA; detection
129 range: 0.5-20 μm diameter particles) was used to measure aerosol. A custom 3D-printed funnel
130 (RAISE3D Pro2 Printer, 3DGBIRE, Chorley, UK) made from polylactide, with a maximum diameter of
131 150 mm, cone height of 90 mm and a 10 mm exit port, was attached to the APS inlet using
132 conductive silicone sampling tubing (TSI, 3001788), approximately 0.90 m long and 4.80 mm in
133 diameter. The experimental set-up is shown in **Supplementary Fig. 1**. The APS was set to sample
134 aerosol number concentration once per second. Further detail on the environment and instruments
135 used can be found in the **Supplemental Material**.

136

137 *Baseline patient measurements*

138 Baseline readings were taken from each participant including tidal breathing at rest (60 s), counting
139 out loud (60 s) and three voluntary coughs. The funnel was positioned at source (as close to the
140 mouth as possible), with the patient seated upright [20]. Baseline characteristics of patients were
141 reported using median and range for continuous data, alongside counts and percentages for
142 categorical data, stratified by specialty type. To assess differences among specialties, age and sex
143 distribution was compared using one-way ANOVA and Fisher's exact test respectively.

144

145 *Patient dental procedure aerosol measurement*

146 We conducted an initial pilot study to investigate the optimum position and orientation for the 3D-
147 printed funnel when sampling dental aerosol. This was determined to be 22 cm from soft tissue
148 nasion to the top of the funnel, at approximately 45 degrees on the patient's left side (11 o'clock
149 position). For every case, a full mouth examination was carried out using a dental mirror, followed
150 by local anaesthetic administration when indicated. Each patient received 3-in-1 syringe air drying
151 (30 s), water (30 s) and then combined air and water (30 s) applied to their all their teeth. When
152 necessary, up to 3-minute intervals between procedural steps were allowed for background reading
153 levels to stabilise. The remainder of the treatment session was dictated by clinical need. A detailed
154 description of the treatments and time-stamped protocols are provided in the **Supplementary**
155 **Methods.**

156

157 *Phantom head control procedure aerosol measurement*

158 To measure aerosol generated by the dental instruments alone, we conducted high fidelity control
159 experiments in triplicate, in a phantom head unit. For phantom head control data, the aerosol
160 number concentration and size distribution were extracted for further analysis. Further detail can be
161 found in the **Supplementary Methods.**

162

163 **Statistical analysis**

164 *Total procedure aerosol number concentration*

165 The aerosol number concentration for each procedure and baseline measurement were compared
166 by calculating particle number concentration detected above background for each patient
167 (irrespective of particle size). As the length of procedure differed across patients, we sampled the


168 mean particle number concentration across the sampling time for each patient and the per patient
169 values were combined to give median and inter-quartile ranges of total aerosol number
170 concentration for each procedure.

171

172 Procedure aerosol size distributions

173 The aerosol size distributions from the phantom head control and patients were compared, with the
174 assumption that if the distributions were the same, all aerosol detected from the patient during the
175 procedure could be explained by the non-salivary contaminated instrument source (represented by
176 the phantom control). For each procedure, mean aerosol number concentrations (dN) for a range of
177 particle size bins (D_p) were calculated by averaging across patients. These were transformed
178 ($dN/d\log(D_p)/\text{cm}^{-3}$) in order to normalise the data, enabling visual comparison of the size
179 distribution in a standardised form typical for reporting aerosol size distributions (Fig. 1). For each
180 procedure, the shape of the phantom head control and patient size distributions were compared
181 visually.

182



183

184 **Figure 1.** Description of the parameters used to describe the average size distribution detected

185 above background. The mode width is described by $\log\sigma$ and the peak of the mode by $D_{p,c}$.

186 Mode amplitude parameter (N) was not compared, as it is highly dependent on sampling variability.

188 In the phantom head control, the model best describing the size distribution (highest r^2) was
189 identified by iteratively altering the number of modes (uni-modal, bi-modal, or tri-modal) using
190 **Supplementary Equation S1**. Once fitted, mode parameters (N , $D_{p,c}$, $\log(\sigma)$) were compared
191 between patient and phantom head control as illustrated in **Fig. 1**. Aerosol size distributions
192 inherently provide a robust fingerprint of source with different aerosol sources yielding different log-
193 normal distributions with different mean sizes and breadths/standard deviations. This enabled us to
194 attribute aerosols to specific sources e.g., phantom head or patient [21].

195

196 To further assess if size distributions between the phantom head control and patient differed other
197 than by chance, a two-sided unpaired t-test was used to compare the mode widths ($\log(\sigma)$) and
198 peak positions ($D_{p,c}$). The mode amplitude parameter (N) was not compared as it is highly dependent
199 on variabilities in sampling efficiency, for example direction of the exhaled airflow, which is not
200 comparable across phantom head and patient sampling. After accounting for multiple parameters
201 compared across instruments ($n=26$), a Bonferroni adjusted p -value of 0.002 was used.

202

203 **Results**

204 Forty-one patients were included in the study with a median age of 47 years (range: 18–75)
205 (**Supplementary Tables 1 & 2**). The mean background aerosol number concentration across patients
206 was 0.18 (+/- SD 0.12) particles cm^{-3} (180 particles per litre). In total, fifteen dental procedures were
207 assessed during periodontal, orthodontic, and oral surgery treatments. Of these, examination with a
208 dental probe, hand scaling, local anaesthetic delivery, routine extraction (with forceps and/or
209 elevator), raising a soft tissue flap, orthodontic bracket removal, alginate impression taking, 3-in-1
210 water only, and suturing did not produce any aerosol (**Supplementary Table 3**). For the other six
211 procedures where aerosol was detected, the percentage of total procedure time that aerosol was

212 observed was 12.7% for ultrasonic scaling, 24.8% for 3-in-1 air only, 75.3% for 3-in-1 air + water,
213 40.1% for high-speed drilling, 49.9% for slow speed drilling and 55.6% for surgical drilling (**Table 1**).

214

215 **Table 1.** Dental procedures for which aerosol was detected above background.

Procedure	Number of patients*	Total sampling time for procedure (s)	Time aerosol detected above background (s)	Percentage time aerosol detected above background (%)
Ultrasonic scaling	12	12,272	1,559	12.7
3-in-1 air only	35	801	199	24.8
3-in-1 air + water	33	772	581	75.3
High speed drilling	15	3,849	1,543	40.1
Slow speed drilling	15	3,324	1,632	49.9
Surgical drilling	9	568	316	55.6


* Some procedures were conducted in more than one patient

216

217 *Aerosol number concentrations from dental procedures*

218 The aerosol number concentration for each procedure is shown in **Fig. 2**. Participant breathing and
219 speaking had similar number concentrations and size distributions to background aerosol, indicating
220 the background dominated the signal for these activities. High speed drilling produced 10-fold more
221 aerosol (median 118.38 cm⁻³) compared to the other five procedures (ultrasonic scaling, surgical
222 drilling, 3-in-1 syringe air/ air + water, and slow speed drilling), which were comparable with median
223 number concentrations of approximately 10 cm⁻³.

224



225


226 **Figure 2.** Box and whiskers plot of total aerosol number concentration for baseline measurements
227 (orange) and dental procedures for which aerosol was detected above mean background value
228 (green). The aerosol number concentration is reported on a logarithmic scale.

229

230 *Aerosol size distributions from dental procedures*

231 Aerosol size distributions inherently provide a robust fingerprint of source. Different aerosol sources
232 yield different log-normal distributions with different mean sizes and breadths. Therefore, a detailed
233 analysis and comparison of size distributions from patient procedures and phantom head controls
234 enables identification of potential sources of salivary aerosol. The aerosol size distributions detected
235 from patient procedures are shown alongside phantom head controls on a linear scale
236 (**Supplementary Fig. 2**) and a logarithmic scale (**Supplementary Fig. 3**). Distributions show that,

237 within each procedure, the patient and phantom head have the same number of modes with similar
238 mode widths (log) and peaks ($D_{p,c}$) but different mode heights (N). Fitted size distributions for
239 ultrasonic scaling, 3-in-1 air + water and slow speed drilling are shown in **Fig. 3**, the remaining
240 procedure fits are presented in **Supplementary Fig. 4**.



241

242 **Figure 3.** Size distribution data for ultrasonic scaling (a, b), 3-in-1 (c, d) and slow-speed drilling (e, f).

243 Mode 1 (red line), mode 2 (green line), mode 3 (dark blue line) and cumulative bi- or tri-modal fit

244 (blue line). 95% confidence band is shown as the red shaded area.

245

246 **Table 2.** A comparison of aerosol size distribution parameters between the phantom control and
 247 patient data.

Procedure	Mode	Parameter	Phantom Mean (95% CI)	Patient Mean (95% CI)	Difference in means (95% CI)	P-value*
Ultrasonic scaling	1	Log(σ)	0.10 (0.05, 0.15)	0.29 (-0.24, 0.83)	0.19 (-0.90, 1.29)	0.74
		$D_{p,c}$ (μm)	0.73 (0.67, 0.79)	0.62 (0.44, 0.80)	0.11 (-0.26, 0.48)	0.57
	2	Log(σ)	0.18 (0.11, 0.25)	0.15 (-0.02, 0.31)	0.03 (-0.31, 0.37)	0.87
		$D_{p,c}$ (μm)	1.66 (1.41, 1.90)	1.82 (1.40, 2.24)	0.17 (-0.70, 1.03)	0.71
Surgical drilling	1	Log(σ)	0.13 (0.03, 0.23)	0.18 (-3.13, 3.49)	0.05 (-5.18, 5.29)	0.98
		$D_{p,c}$ (μm)	0.41 (0.29, 0.53)	0.23 (-5.12, 5.58)	0.18 (-8.28, 8.64)	0.97
	2	Log(σ)	0.14 (0.10, 0.18)	0.20 (-0.12, 0.53)	0.06 (-0.45, 0.58)	0.81
		$D_{p,c}$ (μm)	0.78 (0.56, 1.00)	0.55 (-1.29, 2.39)	0.23 (-2.68, 3.15)	0.88
3-in-1 air + water	1	Log(σ)	0.08 (0.07, 0.08)	0.09 (0.08, 0.09)	0.01 (-0.01, 0.03)	0.34
		$D_{p,c}$ (μm)	0.65 (0.64, 0.66)	0.65 (0.64, 0.65)	0.00 (-0.02, 0.02)	0.74
	2	Log(σ)	0.22 (0.19, 0.25)	0.18 (0.15, 0.20)	0.04 (-0.04, 0.12)	0.35
		$D_{p,c}$ (μm)	0.86 (0.70, 1.02)	1.16 (1.07, 1.25)	0.30 (-0.01, 0.62)	0.07
3-in-1 air only	1	Log(σ)	0.15 (0.00, 0.31)	0.07 (0.04, 0.11)	0.08 (-0.03, 0.20)	0.16
		$D_{p,c}$ (μm)	0.52 (0.32, 0.71)	0.59 (0.54, 0.65)	0.08 (-0.07, 0.23)	0.32
	2	Log(σ)	0.36 (0.00, 0.73)	0.28 (0.12, 0.44)	0.08 (-0.26, 0.42)	0.65
		$D_{p,c}$ (μm)	0.72 (-1.03, 2.48)	0.47 (-0.08, 1.03)	0.25 (-1.17, 1.67)	0.73
High speed drilling	1	Log(σ)	0.16 (0.15, 0.17)	0.12 (0.12, 0.13)	0.04 (0.02, 0.05)	1.10×10^{-5}
		$D_{p,c}$ (μm)	0.66 (0.65, 0.67)	0.65 (0.65, 0.66)	0.01 (-0.01, 0.02)	0.29
	2	Log(σ)	0.19 (0.16, 0.21)	0.18 (0.17, 0.20)	0.00 (-0.03, 0.04)	0.86
		$D_{p,c}$ (μm)	1.68 (1.59, 1.77)	1.41 (1.36, 1.47)	0.26 (0.12, 0.40)	1.81×10^{-3}
		Log(σ)	0.09 (0.03, 0.15)	0.07 (-0.10, 0.24)	0.03 (-0.38, 0.43)	0.90
		$D_{p,c}$ (μm)	4.32 (3.65, 5.00)	4.79 (2.92, 6.67)	0.47 (-3.97, 4.90)	0.84
Slow speed drilling	1	Log(σ)	0.34 (0.27, 0.40)	0.17 (0.16, 0.18)	0.17 (0.13, 0.20)	5.54×10^{-8}
		$D_{p,c}$ (μm)	1.00 (0.60, 1.40)	0.65 (0.64, 0.67)	0.35 (0.16, 0.53)	1.65×10^{-3}
	2	Log(σ)	0.19 (0.16, 0.22)	0.16 (0.14, 0.18)	0.03 (-0.01, 0.07)	0.22
$D_{p,c}$ (μm)		2.11 (2.01, 2.20)	1.55 (1.48, 1.63)	0.56 (0.40, 0.72)	1.89×10^{-6}	
	3	Log(σ)		0.11 (0.08, 0.14)		
		$D_{p,c}$ (μm)		3.48 (3.09, 3.86)		

P<0.002 is the Bonferroni adjusted equivalent of p<0.05

248

249 We assessed how aerosol size distributions from patients may differ from the phantom head control
 250 for the same procedure. Such assessment enables identification of sources of salivary aerosol,
 251 relative to the non-salivary aerosol generated by the dental instrument. For ultrasonic scaling, bi-
 252 modal fits to the patient and phantom head control data show a high level of agreement for the
 253 mode width (log(σ)) (Mode 1: $p= 0.74$; Mode 2 $p= 0.87$) and peak ($D_{p,c}$) (Mode 1: $p= 0.57$; Mode 2: $p=$
 254 0.71) between both patient and phantom head control (**Fig. 3a,b** and **Table 2**). This indicates that

255 the phantom head control data (instrument source) may account for all the aerosol seen during
256 ultrasonic patient procedures. Similarly, data from surgical drilling, 3-in-1 air + water and air alone
257 could be represented by bi-modal fits, with shape of size distribution curves similar for both the
258 patient and phantom head controls. Model parameters were similar again, suggesting the aerosol
259 detected arose from the dental instrument source (**Fig. 3c,d, Supplementary Fig. 4a-d, Table 2**).

260

261 For high-speed drilling, both the phantom head control and patient data fit can be represented by a
262 tri-modal fit, with similar shaped size distribution curves (**Supplementary Fig. 4e,f**). There was
263 statistical evidence passing the multiple testing threshold, that the size distribution modes were
264 different between the phantom head control and the patients ($\log(\sigma)$) (Mode 1: $p=1.10 \times 10^{-5}$) and
265 peak ($D_{p,c}$) (Mode 2: $p=1.81 \times 10^{-3}$), suggesting the patient aerosol size distribution may not be
266 completely explained by instrument aerosol (**Table 2**). Slow speed drilling phantom head control
267 data were best represented by a bi-modal fit, whereas three modes were required for the patient
268 data. These aerosol size distributions are different from each other, with clear divergence below 3
269 μm and above 7 μm particle sizes (**Fig. 3e,f**). This difference in size distributions was reinforced as
270 the parameters showed strong evidence for a difference in Mode 1 (mean difference $\log(\sigma) = 0.17$
271 (95% confidence intervals (95%CI), 0.13, 0.20), $p=5.54 \times 10^{-8}$; $D_{p,c} = 0.35$ (95%CI 0.16, 0.53), $p=1.65 \times$
272 10^{-3}) and Mode 2 (mean difference $\log(\sigma) = 0.03$ (95%CI -0.01, 0.07), $p=0.22$; $D_{p,c} = 0.56$ (95%CI 0.40,
273 0.72), $p=1.89 \times 10^{-6}$) (**Table 2**). Therefore, for slow speed drilling the phantom head control data do
274 not fully explain what was observed during patient procedures.

275

276 Discussion

277 This study explored aerosol generation during fifteen different dental procedures at source (as close
278 to the patient as possible). Only six procedures generated aerosol detectable above background. Our
279 background particle concentration was very low (0.18 cm^{-3}) and of similar magnitude to that

280 generated by a person speaking but less than that generated by a person coughing, enabling
281 confident detection of aerosol produced during dental procedures.

282

283 Of the six procedures that generated detectable aerosol, the size distributions observed in patients
284 closely matched those observed in phantom head controls for four of them: ultrasonic scaling, 3-in-1
285 air/ air + water and surgical drilling. In other words, we did not detect additional aerosol beyond that
286 generated by the dental instrument alone, which is a non-contaminated source. Dental instrument
287 aerosol could in principle be contaminated through impaction and resuspension in the mouth or
288 through coalescence with contaminated aerosol in the oral cavity. However, aerosol coalescence
289 rates within the dental aerosol plume are too small to be significant. For example, coalescence of 1
290 μm diameter particles at 100 cm^{-3} concentration proceeds with a coagulation coefficient equal to 3.4
291 $\times 10^{-16}\text{ m}^3\text{ s}^{-1}$, reducing the particle concentration to only 99.999 cm^{-3} in 100 s. Even for coalescence
292 of dissimilar size particles (e.g., 100 nm particles with 1 μm particles), coalescence is so inefficient
293 that the concentration changes by $<0.1\%$. Therefore, if aerosol from the dental instruments cannot
294 pick up patient biological aerosol by coalescence in the oral cavity (either respirable or resulting
295 from the dental procedure), the only remaining alternative is that the aerosol from the dental
296 instrument deposits in the oral cavity and then, having mixed with salivary components and
297 potentially infectious virus, is re-suspended by the instrument. This process would generate an
298 entirely new source of aerosol and be identifiable by the emergence of an additional mode in the
299 size distributions. Because the patient and phantom head size distributions match well for these
300 procedures, it is unlikely the measured aerosol is contaminated by patient biological material unless
301 the new aerosol is generated at a concentration low enough not to be resolved from the size
302 distribution of the instrument-generated aerosol. By contrast, with both high and slow speed drilling
303 there were differences observed between the phantom head and patient aerosol size distributions.
304 The presence of this unexplained aerosol suggests the generation of salivary aerosol and
305 consequently the potential for viral transmission.

306

307 Our study in patients supports findings from previous phantom head studies. Din et al. (2021),
308 showed that orthodontic debonding using a high speed drill led to the most significant increase in
309 particles, while combined use of the 3-in-1 air-water syringe did not result in any detectable increase
310 in the aerosol levels. Similarly, Allison et al. (2021) and an N-of-one human volunteer study [22]
311 found that ultrasonic scaling produces mainly instrument-generated aerosol. Similar to these groups,
312 we found the amount of aerosol generated by ultrasonic scaling was low in comparison to high
313 speed drilling (at least 10 times less) and intermittent, with no detectable aerosol for the majority of
314 the time the instrument was in use. This may reflect the non-continuous use of dental instruments,
315 that aerosol does not always escape the oral cavity, that aerosol is mitigated by use of high-volume
316 suction, or that there is directionality to the generated aerosol plume, which cannot be continuously
317 sampled.

318

319 Our study characterises aerosol generation during dental procedures but did not test for the
320 presence of SARS-CoV-2, although aerosols and droplets are the vehicles that transmit SARS-CoV-2.
321 Observation of increased aerosol generation does not confirm the potential for pathogen
322 transmission, and it is possible we were unable to identify potential salivary aerosol if the patient
323 size distribution were altered minimally from the phantom control or procedure generated new
324 aerosol at such a low concentration it could not be differentiated from that generated by the dental
325 instrument. While some air sampling studies have detected viable SARS-CoV-2, others have not, and
326 this remains technically challenging [23-25]. The use of time-of-flight mass spectrometry (MALDI-
327 TOF), fluorescein dye or salivary enzyme markers could be useful in determining if unexplained
328 aerosol contains biological material from the patient. For dental instrument-generated aerosol to
329 increase the risk of SARS-CoV-2 transmission, it must interact with saliva containing the virus, be of a
330 size distribution that can contain SARS-CoV-2, withstand irrigant dilution, and ultimately go on to
331 interact with a susceptible host.

332

333 In this work, the background aerosol level was low for a typical dental surgery (0.18 cm^{-3}), but still 50
334 times higher than can be achieved in a laminar flow theatre setting (Brown et al. 2021). Very low
335 levels of aerosol (e.g., from breathing or speaking) were not clearly resolved [19, 20]. Nonetheless,
336 we accurately measured aerosol size distributions to identify differences in patient data compared
337 to phantom head controls. Many factors that are uncontrollable in a clinical setting (e.g., patient
338 movement, differences in use of instruments by clinicians, specific tooth or quadrant treated) will
339 affect the aerosol number concentration, but these would minimally affect size distributions. Both
340 sets of experiments are comparable because the set-up for phantom head controls and patient
341 measurements were the same, including relative humidity and temperature.

342

343 While we investigated a wide range of dental procedures, it is not clear if these results can be
344 extrapolated to the same instrument being used for a different purpose (e.g., cutting a cavity using a
345 high speed drill) or different instruments performing the same procedure (e.g., piezo surgery instead
346 of surgical drilling). This study limited itself to aerosols in the $0.5\text{--}20 \mu\text{m}$ diameter size range, which
347 includes respirable aerosol. The removal of aerosol in this size range is mainly governed by room
348 ventilation. The studied procedures may generate larger droplets, which tend to behave more
349 ballistically [26]. Particles smaller than those studied here ($<0.5 \mu\text{m}$ diameter) are less likely to
350 harbour the virus [18]. Evaporation may occur between aerosol generation and measurement,
351 potentially altering the aerosol size distribution [27].

352

353 It has been suggested that the use of the term AGP should be reconsidered [28]. For instance,
354 coughing can occur during any dental procedure (**Supplementary Figure 5**) and may pose a higher
355 risk of viral transmission than many AGPs because coughing generates orders of magnitude higher
356 aerosol number concentrations than many AGPs [29-31]. The potential for viral transmission may be

357 governed primarily by the proximity of the dental care professional to the patient, given the
358 inevitable exposure to short range aerosol and droplet transmission of respiratory aerosol, rather
359 than by specific dental instruments or procedure. This study provides further evidence for sources of
360 aerosol generation during common dental procedures, enabling a more holistic approach to risk
361 assessment [28].

362

363 **Author Contributions**

364 T. Dudding, S. Sheikh, F. Gregson, J. Haworth, S. Haworth, A.J. Ireland, J.P. Reid, B.R. Bzdek and M.
365 Gormley contributed to conception and study design. T. Dudding, S. Sheikh, J. Haworth, and M.
366 Gormley were responsible for data acquisition and curation. T. Dudding, S. Sheikh, F. Gregson, and
367 M. Gormley drafted the initial manuscript and all authors contributed to subsequent drafts. Data
368 analysis was carried out by T. Dudding, S. Sheikh, F. Gregson, J. Haworth, S. Haworth, J.P. Reid and
369 B.R. Bzdek. Data interpretation and critical revisions were completed by T. Dudding, S. Sheikh, F.
370 Gregson, J. Haworth, S. Haworth, B.G. Main, A.J. Shrimpton, F.W. Hamilton, A.J. Ireland, N.A.
371 Maskell, B.R. Bzdek, J.P. Reid and M. Gormley and all authors helped write the final version of the
372 manuscript. This study is part of a wider collaboration represented by the AERATOR Group who were
373 responsible for the grant idea, writing and funding of this work. All authors gave final approval and
374 agree to be accountable for all aspects of the work.

375

376 **Acknowledgements**

377 Firstly, we would like to thank the patients who volunteered to take part in this study and for
378 recognising the value of this work. We thank all the management, administrative and clinical staff at
379 Bristol Dental Hospital and School for their logistical support with this study. In particular Ms Sarah
380 Bain, Dr Julie Weeks, Ms Sarah Constant, Ms Becki Bullock, Ms Kuldip Bhakerd, Ms Ceit Scott and Mr
381 James Tubman. We also extend our thanks to the nursing staff and clinicians who assisted with the

382 patient procedures, including Dr Alex Gormley, Dr Charlotte Richards and the BSc Dental Hygiene
383 and Therapy students. We would like to thank the Medical Equipment Management Organisation
384 Clinical Engineering (MEMO) team at University Hospitals Bristol and Weston Foundation Trust, Mr
385 John Woods (Nuview Ltd.) for sourcing and supplying the Jade SCA5000C Air Purification units
386 (Surgically Clean Air®, Toronto, Ontario) and Ms Adele Carter, for her assistance with arranging the
387 phantom head simulation equipment. Finally, we acknowledge the AERATOR Group who helped
388 write the grant to obtain the funding for this work.

389

390 *AERATOR Group members who contributed to this work include: Alice Milne, James Murray,
391 Johannes Keller, Jules Brown, Andrew Shrimpton, Anthony Pickering, Timothy Cook, Mark Gormley,
392 David Arnold, George Nava, Jonathan Reid, Bryan R Bzdek, Sadiyah Sheikh, Florence Gregson, Fergus
393 Hamilton, Nick Maskell, James Dodd, Ed Moran.

394

395 **Declaration of Conflicting Interests**

396 The authors declared no potential conflicts of interest with respect to the research, authorship,
397 and/or publication of this article. No corporate funding was received, or any material transfer
398 agreements made with any company or sponsor.

399

400 **Funding**

401 The AERosolisation And Transmission Of SARS-CoV-2 in Healthcare Settings (AERATOR) study was
402 funded by National Institute of Health Research (NIHR) (Award ID: COV0333) and has been awarded
403 urgent public health status (IRAS-Number: 288784; CPMS-ID: 47097). T. Dudding and S. Haworth are
404 supported by NIHR through the Academic Clinical Fellowship scheme. M. Gormley is supported by a
405 Wellcome Trust GW4-Clinical Academic Training PhD Fellowship. This research was funded in part,
406 by the Wellcome Trust [Grant number 220530/Z/20/Z]. For the purpose of open access, the author

407 has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from
408 this submission. Bryan R. Bzdek is supported by the Natural Environment Research Council
409 (NE/P018459/1).

410

411 **References**

- 412 1. World Health Organization. World Health Organization. Infection prevention and control
413 during health care when novel coronavirus (nCoV) infection is suspected 2020 [Available from:
414 <https://www.who.int/publications/i/item/10665-331495>.
- 415 2. Wilson N, Corbett S, Tovey E. Airborne transmission of covid-19. *BMJ*. 2020;370:m3206.
- 416 3. Jeong HW, Kim S-M, Kim H-S, Kim Y-I, Kim JH, Cho JY, et al. Viable SARS-CoV-2 in various
417 specimens from COVID-19 patients. *Clinical microbiology and infection : the official publication of*
418 *the European Society of Clinical Microbiology and Infectious Diseases*. 2020;26:1520-4.
- 419 4. Zhu J, Guo J, Xu Y, Chen X. Viral dynamics of SARS-CoV-2 in saliva from infected patients. *The*
420 *Journal of infection*. 2020;81:e48-e50.
- 421 5. Conway DI, Culshaw S, Edwards M, Clark C, Watling C, Robertson C, et al. SARS-CoV-2
422 Positivity in Asymptomatic-Screened Dental Patients. *J Dent Res*. 2021;100:583-90.
- 423 6. Public Health England. Public Health England. COVID-19: Infection prevention and control
424 dental appendix 2020 [Available from:
425 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/954834/Infection_prevention_and_control_guidance_dental_appendix.pdf)
426 [/954834/Infection_prevention_and_control_guidance_dental_appendix.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/954834/Infection_prevention_and_control_guidance_dental_appendix.pdf).
- 427 7. Ehtezazi T, Evans DG, Jenkinson ID, Evans PA, Vadgama VJ, Vadgama J, et al. SARS-CoV-2:
428 characterisation and mitigation of risks associated with aerosol generating procedures in dental
429 practices. *Br Dent J*. 2021:1-7.

- 430 8. Szymańska J. Dental bioaerosol as an occupational hazard in a dentist's workplace. *Ann Agric*
431 *Environ Med.* 2007;14:203-7.
- 432 9. Shahdad S, Patel T, Hindocha A, Cagney N, Mueller J-D, Seoudi N, et al. The efficacy of an
433 extraoral scavenging device on reduction of splatter contamination during dental aerosol generating
434 procedures: an exploratory study. *British Dental Journal.* 2020.
- 435 10. Public Health England. Public Health England. COVID-19: Guidance for maintaining services
436 within health and care settings: Infection prevention and control recommendations. 2021 [Available
437 from:
438 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/954690/Infection_Prevention_and_Control_Guidance_January_2021.pdf)
439 [/954690/Infection_Prevention_and_Control_Guidance_January_2021.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/954690/Infection_Prevention_and_Control_Guidance_January_2021.pdf).
- 440 11. Innes N, Johnson IG, Al-Yaseen W, Harris R, Jones R, Kc S, et al. A systematic review of
441 droplet and aerosol generation in dentistry. *Journal of dentistry.* 2021;105:103556-.
- 442 12. Leggat PA, Kedjarune U. Bacterial aerosols in the dental clinic: a review. *Int Dent J.*
443 2001;51:39-44.
- 444 13. Harrel SK, Molinari J. Aerosols and splatter in dentistry: a brief review of the literature and
445 infection control implications. *J Am Dent Assoc.* 2004;135:429-37.
- 446 14. Dawson M, Soro V, Dymock D, Price R, Griffiths H, Dudding T, et al. Microbiological
447 assessment of aerosol generated during debond of fixed orthodontic appliances. *Am J Orthod*
448 *Dentofacial Orthop.* 2016;150:831-8.
- 449 15. Allison JR, Currie CC, Edwards DC, Bowes C, Coulter J, Pickering K, et al. Evaluating aerosol
450 and splatter following dental procedures: Addressing new challenges for oral health care and
451 rehabilitation. *Journal of Oral Rehabilitation.* 2021;48:61-72.

- 452 16. Veena HR, Mahantesha S, Joseph PA, Patil SR, Patil SH. Dissemination of aerosol and splatter
453 during ultrasonic scaling: A pilot study. *Journal of Infection and Public Health*. 2015;8:260-5.
- 454 17. Din AR, Hindocha A, Patel T, Sudarshan S, Cagney N, Koched A, et al. Quantitative analysis of
455 particulate matter release during orthodontic procedures: a pilot study. *British Dental Journal*.
456 2020;<https://doi.org/10.1038/s41415-020-2280-5>.
- 457 18. Fennelly KP. Particle sizes of infectious aerosols: implications for infection control. *Lancet*
458 *Respir Med*. 2020;8:914-24.
- 459 19. Gregson FKA, Sheikh S, Archer J, Symons HE, Walker JS, Haddrell AE, et al. Analytical
460 challenges when sampling and characterising exhaled aerosol. *Aerosol Science and Technology*.
461 2021;1-16.
- 462 20. Gregson FKA, Watson NA, Orton CM, Haddrell AE, McCarthy LP, Finnie TJR, et al. Comparing
463 aerosol concentrations and particle size distributions generated by singing, speaking and breathing.
464 *Aerosol Science and Technology*. 2021;55:681-91.
- 465 21. Hinds WC. *Aerosol Technology: Properties, Behavior, and Measurement of Airborne*
466 *Particles*: John Wiley & Sons; 1999. 262-3 p.
- 467 22. Yang M, Chaghtai A, Melendez M, Hasson H, Whitaker E, Badi M, et al. Mitigating saliva
468 aerosol contamination in a dental school clinic. *BMC Oral Health*. 2021;21:52.
- 469 23. Greenhalgh T, Jimenez JL, Prather KA, Tufekci Z, Fisman D, Schooley R. Ten scientific reasons
470 in support of airborne transmission of SARS-CoV-2. *Lancet*. 2021;1;397(10285):1603-1605.
- 471 24. Borges JT, Nakada LYK, Maniero MG, Guimarães JR. SARS-CoV-2: a systematic review of
472 indoor air sampling for virus detection. *Environmental Science and Pollution Research*. 2021;25:1–
473 14.

- 474 25. Meethil AP, Saraswat S, Chaudhary PP, Dabdoub SM, Kumar PS. Sources of SARS-CoV-2 and
475 Other Microorganisms in Dental Aerosols. *J Dent Res*. 2021: 220345211015948.
- 476 26. Micik RE, Miller RL, Mazarella MA, Ryge G. Studies on dental aerobiology. I. Bacterial
477 aerosols generated during dental procedures. *J Dent Res*. 1969;48:49-56.
- 478 27. Walker JS, Archer J, Gregson FKA, Michel SES, Bzdek BR, Reid JP. Accurate Representations of
479 the Microphysical Processes Occurring during the Transport of Exhaled Aerosols and Droplets. *ACS*
480 *Central Science*. 2021;7:200-9.
- 481 28. Hamilton F, Arnold D, Bzdek BR, Dodd J, Reid J, Maskell N. Aerosol generating procedures:
482 are they of relevance for transmission of SARS-CoV-2? *Lancet Respir Med*. 2021;S2213-
483 2600(21)00216-2.
- 484 29. Gregson FKA, Shrimpton AJ, Hamilton F, Cook TM, Reid JP, Pickering AE, et al. Identification
485 of the source events for aerosol generation during oesophago-gastro-duodenoscopy. *Gut*.
486 2021:gutjnl-2021-324588.
- 487 30. Hamilton F, Gregson F, Arnold D, Sheikh S, Ward K, Brown J, et al. Aerosol emission from the
488 respiratory tract: an analysis of relative risks from oxygen delivery systems. *Thorax*. 2021;
489 doi.org/10.1136/thoraxjnl-2021-217577.
- 490 31. Sheikh S, Hamilton FW, Nava GW, Gregson FKA, Arnold DT, Riley C, et al. Are Aerosols
491 Generated During Lung Function Testing in Patients and Healthy Volunteers? Results From the
492 AERATOR Study. *Thorax*. 2021; [doi: 10.1136/thoraxjnl-2021-217671](https://doi.org/10.1136/thoraxjnl-2021-217671).
- 493