# Technology in Palliative Care (TIP): the identification of digital priorities for palliative care research using a modified Delphi method

## \*Amara Callistus Nwosu

Lancaster Medical School, Lancaster University, Lancaster, United Kingdom.

Marie Curie Hospice Liverpool, Liverpool, United Kingdom.

Liverpool University Hospitals NHS Foundation Trust, Liverpool, United Kingdom.

### **Tamsin McGlinchey**

Palliative Care Unit, University of Liverpool, United Kingdom.

## **Justin Sanders**

Dana-Farber Cancer Institute, Boston, MA, USA.

Ariadne Labs, Brigham and Women's Hospital and Harvard T.H. Chan School of Public Health, Boston, MA, USA.

Harvard Medical School, Boston, MA, USA.

### Sarah Stanley

Marie Curie Hospice Liverpool, Liverpool, United Kingdom.

### **Jennifer Palfrey**

Princess Alice Hospice, Surrey, United Kingdom.

## Patrick Lubbers

Netherlands Comprehensive Cancer Organization (IKNL), Utrecht, Netherlands.

# Laura Chapman

Marie Curie Hospice Liverpool, Liverpool, United Kingdom.

## Anne Finucane

Clinical Psychology, University of Edinburgh, Edinburgh, United Kingdom.

## **Stephen Mason**

Palliative Care Unit, University of Liverpool, Liverpool, United Kingdom.

WORD COUNT: 3675

# **KEY WORDS**

Terminal care; supportive care; quality of life; symptoms and symptom management

# **Corresponding author**

\*Amara Callistus Nwosu, Lancaster Medical School, Lancaster University, Lancaster,

LA1 4YG, United Kingdom.

Email: <u>a.nwosu@lancaster.ac.uk</u>

#### ABSTRACT

#### Background

Developments in digital health (describing technologies which use computing platforms, connectivity, software, and sensors for health care and related purposes) has the potential to transform the delivery of health and social care to help citizens manage their own health. Currently, we lack consensus about digital health research priorities in palliative care and lack theories about how these technologies might improve care outcomes. Global palliative care need is expected to increase due to the consequences of an ageing population; therefore, it is important for healthcare leaders to identify innovations to ensure that an increasingly frail population have appropriate access to palliative care services. Consequently, it is important to articulate research priorities as the first step to determine how we should allocate finite resources to a field saturated with rapidly developing innovations.

## Aims

To identify research priority areas for digital health in palliative care.

#### Methods

We selected the digital health trends, most relevant to palliative care, from a list of emerging trends reported by the 'Future Today Institute'. We conducted a modified Delphi process and consensus meeting with palliative care experts to identify research priorities. We used the views of public representatives to gain their perspectives of the agreed priorities.

### Results

One hundred and three experts (representing 11 countries) participated in the 1st Delphi round. Fifty-five participated in the 2<sup>nd</sup> round (53% of 1<sup>st</sup> round). Eleven experts attended the final consensus meeting. We identified 16 priorities areas, which were summarised into eight themes. These themes were: big data, mobile devices, telehealth and telemedicine, virtual reality, artificial intelligence, the smart home, biotechnology and digital legacy.

## Conclusions

The identified priorities in this paper represent a wide range of important emerging areas in field of digital health, personalised medicine, and data science. Humancentred design and robust governance systems should be considered in future research. It is important that the barriers and risks of using these technologies in palliative care are properly addressed to ensure that these tools are used meaningfully, wisely and safely and that do not cause unintentional harm.

# Technology in Palliative Care (TIP): the identification of digital priorities for palliative care research using a modified Delphi method

#### BACKGROUND

Developments in digital health (describing technologies which use computing platforms, connectivity, software, and sensors for health care and related purposes) has the potential to transform the delivery of health and social care to help citizens manage their own health.<sup>1-3</sup> Currently, we lack consensus about digital health research priorities in palliative care and lack theories about how these technologies might improve care outcomes. Therefore, it is important to articulate research priorities as the first step to determine how we should allocate finite resources to a field saturated with rapidly developing innovations. Global palliative care need is expected to increase due to the consequences of an ageing population; therefore, it is important for healthcare leaders to identify innovations to ensure that an increasingly frail population have appropriate access to palliative care services.<sup>4</sup> Research demonstrates that, when used well, digital health initiatives improve healthcare delivery and access,<sup>5-15</sup> and the World Health Organisation (WHO) promotes that digital health should be an integral part of health priorities as a means to improve health on a global scale.<sup>16 17</sup> To date, many barriers have prevented the meaningful use of digital health in palliative care;<sup>18</sup> these barriers include expense, inter-operability issues, data privacy and security concerns, lack of effectiveness, equity, and the concern that technology will reduce face-to-face consults between patients and clinicians.<sup>19 20</sup>

Strategic forethought (futurism) can help palliative care leaders to recognise emerging trends, to test, plan and use these innovations in practice.<sup>21</sup> Consequently, this study aims to identify digital health research priorities and to theorize how innovations in emerging technologies can improve palliative care.

#### AIM

To identify research priority areas for technology in palliative care.

### **METHODS**

We used a Delphi process, informed by the guidance on conducting and reporting Delphi studies (Guidance on Conducting and REporting DElphi Studies - CREDES<sup>22</sup>) in palliative care, to establish the opinion of palliative care experts. A Delphi process can be used as a consensus-based, forecasting process, enabling anonymous expert contributions to predict phenomena.<sup>23 24</sup> We chose to use the Delphi method for its potential to achieve consensus in areas of uncertainty.<sup>25-28</sup> We conducted two rounds of Delphi questionnaire, followed by a consensus meeting and a public engagement workshop to establish final consensus on research priorities for digital technology in palliative care. Data collection took place between November 2018 and September 2019.

### Identification of technology trends from the Future Today Institute

We selected technology trends most relevant to palliative care from a list of emerging technology trends reported by the Future Today Institute (FTI - <u>https://futuretodayinstitute.com</u>). The FTI is a multi-professional organisation that uses data-driven applied research to develop models that forecast risk and opportunity across several disciplines, which are mapped into technology trends.

The 2018 trend list included 225 emerging trends, which were stratified into 19 categories (Appendix: Future Today Institute 2018 Trends list).

#### Selection of technology trends for palliative care

We developed criteria to select the FTI trends, based on recommendations from a UK-based policy report, which reported public and professional views on new types of healthcare data.<sup>29</sup> We developed the following statement to select FTI trends for inclusion: 'Trends should involve analysis or use data generated by a patient, caregiver or healthcare professional with potential use in palliative care'. Two authors (ACN and TMc) reviewed all 225 FTI trends. We included 95 (42.2.%) of the trends. We then combined and simplified similar trends to reduce the number to 32 (Figure 1 - Flow diagram to outline study process for identifying research priority areas). To ratify the validity of the trends for palliative care, we conducted a focused literature review to identify examples where these technologies have been used in healthcare. We used an Excel spreadsheet to collate this data for reference.

# Figure 1: Flow diagram to outline study process for identifying research priority areas

### Delphi Questionnaire Development

We developed 32 items for inclusion in the Delphi questionnaire, which reflected the 32 trends identified from the Future Today Institute Report (see Figure 1 - Flow diagram to outline study process for identifying research priority areas). We used Google Forms (<u>https://www.google.co.uk/forms/about/</u>) to develop the survey. We designed the questionnaire to collect demographic information (geographic location,

age, occupation), and individuals' rating of importance for each item via a 5-point Likert scale (1 = low priority, 5 = high priority). To ensure that the survey questions were appropriate, we conducted a local pre-study pilot of the questionnaire and supporting materials (Appendix – 'Delphi Questionnaire' and 'Scoping review').

#### **Participant Recruitment and Consent**

We solicited a convenience sample of palliative care professionals with expressed interest in technological innovation; we used professional networks, social media and email to contact individuals (Appendix - Summary of the networks used to invite palliative care professionals to participate). Consenting participants accessed the study material online to complete an electronic consent form and the 1st round Delphi questionnaire. We invited participants who completed the 1<sup>st</sup> round questionnaire to participate in the 2nd round.

#### **Ethical approval**

This study was approved by the University of Liverpool Ethics Committee (study approval number 3564).

#### Data collection and analysis

Quantitative statistical analyses of participants ratings were undertaken with the statistical software package SPSS 22.0. We used the Interquartile Range (IQR) to determine the level of agreement on the five-point scales for each 'area' on the questionnaire. The justification for the levels of agreement were based on thresholds previously used in palliative care Delphi studies, which used a 5-point Likert scale to determine agreement (Appendix: Interquartile Range to be used to guide the Level of

Agreement for Delphi responses).<sup>22 30</sup> We emailed a summary of the 1<sup>st</sup> round Delphi results to each participant. The email included the following information: (i) a summary of how the participant rated each item in the first Delphi round, and (ii) a summary of all participants' responses for each item (pooled level of agreement). We provided this information so participants could consider whether they wished to rank items differently in the 2nd Delphi round, based on the ranking data generated by other participants.

## **Round 2 Delphi questionnaire**

We provided participants with an electronic link to access the 2<sup>nd</sup> round Delphi questionnaire. We asked participants to answer the same questions that were included in the first-round questionnaire. Participants were required to complete the questionnaire within 4 weeks. We analysed responses from the 2nd questionnaire by IQR to provide a final list of items according to their level of agreement.

## Final consensus meeting and voting

We organised a consensus meeting to agree the trend list as the final stage of the Delphi process.<sup>22</sup> We invited all participants to attend the meeting at the University of Liverpool, UK. We divided participants into two groups. We attempted to ensure the groups were similar by allocating individuals according to their gender, experience and occupation. We provided participants with the Delphi results, via (i) an oral presentation and (ii) a written summary. ACN and TMc acted as group facilitators and ACN chaired the meeting. We facilitated group discussion and voting to determine which items were included or excluded from the final list. Each item was

discussed and debated, and a 'raised-hand' vote was undertaken within each group to determine if each item was included or excluded from the final list.

After voting, we compared the outcomes for both groups. We included items if both groups voted for their inclusion. Similarly, we excluded items if both groups voted for exclusion. When the groups disagreed (i.e., one group voting for inclusion and the other voting for exclusion), we facilitated debate with both groups together, which was followed by rounds of voting until consensus was achieved.

## Public engagement workshop

Following the Consensus meeting, we conducted a public engagement workshop with lay representatives to determine their views on the agreed priorities. We used volunteer coordinators from Marie Curie Hospice Liverpool and Liverpool University Hospitals NHS Foundation Trust, to invite palliative care volunteers (by telephone and email).

### RESULTS

### Round 1 Delphi Questionnaire

Round 1 included 103 people participants (Table 1 - Demographics of study participants). The median age of participants was 45 years. Most participants were female (n = 65, 63.1%) and had a clinical background (n = 74, 72%). Participants represented 11 countries, most commonly the United Kingdom (n = 88, 85.4%). Most of the items (n = 25, 78%) achieved a median priority rating of 4 or 5 (Appendix: Level of agreement for each 'priority area' following both Delphi rounds), which suggests that participants considered most items were important.

### Round 2 Delphi

Fifty-five (53%) of the round 1 participants completed the round 2 questionnaire. The median age was 44 years, which was similar to round 1. More women than men completed the questionnaire (n = 32, 58.2%). The distribution of occupations was similar across both rounds. Fewer countries (n = 8) were represented among the final sample. The final IQR analysis (Appendix: Level of agreement for each 'priority area' following both Delphi rounds) demonstrates that most items (n = 21, 65.6%) had *low* levels of agreement, with two (6.3%) and nine (28.1%) items achieving *moderate* and *high* levels of agreement respectively.

### Consensus meeting and final list of priorities

Eleven people participated in the consensus meeting (10.7% of total participants and 20% of second round participants). The median age of participants was 44, and most were female (n = 7, 63.6%). All participants were based in the UK and were mostly from clinical (n = 6, 55%) or academic backgrounds (n = 4, 36%). The debate resulted in agreement, rejection, modification (rewording and combination) of trends, and the addition of a new item, digital legacy (Appendix: Voting outcomes for consensus meeting). We classified the priorities into eight themes which were: big data, mobile devices, telehealth/telemedicine, virtual reality, artificial intelligence, the smart home, biotechnology and digital legacy (Table 2 - Final list of priorities).

### Public Engagement Event

We conducted the public engagement event at Marie Curie Hospice Liverpool, UK, which was attended by six lay representatives, two staff members (nurse and doctor) and a medical student. In this meeting, we presented the Delphi outcomes and we

facilitated round table discussions to explore attendees' views on the agreed priorities. We asked attendees to identify areas within these themes that they wanted researchers to study further.

Our public representatives recommended that future research should: (1) ensure a human centre co-design approach to ensure that technologies are designed according to the needs of individuals and (2) that appropriate governance processes should be in place to evaluate efficacy, effectiveness and ethical issues of current and future digital health tools and systems.

## Table 1: Demographics of study participants

## Table 2: Final list of priorities

# DISCUSSION

## Summary of main findings

This is first study to identify digital health research priorities for palliative care and provides guidance for researchers, funders and policy makers to consider areas for future research and development. We identified 16 research priority areas for technology in palliative care, representing 8 themes of big data, mobile devices, telehealth, virtual reality, artificial intelligence, the smart home, biotechnology and digital legacy.

#### Contribution and strengths of this paper

The outcomes of our detailed analysis (involving a modified Delphi process and patient engagement workshop) indicates further digital health research is needed to identify how to best use technology to support palliative care. Our paper is the first priority-setting paper for palliative care digital health and provides a foundation for digital health focused palliative care research.

#### Telehealth and telemedicine

Prior to the novel coronavirus disease 2019 (COVID19) pandemic, researchers highlighted the potential to use telehealth (i.e., technology to support remote clinical access), and telemedicine (i.e., technology to support remote clinical care delivery) in palliative care. These technologies are increasingly used in palliative care;<sup>31 32</sup> however, many are un-evaluated for use in real-world settings.<sup>19 33</sup> Beyond the pandemic, researchers can consider how these technologies can improve palliative care access (e.g. for remote communities, hard to reach groups) to support new models of care (e.g. tele-palliative care clinics). It is important to consider different issues with this technology, such as reasons for use, managing distress, privacy and security issues, quality of communication and care, equity of access and barriers and facilitators to implementation.

### Artificial Intelligence

Artificial intelligence (AI) is often used as an umbrella term to describe a number of processes (e.g. Machine Learning, Natural Language Processing (NLP), deep learning, neural networks).<sup>34</sup> Clinicians and researchers are increasingly using AI to predict survival,<sup>35-38</sup> classify pain severity,<sup>39 40</sup> identify quality indicators,<sup>41 42</sup> and to

identify serious illness conversations from electronic healthcare records.<sup>43</sup> However, most of these studies are exploratory and do not provide recommendations for clinical practice.<sup>18</sup> Therefore, researchers should explore how different AI techniques can support palliative care research and practice, with consideration to the ethical issues associated with these methods.

#### Big data

Big data describes the large amounts of (previously unmanageable) data, which can now be processed by modern-day computer analysis techniques. Researchers have reported opportunities to use routine data to support palliative care decisions for populations and inidviduals.<sup>44 45</sup> Currently, there is no consensus for how nontraditional sources of big-data can be used meaningfully in palliative care. For example, patient-generated data (e.g., from mobile devices and wearables) can potentially be used to evaluate quality-of-life. Further, open source genomic databases may provide opportunities to better study the interplay between genetics and health, to inform how services use data to plan care for people with specific illnesses. Social media, and other forms of online data, are increasingly used to support public and professional communication, and to gain insight on the public attitudes to palliative care.<sup>46-49</sup> Consequently, researchers should identify what data to collect, and how to best use both traditional and non-traditional sources of palliative care big data.<sup>18 50 51</sup>

#### Mobile devices and wearables

Many studies have described how mobile devices and wearables can support palliative care (e.g. remote monitoring of physical activity and symptoms, to deliver

wellbeing activity, for documentation of advance care planning, education access/delivery and guideline access).<sup>52-56</sup> The capabilities of these devices to collect and store data is increasing; therefore, it is important to determine how this data can be meaningfully used.<sup>57 58</sup> Researchers have previously described how patient-reported outcomes (PRO) can improve palliative care patients,<sup>59-61</sup> however, further work is needed to explore how this technology can best support PRO collection (and use) in real world settings.<sup>62 63</sup> It is important to examine how mobile devices are designed to meet the requirements of users with palliative care needs.<sup>64</sup> Furthermore, studies should provide more information of how mobile devices can help patients to record their care preferences (e.g., advance care planning).<sup>65 66</sup>

#### Virtual reality

Virtual reality (VR) is a human-computer interface technology that uses visual graphics, sounds and other sensory input to create an interactive computer world.<sup>67</sup> Previous studies have described the potential to use VR to support psycho-social symptoms and wellbeing; however, most work is unevaluated so further research is needed.<sup>68-71</sup> We recognise the potential of VR to support palliative care education;<sup>72</sup> <sup>73</sup> however, the Consensus group did not identify this as a current priority. Following our study, we recognize that COVID19 pandemic has accelerated the use of virtual learning environments for medical education,<sup>74</sup> particularly with the potential to use VR for communication skills training.<sup>72</sup> Consequently, it is possible that VR for education would rate higher as a palliative care digital health priority if this study were repeated.

#### The smart home

A smart home describes a living environment where sensor-based systems and internet-connected devices (the Internet of Things) are used for remote monitoring and automation of appliances, such as lighting and heating.<sup>75</sup> Previous studies illustrate how various technologies have been used to support care for people who experience a decline in physical function (e.g. virtual assistants and supportive robotics), which highlights the wider role that these technologies may have in palliative care practice.<sup>76</sup> Consequently, future work should explore the usefulness of smart home technologies in supporting physical function, and the legal, privacy and ethical issues associated with these developments.<sup>51 3 65 75-77</sup>

#### Biotechnology

Biotechnology involves the combination of technology with living things.<sup>78</sup> Palliative care developments in this area include use biomarkers to predict survival.<sup>79 80</sup> Other studies have identified biomarkers to predict constipation<sup>81</sup> and delirium,<sup>82 83</sup> and highlighted potential to personalise cancer pain to an individual's genetics.<sup>84-86</sup> Consequently, it is possible to imagine future scenarios where technologies are used for early identification (and prediction) of clinical issues, facilitating personalised treatment for the individual (e.g. early identification and management of pathological fracture).

#### Digital legacy

A digital legacy is the digital information available about someone after death, such as social media profiles, photos, videos and gaming profiles.<sup>87</sup> The volume of digital information generated by citizens is increasing, which creates new challenges after

death.<sup>88</sup> The increasing use of cloud storage and social media is contributing to uncertainty of data ownership, which creates difficulties for caregivers to manage the digital legacy of the deceased. Studies demonstrate that healthcare professionals can positively support their patients to manage their digital legacy.<sup>87 89 90</sup> However, digital legacy is not routinely discussed in clinical practice, which means that we generally do not know how individuals want their data to be managed after death.<sup>91</sup> Therefore, we believe that researchers should explore how patients and caregivers can be supported to manage their digital legacy after death, with exploration on the different methods and materials that can be used.

Relation to previous work in this area and areas of interest following the novel COVID19 pandemic

Our study is synergistic with previous work, which has been conducted across the theme areas.<sup>19 33</sup> We acknowledge that our study pre-dates the pandemic and it is possible that the priorities we identified may now have shifted. However, we believe our research findings are valid as the digital health innovations adopted during the pandemic were in sync with our priority list. (Appendix: Examples of technologies used in palliative care during the COVID19 pandemic).<sup>32 33</sup> Telehealth was commonly used during the pandemic, with many palliative care services using this to provide remote clinical support,<sup>92-104</sup> to communicate<sup>105</sup> and for education.<sup>106</sup> Technologies were used to maintain connection, and to develop communities of palliative care practice.<sup>107 108</sup> VR was used to provide psychological care and symptom management.<sup>109 110</sup> In general, the findings these studies are positive about the potential of digital health to improve palliative care; however, its rapid

implementation during the pandemic has created a number of challenges (e.g., technical issues, data security and wellbeing considerations) which, require further evaluation.<sup>105</sup>

#### Limitations

It is possible that recent developments were not reflected in the priority list due to ongoing advancement of healthcare technologies. For example, the FTI trends list is now in its 2021 version and includes new trends such as, home medical laboratory tests and remote metabolic monitoring. Therefore, it is possible that relevant areas are absent from this analysis. Also, a weakness of digital health research is the rapid change associated with technology, which may cause the findings of this study to lose relevancy overtime.

Our decision to reduce the number of trends from 95 to 32 items, has broadened the focus of the list, which means it is possible that more specific and technical areas were not explored in greater depth (e.g., faceprints, voiceprints, chatbots etc etc). It is also possible that our Delphi participants will have different views on priority of some areas post COVID19, due to the resulting increase of clinical applications of digital health. Questionnaires were mostly completed by participants arising from English-speaking countries, meaning that the experience of non-English speaking populations may not be reflected. It is possible, due to the novel nature of some areas, that participants gave more priority to familiar areas and therefore, less priority to unfamiliar areas. The final priority list may not represent non-UK healthcare systems, as the consensus meeting was only attended by UK residents. Furthermore, people from different professional backgrounds (including cultures and

settings) may assign different levels of priority to trends, due their experience, workrequirements and personal beliefs. As most participants were clinically-focused, it is possible that the priorities were orientated to clinical-utility, rather than methodology.

#### Relevance to research, practice and policy

Decision-makers should ensure that technology is relevant to the needs of the palliative care user, as these requirements will influence the design, use and function of systems. For example, healthcare professionals will generally use technology to access patient data and communicate with other professionals, whereas patients may wish to access their own health data and to contact healthcare services. Further research is needed to develop specific use-cases for these scenarios, to ensure that the technology can be used meaningfully to achieve the intended outcomes. Furthermore, as the user requirements of people with palliative care needs may differ from the general population<sup>111</sup> and because we lack resources for wide-spread implementation of all technologies currently, it is important that digital health studies provide the data needed for determine best practice, and to help identify the barriers and facilitators for adoption.

Researchers should use appropriate methodologies to explore these questions and should study associated areas such as ethical issues, data security, and design. It is important that researchers work with the public, as the comments of the lay representatives in our study (from both the consensus meeting and the public engagement workshop) highlighted concerns of how personal health information is collected, stored, and used. Policymakers should consider issues related to governance and ethics of current and future digital interventions and systems. From a design perspective, we suggest that palliative care professionals work

collaboratively with creative industries (e.g., designers, developers and engineers) to ensure that designed technologies fulfil the user requirements for specific palliative care use-cases.

## Conclusion

The identified priorities in this paper represent a wide range of important emerging areas in the field of digital health, personalised medicine, and data science. Humancentred design and robust governance systems should be considered in future research. Transdisciplinary studies using a range of methodologies are required to further study this priority list. It is important that the barriers and risks of using these technologies in palliative care are properly addressed to ensure that these tools are used meaningfully, wisely and safely and that we do not cause unintentional harm.

## ACKNOWLEDMENTS

This research was supported by the following funding streams: Liverpool Clinical Commissioning Group (CCG) Research Capability Funding (researcher salary), £20,970. Wellcome Trust, Public Engagement Grants Scheme (Public Engagement Event costs), £360. Liverpool Clinical Commissioning Group (consensus meeting costs) £807. The posts of ACN and SS were funded by Marie Curie: https://www.mariecurie.org.uk/

# **AUTHOR CONTRIBUTIONS**

The author's responsibilities were as follows:

Study design: ACN, TMc

Data collection: ACN, TMc

Paper writing: ACN, TMc, JS, JP, LC, SS

Critique and review of the final manuscript: ACN, TMc, SS, LC, JP, AF, PL, JS, SM

## **COMPETING INTEREST STATEMENT**

The authors declare no competing interests.

## ETHICS

This study was approved by the University of Liverpool Ethics Committee (study approval number 3564).

# DATA AVAILABILITY

The authors declare that the data supporting the findings of this study are available within the paper and its supplementary information files.

## LIST OF ABBREVIATIONS USED

- AI: Artificial Intelligence
- EHR: Electronic Health Records
- ePROM: electronic patient-reported measures
- FTI: Future Today Institute
- ML: Machine learning

NHS: National Health Service
NLP: Natural Language Processing
PACD: Physical Activity Change Detection
PGHD: Patient-Generated Health Data
PROMs: Patient reported outcome measures
VR: Virtual Reality

# REFERENCES

- Topol E. Preparing the Healthcare Workforce to Deliver the Digital Future the Topol Review: An Independent Report on Behalf of the Secretary of State for Health and Social Care. NHS Health Education: London, UK 2019
- Department of Health Social Care. The future of healthcare: our vision for digital, data and technology in health and care: DHSC London, 2018.
- Haghi M, Thurow K, Stoll R. Wearable Devices in Medical Internet of Things: Scientific Research and Commercially Available Devices. *Healthcare informatics research* 2017;23(1):4-15. doi: 10.4258/hir.2017.23.1.4 [published Online First: 2017/03/07]
- 4. Bone AE, Gomes B, Etkind SN, et al. What is the impact of population ageing on the future provision of end-of-life care? Population-based projections of place of death. *Palliative Medicine* 2017;32(2):329-36. doi:

10.1177/0269216317734435

 Barrett M, Combs V, Su JG, et al. AIR Louisville: addressing asthma with technology, crowdsourcing, cross-sector collaboration, and policy. *Health Affairs* 2018;37(4):525-34.

- Morawski K, Ghazinouri R, Krumme A, et al. Association of a smartphone application with medication adherence and blood pressure control: the MedISAFE-BP randomized clinical trial. *JAMA internal medicine* 2018;178(6):802-09.
- 7. Wild SH, Hanley J, Lewis SC, et al. Supported telemonitoring and glycemic control in people with type 2 diabetes: the telescot diabetes pragmatic multicenter randomized controlled trial. *PLoS medicine* 2016;13(7):e1002098.
- Koehler F, Koehler K, Prescher S, et al. Mortality and morbidity 1 year after stopping a remote patient management intervention: extended follow-up results from the telemedical interventional management in patients with heart failure II (TIM-HF2) randomised trial. *The Lancet Digital Health* 2020;2(1):e16e24.
- de Jong MJ, van der Meulen-de AE, Romberg-Camps MJ, et al. Telemedicine for management of inflammatory bowel disease (myIBDcoach): a pragmatic, multicentre, randomised controlled trial. *The Lancet* 2017;390(10098):959-68.
- Denis F, Yossi S, Septans A-L, et al. Improving survival in patients treated for a lung cancer using self-evaluated symptoms reported through a web application. *American journal of clinical oncology* 2017;40(5):464-69.
- 11. Müller KI, Alstadhaug KB, Bekkelund SI. A randomized trial of telemedicine efficacy and safety for nonacute headaches. *Neurology* 2017;89(2):153-62.
- Rono HK, Bastawrous A, Macleod D, et al. Smartphone-based screening for visual impairment in Kenyan school children: a cluster randomised controlled trial. *The Lancet Global Health* 2018;6(8):e924-e32.

- Freeman D, Sheaves B, Goodwin GM, et al. The effects of improving sleep on mental health (OASIS): a randomised controlled trial with mediation analysis. *The Lancet Psychiatry* 2017;4(10):749-58.
- 14. Kollins SH, DeLoss DJ, Cañadas E, et al. A novel digital intervention for actively reducing severity of paediatric ADHD (STARS-ADHD): a randomised controlled trial. *The Lancet Digital Health* 2020;2(4):e168-e78.
- 15. Craig TK, Rus-Calafell M, Ward T, et al. AVATAR therapy for auditory verbal hallucinations in people with psychosis: a single-blind, randomised controlled trial. *The Lancet Psychiatry* 2018;5(1):31-40.
- 16. Mariano B. Towards a global strategy on digital health. *Bulletin of the World Health Organization* 2020;98(4):231.
- 17. World Health Organization. Global strategy on digital health 2020–2025: World Health Organization, 2020.
- 18. Nwosu AC, Collins B, Mason S. Big Data analysis to improve care for people living with serious illness: The potential to use new emerging technology in palliative care. *Palliative Medicine* 2018;32(1):164-66. doi:

10.1177/0269216317726250

- Hancock S, Preston N, Jones H, et al. Telehealth in palliative care is being described but not evaluated: a systematic review. *BMC palliative care* 2019;18(1):114.
- 20. Nwosu AC, Sturgeon B, McGlinchey T, et al. Robotic technology for palliative and supportive care: Strengths, weaknesses, opportunities and threats. *Palliat Med* 2019;33(8):1106-13. doi: 10.1177/0269216319857628 [published Online First: 2019/06/30]

- 21. Archibald MM, Barnard A. Futurism in nursing: Technology, robotics and the fundamentals of care. *Journal of Clinical Nursing* 2018;27(11-12):2473-80.
- 22. Jünger S, Payne SA, Brine J, et al. Guidance on Conducting and REporting DElphi Studies (CREDES) in palliative care: Recommendations based on a methodological systematic review. *Palliative Medicine* 2017;31(8):684-706. doi: 10.1177/0269216317690685
- 23. Rowe G, Wright G. Expert opinions in forecasting: the role of the Delphi technique. Principles of forecasting: Springer 2001:125-44.
- 24. Ono R, Wedemeyer DJ. Assessing the validity of the Delphi technique. *Futures* 1994;26(3):289-304.
- 25. von der Gracht HA. The Delphi technique for futures research. *The Future of Logistics: Scenarios for 2025* 2008:21-68.
- 26. Dalkey NC, Brown BB, Cochran S. The Delphi method: An experimental study of group opinion: Rand Corporation Santa Monica, CA 1969.
- 27. Avella JR. Delphi panels: Research design, procedures, advantages, and challenges. *International Journal of Doctoral Studies* 2016;11(1):305-21.
- 28. Hsu C-C, Sandford BA. The Delphi technique: making sense of consensus. *Practical assessment, research & evaluation* 2007;12(10):1-8.
- 29. The Academy of Medical Sciences. Our data-driven future in healthcare. Website of the Academy of Medical Sciences, 2018.
- 30. Jünger S, Payne S, Brearley S, et al. Consensus building in palliative care: a Europe-wide delphi study on common understandings and conceptual differences. *J Pain Symptom Manage* 2012;44(2):192-205. doi: 10.1016/j.jpainsymman.2011.09.009 [published Online First: 2012/06/19]

31. Wherton J, Shaw S, Papoutsi C, et al. Guidance on the introduction and use of video consultations during COVID-19: important lessons from qualitative research. *BMJ Leader* 2020:leader-2020-000262. doi: 10.1136/leader-2020-000262

32. Etkind SN, Bone AE, Lovell N, et al. The Role and Response of Palliative Care and Hospice Services in Epidemics and Pandemics: A Rapid Review to Inform Practice During the COVID-19 Pandemic. *Journal of pain and symptom management* 2020;60(1):e31-e40. doi:

10.1016/j.jpainsymman.2020.03.029 [published Online First: 2020/04/08]

- 33. Finucane A, O'Donnell H, Lugton J, et al. Digital Health Interventions in Palliative
  Care: A Systematic Meta-Review and Evidence Synthesis. *medRxiv*2020:2020.09.16.20195834. doi: 10.1101/2020.09.16.20195834
- 34. Chen M, Decary M. Artificial intelligence in healthcare: An essential guide for health leaders. *Healthc Manage Forum* 2020;33(1):10-18. doi: 10.1177/0840470419873123 [published Online First: 2019/09/26]
- 35. Einav L, Finkelstein A, Mullainathan S, et al. Predictive modeling of U.S. health care spending in late life. *Science* 2018;360(6396):1462-65. doi: 10.1126/science.aar5045 [published Online First: 2018/06/30]
- 36. Makar M, Ghassemi M, Cutler DM, et al. Short-term Mortality Prediction for Elderly Patients Using Medicare Claims Data. *Int J Mach Learn Comput* 2015;5(3):192-97. doi: 10.7763/ijmlc.2015.V5.506 [published Online First: 2015/06/01]
- 37. Sahni N, Simon G, Arora R. Development and Validation of Machine Learning Models for Prediction of 1-Year Mortality Utilizing Electronic Medical Record Data Available at the End of Hospitalization in Multicondition Patients: a

Proof-of-Concept Study. J Gen Intern Med 2018;33(6):921-28. doi:

10.1007/s11606-018-4316-y [published Online First: 2018/02/01]

38. Avati A, Jung K, Harman S, et al. Improving palliative care with deep learning. BMC medical informatics and decision making 2018;18(Suppl 4):122. doi:

10.1186/s12911-018-0677-8 [published Online First: 2018/12/13]

 Heintzelman NH, Taylor RJ, Simonsen L, et al. Longitudinal analysis of pain in patients with metastatic prostate cancer using natural language processing of medical record text. *J Am Med Inform Assoc* 2013;20(5):898-905. doi: 10.1136/amiajnl-2012-001076 [published Online First: 2012/11/13]

- 40. Lodhi MK, Stifter J, Yao Y, et al. Predictive Modeling for End-of-Life Pain
  Outcome using Electronic Health Records. *Adv Data Min* 2015;9165:56-68.
  doi: 10.1007/978-3-319-20910-4\_5 [published Online First: 2016/08/09]
- Lodhi MK, Ansari R, Yao Y, et al. Predictive Modeling for Comfortable Death Outcome Using Electronic Health Records. *Proc IEEE Int Congr Big Data* 2015;2015:409-15. doi: 10.1109/BigDataCongress.2015.67 [published Online First: 2015/06/01]
- 42. Lindvall C, Lilley EJ, Zupanc SN, et al. Natural Language Processing to Assess End-of-Life Quality Indicators in Cancer Patients Receiving Palliative Surgery. *J Palliat Med* 2019;22(2):183-87. doi: 10.1089/jpm.2018.0326 [published Online First: 2018/10/18]
- 43. Chan A, Chien I, Moseley E, et al. Deep learning algorithms to identify documentation of serious illness conversations during intensive care unit admissions. *Palliat Med* 2019;33(2):187-96. doi: 10.1177/0269216318810421 [published Online First: 2018/11/15]

- 44. Tanuseputro P. Delivering care to those in need: Improving palliative care using linked data. *Palliat Med* 2017;31(6):489-91. doi: 10.1177/0269216317704629
  [published Online First: 2017/04/26]
- 45. Nwosu AC, Collins B, Mason S. Big Data analysis to improve care for people living with serious illness: The potential to use new emerging technology in palliative care. *Palliat Med* 2017:269216317726250. doi:

10.1177/0269216317726250 [published Online First: 2017/08/15]

- 46. Selman LE, Chamberlain C, Sowden R, et al. Sadness, despair and anger when a patient dies alone from COVID-19: A thematic content analysis of Twitter data from bereaved family members and friends. *Palliat Med* 2021:2692163211017026. doi: 10.1177/02692163211017026 [published Online First: 2021/05/22]
- 47. Nwosu AC, Debattista M, Rooney C, et al. Social media and palliative medicine: a retrospective 2-year analysis of global Twitter data to evaluate the use of technology to communicate about issues at the end of life. *BMJ Support Palliat Care* 2015;5(2):207-12. doi: 10.1136/bmjspcare-2014-000701 [published Online First: 2014/09/04]
- Parker Oliver D, Washington K, Gage LA, et al. The promise of secret Facebook groups for active family caregivers of hospice patients. *J Palliat Med* 2014;17(11):1199-200. doi: 10.1089/jpm.2014.0311 [published Online First: 2014/10/15]
- Selman LE, Sowden R, Borgstrom E. 'Saying goodbye' during the COVID-19 pandemic: A document analysis of online newspapers with implications for end of life care. *Palliat Med* 2021:2692163211017023. doi: 10.1177/02692163211017023 [published Online First: 2021/05/22]

- 50. Rajaram A, Morey T, Dosani N, et al. Palliative Care in the Twenty-First Century: Using Advanced Analytics to Uncloak Insights from Big Data. *J Palliat Med* 2019;22(2):124-25. doi: 10.1089/jpm.2018.0609 [published Online First: 2019/02/02]
- 51. Harris J, Cheevers K, Armes J. The emerging role of digital health in monitoring and supporting people living with cancer and the consequences of its treatments. *Current opinion in supportive and palliative care* 2018;12(3):268-75. doi: 10.1097/spc.0000000000000362 [published Online First: 2018/06/22]
- 52. Nwosu AC, Mason S. Palliative medicine and smartphones: an opportunity for innovation? *BMJ Support Palliat Care* 2012;2(1):75-7. doi:

10.1136/bmjspcare-2011-000151 [published Online First: 2012/03/01]

- 53. Bienfait F, Petit M, Pardenaud R, et al. Applying M-Health to Palliative Care: A Systematic Review on the Use of M-Health in Monitoring Patients With Chronic Diseases and its Transposition in Palliative Care. Am J Hosp Palliat Care 2020;37(7):549-64. doi: 10.1177/1049909119885655 [published Online First: 2019/11/28]
- 54. Weekly T, Walker N, Beck J, et al. A Review of Apps for Calming, Relaxation, and Mindfulness Interventions for Pediatric Palliative Care Patients. *Children* (*Basel*) 2018;5(2) doi: 10.3390/children5020016 [published Online First: 2018/01/27]
- Meghani SH, MacKenzie MA, Morgan B, et al. Clinician-Targeted Mobile Apps in Palliative Care: A Systematic Review. *J Palliat Med* 2017;20(10):1139-47. doi: 10.1089/jpm.2017.0070 [published Online First: 2017/05/31]

- Phongtankuel V, Adelman RD, Reid MC. Mobile health technology and home hospice care: promise and pitfalls. *Prog Palliat Care* 2018;26(3):137-41. doi: 10.1080/09699260.2018.1467109 [published Online First: 2018/12/07]
- 57. Maggi N, Douglas Magnoni L, Ruggiero C, et al. Information Technology System Including Patient Generated Health Data for Cancer Clinical Care and Research. *Studies in health technology and informatics* 2019;261:289-93. [published Online First: 2019/06/04]
- 58. Sayeed R, Gottlieb D, Mandl KD. SMART Markers: collecting patient-generated health data as a standardized property of health information technology. NPJ Digit Med 2020;3:9. doi: 10.1038/s41746-020-0218-6 [published Online First: 2020/01/30]
- 59. Dudgeon D. The Impact of Measuring Patient-Reported Outcome Measures on Quality of and Access to Palliative Care. *J Palliat Med* 2018;21(S1):S76-s80. doi: 10.1089/jpm.2017.0447 [published Online First: 2017/12/29]
- 60. Johnston B, Flemming K, Narayanasamy MJ, et al. Patient reported outcome measures for measuring dignity in palliative and end of life care: a scoping review. *BMC Health Serv Res* 2017;17(1):574. doi: 10.1186/s12913-017-2450-6 [published Online First: 2017/08/24]
- 61. Kavalieratos D, Corbelli J, Zhang D, et al. Association Between Palliative Care and Patient and Caregiver Outcomes: A Systematic Review and Metaanalysis. *Jama* 2016;316(20):2104-14. doi: 10.1001/jama.2016.16840 [published Online First: 2016/11/29]
- 62. Benze G, Nauck F, Alt-Epping B, et al. PROutine: a feasibility study assessing surveillance of electronic patient reported outcomes and adherence via

smartphone app in advanced cancer. *Ann Palliat Med* 2019;8(2):104-11. doi: 10.21037/apm.2017.07.05 [published Online First: 2017/11/22]

- 63. Abernethy AP, Ahmad A, Zafar SY, et al. Electronic patient-reported data capture as a foundation of rapid learning cancer care. *Medical care* 2010;48(6 Suppl):S32-8. doi: 10.1097/MLR.0b013e3181db53a4 [published Online First: 2010/05/18]
- 64. Mandel JC, Kreda DA, Mandl KD, et al. SMART on FHIR: a standards-based, interoperable apps platform for electronic health records. *J Am Med Inform Assoc* 2016;23(5):899-908. doi: 10.1093/jamia/ocv189 [published Online First: 2016/02/26]
- 65. Pavic M, Klaas V, Theile G, et al. Feasibility and Usability Aspects of Continuous Remote Monitoring of Health Status in Palliative Cancer Patients Using Wearables. Oncology 2020;98(6):386-95. doi: 10.1159/000501433 [published Online First: 2019/07/25]
- 66. Nwosu AC, Quinn C, Samuels J, et al. Wearable smartwatch technology to monitor symptoms in advanced illness. *BMJ supportive & palliative care* 2018;8(2):237-37.
- 67. Chirico A, Lucidi F, De Laurentiis M, et al. Virtual Reality in Health System: Beyond Entertainment. A Mini-Review on the Efficacy of VR During Cancer Treatment. *Journal of cellular physiology* 2016;231(2):275-87. doi: 10.1002/jcp.25117 [published Online First: 2015/08/05]
- 68. Johnson T, Bauler L, Vos D, et al. Virtual Reality Use for Symptom Management in Palliative Care: A Pilot Study to Assess User Perceptions. *Journal of Palliative Medicine* 2020

69. Niki K, Okamoto Y, Maeda I, et al. A Novel Palliative Care Approach Using Virtual Reality for Improving Various Symptoms of Terminal Cancer Patients: A Preliminary Prospective, Multicenter Study. *J Palliat Med* 2019;22(6):702-07. doi: 10.1089/jpm.2018.0527 [published Online First: 2019/01/25]

- 70. Austin P, Lovell M, Siddall P. The Efficacy of Virtual Reality for Persistent Cancer
  Pain: A Call for Research. *J Pain Symptom Manage* 2019;58(4):e11-e14. doi:
  10.1016/j.jpainsymman.2019.07.009 [published Online First: 2019/07/20]
- 71. Hsieh W-T. Virtual reality video promotes effectiveness in advance care planning. *BMC palliative care* 2020;19(1):1-10.
- 72. Evans L, Taubert M. State of the science: the doll is dead: simulation in palliative care education. *BMJ Support Palliat Care* 2018 doi: 10.1136/bmjspcare-2018-001595 [published Online First: 2018/09/27]
- 73. Lee AL, DeBest M, Koeniger-Donohue R, et al. The feasibility and acceptability of using virtual world technology for interprofessional education in palliative care: a mixed methods study. *Journal of interprofessional care* 2019:1-11. doi: 10.1080/13561820.2019.1643832 [published Online First: 2019/08/23]
- 74. Clabburn O, Groves KE, Jack B. Virtual learning environment ('Ivy Street') for palliative medicine education: student and facilitator evaluation. *BMJ* Supportive & Palliative Care 2020;10(3):318-23.
- 75. Liu L, Stroulia E, Nikolaidis I, et al. Smart homes and home health monitoring technologies for older adults: A systematic review. *Int J Med Inform* 2016;91:44-59. doi: 10.1016/j.ijmedinf.2016.04.007 [published Online First: 2016/05/18]
- 76. Pavic M, Klaas V, Theile G, et al. Mobile Health Technologies for Continuous Monitoring of Cancer Patients in Palliative Care Aiming to Predict Health

Status Deterioration: A Feasibility Study. *J Palliat Med* 2020;23(5):678-85. doi: 10.1089/jpm.2019.0342 [published Online First: 2019/12/25]

77. Pantelopoulos A, Bourbakis NG. A survey on wearable sensor-based systems for health monitoring and prognosis. *IEEE Transactions on Systems, Man, and Cybernetics, Part C (Applications and Reviews)* 2010;40(1):1-12.

78. Di Sanzo M, Cipolloni L, Borro M, et al. Clinical Applications of Personalized Medicine: A New Paradigm and Challenge. *Curr Pharm Biotechnol* 2017;18(3):194-203. doi: 10.2174/1389201018666170224105600 [published Online First: 2017/02/28]

- 79. Reid VL, McDonald R, Nwosu AC, et al. A systematically structured review of biomarkers of dying in cancer patients in the last months of life; an exploration of the biology of dying. *PloS one* 2017;12(4):e0175123.
- 80. Coyle S, Scott A, Nwosu AC, et al. Collecting biological material from palliative care patients in the last weeks of life: a feasibility study. *BMJ open* 2016;6(11):e011763. doi: 10.1136/bmjopen-2016-011763 [published Online First: 2017/02/12]
- 81. Kim J-E, Lee Y-J, Kwak M-H, et al. Metabolomics approach to serum biomarker for loperamide-induced constipation in SD rats. *Laboratory animal research* 2014;30(1):35.
- 82. Amgarth-Duff I, Hosie A, Caplan G, et al. Toward best practice methods for delirium biomarker studies: an international modified Delphi study. *International journal of geriatric psychiatry* 2020;35(7):737-48.
- 83. DeMarshall C, Oh E, Kheirkhah R, et al. Detection of early-stage Alzheimer's pathology using blood-based autoantibody biomarkers in elderly hip fracture repair patients. *PloS one* 2019;14(11):e0225178.

84. Fladvad T, Klepstad P, Langaas M, et al. Variability in UDP-

glucuronosyltransferase genes and morphine metabolism: observations from a cross-sectional multicenter study in advanced cancer patients with pain. *Pharmacogenetics and genomics* 2013;23(3):117-26. doi:

10.1097/FPC.0b013e32835ce485 [published Online First: 2013/01/02]

85. Barratt DT, Bandak B, Klepstad P, et al. Genetic, pathological and physiological determinants of transdermal fentanyl pharmacokinetics in 620 cancer patients of the EPOS study. *Pharmacogenetics and genomics* 2014;24(4):185-94. doi: 10.1097/fpc.000000000000022 [published Online First: 2014/01/29]

86. Klepstad P, Fladvad T, Skorpen F, et al. Influence from genetic variability on opioid use for cancer pain: a European genetic association study of 2294 cancer pain patients. *Pain* 2011;152(5):1139-45. doi:

10.1016/j.pain.2011.01.040 [published Online First: 2011/03/15]

- 87. DeSanto-Madeya S, Tjia J, Fitch C, et al. Feasibility and Acceptability of Digital Legacy-Making: An Innovative Story-Telling Intervention for Adults With Cancer. American Journal of Hospice and Palliative Medicine® 2020:1049909120971569.
- Taubert M, Watts G, Boland J, et al. Palliative social media. *BMJ Supportive & Palliative Care* 2014;4(1):13-18.
- 89. Clabburn O, Knighting K, Jack BA, et al. The use of digital legacies with people affected by motor neurone disease for continuing bonds: An interpretative phenomenological analysis study. *Palliative medicine* 2019;33(7):812-22.
- 90. Taubert M, Norris J, Edwards S, et al. Talk CPR-a technology project to improve communication in do not attempt cardiopulmonary resuscitation decisions in palliative illness. *BMC palliative care* 2018;17(1):1-12.

91. Coop H, Marlow C. Do we prepare patients for their digital legacy? A survey of palliative care professionals. *Palliat Med* 2019;33(1):114-15. doi:

10.1177/0269216318802748 [published Online First: 2018/09/28]

92. Calton B, Abedini N, Fratkin M. Telemedicine in the Time of Coronavirus. *J Pain Symptom Manage* 2020;60(1):e12-e14. doi:

10.1016/j.jpainsymman.2020.03.019 [published Online First: 2020/04/03]

- 93. Grewal US, Terauchi S, Beg MS. Telehealth and Palliative Care for Patients With Cancer: Implications of the COVID-19 Pandemic. *JMIR Cancer* 2020;6(2):e20288. doi: 10.2196/20288 [published Online First: 2020/10/14]
- 94. Ritchey KC, Foy A, McArdel E, et al. Reinventing palliative care delivery in the era of covid-19: how telemedicine can support end of life care. American Journal of Hospice and Palliative Medicine® 2020;37(11):992-97.
- 95. Bettini EA. COVID-19 Pandemic Restrictions and the Use of Technology for Pediatric Palliative Care in the Acute Care Setting. *J Hosp Palliat Nurs* 2020;22(6):432-34. doi: 10.1097/njh.000000000000694 [published Online First: 2020/09/15]
- 96. Chávarri-Guerra Y, Ramos-López WA, Covarrubias-Gómez A, et al. Providing Supportive and Palliative Care Using Telemedicine for Patients with Advanced Cancer During the COVID-19 Pandemic in Mexico. *The oncologist* 2020
- 97. Mackey RM, Yeow ME, Christensen AR, et al. Reconnecting: Strategies for Supporting Isolated Older Adults during COVID-19 through Tele-palliative Care. *Clinical gerontologist* 2020:1-8.

- 98. Harris DA, Archbald-Pannone L, Kaur J, et al. Rapid Telehealth-Centered Response to COVID-19 Outbreaks in Postacute and Long-Term Care Facilities. *Telemedicine and e-Health* 2021;27(1):102-06.
- 99. Flores S, Abrukin L, Jiang L, et al. Novel Use of Telepalliative Care in a New York City Emergency Department During the COVID-19 Pandemic. *The Journal of emergency medicine* 2020;59(5):714-16.
- 100. Lu Y, Xie D, Zhang X, et al. Management of intractable pain in patients with implanted spinal cord stimulation devices during the COVID-19 pandemic using a remote and wireless programming system. *Frontiers in neuroscience* 2020;14
- 101. Sansom-Daly UM, Bradford N. Grappling with the 'human'problem hiding behind the technology: Telehealth during and beyond COVID-19. *Psychooncology* 2020
- 102. Samara J, Liu W-M, Kroon W, et al. Telehealth Palliative Care Needs Rounds During a Pandemic. *The Journal for Nurse Practitioners* 2021
- 103. Chua IS, Jackson V, Kamdar M. Webside Manner during the COVID-19
  Pandemic: Maintaining Human Connection during Virtual Visits. *J Palliat Med* 2020;23(11):1507-09. doi: 10.1089/jpm.2020.0298 [published Online First: 2020/06/12]
- 104. Lally K, Kematick BS, Gorman D, et al. Rapid Conversion of a Palliative Care
  Outpatient Clinic to Telehealth. JCO Oncol Pract 2021;17(1):e62-e67. doi:
  10.1200/op.20.00557 [published Online First: 2020/12/12]
- 105. Crosby B, Hanchanale S, Stanley S, et al. Evaluating the use of video communication technology in a hospital specialist palliative care team during the COVID-19 pandemic. *AMRC Open Research* 2021;3(5):5.

- 106. Lal A, Bell G, Curseen K, et al. Teaching Telepalliative Care: An Elective Rotation for Medical Students during the COVID-19 Pandemic. *Journal of Palliative Medicine* 2020
- 107. Abel J, Taubert M. Coronavirus pandemic: compassionate communities and information technology. *BMJ Supportive & amp; Palliative Care* 2020;10(4):369-71. doi: 10.1136/bmjspcare-2020-002330
- 108. Mills J, Li C, Fullerton S, et al. Staying connected and informed: Online resources and virtual communities of practice supporting palliative care during the novel coronavirus pandemic. *Progress in Palliative Care* 2020;28(4):251-53. doi: 10.1080/09699260.2020.1759876
- 109. Niki K, Okamoto Y, Ueda M. Response to Wang et al., Virtual Reality as a Bridge in Palliative Care during COVID-19 (DOI: 10.1089/jpm.2020.0212). J Palliat Med 2020;23(7):892-94. doi: 10.1089/jpm.2020.0261 [published Online First: 2020/05/13]
- 110. Wang SS, Teo WZ, Teo WZ, et al. Virtual reality as a bridge in palliative care during COVID-19. *Journal of palliative medicine* 2020;23(6):756-56.
- 111. Etkind SN, Bone AE, Gomes B, et al. How many people will need palliative care in 2040? Past trends, future projections and implications for services. *BMC Medicine* 2017;15(1):102. doi: 10.1186/s12916-017-0860-2

# LEGENDS

Figure 1: Flow diagram to outline study process for identifying research priority areas


## Table 1: Demographics of study participants for the Delphi questionnaires and

### consensus meeting

## Table 2: Final list of priorities

## APPENDICES

- Future Today Institute 2018 Trends List.
- Technology in Palliative Care study scoping review.
- Delphi questionnaire (Google forms).
- Voting outcomes for consensus meeting.
- Descriptive Statistics and level of agreement for each 'priority area'.
- Summary of the networks used to invite palliative care professionals to participate.
- Interquartile Range to be used to guide the level of agreement for Delphi responses.
- Level of agreement for each 'priority area' following both Delphi rounds.
- Examples of technologies used in palliative care during the COVID19 pandemic.



Delphi Questionnaires

Review of selected trends for simplification and reduction N= 95

Trends included in Delphi questionnaire and consensus meeting 16 Healthcare significance but limited significance to palliative care

