

Title- Awareness regarding Anemia in Antenatal women attending tertiary care hospitals in South India

Author:

Mehul Sinha¹

1- Kasturba Medical College, Mangalore

Correspondence email- mehul666666@gmail.com

Conflict of interest- None

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

ABSTRACT

Lack of adequate nutrition during pregnancy is one of the major risk factors for the development of anemia which further leads to a high-risk pregnancy. Anemia in pregnancy accounts for one-fifth of maternal deaths and is a major factor responsible for low birth weight. The aim of the study was to evaluate the awareness about anemia among the antenatal women attending tertiary care hospitals in South India and also to find out the methods adopted by the women to prevent and correct their anemia. It was a cross-sectional type of study conducted on pregnant women who gave consent. A majority of the pregnant women were aware of the term anemia and the most common source of their knowledge was their treating physician, followed by ASHA/Anganwadi worker and friends/family. Half of the participants in the study were well aware that anemia was a problem related to blood and a majority knew that it is most common in pregnant women. Around half of the study population knew the correct value of minimum Hb level in a pregnant woman and even a lesser percentage of women knew about the consequence of severe anemia during delivery. Most of the pregnant women were including iron-rich foods normally in their diet. A majority of the respondents noticed a health benefit after taking the iron and folic acid tablets. Thus, there is a need to endorse the iron and folic acid (IFA) tablets and educate the women about the anemia in pregnancy so that the health benefits of the free supplementation program of the government percolates down to even the lower strata of the society.

Keywords: Pregnant women, Folic acid, anemia, nutritional status, tertiary care center

INTRODUCTION

Pregnancy is a normal physiological state of the body in which there is an increase in body's demands to cope up with the growth and development of fetus within mother's body. Therefore, maternal nutrition is an area of major concern which consequently affects the health of both the mother and the fetus. Lack of adequate nutrition during pregnancy is one of the major risk factors for the development of anemia which further leads to a high-risk pregnancy. Anemia in pregnancy accounts for one fifth of maternal deaths and is a major factor responsible for low birth weight. In India, 16% maternal deaths are attributed to anemia ⁽¹⁾. Dietary habits play an important role in Nutritional Iron Deficiency Anemia.

Administration of Iron and Folic acid tablets during pregnancy have improved the anemic states in pregnant women. India became the first developing country to take up a National Nutritional Anemia Prophylaxis Program (NNAP) to prevent anemia among pregnant women. NNAPP was initiated in 1970 during the fourth 5-year health plan with the aim of reducing the prevalence of anemia to 25%. The Government of India recommends a minimum dose of total 100 iron and folic acid tablets to be prescribed during pregnancy. ⁽²⁾ According to DLHS-4 (2012-2013 survey), only 47.3% of pregnant women consumed 100 or more iron and folic acid tablets in Karnataka. ⁽²⁾ This number further decreases in the district of Dakshin Kannada - According to NFHS-4, only 44.4% of mothers consumed iron and folic acid tablets during their pregnancy. ⁽²⁾ Although iron and folic acid supplementation is given free of cost in public healthcare systems, then also there is high prevalence of anemia in pregnant women. Etiological factors include lack of awareness about anemia, non-compliance and other socio demographic factors.

Unfavourable sociodemographic factors are the major barriers to the efforts in place for the prevention of anemia during pregnancy. Educating the women only will not produce any desirable change but increasing the degree of literacy of the family will definitely help to solve this problem. The educational status of the husbands and the women are equally important factors as it makes the couple receptive to the advice given by the health staff. Therefore, there is a need for dietary counselling and nutritional education in antenatal clinics to tackle the issue of anemia in pregnancy with missionary zeal, innovative approach, and evidence-based interventions.

While women frequently recognize symptoms of anemia, they do not know the clinical term for anemia. Half of women in all countries consider these symptoms to be a priority health concern that requires action and half do not. Those women who visit prenatal health services are often familiar with iron supplements, but commonly do not know why they are prescribed.

AIMS AND OBJECTIVES

- To evaluate the awareness about anemia among the antenatal women attending tertiary care hospitals in South India
- To find out the proportion of women having anemia
- To find out the methods adopted by the women to prevent and correct their anemia

MATERIALS AND METHODS

Background Information: Mangalore, a city in the coastal belt of the State of Karnataka, in India, has a population of about 6,23,841 as per 2011 census of India. The present study will be carried out in tertiary care hospitals of Mangalore, Karnataka, India. Known for its good medical facilities, patients come here from different parts of India. The literacy rate of Mangalore is about 82.2% according to 2001 census data which includes a high female literacy rate also.

Study design: Cross sectional study

Study type: Analytical study

Study settings: Tertiary care hospitals affiliated with Kasturba Medical College (KMC), MAHE, Mangalore – KMCH, Attavar and Lady Goschen Hospital.

Study population: Antenatal Women attending tertiary care hospitals affiliated with Kasturba Medical College, Mangalore

Inclusion criteria : Pregnant women who give consent.

Exclusion criteria :

- Pregnant women who didn't give consent.
- Patients who have been diagnosed with hemolytic anemia.

Study duration : 13th march - 31st march 2019

Sample size calculation :

With proportion being 35.8% ⁽¹¹⁾, 7% of absolute precision, 95% confidence level, the sample size comes to 205, using the following formula.

$$N = \frac{Z^2 pq}{d^2}$$

Where N is sample size, Z=1.96, p is proportion = 35.8, q is (1-p) = 64.2, d = 7%.

Sampling Strategy : Convenient, Non random sampling

Data Collection Methodology :

Institutional Ethics Committee (IEC) clearance will be obtained prior to commencement of the study. Permission will be obtained from the Medical Superintendent of the above mentioned hospitals. The data will be collected using a semi-structured questionnaire designed after literature review.

The questionnaire consists of 3 parts :

1. Patient Particulars
2. Questions to assess knowledge about anemia
3. Practices followed to prevent or control anemia

The patients will be selected by non-random sampling and will be explained the nature and the purpose of the study in a manner comprehensible by them. A written informed consent will be taken from those willing to participate.

Data Analysis: Data analysis will be done using descriptive statistics. The collected data will be entered in and analyzed using SPSS (Statistical Package for Social Sciences) version 25.0. If they are willing to participate in the study, a written informed consent (Annexure I) will be obtained and they will be interviewed which includes filling up of a semi structured questionnaire (Annexure II). Chi square test will be used to correlate the association between the factors and P value <0.05 will be considered statistically significant.

RESULTS

TABLE NO: 1. SOCIO DEMOGRAPHIC PROFILE OF STUDY SUBJECTS

Age	N=205	Percentage (%)
18-23	47	22.9
24-29	104	50.7
30-35	50	24.3
>36	4	1.9
Socioeconomic status		
Upper middle class	5	5.2
Lower middle class	52	52.2
Upper lower class	62	30.2
Lower class	83	40.4
Occupation of respondents		
	N=205	Percentage (%)
Housewife	158	77.07
Lecturer	9	4.3
Hospital setting environment	7	3.4
Semi-skilled	10	4.8
Shop/cleric/private/corporate	21	10.2
Educational status of respondents		
	N=205	Percentage (%)
Illiterate	44	21.4
Primary school	26	12.6
Secondary school	52	25.3
Senior secondary	45	21.9
Graduate	38	18.5
Type of family		
Nuclear	134	65.3
Joint	71	34.6

TABLE NO: 2. OBSTETRIC HISTORY OF ANTENATAL WOMEN

Obstetric status	N=205	Percentage
Primigravida	69	33.6
Multigravida	136	64.6
abortion	42	20.4
Gestational Age		
1 st Trimester	28	13.6
2 nd Trimester	32	15.6
3 rd Trimester	145	70.7

Trimester in which registered for ANC	N=205	Percentage (%)
1st trimester	200	97.5
2 nd Trimester	5	2.4
3 rd trimester	0	0
Age at Marriage		
17-20	44	21.4
21-24	102	49.7
25-28	45	21.9
29-32	10	4.8
>32	4	1.9
Complications during pregnancy		
	N=205	Percentage
Intrauterine death	7	3.4
Pre-eclampsia	1	0.4
C section	2	0.8
No complication	192	93.6
Hb status		
Moderate anemia (7-9.9gm%)	135	65.8
Mild anemia (10-10.9gm%)	32	15.6
Normal (>11gm%)	23	11.2

TABLE NO: 3. AWARENESS REGARDING ANEMIA

Awareness about the term anemia	N = 205	Percentage
Aware	139	67.8
Not aware	66	32.6
Source of information on anemia		
Treating physician	96	46.8
Anganwadi/ASHA worker	42	20.4
Internet/social media	11	5.3
Friends/family	31	15.1
College/school	19	9.2
Not applicable	6	2.9
Perception of Anemia	N=205	Percentage (%)
Natural state of pregnancy	64	31.2
Increased Body weight	16	7.8
Problem related to blood	113	55.1
Increase in Blood sugar	12	5.8
Who is at high risk of being an anemic?		
Male Adolescent	8	3.9
Pregnant women	165	81.6
Disabled patients	12	5.8
Diabetics	20	9.7
Perception of Anemia	N=205	Percentage (%)
Natural state of pregnancy	64	31.2
Increased Body weight	16	7.8
Problem related to blood	113	55.1
Increase in Blood sugar	12	5.8
What should the minimum Hb level be in a pregnant woman?	N=205	Percentage (%)
Correct response	65	31.7
Wrong response	52	25.3
Don't know	86	42.5
What is the consequence of anemia during delivery?		
Death of baby	30	14.6
Difficulty in delivery	73	35.6
Post-partum haemorrhage	56	27.3
Don't know	46	22.4
Does pregnancy create a greater demand of Iron?		
Yes	164	80
Does repeated pregnancy causes anemia?		
Yes	125	60.9
Are citrus fruits a rich source of vitamin C which prevents anemia?		
Yes	145	70.7

TABLE NO: 4. CAUSES, SIGNS AND SYMPTOMS OF ANEMIA

Is anemia communicable disease?	N=205	Percentage (%)
Yes	64	31.2
Can infections cause anemia?		
Yes	133	65.8
Can consumption of contaminated food and water cause anemia?		
Yes	113	55.1
Can heavy blood loss due to menstruation cause anemia?		
Yes	149	72.6
How would you identify an anemic person?		
She appears pale	129	62.9
She suffers from seizures	18	8.7
Red face with flushing	40	19.5
Bluish discoloration of extremities	18	8.7
Do anemic patients become breathless easily?		
Yes	110	54.4
Do anemic patients normally experience fatigue and tiredness?		
Yes	177	86.3
Excessive vomiting and diarrhea are signs of anemia?		
Yes	111	54.1
Burning urination with increased frequency is a sign of anemia?		
Yes	81	39.5

TABLE NO: 5. PREVENTION OF ANEMIA

Can anemia be prevented?	N=205	Percentage (%)
Yes	166	89.5
No	39	10.5
How do you prevent/ correct anemia?		
By taking Iron and folic acid tablets and iron rich diet	117	57
By taking calcium supplements	21	10.2
By taking insulin injection	16	7.8
Not sure / Don't know	51	24.8
Can regular exercise prevent anemia?		
Yes	138	67.3
No	67	32.7
Name few iron rich foods		
Green leafy vegetables	61	29.7
Dry fruits	14	6.8
Non-Vegetarian	28	13.6
Milk	10	4.8
Fruits	97	47.3
Cereals and Pulses	15	7.3
Do you normally include the above food items in your diet?		
Yes	155	75.6
Have you taken deworming tablets previously?		
Yes	99	48.2
Did you change your normal dietary pattern during pregnancy?		
Yes	67	32.6
If so, what were the changes?		
Decreased intake of carbohydrates and fats	8	3.9
Increased vitamin and iron rich food	13	6.3
Increased protein rich food	10	4.8
Increased fruits and Vegetables	36	17.5
No change	138	67.3

TABLE NO: 6. PRACTICES FOLLOWED TO PREVENT OR CORRECT ANEMIA

Respondents noticed a health benefit's of taking the tablets and therefore continued taking them	N=205	Percentage (%)
Noticed a benefit	157	76.5
Did not notice a benefit	48	23.2
Respondents increased the intake of Iron rich food/ fortified foods to control anemia		
Increased	163	79.5
Did not increase	42	20.5
Respondents took their Iron and folic acid (IFA) supplements as per prescription (regularly)		
Yes	165	80.4
Respondents started including more non-veg food in their diet following a fall in their Hb levels		
Yes	103	50.2
Respondents continued to take the preventive supplementation regimen after the completion of their 3 months of therapeutic supplementation		
Continued	130	63.4
Did not continue	75	36.6
Respondents came for regular follow ups as instructed by their physician		
Yes	184	89.7

No	6	10.3
----	---	------

DISCUSSION

Most of the women who participated in the study belonged to the lower class of economic status with housewife as their occupation. Majority of the women who participated in the study had basic level of education and only one –fifth of them were illiterate, living in a nuclear family. Most of the antenatal women were multi gravida who were in their 3rd trimester of pregnancy. Our study group included most of the women who were moderately anemic. A majority of the pregnant women were aware of the term anemia and the most common source of their knowledge was their treating physician, followed by ASHA/Anganwadi worker and friends/family. Half of the participants in the study were well aware that anemia was a problem related to blood and a majority knew that it is most common in pregnant women. Around half of the study population knew the correct value of minimum Hb level in a pregnant woman and even a lesser percentage of women knew about the consequence of severe anemia during delivery.

Most of the respondents knew that pregnancy creates a demand of iron. Almost two-thirds women knew that anemia is a non-communicable disease which is caused due to heavy blood loss during menstruation. Majority of the patients knew the clinical signs and symptoms of anemia. Almost more than half the respondents knew that anemia can be prevented by increasing the intake of iron rich foods like green leafy vegetables and fruits in their diet.

Half of the women thought that anemia can also be prevented by exercise which is a myth. Most of the pregnant women were including iron rich foods normally in their diet. A majority of the respondents noticed a health benefit after taking the Iron and folic acid tablets. Half of the pregnant women also started including more nonveg food items in their diet after a fall in their Hb levels. Most of the women took their IFA tablets very regularly to prevent anemia. More than half the population continued the preventive dose of IFA tablets even after the prophylactic dose of 3 months due to its health benefit. Almost all of the Antenatal women in their 3rd trimester who were registered under the ANC program visited their physician regularly for follow ups. Therefore, there is a positive correlation between the education level of women and the awareness regarding anemia in Antenatal women.

CONCLUSION

While women frequently recognize symptoms of anemia, they do not know the clinical term for anemia. Half of women in all countries consider these symptoms to be a priority health concern that requires action and half do not. Those women who visit prenatal health services regularly are often familiar with iron supplements and their health benefits. Thus, there is a need to endorse the IFA tablets and educate the women about the anemia in pregnancy so that the health benefits of the free supplementation program of the government percolates down to even the lower strata of the society.

REFERENCE

- 1 - Government of India (sample registration system). Maternal mortality in India: 1997-2003, Trends, causes and risk factors. Register General India, New Delhi in collaboration with Centre for Global Health Research University of Toronto, Canada. -
- 2 - Agarwal DK, Agarwal KN, Roychaudhary S. Targets in national anaemia prophylaxis program for pregnant women. Indian Pediatric Journal 1988;25:319-22
- 3 - Lokare PO, Karanjekar VD, Gattani PL, Kulkarni AP. A study of prevalence of anemia and sociodemographic factors associated with anemia among pregnant women in Aurangabad city, India. Ann Nigerian Med 2012;6:30-4
- 4-Taye B, Abeje G, Mekonen A. Factors associated with compliance of prenatal iron folate supplementation among women in Mecha district, Western Amhara: a cross-sectional study. Pan African Medical Journal. 2015;20(1).
- 5-Stoltzfus RJ, Dreyfuss ML. Guidelines for the use of iron supplements to prevent and treat iron deficiency anemia. Washington, DC: Ilsi Press; 1998.
- 6-Gedefaw L, Ayele A, Asres Y, Mossie A. Anaemia and associated factors among pregnant women attending antenatal care clinic in Walayita Sodo town, Southern Ethiopia. Ethiopian journal of health sciences. 2015;25(2):155-64.
- 7- Brabin JB, Hakimi M, Pelletier D. An Analysis of Anemia and Pregnancy-Related Maternal Mortality. The Journal of Nutrition 2001;131(2):604-15
- 8- Shitie D, Zewde T, Molla Y. Anemia and other hematological profiles of pregnant women attending antenatal care in Debre Berhan Referral Hospital, North Shoa, Ethiopia. BMC Research Notes 2018;11:704
- 9- Vijayaraghavan K, Brahmam GNV, Nair KM et al. Evaluation of national nutritional anemia prophylaxis program. Indian J Pediatr 1990; 57(2): 183-9.
- 10- Seck BC, Jackson RT. Determinants of compliance with iron supplementation among pregnant women in Senegal. Public health nutrition. 2008 Jun;11(6):596-605.

WITHDRAWN
see manuscript DOI for details