

Developing a co-produced, systems-informed, sexually transmitted infection contact tracing intervention for gay and bisexual men who have sex with men and their ‘one-off’ sexual partners.

Paul Flowers* <http://orcid.org/0000-0001-6239-5616> Email: paul.flowers@strath.ac.uk

Sarah Lasoye <https://orcid.org/0000-0002-2408-3555> Email: s.lasoye@ucl.ac.uk

Jean McQueen <https://orcid.org/0000-0001-5330-3564> Email: Jean.McQueen@gcu.ac.uk

Melvina Woode Owusu : Email: m.woode-owusu@ucl.ac.uk

Merle Symonds <https://orcid.org/0000-0001-5793-8946> Email: merle.symonds2@nhs.net

Claudia Estcourt <https://orcid.org/0000-0001-5523-5630> Email: Claudia.Estcourt@gcu.ac.uk

Key words: contact tracing, complex intervention, sexually transmitted infections, intervention development

This work presents independent research funded by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research Programme (reference number RP-PG-0614-20009).

The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. The funders had no role in study design, collection, management, analysis and interpretation of data; writing of the report and the decision to submit the report for publication.

RP-PG-0614-20009

Corresponding author: Paul Flowers

Postal address: Professor of Health Change, School of Psychological Sciences & Health, Graham Hills Building, 40 George Street, Glasgow, G1 1QE Email: Paul.Flowers@strath.ac.uk

Telephone: +44 (0)141 548 2700

Fax: +44 (0)141 548 2721

Contributions to date:

Paul Flowers: funding acquisition, methodological lead, data collection and analysis, contributions to first draft,

Sarah Lasoye: contributions to first draft

Jean McQueen: data collection and analysis, contributions to second draft

Melvina Woode Owusu : Supporting data collection, contributions to first draft

Merle Symonds: funding acquisition, contributions to second draft

Claudia Estcourt: principal investigator, funding acquisition, contributions to first draft

Developing a co-produced, systems-informed, sexually transmitted infection contact tracing intervention for gay and bisexual men who have sex with men and their ‘one-off’ sexual partners

Abstract

Objective Gay and bisexual men who have sex with men (GBMSM) bear a disproportionate burden of sexually transmitted infections (STIs). Most STIs are asymptomatic and people infected won't know to seek care unless they are told about their exposure. Contact tracing, is the process of identifying and contacting sex partners of people with STIs for testing and treatment. Contact tracing is sometimes particularly challenging amongst GBMSM because of the kinds of sexual relationships which GBMSM enjoy. These include one-off partners who are particularly important for transmission dynamics as they contribute disproportionately to onwards transmission. The effectiveness of contact tracing interventions within sexual health are patterned by sexual-partner type. Contact tracing and management for ‘one-off’ partners is an on-going public health challenge. Low motivation amongst index patients, high resource burden on health care professionals and problems with contactability are key barriers to contact tracing. Using insights from complex adaptive systems thinking and behavioural science, we sought to develop an intervention which addressed both the upstream and down-stream determinants of contact tracing and change the system in which many inter-dependent contact tracing behaviours are embedded.

Setting UK community-recruited GBMSM, stakeholders, sexual hcp, dating app providers **Method** Using the MRC complex intervention framework and insights from the INDEX study, a three-phase intervention development process was adopted to specify intervention content. Phase one consisted of an inter-professional and community-member stakeholder event (n=45) where small mixed groups engaged in exploratory systems-mapping and the identification of ‘hot spots’ for future intervention. Phase two used a series of focus groups with GBMSM (n=28) and interviews with representatives from key dating app providers (DAPs) (n=3) to further develop intervention ideas using the theoretical domains framework, the behaviour change wheel and the behaviour change technique taxonomy. In Phase 3 we again worked with key stakeholders expert health care professionals (HCPs) (n=5) and key workers from community-based organisations (CBOs) (n=6) to hone the intervention ideas and develop programme theory using the APEASE criteria.

Results The co-produced intervention levers change simultaneously across the system within which contact tracing is embedded. Multiple change-agents (i.e., GBMSM, CBOs, HCPs) work together, sharing an overall vision to improve sexual health through contact tracing. Each make relatively modest changes that over time, synergistically combine to produce a range of multiple positively-reinforcing feedback loops to engender sustainable change around contact tracing. Agreed intervention elements included: a co-ordinated, co-produced mass and social media intervention to tip cultural norms and beliefs of GBMSM towards enabling more contact tracing and to challenge enduring STI- and sex-related stigmas; complementary CBO-co-ordinated, peer-led work to also focus on reducing STI stigma and enabling more contact tracing between one-off partners; priming GBMSM at the point of STI diagnosis to prepare for contact tracing interactions and reduce HCP and sexual health-service burden; changes to SHS environments and HCP-led interactions to systematically endorse contact tracing; changing national audits and monitoring systems to directly address one-off partner targets; delivering bespoke training to HCPs and CBO staff on one-off partners and the social and cultural context of GBMSM; DAPs active involvement in mass and social media promoting appropriate contact tracing messaging.

Conclusion Our combination of multiple data sources, theoretical perspectives and diverse stakeholders have enabled us to develop an expansive, complex intervention that is firmly based in the

priorities of those it will affect, and which has a solid theoretical foundation. Future work will assess if and how it will be possible to evaluate it. The resulting intervention is profoundly different than other ways of enhancing contact tracing, as it simultaneously addresses multiple, multi-levelled, upstream and social determinants of contact tracing.

Developing a co-produced, systems-informed, sexually transmitted infection contact tracing intervention for gay and bisexual men who have sex with men and their ‘one-off’ sexual partners

Introduction

Contact tracing and the management of those potentially exposed to infectious disease has long been a cornerstone of public health. Its primary logic is simple: once having identified someone with an infectious disease *focus* further efforts to reduce onwards transmission where transmission is most likely to have occurred. Depending on the nature of the infectious disease, its transmission routes, and our ability to treat it, further focussed efforts may include testing, treatment, a range of behaviours to reduce onwards transmission (e.g., testing, self-isolation and physical distancing) and of course more contact tracing.

Across a range of infectious diseases, the effectiveness of contact tracing efforts remain largely unknown (Fitzgerald and Bell, 1998). Because of the ubiquitous nature of contact tracing as a routine intervention in public health, evaluations of its effectiveness are relatively rare (see however Saurabh and Prateek, 2017; Swanson *et al.*, 2018; Juneau *et al.*, 2020) although many modelling studies do exist (Müller and Kretzschmar, 2021). However, the COVID-19 pandemic has fuelled a recent explosion of interest in contact tracing as its urgent and rapid implementation quickly highlighted major problems in the COVID-19 context. Within England, for example, contact tracing has been particularly problematic (Department of Health and Social Care, 2020; Limb, 2020). We would argue one key problematic aspects of contact tracing *per se*, as has been seen with contact tracing COVID-19 in particular, relates to the lack of attention to its psychosocial and sociocultural aspects. On a superficial level, contact tracing can of-course be thought of simply as the focusing of public health efforts on the contacts of an index patient to reduce onwards transmission. However, this somewhat deracinated perspective overlooks the complexity of the social worlds in which both the index patient and their contacts are embedded. Stepping away from the specific interactions between an index patient and their contacts opens new opportunities for intervention. A focus on the patterning of factors, which drive interactions within communities and networks that are associated with onwards transmission represents a complementary and viable alternative. In contact tracing interventions across varied infectious diseases, little attention has been paid to health inequities and the need to target and tailor contact tracing interventions to particular communities and their life circumstances (Mathevet *et al.*, 2021). Equally, until recently, very little research has explored the factors shaping particular communities’ engagement with contact tracing. However, a recent rapid review highlights major barriers and facilitators to engaging in contact tracing across a range of infectious diseases. It suggests key barriers relate to: mistrust and apprehension, fear of stigmatisation, specific challenges relating to mode of its delivery (i.e., digital vs otherwise) and privacy concerns (Megnin-Viggars *et al.*, 2020). In contrast, the rapid review notes that key facilitators of effective contact tracing interventions relate to the co-production of contact tracing systems, perceptions of collective responsibility, perceived personal benefit, and perceptions of the contact tracing intervention as efficient, rigorous and reliable.

It follows that if we want to enhance contact tracing interventions for infectious diseases *per se* and for STIs in particular, we need to systematically engage much more with their meaning and implementation, and directly address their intersection with the lives and social worlds of those they

directly affect. This paper pursues this goal and seeks to begin the process of developing an intervention to reduce onwards transmission of STIs by improving contact tracing.

Sexually transmitted infections, GBMSM and contact tracing

Globally, more than one million sexually transmitted infections (STIs) are acquired each day, and result in serious negative health impacts (Rowley *et al.*, 2019). Reducing the prevalence and rates of transmission of STIs among GBMSM has become a key area of focus for public health. GBMSM bear a particular burden of sexual ill-health. It is highly likely that the on-going, ever-transforming, social worlds of GBMSM have been shaped by the cumulative effects of homophobia, heterosexism and a lack of a salutogenic intergenerational culture. Unlike other communities in which the rhythm and cycle of childbirth, parenting, grandparenting and extended families and local communities all enable intergenerational culture to be shared across time, for GBMSM it is primarily, although not exclusively, through sex that culture is sustained. Again, although it can never be empirically demonstrated, these factors may be associated with the on-going presence of hyper sexualised subcultures amongst GBMSM. These subcultures are characterised by high numbers of disassortative sexual mixing which in turn are associated with higher risks of STI transmission. Arguably, increases in the use of digitally mediated sexual mixing has further contributed to these issues. Although some of the most profound behaviour change ever to have been recorded was seen in these communities (Stall *et al.*, 2005 in response to the impact of HIV/AIDS), in recent years the incidence of STIs other than HIV has been increasing. Within the UK, GBMSM account for a disproportionate number of syphilis and chlamydia diagnoses (*Public Health England 2019*; Health Protection Scotland 2019). Addressing the rise of STI transmission among GBMSM is of increasing importance, especially given growing evidence of emerging antimicrobial resistance in key pathogens *Neisseria gonorrhoea* and *Mycoplasma genitalium* (Wi *et al.*, 2017). To address this complex situation WHO have set a target to reduce the incidence of gonorrhoea by 90% globally by 2021 (World Health Organisation, 2016). They recommend combining *education, behaviour change, development of vaccines, new medications and partner management to notify and treat the sex partners of individuals diagnosed with STIs*. Partner management is a term used to describe contact tracing within the sexual health field. It is a key component of STI control by breaking chains of transmission. Contact tracing for GBMSM remains poorly understood, and effective approaches specifically for one-off sex partners in particular are largely unknown (Wang *et al.*, 2016).

Approach to intervention development for contact tracing amongst GBMSM and their one-off partners for STIs

As part of the LUSTRUM programme of research which aims to improve contact tracing interventions for STIs & HIV we have used psychosocial and social insights to optimise and then trial a contact tracing intervention known as accelerated partner therapy (APT, see for example, (Estcourt *et al.*, 2020; Pothoulaki *et al.*, 2020). In typically contact tracing in the UK a health care professional asks the person with the STI (known as the 'index' patient) to 1) think back to all the people they have had sex with in the last 3 months, 2) inform those sex partners that they need to get tested and treated (at the sexual health clinic). Alternatively, the health care professional offers to inform the sex partners for the index patient, usually without disclosing the index patient's identity. In relation to LUSTRUM, and APT where the index patient takes the testing and treatment to their partners, we theorised that this particular contact tracing intervention would work well when index patients and their partners are in types of relationship where sex is likely to occur again, and affective ties motivate sexual partners to engage in contact tracing (Estcourt *et al.*, 2021). APT, for example, even harnesses people's on-going contact to actually deliver testing and

treatment directly. In contrast, and perhaps more akin to the affective elements of much contact tracing within the recent COVID-19 pandemic, when sex is unlikely to occur again, and when there is no affect to motivate contact tracing and enable further focussed public health efforts, contact tracing initiatives are much harder to gain traction. Actively being part of contact tracing is only associated with costs (e.g., self-isolation in COVID). For GBMSM and their one-off partners the costs of engaging in contact tracing with one-off partners stem the lives and social worlds of those affected, for example, cultural factors such as the on-going negative impacts of homophobia and heterosexism which in turn shape STI-related stigma and negative attitudes towards sex (Flowers et al., 2021). At the individual-level of particular GBMSM, issues such as reputational damage and perceptions of culpability and blame are also important considerations (Flowers, Duncan and Frankis, 2000; Flowers et al., 2021). Equally, from the perspectives of those who provide health services, it can be hard to justify the considerable resources required to achieve good contact tracing outcomes in harder to reach contacts, despite them contributing disproportionately to community transmission.

Given the importance of one-off sexual contacts to onward transmission and the difficulties in achieving good PN outcomes in this group we did an intervention development study that aims to improve contact tracing amongst GBMSM and their 'one-off' partners. Particularly mindful of the psychosocial and sociocultural factors that shape contact tracing *per se* and our insights into the specifics of contact tracing with 'one-off partners', we were keen to use the best available approaches to intervention development. Recent insights into the science of intervention development from the INDEX study (O'Cathain *et al.*, 2019) and the updated MRC Complex Intervention guidance (Medical Research Council, 2019) highlight the centrality of working with stakeholders from the start of the intervention development process. They also highlight the need to theorise interventions and their causal mechanisms in relation to the specific contexts in which they will be delivered. Recent studies in the science of intervention development also demand more focussed attention on long term sustainability and attempts to achieve intervention implementability early on within the intervention development process. In this way, we can avoid the oft reported 'implementation gap' associated with apparently successful interventions, showing promise within trials, later failing to translate into routine practice (Murray *et al.*, 2010; May, 2013).

In relation to using theory within intervention development we used a combination of theoretical approaches. Complex adaptive systems (CAS) perspectives are heralded as being particularly useful approaches to seemingly intractable public health problems (Wilson, Holt and Greenhalgh, 2001; Rutter *et al.*, 2017). They are particularly good at enabling an overview of a problem and generating *novel* solutions to potential interventions that are not limited by our typical constrained and linear understanding of cause and effect. They encourage us to think wholistically and systemically and deal with complexity head on. These approaches focus our attention on ways to intervene which are not overly simplified and instead highlight the need to address upstream (i.e., distal) and often multi-faceted causal influences (e.g. interdependent and multi-levelled drivers which shape system behaviour in ways that are challenging to reduce). As such we sought to use these systems perspectives at the start of our intervention development process. However, we would argue that although CAS perspectives are useful for imagining new ways of intervening, they are far less useful for actually operationalising the intervention ideas they can generate. Therefore, we also drew upon a range of theories and approaches from implementation science that assist with the clear specification and operationalisation of intervention ideas. (for more details see the intervention development protocol (McQueen *et al.*, 2020). In this way we drew upon a range of other theoretical frameworks that focus on implementation *per se* and ways of changing implementation behaviour.

Aim: To develop a co-produced, systems-informed, multi-level, contact tracing intervention for GBMSM with bacterial STIs and their “one-off” partners.

Research Questions

1. What did diverse stakeholders tell us about where best to intervene within the system that drives contact tracing?
2. What can we learn through behaviourally informed qualitative work to suggest particular evidence-based and theoretically-informed future intervention content
3. What is the final suggested intervention content after further stakeholder engagement?

Methods

A full account of the design and details of each phase of the research is detailed within the initial protocol (McQueen *et al.*, 2020). Here we provide a brief overview.

Design

Overall a three-phase design was used (See Figure 1). The first phase consisted of a major stakeholder event, an appraisal of the literature, and discussions with experts in the field. Initial ideas for intervention components generated at this phase were explored and developed further within Phase 2 and Phase 3.

Phase 1 *What did diverse stakeholders tell us about where best to intervene within the system that drives contact tracing?*

The stakeholder event aimed to work with important stakeholders to map the system which shapes contact tracing and begin to identify potential areas and methods of future intervention.

Data collection: Within the stakeholder workshop a series of small mixed groups were convened and tightly facilitated to ensure the problem of contact tracing for one-off partners was seen holistically and explored from multiple vantage points simultaneously. Small groups consisted of GBMSM, those working in CBOs, sexual health care professionals, service commissioners, sexual health researchers and those working in the dating app industry. Topics of discussion included the overall system which drives contact tracing, and an exploration of the full range of key organisations, communities, professionals, individuals, social processes, dynamics, and interactions involved and connected within the system. Throughout, in the context of GBMSM and their one-off partners, participants were encouraged to consider what currently worked to facilitate contact tracing and conversely what did not work well and why. Participants were also asked to think expansively and creatively to consider if improvements to contact tracing for GBMSM and their one-off partners were possible, and if so where and with whom this might take place and why they thought this might be important. Equally participants were asked to note where and why improvements were unlikely to ever be possible. Discussions were not audio recorded. Data was collected in the forms of visualisations from participants during systems-mapping and detailed notes were collected by scribes during facilitated group-discussion

Data Analysis: The stakeholder workshop data analysis focussed on systematically appraising the visualisations, detailed notes and sticky labels. PF and JMcQ considered *how* participants mapped the system shaping current and potential contact tracing and extracted a series of heterogeneous potential ‘hot spots’ where future intervention could usefully focus. An informal rapid review of the available published literature was also conducted at this time to assess the nature and relative success of existing interventions in this field. These ideas were examined in relation to workshop insights. PF and JMcQ began the process of considering potentially useful intervention ideas for future interventions to improve PN with MSM and their one-off partners. Through presentations and discussions with the wider team they were able to formulate working hypotheses about potentially useful intervention ideas. In this way a very basic and provisional programme theory was being formulated.

Phase 2 *What was learned through behaviourally informed qualitative work to suggest particular evidence-based and theoretically-informed future intervention content?*

We further explored the nascent intervention ideas with samples of people who would be affected by the intervention. Analytic tools from implementation science were used to understand barriers and facilitators to the implementation of the emerging intervention ideas developed from Phase 1 and to specify ways of optimising these ideas to secure an acceptable, feasible, sustainable future intervention.

Data collection: Focus groups with GBMSM, health care professionals (HCPs) and Dating App providers (DAPs) were planned to further develop initial intervention ideas. This sampling strategy and the topic guides for both the interview and focus group participants were informed by the initial intervention ideas generated by stakeholders. Participants were asked to discuss the emerging intervention ideas, whether and how they could be implemented, and the barriers and facilitators to implementation. All interview and focus group data were digitally recorded and transcribed.

Data Analysis: The data were analysed firstly to assess broad acceptability and whether participants thought the implementation of the intervention idea was actually feasible. Subsequently deductive thematic analysis was then undertaken to detail barriers and facilitators to implementing the intervention ideas deemed acceptable and amenable to change. Further data analysis then consisted of using the Theoretical Domains Framework (TDF) to theorise important barriers to the uptake or delivery of intervention ideas. Then the Behaviour Change Wheel (BCW) approach, incorporating the Behaviour Change Technique Taxonomy (BCTT), was used to specify potential intervention content in granular detail. Analyses were led by JMcQ and audited by GV and PF - both trained within these approaches (Michie, van Stralen and West, 2011; Michie *et al.*, 2013; Michie, Atkins and West, 2014; Atkins *et al.*, 2017).

Phase 3 *What is the final suggested intervention content after further stakeholder engagement?*

Although our original aim had been to work with a similarly diverse range of interprofessional stakeholders that had contributed to Phase 1, this was no longer possible because of the impact of COVID 19 which precluded the conducting of research which involved HCPs as participants.

Data collection: Instead, the project relied on working with staff from community-based organisations (CBOs) and working with sexual health experts from within the research team itself (primarily HCPs). This phase involved a series of four virtual workshops where participants were presented with both an overall high-level account of the emerging intervention ideas, as well as specific and granular detail generated during Phase 2. Phase 3 participants were asked to consider specific intervention ideas in relation to the APEASE criteria (APEASE=Affordability, Practicability, Effectiveness and cost-effectiveness, Acceptability, Side-effects and safety, Equity) and comment on whether content should be kept, modified or abandoned. All data was digitally recorded and transcribed.

Data Analysis: JMcQ and PF appraised data collected within these virtual workshops and iteratively worked with the data to identify content from the developing intervention which could be kept, abandoned or modified. Colour coding (green, amber, red) was then applied to the detailed intervention content outlined in Phase 2 to result in a final list of intervention content ([Appendix 1](#)).




Figure 1: Diagram presenting the components and process of each phase of the intervention development study.

Results

Phase 1: *What did diverse stakeholders tell us about where best to intervene within the system that drives contact tracing?*

In November 2019 the stakeholder event recruited 45 diverse participants representing a broad spectrum of agents within the system that drives contact tracing. These included GBMSM with no professional interest in the topic (20%), public health / sexual health service commissioners (17%), representatives of community-based organisations (9%), Health advisors and nurses (17%), sexual health researchers (11%), medical consultants from sexual health (20%) and representatives from DAPS (6%). A substantial minority of the stakeholders were both professionals and GBMSM.

Together, stakeholders highlighted how drivers of effective contact tracing were complex and distributed across a complex system. Figure 2 provides a schematic overview of the systemic drivers effecting contact tracing, collating stakeholder input from the initial engagement event. The drivers of contact tracing are multi-levelled and considerably inter-dependent. There is a clear balance of connected drivers that both enable and constrain contact tracing.




Figure 2 Schematic overview of the diverse systemic drivers effecting contact tracing, including pathogenic (red) and salutogenic (green) influences collated from the initial stakeholder event.

Stakeholders described how many critically important drivers of effective contact tracing tended to be distal to the actual act of notifying partners (e.g., the legacy of inadequate sex and relationships

education). Current contact tracing approaches with their primary focus on proximal determinants (e.g, using index patient affect to motivate patient referral) were understood to be necessarily limited when applied to one-off partners (e.g., practical and resources-related barriers) particularly within the age of digital dating apps (e.g., blocking)

Table 1 below summarises the findings of the stakeholder workshop and what we learned about the drivers of contact tracing.. It should be noted that, when compared to Figure 2, the tabularisation of these issues minimises the sense of how interlinked and interconnected the drivers of contact tracing are. A full narrative account of the drivers is available within supplementary file A.

Table 1- The multi-levelled and systematic drivers of contact tracing

The distal drivers of contact tracing
<i>Negative drivers (factors limiting contact tracing with one-off partners)</i>
The socioeconomic context of sexual health service provision
Overstretched sexual health services
The business models of dating apps
Elements of contemporary and historical culture
The on-going impact of heterosexism
The on-going impact of homophobia
History of pathologizing and blaming GBMSM
Legacy of inadequate sex and relationship education
High levels of sex and STI-related stigma
<i>Positive drivers (factors facilitating contact tracing with one-off partners)</i>
A distributed approach to improving drivers of contact tracing that did not over burden any single stakeholder
The agency and involvement of GBMSM throughout
Collective and co-ordinated efforts across the wider system
Saturate the system with clear signals about cultural value of contact tracing
Working with communities of GBMSM to enhance existing salutogenic aspects
Use existing assets: presence and history of existing norms about contact tracing, community resilience in relation to HIV
Harness peer interactions to drive improved contact tracing
Use all available infrastructures (websites, dating apps, broad range of health services)
Compensatory sexual health education
New cultural norms about contact tracing
New cultural expectations about contact sharing
Co-produce mass and social media intervention elements to change GBMSM cultures and norms
Ensure dating apps promote and endorse norms to facilitate contact tracing
Change dating app functionality and features
The proximal drivers of contact tracing
<i>Negative drivers (factors limiting contact tracing with one-off partners)</i>
Sexual health services experienced as unwelcoming to GBMSM
Interactions between HCPs and GBMSM experienced as judgemental and unhelpful
<i>Positive drivers (factors facilitating contact tracing with one-off partners)</i>
Opportunities for HCPs to reflect on current and optimal practice
Data driven approaches to understand current success and failure
Opportunities to share good practice within and across services
Training to ensure HCP cultural competence in relation to GBMSM
Scripts for HCPs to motivate GBMSM to engage in contact tracing

Work with GBMSM to anticipate and prepare for contact tracing interactions within sexual health services
Enhanced use of digital technology within contact tracing interactions (smart phones, dating apps)
Work with key people organising sex parties, sex clubs for GBMSM
Work around contact tracing within public sex environments

The selection of intervention ideas

Using the APEASE criteria (i.e., Affordability, Practicability, Effectiveness/cost-effectiveness, Acceptability, Side-effects/safety, Equity) and consideration of which elements of the system were actually amenable to change, Phase 1 ended with an appraisal and selection of key areas to explore in greater depth and detail during Phase 2 (see Figure 3 for an overview). Our stakeholders' system mapping had highlighted that future interventions must be *multi-levelled* (i.e. addressing both distal and proximal determinants of contact tracing). It also suggested that they must be *multi-agent* in order to share the burden of activity across the whole system whilst simultaneously normalising contact tracing in new ways. These core principles have the potential of realising sustainable change by re-organising the system itself to in turn improve contact tracing. These key principles were taken forwards to the subsequent phases.

Although acknowledging their fundamental importance in shaping contact tracing we chose not to further examine the following because they were not amenable to change within the intervention being developed here: inadequate sex and relationships education, the need for compensatory lifelong health and well-being education for GBMSM to foster better care of the self and the other, focussed efforts to reduce heterosexism and homophobia, attempts to build the resilience of GBMSM, the business models of the dating app industry, the economic drivers of health services challenges that stretch sexual health services. Although we had chosen to remove these elements as they were not particularly amenable to change, we were also mindful of the potential of these elements to re-emerge within Phase 2 and Phase 3 as we learned more from our participants and stakeholders.

We chose to focus on how communities of GBMSM, dating apps providers and sexual health services could work together in new ways and which complementary and co-ordinated activities could mutually reinforce each other. Such activities, included, targeted mass and social media with impactful messaging that changed aspects of gay culture that constrained contact tracing yet celebrated aspects that enabled it, dating app endorsement of contact tracing and hosting of media messaging, changing app functionality to enable GBMSM to take part in contact tracing, changing peer interactions to exact social influence echoing the key messages of targeted mass and social media, enhancing the ways HCPs worked with GBSMS though using technology within contact tracing interactions, through providing consistent and multiple messages through interactions and changes to the service environment to make contact tracing more salient, through providing culturally competent services able to support GBMSM and their contact tracing with one-off partners, through using motivational and culturally competent scripts to enable contact tracing with one-off partners and through using data driven approaches to enable services to learn more about their relative success and failures in contact tracing through (e.g. audit, monitoring, service reviews).




Figure 3 Selection of key areas to examine further within Phase 2

Phase 2 *What was learned through behaviourally informed qualitative work to suggest particular evidence-based and theoretically-informed future intervention content?*

The selected preliminary intervention components from Phase 1 were re-considered and further developed through a series of focus group discussions and interviews with key stakeholders. Critically, because of the impact of COVID-19 on HCP and NHS staff it was not possible to collect data from diverse HCPs as had been intended. As such the proposed intervention ideas generated in Phase 1 did not benefit from the critical appraisal of HCPs. For some of the intervention ideas relating to the SHS this was a particular loss. Developing new ways of working for sexual health clinics, for example, or addressing issues of cultural competency, harnessing insights into current good practice, were unable to be further developed by diverse HCPs in this phase. Focus groups with GBMSM were facilitated in Glasgow, London and Leeds. Two further focus groups were facilitated virtually (n=28). Telephone interviews were held with DAPs (n=3).

Overall within this phase, and bearing in mind the caveats concerning the lack of input from sexual health services and HCPs, there was on-going support for key intervention elements from Phase 1.

In the paragraphs below we illustrate how participants thought about acceptability and whether intervention ideas were amenable to change. The participants also indicate key barriers and facilitators to their implementation.

Participants agreed with the principle of needing to work across the system in different ways at the same time ‘all these different tools, all these different ways, you know, to make it easier to just inform people’ (GBMSM virtual 1) as was the need not to overburden any particular stakeholder ‘we need to find a way that we’re not heavily just relying everything on health care professionals, but also creating a culture of ownership, responsibility, and awareness, in the LGBT community. (GBMSM virtual 1). Some of the later data collection events also stressed the timeliness of the project and talked about how COVID-19 had done a huge amount of groundwork for the current project ‘what makes this conversation, particularly easy for everyone to understand now, is the experience of being locked down, the test and trace discussion around Covid-19 and being able to illustrate how that connects to STIs. So, everyone is in a different place right now than if we’d been having this conversation 12 months ago. So, it’s linking to that Covid-19 experience, must make selling this concept a lot easier’.(DAP2). Against this positive support for the intervention ideas as a whole, there was some mention of extant challenges. It was felt important to acknowledge the on-going and longstanding dynamics that negatively shape cross-stakeholder partnership work ‘And I think that, you know, as we talk about contact tracing and then we think about the sort of history of partner notification, there’s a really bad relationship there based on behaviour of public health in terms of gay men and gay men sexuality [...] There’s a history there and they’re not approaching this from a sort of neutral standpoint....bias and homophobia. I think that there are ways that we can also sort of work more collaboratively on some of the messaging and work more collaboratively on, you know, frankly repairing some of that relationship damage (DAP1)

In relation to what GBMSM said about changing their cultures and communities to normalise contact tracing there was widespread agreement and enthusiasm for progressing these areas of future intervention. The pre-COVID focus groups in particular highlighted a lack of knowledge and understanding about contact tracing amongst some GBMSM. They also highlighted a lack of consistent norms about what men should and could do about notification. For some GBMSM this was the norm but for others it clearly wasn’t. However there was a clear sense of the need to address these issues within future intervention. ‘I think what we need, first of all, is more visible campaigning, promotion, advertising, social media, that it becomes cool to say it, it just becomes cool, it makes a person who brings this up more cool, than somebody who doesn’t bring it up. So what we need is some kind of social media campaigning that normalises people discussing openly with each other’. (GBMSM Virtual1). There was also consensus and agreement that peer-focussed work between GBMSM was important ‘the facts are that people have sex, everyone should notify each other if they get something. That’s It.’ (GBMSM Glasgow).There was clear agreement to progress ideas such as targeted mass and social media including impactful messaging to normalise contact tracing.

Across the focus groups there was considerable heterogeneity of experience about contact tracing. From the pre-COVID focus groups in particular, there was a clear acknowledgment that it was not always a central feature of GBMSMs culture and that there was a lack of consistent knowledge about what contact tracing is, and what should happen around it, ‘If everyone had heard the same information, we’d probably be notifying our partners a lot more, but because of inconsistency in the information being given, it just isn’t there. Not everyone’s at the same level of access to it’.(GBMSM Leeds). Filling the basic knowledge gap around contact tracing amongst GBMSM was thought to be an important first step in progressing contact tracing interventions ‘I think normalising the words [about contact tracing] will normalise talking about it, will normalise telling each other about it. (GBMSM, Glasgow). Across the focus groups here was a shared sense of the high value and cultural acceptance of earlier STI and HIV mass media interventions. Discussion of these effective interventions focused on ideas about potential essential requirements for future contact tracing interventions, for example: ‘it’s got to be a very clever strapline, and a very kind of, immediately known, immediately remembered, and everyone knows exactly what it’s about. So yeah, you’re right, it’s got to be something short, but clear’. One focus group went as far as outlining a potential key message ‘Three Ts – test, treat, tell. (GBMSM, Virtual focus group 2). Equally, when considering the knowledge gaps around contact tracing, participants talked about how important it was to model what contact tracing was like: ‘maybe showing a video of what it’s like to tell someone, actually seeing

something like that, I think that would be...actually, if we're talking about the fear of it, let's actually have something that shows that experience'. (GBMSM, Virtual 1).

In relation to what DAPs told us about implementing the earlier intervention ideas, there was consensus across the interviews regarding the need for working in partnership across the system, there was also strong agreement with the need for culturally appropriate social and mass media. New suggestions emerged concerning associated merchandising 'do something that people are willing to wear'. There was also a clear acknowledgement of the need for targeted messaging that genuinely celebrated the diversity of GBMSM communities and used culturally appropriate figures 'porn stars and sex workers and their perspective on sexual health issues as opposed to a professional talking about it in a dry sense of the word. (DAP2). DAPs highlighted the need to think about the diversity of GBMSM within interventions and issues of visual representation 'I feel like every single gay person no matter where they fit in the community has to be able to look at that and realise that, okay, this is affecting me as well. So, it can't just be, you know, white guy who basically dresses in H and M head to toe and is saying, oh, you should tell your partner [inaudible 16.09], you know, head to toe in tattoos and blah blah'.

DAPs also highlighted the constraints they felt as small organisations with limited capacity to work beyond their primary remit as businesses 'We are a team of four probably I think it's everyone, I mean we're a small organisation so there's only so many hours in the day' (DAP2). However, there was support for the idea of signing up to a code of conduct relating to contact tracing but also an acknowledgement of its practical problems 'I mean the challenge would be agreeing what are the standards that we're going to offer to users and what's the technology required to get there'. In contrast to the relative enthusiasm for being involved in messaging and media interventions, changing functionality to enable contact tracing was not seen as a viable option, not only would the process be complex but 'Its around the right to block someone. And I think that would kind of violate it. I'd probably say most dating apps would say no'.

In relation to what GBMSM said about SHS and innovations in contact tracing ('new ways of working') many highlighted the potential of innovations within services, where future changes could 'start to feel you're being treated more like a human being' (GBMSM, Virtual focus group 2). For some GBMSM there was appreciation of the pressure facing many services and how this limited innovations in contact tracing. Time invested in one patient, receiving intense motivational counselling for contact tracing for one-off partners, for example, was understood to essentially take time away from others 'that means somebody else is waiting 20 minutes more just to go in and get tested' (GBMSM, London). However, they also perceived other opportunities within the services that could be harnessed for contact tracing 'you're waiting in the waiting room, the TV screen, they can do it (educate) that way as well, while you're waiting you just look at the screen for helpful messages tips' (GBMSM, London). Equally, some of the interaction time within services could be reduced if GBMSM prepared and shared information before arriving in the service, saving what was perceived to be valuable HCP time, 'if they sent you a link or something where you can send them back the number or a Grindr profile or whatever'. (GBMSM, London). Overall however, GBMSM felt that HCPs within some services could do more to engage GBMSM to do more in relation to contact tracing 'I think, the attitude of the people at those more progressive, and welcoming clinics, are a bit more supportive in terms of the actual, like, practical need for support, to do the disclosure bit' (GBMSM, Virtual focus group 2)

Our analysis went beyond the analysis of acceptability and barriers and facilitators to implementation outlined above. Because of the size of our comprehensive analyses of data collected in this phase we do not present the details of our theorisation of them using the TDF. Equally, we cannot show here how we used the BCW and the BCTT to generate highly specific evidence-based and theoretically informed ways in which the Stakeholders initial ideas could be operationalised in ways that systematically reduced barriers to implementation yet harnessed perceived facilitators. However, for the interested reader the full analyses underpinning these aspects of Phase 2 are presented within the supplementary files (Tables 1-7).




Figure 4 An overview of intervention ideas after detailed consideration of how to implement them and taken to Phase 3

As a very overarching summary of this work, our deductive analysis generated 17 barriers and 15 facilitators. We generated 83 potential intervention functions (Environmental restructuring, education, enablement, persuasion, modelling, training and incentivisation) and 141 potential BCTs. The full analyses underpinning this Phase of the research are available as supplementary files (Tables 1-7).

In relation to broad changes where our participants noted intervention ideas were not acceptable and/or not amenable to change the overall span of intervention ideas was reduced. This primarily included the abandoning of attempts to change dating app functionality and the use of technologies within sexual health services. Figure 4 shows an overview of the remaining intervention ideas that were taken forwards to Phase 3.

Phase 3 *What is the final suggested intervention content after further stakeholder engagement?*

A second large stakeholder engagement event mirroring that at the start of the project had originally been planned to take place, wherein intervention components developed and specified at the end of Phase 2 would be finalised for inclusion. However, due to the outbreak of COVID-19, this approach was no longer possible due to reduced stakeholder availability and the severe limitations of online facilitation for complex multi-stakeholder engagement. Consequently, Phase 3 was re-assessed, and an alternative scaled-down approach to refining the intervention’s final components through stakeholder engagement was agreed. This consisted of two stakeholder groups with community-based organisations with expertise in the field of sexual health and GBMSM (n=9, representing four

organisations) and two stakeholder events with diverse sexual health clinicians drawn from the wider LUSTRUM research team (n=5). Using the materials generated at the end of Phase 2 these stakeholders firstly discussed the general principles shaping the emerging intervention. Subsequently, they discussed the range of individual intervention components that Stage 2 had generated in detail. Throughout the APEASE criteria were used to facilitate rounded discussions about whether intervention ideas should be kept, modified or rejected.

In relation to the idea of working in targeted mass and social media interventions with impactful messaging, all stakeholders explored earlier intervention ideas such as *'test, trace, tell and treat'* with other messaging such as *'Identify, Inform, Invest, Improve'* (Expert clinicians, 2) wherein people Identify their contacts, Inform their contacts, Invest in a check-up and together this Improves your sexual health and those of the community. Echoing Phase 2 community-based stakeholders here also emphasised the need to represent the diversity of GBMSM communities and ensure materials depicted a diversity of experiences and perspectives. There was agreement about the need to educate all GBMSM about contact tracing *'there should be like some sort of educational awareness in terms of telling people or educating them how to respond if somebody tells you about, if they've been diagnosed with an STI'* (CBO stakeholder group1) and indeed to challenge STI-related stigma and blame *'extinguishing blame I think is definitely the central thing that we need to out'* (CBO stakeholder group2). Community-based stakeholders noted the need for positive focus *'the positive effects of like feeling a sense of responsibility to your community'* (CBO stakeholder group 2). There was also support for the idea of comparing two alternative story lines, one depicting effective contact tracing and one showing an example of contemporary blame culture, *'I think any campaign should talk about why there's a benefit to you on both sides. So, it might be the quote is, 'that dirty bastard!' and you can turn round and say, actually, 'I think it was great, because he got in contact with me, and he was looking after my health, so I was able to get treatment'. So, you switch it, you know, it's kind of a solutions focus, you make the negative into a positive. And you make it about how it's benefitting you and that person, and vice versa'* (CBO stakeholder group, 2).

From the community-based organisations there was consensus regarding the need to empower GBMSM to be at the heart of driving new interventions to improve contact tracing. There was support for the idea of peer driven work in this regard although caveats around appropriate funding to support such activity were clearly expressed. It was felt that volunteers were not an appropriate resource for these activities. Clear ideas about working with and celebrating community action were expressed *'one thing that really frustrates me more than anything about when we make advances, like PREP and U=U, is you'll have the clinicians shouting how we've done it, you'll even have NGOs shouting about we've done it, and to be frank, the only people that have actually done this and brought down the HIV rate and things like that, is the community, because they have taken action to do that. And for me, that's the really important thing to get across, that any intervention is, bring the community in, because they're the actual people that make that intervention work'* (CBO stakeholder group, 1). In turn galvanising community action was seen as a potential mechanism that could overcome some of the enduring barriers to delivering contact tracing for one-off partners amongst HCPs and sexual health services. Expert clinicians felt that GBMSM actively asking for help from HCPs in relation to seeking assistance with protecting their communities was seen as a very powerful way of activating overstretched HCPs (e.g., paraphrasing some of the recent community activism around PrEP, *'I want PN now'*).

Expert clinicians also agreed about the value of working with SHS and HCPs to learn from insights into their own successes and failures. Indeed it was felt that this mechanism not only could deliver on-going improvements but that it could also assure a consistency of contact tracing approaches across varied and deeply heterogeneous services. However, it was agreed that nationally co-ordinated guidance was a probably a more effective and efficient means to achieve these ends than attempts to host 'insight-sharing' initiatives within cross professional cross service platforms for example. Finally, all stakeholders agreed on the need to improve HCPs cultural competencies and provide services which could accommodate and facilitate contact tracing for GBMSM and their one-off partners. However expert clinicians acknowledged this might require approaches such as using secret shoppers and giving feedback in non-confrontational ways. Other suggested approaches included making more use of community-based organisations *'the third sector, have a great opportunity to build a culture where [contact tracing] is acceptable, and I'm not necessarily sure that the statutory sector can do that'* (CBO)

Key intervention elements that expert clinician stakeholders suggested should be abandoned included modifying the clinical environment to reinforce contact tracing messages. Stakeholders with expertise in SHS suggested this may not work for the diverse patients using the service from a range of backgrounds and for very diverse reasons




Figure 5 An overview of intervention ideas after final appraisal and modification with stakeholder groups

Final granular recommendations concerning *how* the acceptable interventions ideas that have been through multiple APEASE processes could be implemented are presented within Supplementary files

(Tables 1-7). In addition, we provide a narrative summary of our agreed core principles and components and this can be found in Appendix 1.

Discussion

The current study sought to follow contemporary guidance about intervention development (O’Cathain et al., 2019) and move the field forwards by systematically working with a range of stakeholders to develop a novel contact tracing intervention for GBMSM with STIs and their one-off partners that could perhaps overcome the multiple and multi-level barriers that have prevented the development of effective interventions in this area of contact tracing to date. As such this paper makes a novel contribution because of its use of diverse stakeholders, its use of diverse theoretical lenses and its focus on the psychosocial and socio-cultural determinants of contact tracing in addition to those that are more service oriented. This approach contrasts sharply with the recent glut of poorly conceived, atheoretical, app-based interventions that attempt to address contact tracing in the context of COVID-19 primarily through the development of Apps (Colizza et al., 2021). Overall, within the study we have worked with diverse stakeholders to see the problems and potential solutions to contact tracing with one-off partners. Across three phases of the project, we have iteratively honed a long-list of potential intervention ideas into a much shorter list of intervention ideas that are both acceptable and implementable and if they were to be operationalised may stand a chance of delivering sustainable change.

Our study has foregrounded the insights of key stakeholders and understands the drivers effecting contact-tracing amongst GBMSM and their one-off partners to relate to a complex and interconnected system. Through iterative data analysis, participant knowledge gleaned from system-mapping, exploration, and critical interviews and discussion, we have been able to refine the core components and content of a wholistic future intervention. To summarise refined intervention components focus on collective action across the system and final recommendations underscore the collaborative cross-sector nature of future interventions.

Our analysis suggested critical areas for future intervention included enabling and empowering GBMSM communities to do more about contact tracing and STIs with their one-off partners. Agreed activities in this area included peer-led initiatives, mass and social media with targeted impactful messaging. The content of such messaging could challenge STI-related stigma and raising awareness of contact tracing amongst GBMSM whilst also articulating norms and scripts of how to tell one-off partners about their risk of exposure and how to react appropriately. Working in partnership with CBOs and DAPs throughout this endeavour should enable the seamless promotion of these intervention elements across the social and sexual environments in which GBMSM meet. In addition to moderating some of these wider upstream and cultural determinants of contact tracing, it was also felt that enabling GBMSM to prepare themselves for contact tracing interactions with HCPs was also a good idea. This capitalised on the agency and responsibility of GBMSM. It was felt that this enabled them to seek support and help from SHS and HCPs rather than expecting overstretched HCPs and SHS to deliver this kind of work. DAP in-put, HCP guidance.

In relation to remaining uncertainties pertaining to the intervention and potential next steps it is important to acknowledge that the current study has only begun the process of intervention development and has successfully outlined potential intervention content. Our major investment in these first steps of effective intervention development were designed as a solution to the on-going

problem of rapidly conceived interventions failing to gain traction with the communities and services they are intended to work within (e.g. Chalmers et al., 2014; Walugembe et al., 2019). As a result of this emphasis, key questions relating to how the emerging interventions could be funded, how it could be evaluated and the feasibility of successfully collecting and analysing such data all remain to be answered.

Limitations and strengths

The outbreak of COVID-19 in the UK in early 2020 severely affected the timeline and planned operations of the key phases of this study. Disruptions to Phase 2 in particular impacted negatively on the project as whole. This negativity relates primarily to the loss of HCP perspectives in appraising and further developing intervention ideas within Phase 2. Given the centrality of diverse perspectives and the idea of problem solving from multiple vantage points simultaneously, the loss of the projects ability to systematically hone intervention ideas using sexual health services and HCPs perspectives remains a limitation. Our work with expert clinical stakeholders from within the project team did of course help compensate for this loss. As a result, some aspects of potential intervention ideas concerned with SHS and HCPs have only been illuminated by GBMSM adopting a ‘patient perspective’ and our expert clinical stakeholders within Phase 3. As such issues such as ‘using more technology such as mobile phones within clinical interactions’, ‘using techniques such as motivational interviewing or scripts to assist with contact tracing with one-off partners’, ‘the need for more culturally competent services’, and ‘ways of overcoming limited capacity’ could potentially have been assessed and modified differently if we had been able to recruit diverse HCPs working within the field. Similarly, although we did manage to recruit some DAPS to the project, our attempts at recruitment struggled in the context of this sector adapting to the impacts of COVID-19. Again, this represents a limitation to the overall design. A final limitation relates to Phase 3 and the impact of COVID on our stakeholder work. Ideally, this would have mirrored Phase 1 with an interactive mixed small groups of professionals working with the intervention ideas. Once more, given COVID-19 restrictions this was not possible. Further limitations include our sole reliance on qualitative approaches, if time and resource had permitted a more expansive mixed methods design would have enhanced the project, tools like using the Delphi approach could have been used.

Strengths of the project include its systematic use of stakeholder input which has ensured the intervention ideas are grounded in the priorities of those the potential intervention will affect. Our use of diverse stakeholders ensured that the potential intervention did not either overburden one part of the system and indeed that it did not shunt the problem elsewhere within the system. Other key strengths included the studies pluralistic approach to using theory. Where complex adaptive systems theory was useful, we used it to see the nature of problem the intervention was trying to solve differently. This enabled us to see the importance of intervening upstream as well as well as downstream. However, where complex adaptive systems theory is weak, we used behaviour change theory to systematically understand and then specify ways to ensure the implementation of the potential intervention ideas our stakeholders had generated. The overall approach has provided a number of lenses and levels in which to see the intervention we have generated. This varies from visual overviews (Figure 2-5), narrative summaries (appendix) to very granular accounts of intervention content (Tables 1-7).

Conclusion

We have undertaken a process of complex intervention development which foregrounds the insights of key stakeholders and understands the drivers effecting contact-tracing amongst GBMSM to be a complex and interconnected system. Through iterative data analysis, participant knowledge gleaned from system-mapping, exploration, and critical interviews and discussion, we have been able to refine the core components and content of a holistic intervention. Final recommendations underscore the collaborative cross-sector nature of potential interventions to intervene in this area. Our suggested intervention ideas are multi-level (upstream and downstream intervention elements), multi-agent (GBMSM, CBOs, HCPs, DAPs) and consist of co-ordinated and complementary activities including mass and social media elements, peer-led initiatives, the priming of individual GBMSM to prepare for interactions within SHS, and HCP training assuring a standard level of cultural competency as well as adequate national guidance and monitoring to enable SHS to appraise and constantly improve their contact tracing outcomes.

References

Atkins, L. *et al.* (2017) 'A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems', *Implementation Science*. BioMed Central Ltd., 12(1), pp. 1–18. doi: 10.1186/s13012-017-0605-9.

Department of Health and Social Care (2020) *Weekly NHS Test and Trace bulletin, England: 16 to 22 July 2020*.

Colizza V, Grill E, Mikolajczyk R, Cattuto C, Kucharski A, Riley S, Kendall M, Lythgoe K, Bonsall D, Wymant C, Abeler-Dörner L. Time to evaluate COVID-19 contact-tracing apps. *Nature Medicine*. 2021 Mar;27(3):361-2.

Estcourt, C. S. *et al.* (2020) 'Accelerated partner therapy (APT) partner notification for people with Chlamydia trachomatis: protocol for the Limiting Undetected Sexually Transmitted infections to RedUce Morbidity (LUSTRUM) APT cross-over cluster randomised controlled trial', *BMJ Open*, 10, p. 34806. doi: 10.1136/bmjopen-2019-034806.

Estcourt CS, Flowers P, Cassell JA, Pothoulaki M, Vojt G, Mapp F, Woode-Owusu M, Low N, Saunders J, Symonds M, Howarth A. Going beyond 'regular and casual': development of a classification of sexual partner types to enhance partner notification for STIs. *Sexually transmitted infections*. 2021 Apr 29.

Fitzgerald M, Bell G. Measuring the effectiveness of contact tracing. *International journal of STD & AIDS*. 1998 Nov 1;9(11):645.

Flowers, P., Duncan, B. and Frankis, J. (2000) 'Community, responsibility and culpability: HIV risk-management amongst Scottish Gay Men.', *Journal of Community & Applied Social Psychology*, 10(4), pp. 285–300.

Flowers, Gerressu, MacLeod, McQueen, Vojt, Symonds, Woode Owusu and Esctourt (2021) How can we increase notification of exposure to STIs between gay and bisexual men?: intervention development using stakeholder engagement, qualitative methods and behavioural science. medRxiv 2021 .05.27.21257903; doi: <https://doi.org/10.1101/2021.05.27.21257903>

Juneau, C.-E. *et al.* (2020) 'Effective Contact Tracing for COVID-19: A Systematic Review', *medRxiv*. Cold Spring Harbor Laboratory Press, p. 2020.07.23.20160234. doi: 10.1101/2020.07.23.20160234.

- Limb, M. (2020) 'Covid-19: Sandwell Council in West Midlands sets up contact tracing, citing failures of national scheme', *BMJ (Clinical research ed.)*. NLM (Medline), 370, p. m3065. doi: 10.1136/bmj.m3065.
- Mathevet, I. *et al.* (2021) 'Accounting for health inequities in the design of contact tracing interventions: A rapid review', *International Journal of Infectious Diseases*. Elsevier B.V., pp. 65–70. doi: 10.1016/j.ijid.2021.03.010.
- May, C. (2013) 'Towards a general theory of implementation', *Implementation Science*. BioMed Central, 8(1), pp. 1–14. doi: 10.1186/1748-5908-8-18.
- McQueen, J. M. *et al.* (2020) 'Protocol to develop a partner notification intervention for men who have sex with men', *medRxiv*. Cold Spring Harbor Laboratory Press, p. 2020.10.21.20209049. doi: 10.1101/2020.10.21.20209049.
- Medical Research Council (2019) *Developing and Evaluating Complex Interventions: Draft of updated guidance (Draft for consultation.)*. Available at: [https://www.sphsu.gla.ac.uk/stakeholder-survey-2019/Full complex guidance draft for consultation v1.1 26.03.19.pdf](https://www.sphsu.gla.ac.uk/stakeholder-survey-2019/Full%20complex%20guidance%20draft%20for%20consultation%20v1.1%2026.03.19.pdf) (Accessed: 25 May 2021).
- Megnin-Viggars, O. *et al.* (2020) 'Facilitators and barriers to engagement with contact tracing during infectious disease outbreaks: A rapid review of the evidence', *PLoS ONE*. Public Library of Science. doi: 10.1371/journal.pone.0241473.
- Michie, S. *et al.* (2013) 'The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: Building an international consensus for the reporting of behavior change interventions', *Annals of Behavioral Medicine*. Ann Behav Med, 46(1), pp. 81–95. doi: 10.1007/s12160-013-9486-6.
- Michie, S., Atkins, L. and West, R. (2014) *The Behaviour Change Wheel: A Guide to Designing Interventions*. London, UK: Silverback Publishing. Available at: www.behaviourchangewheel.com.
- Michie, S., van Stralen, M. M. and West, R. (2011) 'The behaviour change wheel: A new method for characterising and designing behaviour change interventions', *Implementation Science*. BioMed Central, 6(1), pp. 1–12. doi: 10.1186/1748-5908-6-42.
- Müller, J. and Kretzschmar, M. (2021) 'Contact tracing – Old models and new challenges', *Infectious Disease Modelling*. KeAi Communications Co., 6, pp. 222–231. doi: 10.1016/j.idm.2020.12.005.
- Murray, E. *et al.* (2010) 'Normalisation process theory: A framework for developing, evaluating and implementing complex interventions', *BMC Medicine*. BioMed Central, 8(1), pp. 1–11. doi: 10.1186/1741-7015-8-63.
- O’Cathain, A. *et al.* (2019) 'Guidance on how to develop complex interventions to improve health and healthcare', *BMJ Open*. BMJ Publishing Group, 9(8). doi: 10.1136/bmjopen-2019-029954.
- Pothoulaki, M. *et al.* (2020) 'The social organisation of sexual relationships: an exploratory qualitative study'. SocArXiv. doi: 10.31235/osf.io/m4fgb.
- Rowley, J. *et al.* (2019) 'Chlamydia, gonorrhoea, trichomoniasis and syphilis: Global prevalence and incidence estimates, 2016', *Bulletin of the World Health Organization*. World Health Organization, 97(8). doi: 10.2471/BLT.18.228486.
- Rutter, H. *et al.* (2017) 'The need for a complex systems model of evidence for public health', *The Lancet*. Lancet Publishing Group, pp. 2602–2604. doi: 10.1016/S0140-6736(17)31267-9.

Saurabh, S. and Prateek, S. (2017) 'Role of contact tracing in containing the 2014 Ebola outbreak: A review', *African Health Sciences*. Makerere University, Medical School, 17(1), pp. 225–236. doi: 10.4314/ahs.v17i1.28.

Swanson, K. C. *et al.* (2018) 'Contact tracing performance during the Ebola epidemic in Liberia, 2014-2015', *PLoS Neglected Tropical Diseases*. Public Library of Science, 12(9). doi: 10.1371/journal.pntd.0006762.

Wang, A. L. *et al.* (2016) 'Optimizing partner notification programs for men who have sex with men: Factorial survey results from South China', *PLoS ONE*. Public Library of Science. doi: 10.1371/journal.pone.0157749.

Walugembe DR, Sibbald S, Le Ber MJ, Kothari A. Sustainability of public health interventions: where are the gaps?. *Health research policy and systems*. 2019 Dec;17(1):1-7.

Wi, T. *et al.* (2017) 'Antimicrobial resistance in *Neisseria gonorrhoeae*: Global surveillance and a call for international collaborative action', *PLoS Medicine*. Public Library of Science, 14(7). doi: 10.1371/journal.pmed.1002344.

Wilson, T., Holt, T. and Greenhalgh, T. (2001) 'Complexity and clinical care', *BMJ*. British Medical Journal Publishing Group, 323(7314), p. 685. doi: 10.1136/bmj.323.7314.685.

World Health Organisation (2016) *Global health sector strategy on sexually transmitted infections 2016-2021. Towards ending STIs*.

Appendix 1: Narrative summary of suggested intervention components

Core principles

- Diverse ‘change agents’ (communities, organisations, professions, individuals) should work together to change the system that drives effective contact tracing for one-off sexual partners. This will require new infrastructure (e.g., meetings, processes) that enable a shared and fair approach to improving such contact tracing
- The ‘burden’ of the intervention should be shared across the system with each change agent using its particular expertise.
- Simultaneously address both the *distal* and *proximal* determinants of effective contact tracing for one-off sexual partners
- Expect system change to be slow and challenging but sustainable in the long-run

Core intervention components (as summarised and narrativised from data tables)

What did we learn from GBMSM about implementing changes relating to the specific content of targeted mass and social media interventions to normalise contact tracing with one-off partners through impactful messaging? (Table 1)

- ❑ **GBMSM should be targeted for interventions to educate them on contact tracing, change behaviours and promote effective contact tracing for all former-partners, especially one-off partners.** Mass media interventions should be developed to promote contact tracing for GBMSM with one-off partners. Messaging about contact tracing and one-off partners should be targeted, short, punchy, easy to remember. Phrases and terms that build on recent developments from COVID and contact tracing might be helpful:- 4T’s Test, Treat, Trace, Tell. This kind of work will enable MSM to tell more one-off partners. Messaging should always be branded with reputable organisations, and preferably co-produced with, and being visibly inclusive of, a diverse range of MSM. The content of these messages should educate men about the value and importance of PN, and the consequences of blame and stigmatisation in relation to STIs.
- ❑ **Public health messages and materials should be developed to educate GBMSM on contact tracing.** In order to implement these intervention ideas a national stakeholder or steering committee, including experienced HCPs, CBOs, GBMSM and other relevant representatives, should co-produce and decide on a range of consistent and clear intervention messages. Intervention messages and materials should be tailored to specific STIs, distributed and made available widely across virtual and physical locations where men meet, and focus on increasing awareness of contact tracing amongst GBMSM. Intervention messages should be clearly branded with reputable, culturally-appropriate organisations, and should build on recent learning from COVID-19 public health messaging. Intervention messages and materials should provide an example of how to tell a one-off partner about their heightened risk of STI acquisition, highlighting the health and wider consequences of not undertaking contact tracing. This could feature a short film or animation. This works as an example to follow and use in practice, and also helps to set norms concerning what to do and

expect from others. Examples would be based on input from experienced HCP, other staff and peer MSM

What did we learn from GBMSM about implementing peer-focussed work between GBMSM to endorse contact tracing with one-off partners (Table 2)

- ❑ A range of peer-led work should be designed and delivered to enable GBMSM to capitalise on existing community norms supportive of contact tracing and also reducing norms which present barriers to contact tracing. Community-based organisations may be best placed to deliver this work. Peer-led initiatives will need funding and are unlikely to be viable through volunteering routes.
- ❑ Training or up-skilling opportunities should be offered to GBMSM on how to talk with other GBMSM about contact tracing and reducing STI-related stigma. Approaches could draw upon the recent activism seen around PrEP and focus this on contact tracing or partner notification (PN), for example, 'I want PN now'. Such approaches could work with classic models of peer education (e.g., key opinion leaders), or 'social influencers' within the social media to change descriptive norms (what people think other people do) and injunctive norms (what people think they should do) around contact tracing with one off partners. Such peer influence could also clearly model how and why to do contact tracing for all partners.
- ❑ Educational materials, including enabling approaches to peer education, should be developed for GBMSM on STIs and the value and importance of contact tracing. To compensate for the failure of mainstream education to provide men with adequate levels of sex education including issues of contact tracing, there is a clear need for a range of intervention activities (e.g. peer interactions), materials and messaging to focus on re-framing issues of contact tracing amongst men. These materials should be developed and branded by reputable organisations such as CBOs and be co-produced alongside diverse GBMSM themselves, enabling culturally appropriate content. Educational materials should focus on celebrating the power of peers to teach each other about contact tracing; what it is, why it should be done, how it should be done and why it should be done for one off partners in particular.
- ❑ Educational materials should provide an example, suggestions of wording for how to tell a one-off partner about their heightened risk of STI acquisition. These would work as examples for others to follow and use in practice but also, simultaneously, helps to set norms concerning what to do and what to expect from others. Educational materials should also directly address the social and sexual cultures of men and reduce perceived negative consequences of talking about STIs to enable more peer conversations about contact tracing to take place.

What did we learn from GBMSM about using the social media to endorse contact tracing with one-off partners (Table 3)

- ❑ **Social media should be utilised as a key component in targeting GBMSM.** Social media could be used within multimodal or multimedia interventions to change norms amongst GBMSM and challenge stigma surrounding contact tracing, and promote particular norms encouraging engagement in contact tracing with one-off partners. Social media could be used to show people contacting one-off partners and, in this way, provide examples of how to do it and reinforce norms that doing it is the right thing to do. Social networks and dating app sites could be used to exchange, repeat and 'share' messaging concerning contact tracing. GBMSM should receive focused, consistent messaging about contact tracing and the health and community impact of effective contact-tracing from key sources such as HCPs, CBOs, dating apps, and physical and virtual hook up sites. Messaging would detail the diverse consequences of PN for all. It may be possible to recruit social influencers from social media to promote PN and harness their networks to promote key messages

What did dating app providers tell us about implementing changes relating to the endorsement of contact tracing within their public facing infrastructure? (Table 4)

- ❑ **DAPs should host educational content on contact tracing on their platforms and website - developed in partnership with CBOs, GBMSM, HCPs and other key stakeholders - and ensure education is inclusive to GBMSM from all backgrounds.** Educational messages can be supported, endorsed and amplified by those with social influence across other social media platforms. DAPs should consider building educational messages from recent developments in COVID-19 contact-tracing phrases and terms familiar to the public.
- ❑ **DAPs should consider collaborations with social influencers to promote contact tracing educational content.** DAPs should ensure content and messaging is linked to relevant national awareness raising campaigns, where possible. DAPs should consider opportunities for partnership and collaboration across stakeholders such as CBOs and public health bodies to harness strengths and share resource burden.
- ❑ **Those aiming to collaborate with DAPs to promote contact tracing should provide DAPs with persuasive information** on how education and advocacy will benefit their future work and positive experience of app users.

What did dating app providers tell us about implementing changes relating to changing dating app functionality? (Table 5)

The intervention development process did not agree on recommendations for DAPs to implement changes to dating app functionality. DAPs were critical of developing in-app functions to enable users to indicate they are positive about contact tracing as this may not represent a genuine statement of intent and may infringe upon users rights. DAPs also indicated that such functionality would only prove effective if other measures such as education and interventions which promote attitudinal change amongst GBMSM are also in place. DAPs also suggested limited impact of functions which would enable anonymous contact tracing messages as uptake of this option is likely to be low amongst GBMSM; research from previous trials shows options for anonymous messaging are unlikely to increase rates of contact tracing due to lack of use. Finally, DAPs were also reluctant to develop functions which would enable users to 'unblock' former sex partners in order to send a message for contact tracing purposes, as this could be considered a violation of a user's 'right to block'.

What did dating app providers tell us about implementing changes relating to signing up to a code of conduct to endorse contact tracing? (Table 6)

The intervention development process did not agree on the recommendation for DAPs to sign up to a code of conduct supporting contact tracing, though the idea was not outrightly rejected. Stakeholders indicated that in order to be successful, a code of conduct would require lead from relevant organisations in development and encouraging commitment from DAPs. The following may be considered to enable this alternative approach.

- ❑ CBOs, GBMSM and public health organisations could lobby for a code of conduct for DAPs to endorse contact tracing. DAPs could work together with organisations to develop standards and co-produce a code of conduct for DAPs to endorse contact tracing.

What have we learned about implementing improved interactions between HCPs and GBMSM to enhance contact tracing for one-off partners? (Table 7)

- ❑ **HCPs and sexual health clinics should form and utilise partnerships with CBOs and GBMSM to deliver culturally competent services.** The delivery of contact tracing should be reframed to consider the roles of CBOs in supporting GBMSM and enabling them to play a more active role. Opportunities for partnership with CBOs should be explored to support over-worked over-burdened clinic staff with contact tracing for GBMSM and their one-off partners. CBO educational initiatives could also promote the value and impact of individuals completing contact tracing themselves, as this helps to reduce stigma, create ownership and removes reliance on HCPs who have limited resources. GBMSM could co-produce materials with CBOs and communities to educate and enable other GBMSM to systematically raise issues of contact tracing with their HCPs
- ❑ **Support GBMSM to prepare for contact tracing interactions with HCPs. HCPs should work to improve GBMSM engagement with contact tracing in-clinics through education, priming, and incentivisation.** Educational messages for GBMSM should be devised by CBOs and HCPs and made widely available to GBMSM through CBO's, DAPs and agreed appropriate clinic-based routes. HCPs should prime GBMSM at the earliest opportunity to enable and motivate them to participate actively in contact tracing, saving valuable clinic time and resources. HCPs should approach all contact tracing related activity in a non-stigmatising way, using guidance and training on how to perform this behaviour. Sexual health clinics should explore the potential role of CBOs in fostering a culture of peer support to persuade GBMSM to contact previous sexual partners in more difficult situations. Sexual health clinics should also consider opportunities to develop an online system for contact tracing, based on functionality and learning. Education for GBMSM should highlight FAQ and educational resources available for GBMSM to view in their own private space as opposed to a busy clinic environment. Educational messages could be linked to how GBMSM get their results to inform and persuade GBMSM to undertake contact tracing, as with similar sexual health interventions such as PrEP.

- ❑ **Education and training should be developed for HCPs on the value and importance of contact tracing, and to underscore promotion of culturally competent contact tracing as a key part of HCPs role.** Promotion of contact tracing should be communicated as a key part of the role of HCPs and built into clinic standards, staff training and induction processes. Clear messages should be targeted to HCPs, developed in collaboration with CBOs, GBMSM and other relevant professional bodies. Clear and consistent messages on the value and importance of contact tracing should be communicated from credible sources such as reputable professional bodies (i.e. BASHH), commissioners, executive team members, senior clinicians, and consultants. Educational initiatives should be developed and delivered to HCPs, focusing on the consequences of poor contact tracing and the potential impact of the process. Educational initiatives should be planned and coordinated with a range of actors, and reinforce the positive public health consequences of contact tracing (i.e. CBOs, DAPs, sexual health organisations, and public health bodies). Educational initiatives should include understanding of GBMSM sexual cultures, to improve contact tracing for GBMSM with one-off partners. Skills-based training should be delivered to HCPs offering advice for consistent and effective contact tracing practice, and focusing on increasing HCPs knowledge and confidence to talk to GBMSM about contact tracing in the context of their sexual cultures.

- ❑ **National networks, professional bodies and senior leaders should enable prioritisation of contact tracing through standardisation of contact tracing approaches, incentivisation, and training for HCPs.** Clinic based consensus could be used to ensure a consistent approach towards contact tracing is adopted, with intervention approaches documented in clinic policies and procedures. Where possible provide prompts and cues these could be used with in-built algorithms on clinic computer system, clinic paperwork, posters around clinic areas, above desks with prompts and reminders about PN to motivate and remind HCPs of their PN role. Specific clinic initiatives and approaches to contact tracing should be standardised and clearly defined with prompts such as a flow chart and predefined questions to minimise demands on HCP time in light of resource constraints. HCPs should use a combination of professional judgement combined with guidance, hints and tips on good practice. Targets and goals related to contact tracing should be used as components of staff supervision and appraisal systems in clinics to incentivise contact tracing amongst HCPs.

- ❑ **Commissioners and senior HCPs should consider in-clinic priority changes in relation to contact tracing and its value for public health.** Links could be made to COVID-19 contact-tracing to stress the importance and value of increasing the percentage of partners contacted (i.e if 10% are reached, versus 50% or 80% of partners). Sexual health clinics should consider the value of all staff completing recognised training such as accredited CPD training with a focus on consistent terminology and phrases that can be used in contact tracing consultations tailored to GBMSM with one-off partners. More experienced staff could offer support and supervision through shadowing, live supervision as a method to boost confidence amongst other staff when asking about sexual cultures and documenting GBMSM partner types.

Table 1 What did we learn from GBMSM about implementing changes relating to the specific content of mass and social media interventions to normalise contact tracing through impactful messaging?

Key component, step, or behaviour of the intervention derived from initial stakeholder event	Barriers to enacting this intervention component	Facilitators to enacting this intervention component	Indicative quote	TDF Domain	Intervention function (BCW)	Potential behaviour change techniques	Suggestions for intervention content
Targeted and impactful mass and social media messages to normalise contact tracing		MSM find it easier to tell their one-off partners when educational messages are short, punchy and easy to remember	Yeah, the classic one that comes to front of mind is the whole HIV campaign, and like, giving HIV the finger. It's just really succinct, and it does the job really well. It communicates that that's the sort of test you have to do, if this is what you're going towards...sorry, I don't want to go off on a tangent on someone else's work. Yeah, it's boiling it down to something pithy. Sorry, and again, it's my job, I do it every day. But no one is going to like really remember something super long-winded. You just have to, and also, nothing too meaningful as well. I don't know what exactly your target age is, but your demographic is quite wide,	Memory, attention and decision processes Knowledge	Environmental restructuring Education	7.1 Prompts/Cues 5.3 Information about Social and environmental consequences 5.1 Information about health consequences 5.6 Information about emotional consequences. 12.2 Restructuring the social environment 9.1 Credible source	Messaging about PN and one-off partners should be targeted short, punchy, easy to remember (7.1). It should be branded with reputable organisations (9.1) and preferably be co-produced with diverse men. (9.1) The content of such branding should relate to <u>educating</u> men about PN (5.3) but also educating men about the <u>consequences of their stigmatising</u> in relation to STIs (5.6,5.1) In relation to <u>educating</u> men about PN catchy phrases and terms that build on recent developments from COVID and contact tracing might be helpful:- 4T's Test, Treat, Trace, Tell. This kind of work will <u>enable</u> MSM to tell more one-off partners (12.2, 7.1)

			<p>and you have to really hone in on a specific group of people that you're trying to communicate with too, is where I would go. Otherwise, you're stabbing in the dark.</p> <p>R4: Yeah, it's got to be a very clever strapline, and a very kind of, immediately known, immediately remembered, and everyone knows exactly what it's about. So yeah, you're right, it's got to be something short, but clear.</p> <p>Yeah, take like the classic, catch it, bin it, kill it, you know. I mean, I don't think something that's as easy to do, you know, but a listing that has transcended time. I don't even know when that came out, was it swine flu, or something else?</p> <p>Three Ts – test, treat, tell. (virtual focus group 2)</p>				
Targeted and impactful mass and social media messages to normalise contact tracing	MSM find it difficult to notify their one-off partners because public health messages on what to do about partner notification, are	MSM find it easy to notify their partner/s because public health messages on partner notification	<i>I hear a lot of people saying, I'm on PrEP, and it really worries me, because I think, that's great for HIV, but not for chlamydia or gonorrhoea, whatever else, so that...I hear that a lot, and I suppose that'd be the thing that...but I guess if there's education</i>	Environmental context and resources Knowledge Persuasion	Environmental restructuring Education Enablement	<p>12.1 Restructuring the physical environment</p> <p>12.2 Restructuring the social environment</p> <p>9.1 Credible source</p>	<p>A national stakeholder/steering committee (9.1) could co-produce and agree on a range of consistent and clear intervention messages (5.1, 5.3). These messages should be tailored to specific STIs wherever possible and co-produced with HCPs, NGOs and MSM themselves. (9.1, 12.5)</p> <p>Intervention messages, and/or materials should be <u>provided and distributed across virtual and physical</u></p>

<p>inconsistent and unclear and do not assist</p>	<p>are consistent and clear</p>	<p><i>done in the background and actually we know that if someone tells you, you've got an STI or an STD, we know that you go and get checked. (Glasgow)</i></p> <p><i>If everyone had heard the same information that I seem to know about partner notification, we'd probably be notifying our partners a lot more, but because of inconsistency in the information being given, it just isn't there. Not everyone's at the same level of access to it.(Leeds)</i></p>			<p>12.5 Adding objects to the environment</p> <p>5.1 Information about health consequences</p> <p>5.3 Information about social and environmental consequences</p> <p>5.2 Salience of consequences</p>	<p>venues where men meet. (12.5, 12.1, 12.2)</p> <p>Intervention messages, and/or materials should focus on <u>increasing awareness</u> of PN and <u>why people should do it (5.1, 5.2, 5.3)</u>, building on recent learning concerning COVID-19 might assist with this task. (5.3, 5.2, 5.1)</p> <p>Intervention messages, and/or materials should be <u>consistent (5.1, 5.3)</u> and <u>clear and highlight that engaging in PN with all partners including one -off partners is the right thing to do(5.2)</u>. They should be clearly branded in relation to reputable organisations that are culturally appropriate. (9.1,)</p>
<p>MSM find it difficult to notify their one-off partners because public health messages on partner notification are not pitched correctly to motivate MSM</p>	<p>MSM find it easy to notify their partners because public health messages on partner notification are pitched correctly to motivate MSM</p>	<p><i>Secondly, yes, or maybe showing a video of what it's like to tell someone, actually seeing something like that, I think that would be...actually, if we're talking about the fear of it, let's actually have something that shows that experience. Whether it's, so a dramatisation, actually a visual...I'm a visual person, I need to see something. To be shown and just read it, it may not work. (Virtual1)</i></p> <p><i>Sure, this was actually something I was going to mention at a point, but posters, message blasts, adverts, slogans that</i></p>	<p>Environment context and resource</p> <p>Knowledge</p> <p>Beliefs about consequences</p> <p>Social influence</p>	<p>Environmental restructuring</p> <p>Enablement</p> <p>Education</p> <p>Persuasion</p> <p>Modelling</p>	<p>9.1 Credible source</p> <p>12.2 Restructuring the social environment</p> <p>3.1 social support (unspecified)</p> <p>12.5 Adding objects to the environment</p> <p>12.1 Restructuring the physical environment</p> <p>5.3 Information about social and environmental consequences</p> <p>5.1 Information about health consequences</p> <p>6.1 Demonstration of a behaviour</p>	<p>Achieve national consistency via stakeholder or steering committees, and with the input of peer MSM, experienced HCPs, NGOs and other staff (9.1). <u>Ensure PN messages are widely available (12.5)</u>, that the same message is expressed in different ways tailored to different MSM groups (3.1). Messaging could focus on normalising testing (12.1) and normalising STIs (12.1), again with input from MSM, to include buzzwords, and what is culturally sensitive and powerful? (9.1, 12.5, 12.2)</p> <p>Intervention messages, and/or materials should be <u>consistent and clear and highlight the health and wider consequences</u> of telling, or not telling, someone (5.1, 5.3, 16.3) about their heightened risk of STI acquisition. This could be achieved through mechanisms like sliding doors/split screen. (16.3, 9.3, 5.3, 5.1)</p> <p>Intervention messages, and/or materials should provide an <u>example of how to tell a one off partner</u></p>

			<p><i>motivate us to tell one off sexual partners about their potential infection. So what I was thinking a minute ago was, an advertising campaign just to normalise the words, just because we don't see chlamydia and gonorrhoea and that sort of thing discussed or mentioned by a lot of people, and I think normalising the words will normalise talking about it, will normalise telling each other about it. (Glasgow)</i></p>			<p>16.3 Vicarious consequences</p> <p>9.3 Comparative imagining of future outcomes</p>	<p><u>(6.1) about their heightened risk of STI acquisition.</u> This could feature a short film or animation. This works as an example to follow and use in practice but also, simultaneously, it helps to set norms (9.3) concerning what to do and expect from others. Examples would be based on input from experienced HCP, other staff and peer MSM. (9.1)</p>
			<p>I know that health care services try to reach out to audience in different forms and shapes. But I think that we need to find a way that we're not heavily just relying everything on health care professionals, but also creating a culture of ownership, responsibility, and awareness, in the LGBT community. So as individuals, we're also doing that, rather than just heavily relying on health care professionals to sort everything out. (virtual 1)</p>	<p>Knowledge</p> <p>Environmental context and resources</p> <p>Behavioural regulation</p>	<p>Enablement</p> <p>Education</p> <p>Training</p>	<p>3.1 Social support (unspecified)</p> <p>5.3 Information about social and environmental consequences</p> <p>5.5 Anticipated regret</p> <p>4.1 Instruction on how to perform a behaviour</p> <p>6.1 Demonstration of behaviour</p> <p>5.6 Information about emotional consequences</p>	<p>Opportunities to empower MSM to take ownership could be explored as this would reduce the burden on HCPs (3.1). NGOs could co-create educational programmes offering tailored MSM specific PN education (5.3, 4.1). This could detail different scenarios for example 'don't care selfishness versus the moral responsibility and more personalised PN' (5.5). Scenarios could be used to educate MSM and outline the different PN approaches and potential outcomes related to STI prevention (6.1, 5.6). Overall aim to empower MSM who have been diagnosed with an STI to complete more PN (3.1).</p>

Green content = Agreed in Phase 3 as final intervention content

Amber content = Content that could be considered for the intervention

Red content = Content to be dropped from the intervention

Table 2 What did we learn from GBMSM about implementing peer-focused work between GBMSM to endorse contact tracing with one-off partners

Key component, step, or behaviour of the intervention derived from initial stakeholder event	Barriers to enacting this intervention component	Facilitators to enacting this intervention component	Indicative quote	TDF Domain	Intervention function (BCW)	Potential behaviour change techniques	Suggestions for intervention content
MSM educate each other on STIs and PN	3.1 MSM find it difficult to 'educate each other' because school-based sex education lacks key factual information about partner notification and is not tailored to MSM	3.2 MSM find it easy to 'educate each other' because school-based sex education covers key factual information about partner notification and is tailored to MSM	<i>It should just be normal, it (the education) should just be the facts are that people have sex, everyone should notify each other if they get something. That's it. I think it should just be normal and if it is factual like that and not kind of sexual, it's about sex but not sexual, then it just becomes the norm (Glasgow)</i>	Environment context and resource Social influence Knowledge Skills	Environmental restructuring Education Training Enablement Persuasion	9.1 Credible source 8.3 Habit formation 4.1 Instructions on how to perform a behaviour 3.2 Social support (practical) 8.6 Generalisation of target behaviour (i.e. PN with all partners including one-offs) 5.3 Information about social and environmental consequences 6.1 Demonstration of a behaviour 7.1 Prompts/cues 12.2 Restructuring the social environment	To compensate for the failure of mainstream education to provide men with adequate levels of sex education (12.1, 3.2) including issues of partner notification, there is a clear need for intervention materials and messaging to focus on issues of PN amongst men (4.1, 5.3). These materials should be branded as having been developed by reputable organisations (9.1) (e.g., NGOs) and preferably be co-produced by men themselves enhancing the normative content. (9.1, 8.6) It may be possible and useful to focus on the power of peers (9.1) to teach each other about PN; what PN is and why it should be done and why it should be done for one off partners in particular. (3.2, 5.3, 9.1, 12.2). Peers could be used to educate MSM about the moral side of PN and how it is the right thing to do (9.1, 3.2, 5.3). Intervention messages, and/or materials, training should provide an example, suggestions of wording (4.1), of how to tell a one off partner about their heightened risk of STI acquisition. These would work as examples for others to follow and use in practice (7.1, 6.1), but also, simultaneously, helps to set norms concerning what to do and what to expect

							<p><i>from others. (8.3))</i></p> <p>Education programmes within schools <u>should include the importance of regular STI testing (12.2) and partner notification, with case studies, scenarios and tips on how to do this well (4.1).</u> This should be part of an inclusive sex and relationship educational package informed by a variety of stakeholders, including young MSM (6.1, 9.1, 8.3)</p>
--	--	--	--	--	--	--	---

Green content = Agreed in Phase 3 as final intervention content

Amber content = Content that could be considered for the intervention

Red content = Content to be dropped from the intervention

Table 3 What did we learn from GBMSM about using the social media to endorse contact tracing with one-off partners

Key component, step, or behaviour of the intervention derived from initial stakeholder event	Barriers to enacting this intervention component	Facilitators to enacting this intervention component	Indicative quote	TDF Domain	Intervention function (BCW)	Potential behaviour change techniques	Suggestions for intervention content
Peer interactions and Social media approaches		5.1 MSM find it easy to notify one-off partners as social media campaigns, and advertising promote PN making it normal practice for MSM to discuss STIs openly	<i>I think what we need, first of all, is more visible campaigning, promotion, advertising, social media, that it becomes cool to say it, it just becomes cool, it makes a person who brings this up more cool, than somebody who doesn't bring it up. The person taking part in discussing with the first potential partner is something cool, rather than uncool. So what we need is some kind of social media campaigning that normalises people discussing openly with each other. (Virtual1)</i>	Social Influence Reinforcement Knowledge Beliefs about consequences	Education Persuasion Environmental restructuring Modelling Enablement Incentivisation	9.1 Credible source 6.2 Social Comparison 6.3 Information about others approval 6.1 Demonstration of a behaviour 12.5 Adding objects to the environment 3.1 social support (unspecified) 5.1 Information about health consequences 5.3 Information about social and environmental consequences. 12.2 Restructuring the social environment	Social media could be used within multimodal or multimedia interventions to <u>change norms (8.3, 6.3)</u> amongst men concerning PN and <u>promote particular norms</u> about engaging in PN with one off partners. This messaging would detail the diverse consequences of PN for all. (5.1, 5.3,6.2, 12.2) It may be possible to <u>recruit social influencers (9.1)</u> from the social media to promote PN and harness their networks to promote key messages. (5.1, 5.3, 12.5, 12.2) Social media could be used to show people taking part in PN (6.1) with one off partners and in this way <u>provide examples of how to do it</u> and reinforce norms that doing it is the right thing to do. (5.1, 5.3, 6.3) Using social networks and dating app sites to <u>exchange, repeat and 'share' messaging (12.5)</u> concerning PN with one off partners will reinforce the message and the norm of doing PN. (3.1, 5.1, 5.3)

							MSM receive focused consistent messaging about PN and the health and community impact (5.1, 5.3) from key sources such as HCPs, NGOs, dating sites, (9.1) physical and virtual hook up sites. These complimentary messages should be consistent, tailored to MSM sexual cultures and reinforced by a range of stakeholders. (5.3, 5.1)
--	--	--	--	--	--	--	--

Green content = Agreed in Phase 3 as final intervention content

Amber content = Content that could be considered for the intervention

Red content = Content to be dropped from the intervention

Table 4 What did Dating app providers tell us about implementing changes relating to promoting the endorsement of contact tracing within their infrastructure ?

Key component, step, or behaviour of the intervention derived from initial stakeholder event	Barriers to enacting this intervention component	Facilitators to enacting this intervention component	Indicative quote	TDF Domain	Intervention function (BCW)	Potential behaviour change techniques	What does the analysis tell us about potentially useful ways forwards in relation to potential intervention content?
DAPs feature PN educational content and message blasts co-created with NGOs within their on-line platforms		DAPS find it easier to feature PN health related messages when they are clear and memorable	I think it's a great idea. I think that one of the things that really sticks in my head, and I totally forgot about this sexually transmitted disease, which is syphilis. There was a great syphilis campaign [inaudible 17.41] long time no syphilis or something like that. And it was on t-shirts and I saw so many people that were, kind of... It was a jokey statement, but a punchy statement that people were willing to kind of dress themselves up with. And I think that that was a really cool way to kind of reach people as well. So, yes, a punchy statement that... I always say with my team, if we do something that people are willing to wear [inaudible 18.13], do you know what I mean, that's a step over and beyond, you know. You're willing to make a statement with your clothing. I think that's a massive thing, basically. So, yes, a punchy statement could work. Anything where there's an aspect of merchandise would be really	Environment, context and resource (DAPS) Professional social role and identity	Environmental Restructuring Education Persuasion	12.2 Restructuring the social environment 7.1 Prompts cues 12.5 Adding objects to the environment 9.1 Credible source 5.6 Information about emotional consequences	DAPS use intervention messages/material on PN developed in partnership with others (e.g. NGOs?) (9.1). This could follow a similar format to other successful campaigns such as 'long time no syphilis' 'U=U', with a clear focus on educating, enabling and empowering MSM (12.5). This should include easily identified branding, promotional material and dedicated websites on importance of PN and a catchy slogan aimed at grabbing attention with information and educational material. (12.5) DAPs display key PN educational messages similar to other sexual health campaigns syphilis, U=U. (12.5). The aim would be to raise awareness, put PN on the radar, get it out there and spoken about widely. (12.2) DAPs use promotional PN resources (co-produced with NGOs MSM) (9.1) through their educational initiatives and FAQ's (12.5) pages with links to the PN website and promotional material offering advice on the options available for contacting previous partners, this should include suggested wording or approaches to PN and how to respond to reduce stigma and blame if a previous partner

			cool, I think, but also, like I said, representation is really important. (DAP3)				contacts them. (7.1, 5.6)
DAPs feature PN educational content and message blasts co-created with NGOs tailored to specific STIs	DAPS find it difficult to feature PN health related messages on their app as they are not pitched right for the gay community		<i>remember we had this horrible, horrible syphilis ad that our health department here in the [place] wanted to run. And it was so ridiculously stigmatising, and fear-based that I told them we wouldn't run it. You know, we do have standards and if we don't want to run it we won't run it. But I think sometimes it's as simple as that, right. It's as simple as saying this is the sort of things that we're... You know, and to that point, it wasn't like I emailed them back and sort of wrote a scheming email and said, you know, don't ever talk to us again. It was more like, this isn't going to work, let's figure out, you know, what's going to work for the audience that you want to reach So, I think that there are ways that we can also sort of work more collaboratively on some of the messaging and work more collaboratively on, you know, frankly repairing some of that relationship damage that's sort of apparent between public health and the community. (DAP1)</i>	Environment, context and resource Social influences Beliefs about capabilities	Environmental restructuring Modelling Enablement Training Education Persuasion	12.2 Restructuring the social environment 12.5 Adding objects to the environment 9.1 Credible source 6.1 Demonstration of a behaviour 5.6 Information about emotional consequences 5.3 Information about social and environmental consequences	NGOs and MSM could work together to create educational initiatives detailing PN options for MSM with one-off partners (9.1) which are inclusive and pitched right for the gay community and freely available to all who want to use them. (12.5, 12.2) DAPs support PN education through their educational initiatives and FAQ's pages which detail advice on the options available for contacting previous partners (12.5) suggested wording or approaches to PN tips on how to respond to reduce stigma and promote a no-blame culture if a previous partner contacts them. (5.6, 5.3, 6.1) NGOs, HCPs, MSM and DAPS work together to devise educational materials, blogs, online educational content which covers specific STIs (9.1, 12.5, 12.2)
DAPs feature educational PN messages supported and endorsed by		DAPS find it easier to educate MSM using health related	<i>we find that we get the best results when we've got interesting people talking about sexual health. So, we do a lot of stuff with porn stars and sex workers and their perspective on</i>	Environment, context and resource Social	Environmental restructuring Education	12.2 Restructure the social environment 7.1 Prompts and	It may be possible to recruit social influencers from the social media to promote PN and harness their networks to promote key messages (9.1, 7.1, 12.2)

<p>those with social influence within their apps</p>		<p>messages if they pitched right for the gay community</p>	<p><i>sexual health issues is what's interesting from a content point of view but also our readers and users could think, oh I can see so and so is talking about the importance of getting tested, I'll read more about that, kind of thing. So, that's often where find the best bits, as opposed to a professional talking about it in a dry sense of the word. (DAP2)</i></p>	<p>influences</p> <p>Professional social role and identity</p>	<p>Persuasion</p>	<p>cues</p> <p>9.1 Credible source</p> <p>5.1 Information about health consequences</p>	<p>Using social networks to <u>exchange, repeat and 'share' messaging</u> concerning PN with one off partners will reinforce the message and the norm of doing PN (5.1).</p> <p>Dating Apps have a dedicated sexual health info/awareness page or section with FAQs on PN with suggestions and tips, these feature scripted and modelled PN examples with appropriate PN wording to encourage MSM to take more responsibility for PN (12.2, 5.1)</p>
<p>DAPs play a part in educating diverse MSM on the value of PN</p>	<p>DAPS find it difficult to feature PN related educational messages as they lack diversity thereby excluding some MSM</p>		<p><i>The problem that I always have with these kind of campaigns is about the diversity of the people that are putting out the message. So, it has to ensure... And I'm not just talking race, I'm not just talking age, I'm not just talking body type. I'm talking about the different cross-sections of the community as well. So, if we're just focusing on gay men or men who have sex with men, it's about making sure... And I'm really glad that you reached out to me because the fetish community aspect, they're very different to the general gay mainstream, do you know what I mean? So, Grindr will have a very different approach than we would have, for example. Say if there was a campaign that was on a higher level, I feel like every single gay person no matter where they fit in the community has to be able to look at that and realise that, okay, this is affecting me as well. So, it can't just</i></p>	<p>Professional social role and identity</p> <p>Beliefs about consequences</p> <p>Environment, context and resource</p>	<p>Environmental restructuring</p> <p>Education</p> <p>Persuasion</p> <p>Enablement</p>	<p>12.2 Restructure the social environment</p> <p>5.3 Information about social and environmental consequences</p> <p>9.1 Credible source</p> <p>5.1 Information about health consequences</p>	<p>Educational programmes aimed at MSM promoting PN with one-off partners should be inclusive, co-created with a diverse group of MSM and NGOs (9.1) to represent different 'tribes' or groups within the MSM community. (5.3, 12.2). Diverse educational messages could be co-created to reach a wide range of MSM (5.1)</p>

			<p><i>be, you know, white guy who basically dresses in H and M head to toe and is saying, oh, you should tell your partner [inaudible 16.09], you know, head to toe in tattoos and blah blah, or whatever it may be, so that it does reach a cross-section of the community, the whole entire thing. So, for me... And that's part of thing that I've been focusing on in the last couple of years, is about [disability 16.25] and representation. So, for me, it has to kind of represent not only a race and body type but also the different tribes, I hate that term, but the different tribes within the community as well. Because otherwise, I would just look at it or I would read it and I would be like, okay, that's great but I don't see myself in it. So, yes.</i></p> <p><i>(DAP3)</i></p>				
		<p>DAPs find it easier to endorse and feature PN health related messages when they are part of a national campaign co-created with NGOs</p>	<p><i>I think if there was a consensus that all of us, in the same way that World AIDS Day there is a consensus they're all going to focus and concentrate on this and highlight such an important issue, then I think that it would be a really good thing to kind of [put 14.16] around. And people would be more conscious of it. Maybe it's a little bit cheesy to kind of do another campaign like that, but I feel like then it would make people be aware at least.</i></p> <p><i>So, that's why, again, I kind of say, you know, the biggie is always HIV and</i></p>	<p>Environment, context and resource</p> <p>Social influences</p> <p>Professional social role and identity</p>	<p>Environmental restructuring</p> <p>Training</p> <p>Enablement</p> <p>Persuasion</p> <p>Modelling</p>	<p>12.2 Restructuring the social environment</p> <p>9.1 Credible source</p> <p>5.3 Information about social and environmental consequences</p> <p>5.1 Information about health consequences</p>	<p>Intervention messaging linked to a national awareness raising campaign could focus on key organisations (e.g., NGOs, DAPs) all consistently saying how blaming people who disclose an STI is not helpful and ultimately damaging to everyone. (9.1,,5.1, 5.3)</p> <ul style="list-style-type: none"> • Provide good examples of doing PN as 'the responsible thing to do' for everyone. (6.1 4.1) • This could normalise responsibility for others, set expectations of being responsible for others within communities. (5.3, 12.2) <p>Ideally such a campaign and accompanied</p>

			<p>AIDS, and I'm wondering... I'm not suggesting that we create a whole day about sexually transmitted things, but to kind of have almost like a worldwide consensus or campaign where we'll mark where we kind of focus on that and bring the attention to it. Because it would also enable apps, I think, generally to kind of focus campaigns around that specific month rather than it be the responsibility of charities or other kind of sexual clinics to kind of carry the torch in a sense. That was a really longwinded way to say that, how can we get the charity to basically provide us with content to send the message out to people about why it's so important that we talk about HIV prevention still to this day. (DAP3)</p>			<p>6.1 Demonstration of a behaviour</p> <p>4.1 Instruction on how to perform a behaviour</p>	<p>messaging would be co-produced by peers, NGO staff and HCPs (9.1)</p>
	<p>DAPS find it easier to run PN health related messages within their app as COVID 19 means they and their users understand the concept of test, protect and contact tracing better</p>	<p>I think as you said at the beginning, what makes this conversation, particularly easy for everyone to understand now, is the experience of being locked down, the test and trace discussion around Covid-19 and being able to illustrate how that connects to STIs. So, everyone is in a different place right now than if we'd been having this conversation 12 months ago. So, it's linking to that Covid-19 experience, must make selling this concept a lot easier. (DAP2)</p>	<p>Environment, context and resource (DAPs)</p> <p>Knowledge (MSM)</p>	<p>Environmental restructuring</p> <p>Enablement</p> <p>Education</p>	<p>12.2 Restructuring the social environment</p> <p>12.5 Adding objects to environment</p> <p>7.1 Prompts and cues</p> <p>5.3 Information about social and environmental consequences</p>	<p>In relation to <u>educating</u> men about PN catchy phrases and terms that build on recent developments from COVID and contact tracing might be helpful:- 4T's Test, Treat, Trace, Tell. This kind of work will <u>enable</u> MSM to tell more one-off partners (12.2, 12.5, 7.1)</p> <p>For men PN health related messages could draw parallels linking PN to COVID19 to reduce the stigma associated with STIs and educate MSM on PN persuading them to prioritise, prepare and undertake. (5.3, 12.2)</p>	
	<p>DAPs find it difficult to</p>		<p>Environment context and</p>	<p>Environmental restructuring</p>	<p>12.2 Restructuring the social</p>	<p>Consider opportunities to create partnership work (NGOs and public health bodies working</p>	

<p>prioritise PN health related messages as they are small businesses which lack resources</p>		<p><i>it's everyone, I mean we're a small organisation so there's only so many hours in the day and if you try to get information out of a charity to do something interesting, they're really stretched because they've got no resources either. So, no-one has got a big marketing team who can go off and create an amazing campaign.(DAP2)</i></p>	<p>resources</p>	<p>Enablement</p>	<p>environment 3.2 Social support practical 5.3 Information about social and environmental consequences</p>	<p>together, options for crowd funding, money from public health) (3.2). Collaboration across stakeholders is likely to be an effective means of harnessing unique strengths and weaknesses and sharing resource burdens. (5.3, 12.2)</p>
<p>DAPS find public health led partner notification difficult as there are issues with trust in public health providers</p>		<p><i>And I think that, you know, as we talk about contact tracing and then we think about the sort of history of partner notification, there's a really bad relationship there based on behaviour of public health in terms of gay men and gay men sexuality. So, it's understandable that people are going to be suspicious and people are going to be doubtful of the intentions. There's a history there and they're not approaching this from a sort of neutral standpoint....bias and homophobia.(DAP1)</i></p>	<p>Beliefs about consequences Environment context and resources</p>	<p>Education Persuasion Environmental restructuring Enablement</p>	<p>5.3 Information about social and environmental consequences 9.1 Credible source 12.2 Restructuring the social environment 9.3 Comparative imaginings of future outcomes</p>	<p>To enable DAPS to work collaboratively with other organisations (9.1) around PN, it would be helpful for those developing interventions to provide DAPS with information, and persuade them, about how it might benefit their business in the longer term (5.3, 9.3)</p> <p>Opportunities to create partnership work across a range of stakeholders are likely to be an effective means of harnessing unique strengths and weaknesses and sharing resource burdens (12.2).</p>

Green content = Agreed in Phase 3 as final intervention content

Amber content = Content that could be considered for the intervention

Red content = Content to be dropped from the intervention

Table 5 What did Dating app providers tell us about implementing changes relating to changing dating app functionality?

Key component, step, or behaviour of the intervention derived from initial stakeholder event	Barriers to enacting this intervention component	Facilitators to enacting this intervention component	Indicative quote	TDF Domain	Intervention function (BCW)	Potential behaviour change techniques	What does the analysis tell us about potentially useful ways forwards in relation to potential intervention content?
	DAPS find it difficult to offer a function on the app profile where MSM indicate they are positive about PN as it may not represent a genuine statement of intent		<i>So, what I'm worried about is even if you say that I'm, you know, welcoming of...tell me whether I've been exposed to something or I will tell people, it's not really a true indication as to whether someone will actually tell someone So, if you don't have on your profile, it's almost like you might be shamed into just putting on your profile and still not going through with it. Do you know what I mean? So, you've kind of ticked all the boxes to say that I am a really good person but when it comes down to it, you're not. Do you know what I mean? I don't know. I think it's an interesting thing. I just worry that it would be received in the wrong way. And I'll tell you another reason why. So, for World AIDS Day, we have a mechanic that basically puts a virtual ribbon on your profile. And once you've made a donation, you could put the ribbon on your profile, or if you want to spread the word about [inaudible 21.36] content is, you could put the [red 21.37] ribbon on your profile.</i>	Beliefs about consequences Environment context and resources	Modelling	5.3 Information on social consequences 9.3 Comparative imaginings of future outcomes 12.5 Adding objects to the environment	MSM indicating they are positive about PN somewhere on their dating profile is only likely to be effective if other measures such as education and interventions which promote attitudinal change are also in place

<p>DAPS offer a function to allow MSM whose partners have blocked them to reconnect</p>	<p>DAPs find it difficult to offer a function allowing previous partners to unblock as people are blocked for a reason</p>	<p>DAP3</p> <p>I think the problem is that say if I blocked you and you'd contracted something and you're trying to get in touch with me, even if you tried to do that, there's something about I don't want to receive anything more from you. So, to kind of take that away is a violation in that member's right to block. So, I think that would be a problem, but, again, there wouldn't be... If it was an anonymous, maybe there's something in... Someone you know. It doesn't really matter, I suppose, if they've been blocked or not, but I think to have received information from that specific person to say that they've contracted something, again, is around the right to block someone. And I think that would kind of violate it. I'd probably say most dating apps would say no. Yes, but I think that if there was some sort of mechanism to go over... But, again, I don't know how much... I don't even know how people store that data, to kind of say that these people don't talk to each other anymore. So, once they've kind of [inaudible 26.05] happened, some people keep it and some people kind of get rid of it. So, it all depends, really (DAP3)</p>	<p>Environmental context and resources</p>	<p>Environmental restructuring</p>	<p>12.1 Restructuring the physical environment</p> <p>12.2 Restructuring the social environment</p>	<p>Not amenable to change</p>
	<p>DAPS find MSM using anonymous text messages for PN difficult as</p>	<p><i>we've partnered years ago with, I want to say GMFO, I think, about something very similar to this. So, I think what we had before was a system that kind of</i></p>	<p>Beliefs about consequences</p>	<p>Environmental restructuring</p> <p>Enablement</p>	<p>9.3 Comparative imaginings of future outcomes</p>	<p>Use of anonymous text messages have limited impact as take up rates are likely to be low amongst MSM, research from previous trials shows they are unlikely to offer increased PN</p>

previously the take up was low			<p><i>plugged into ours which enabled you to... If you had contracted an STI, you could send out basically a private message to people that you may have infected on xxxxx.</i></p> <p><i>And it was all meant to be private and all that kind of stuff, but it didn't really get used, and I'm not really sure why. And I think it's mostly to do with the fact that, as you probably know, there is a lot of shame within the gay community anyway. And I think that as much as maybe it's a private message, it wasn't private message, you can still kind of work out where and who it might have come from. So, I think people were still kind of worried of that aspect of things. So, that's why it didn't really work that well probably, and we actually ended up removing it completely (DAP3)</i></p>			<p>12.1 Restructuring the physical environment</p> <p>12.5 Adding objects to the environment</p>	<p>due to lack of use.</p>
--------------------------------	--	--	--	--	--	--	-----------------------------------

Green content = Agreed in Phase 3 as final intervention content

Amber content = Content that could be considered for the intervention

Red content = Content to be dropped from the intervention

Table 6 What did Dating app providers tell us about implementing changes relating to signing up to a code of conduct to endorse contact tracing ?

Key component, step, or behaviour of the intervention derived from initial stakeholder event	Barriers to enacting this intervention component	Facilitators to enacting this intervention component	Indicative quote	TDF Domain	Intervention function (BCW)	Potential behaviour change techniques	Suggestions for intervention content
DAPs agree and sign up to a code of conduct supporting partner notification		DAPs find it easier to sign up to a code of conduct if the right organisations work together	A code of conduct around partner notification, you would need someone to bring it together, in terms of a charity working in the sector or someone who could get the right people at the table and then you could possibly...I would keep it really simple, I mean the challenge would be agreeing what are the standards that we're going to offer to users and what's the technology required to get there. Everyone would have to deal with their own technology because everyone runs their own platforms. So, it would be fairly broad and fairly general, a statement of intent, as opposed to anything else, I imagine. you could even do it where you can sign up to undertake to deliver opt-in partner notification by 2027 or	Environment context and resources Professional social role and identity Reinforcement	Environmental restructuring, Persuasion Enablement Incentivisation	12.2 Restructuring the social environment 3.2 Social support (practical) 12.5 Adding objects to the environment 1.8 Behavioural contract	NGOs MSM and public health organisations Lobby for code of conduct for dating app providers supporting PN. (3.2, 12.2) Consider the development of an NGO and dating app led code of conduct (12.5). Key actors, i.e. NGOs, DAPs agree standards and co-produce a DAP code of conduct to support partner notification (12.5, 1.8)

			whatever, that kind of thing. So, you commit to being on a road map towards enabling that notification.(DAP2)				
--	--	--	---	--	--	--	--

Green content = Agreed in Phase 3 as final intervention content

Amber content = Content that could be considered for the intervention

Red content = Content to be dropped from the intervention

Table 7 What have we learned about implementing improved interactions between HCPs and GBMSM to enhance contact tracing for one-off partners

Key component, step, or behavioural aspect of the intervention derived from initial stakeholder event	Barriers to enacting this intervention component	Facilitators to enacting this intervention component	Indicative quote	TDF Domain	Intervention function (BCW)	Potential behaviour change techniques	What does the analysis tell us about potentially useful ways forwards in relation to potential intervention content?
MSM play an active role in PN activities for their one-off partners by preparing for PN ahead of their appointment	MSM find it difficult to undertake PN in clinic settings due to perceived lack of appointment time		<i>I find it difficult when I'm in one of these...one of the appointments, at [clinic] for example, and they say...they usually routinely ask you to talk...how many of your partners could we inform? I find that very difficult to say that in ten seconds. So would be good to have, like, a...if they sent you a link or something where you can send them back the number or a Grindr profile or whatever. (London)</i>	Environmental context and resources Knowledge	Enablement Environmental Restructuring Education	3.2 Social support practical 11.2 Reduce negative emotions 12.5 Adding objects to the environment 5.3 Information about social and environmental consequences 5.1 Information about health consequences 7.1 Prompts and cues 5.6 Information about emotional consequences	NGOs should support, educate NGO led educational programmes could include a focus on reducing STI related stigma (11.2, 5.3, 5.1, 5.6) Work should focus on priming MSM to attend sexual health appointments prepared for PN. (3.2). NGOs could support, educate and raise awareness of the importance of PN emphasising that whenever possible MSM should be aware of the importance of being prepared for PN, bringing contacts and/or smart phone to consultations (7.1, 12.5) and a commitment to contact as many partners as possible (3.2).

<p>HCPs motivate GBMSM to engage in contact tracing</p>		<p>MSM find it easier to notify one-off partners when HCPs use counselling skills to motivate and encourage PN behaviours</p>	<p><i>It's them basically saying, well, look, there's always going to be a risk that you can catch an STI, but you're being really responsible here by actually coming to get tested, building up the courage to be tested, and actually by telling the people that you've slept with and had sex with, actually that's not a bad thing, that's a really good thing, because actually you're helping them as well as yourself, so it's kind of getting that person on side. (Glasgow)</i></p> <p><i>Yeah. Why I said that is clinics as with anything in the NHS and everywhere are on a limited budget, limited amount of staff. If they're going to sit and counsel each person for ten minutes, that means somebody else is waiting 20 minutes more just to go in and get tested. So they're paying the price for the other person sitting there needing to be counselled. And I've been in to a clinic where they talk and they talk, and I'm thinking, I don't really want to hear all this. I'm here, I've just had a couple of pints in the pub, (London)</i></p>	<p>Professional social role and identity</p> <p>Environmental context and resources</p> <p>Beliefs about consequences</p> <p>Beliefs about capabilities</p>	<p>Enablement</p> <p>Persuasion</p> <p>Training</p> <p>Education</p>	<p><i>3.1 Social support</i></p> <p><i>3.3 Social support emotional</i></p> <p><i>6.1 Demonstration of a behaviour</i></p> <p><i>9.1 Credible source</i></p> <p><i>5.1 Information about health consequences</i></p>	<p>HCPs could use motivational counselling type skills (3.1) to encourage PN, persuade and support MSM in a non-judgemental way to consider PN and discuss options and MSMs preferences (3.3) on how best to contact one-off partners who may be at risk of an STI. (3.1)</p> <p>The value of peer training for HCPs (9.1) could be used to cascade MI skills. Training could be focused on skills based motivational interviewing for HCAs using scenarios linked to PN outcomes (6.1) and the difference PN makes to health outcomes for MSM with one-off partners (5.1)</p>
---	--	--	--	---	--	--	---

<p>Health care professionals use opportunities in clinic waiting rooms, webpages to educate MSM about PN particularly for their one-off partners</p>	<p>HCP find it difficult to display MSM specific PN messages in waiting areas as this might not be the most appropriate venue</p>	<p>MSM find PN easier to notify one-off partners when key educational messages are displayed in clinic waiting areas</p>	<p><i>when you're waiting in the waiting room, the TV screen, they can do it (educate) that way as well, while you're waiting you just look at the screen for helpful messages tips, it maybe helps somebody helps get the message out (London)</i></p>	<p>Environmental context and resources Beliefs about capabilities Beliefs about consequences</p>	<p>Environmental restructuring Education Enablement Persuasion</p>	<p><i>12.1 Restructuring the physical environment</i> <i>3.2 Social support practical</i> <i>7.1 Prompts/ cues</i> <i>5.1 Information on health consequences</i> <i>5.3 Information on social and environmental consequences</i> <i>12.5 Adding objects to the environment</i></p>	<p>Clinic based initiatives could re-structure clinic waiting areas and webpages (12.1) to include education on the consequences of STIs and importance of partner notification (3.2) to motivate more MSM to play an active part in PN.</p> <p>Educational PN messages with links to health and community consequences (5.1, 5.3) of good PN could be linked to how MSM get their results (12.1) to inform and persuade MSM. referring to Prep (i.e. it can be done because something highly similar has been done). This could include links to an online message or website 'now you have your results what should you do next (3.2)'. Education for MSM should highlight FAQ with an educational resource which could be viewed by MSM in their own private space as opposed to a busy clinic environment. Educational messages with prompts and cues on different PN options and how to tell partners i.e. suggested options to re-contact, suggested wording could be used to foster a culture of peer support (7.1).</p> <p>Widely available PN educational messages could be</p>

<p>Health care professionals make clinics welcoming and easy to use</p>		<p>MSM find it easier to attend sexual health clinics and participate in PN for one off partners if clinic welcoming with a professional but relaxed atmosphere</p>	<p><i>Some like [clinic] is really easy, it's really inviting and welcoming. (Glasgow)</i></p> <p>So, until I was active in my HIV care, I was mainly being seen by health care professionals in hospitals, in GUM clinics, which aren't particularly pleasant places to be. They always seem to be tucked away at the ***-end of the hospital, and you know, everybody knows why you're going there, it's all just this kind of strange subterfuge going on. But the people working there are lovely, it's just the way it feels. And certainly, when I got (clinic name) and started experiencing, like a different setup, just the way they make the waiting rooms look, and the atmosphere, and the ambience, is a little bit less clinical. And so, you start to feel you're being treated more like a human being. (Virtual focus group 2)</p>	<p>Environmental context and resources</p>	<p>Environmental restructuring, Enablement</p>	<p>12.1 Restructuring the physical environment</p> <p>12.2 Restructuring the social environment</p> <p>11.2 Reduce negative emotions</p>	<p>devised by NGOs and HCPs (5.1) these messages could be available to MSM through NGO's, DAPs and agreed appropriate clinic based routes. Standardised educational messages (punchy and easy to remember) (7.1) should be available to motivate and remind MSM on the importance of PN in a variety of formats to reinforce and maximise impact (12.5).</p> <p>HCPs should promote a positive atmosphere being mindful of what the waiting room area looks like. Feels like. All HCPs could take steps to ensure clinical environments are as welcoming as possible (12.1, 11.2) with a focus on promoting a positive atmosphere for both in-person and online appointments where those attending are supported to feel as relaxed as possible (12.2)</p>
---	--	--	--	--	--	---	--

MSM are given clear and consistent messages about PN from sexual health clinic staff		MSM find it easier to notify one-off partners as they get consistent clear messages about PN from sexual health clinics	I think, the attitude of the people at those more progressive, and welcoming clinics, are a bit more supportive in terms of the actual, like, practical need for support, to do the disclosure bit. Because I think, obviously, many people are committed to the disclosure bit, and it's your health care professionals, the way they encourage you, or however. (virtual focus group2)	Environmental context and resources Social Influences	Training, Enablement Environmental restructuring Education	<p><i>5.1 Information about health consequences</i></p> <p><i>3.1 Social support</i></p> <p><i>3.3 Social support emotional</i></p> <p><i>2.7 Feedback on outcomes of behaviour</i></p> <p><i>12.5 Adding objects to the environment</i></p> <p><i>7.1 Prompts/ cues</i></p> <p><i>4.1 Instruction on how to perform a behaviour</i></p> <p><i>12.2 Restructuring the social environment</i></p>	<p>HCPs inform (5.1, 2.7) and prompt (7.1) MSM about the importance of PN at the earliest opportunity ideally when diagnosis revealed or before. HCPs are encouraged through PN related education and guidelines to encourage MSM to participate in PN using a non-judgemental approach (4.1). HCPs offer consistent and clear information around the health benefits and importance of PN (5.1) with all clinic staff offering clear and consistent messages on the public health benefits and variety of PN options available to MSM (3.1).</p> <p>Clinics to explore role of NGOs who could play a part in harder to do PN fostering a culture of peer support to persuade MSM to complete PN. Consider opportunities to develop a system based on functionality and learning from RELAY. (12.5, 12.2)</p>
HCPs and MSM access smart phones together to locate partners	10.1 MSM find it difficult to access their smart phone with HCPs due to fear of being judged and concerns re privacy	10.2 MSM find it easier to notify one-off partners when HCPs and MSM access smart phones together	<i>I don't know. Maybe it's my judgemental side coming out, but if I was sitting down with like...you know, with someone like a healthcare professional at an STI...like at a health clinic, and it's like maybe someone my mum's age, and I'm sitting looking through my phone with her, you know, people in my phone, and all of a sudden a message flashes up or...I</i>	Environmental context and resources Knowledge	Enablement Environmental restructuring Education	<p><i>3.2 Social support practical</i></p> <p><i>7.1 Prompts/ cues</i></p> <p><i>12.5 Adding objects to the environment</i></p> <p><i>3.3 Social support emotional</i></p> <p><i>5.3 Information about social and</i></p>	<p>NGO led educational programmes could focus on priming MSM (3.2, 7.1), to attend sexual health appointments prepared for PN, bringing contacts and/or smart phone to consultations (12.5)</p> <p>Where MSM and HCPs access smart phones together for PN purposes this is likely to be enhanced when HCPs clearly demonstrate a non-judgemental attitude (3.3) and acknowledge and accept the likelihood of high numbers of sexual partners. (3.3)</p>

<p>HCPs record dating app profile names within MSMs case notes to be used for later PN purposes if required</p>	<p>Health care professionals recording dating app profile names would be difficult because profile names change</p>	<p><i>don't know, I've just got this image in my head of this woman like...(Glasgow)</i></p> <p><i>'well I don't have that person's number, but actually, we hooked up on Grindr, so I have that person's profile. So, you need options, like how can, the idea is to make it as easy as possible, and you can, you know, get through to as many people as possible. So that's why I'm thinking of, you know, all these different tools, all these different ways, you know, to make it easier to just inform people. All these things, you know, they might look a bit awkward at first. (virtual 1)</i></p> <p><i>I don't know how this really works in a time where you can easily block someone or change your details or change your name (Glasgow)</i></p> <p><i>You can actually have five profiles per phone because they know the IP address, so they allow you give maximum profiles from one particular phone, using different emails. (Leeds)</i></p>	<p>Environmental context and resources</p>	<p>Enablement</p>	<p><i>environmental consequences</i></p> <p><i>5.1 Information about health consequences</i></p>	<p>For a number of reasons this intervention is not likely to be effective or one we will take forward, important to acknowledge this with any stakeholder presentations on communication.</p>
---	--	---	--	-------------------	--	---

Figure 2 Schematic overview of the diverse systemic drivers effecting contact tracing, including pathogenic (red) and salutogenic (green) influences collated from the initial stakeholder event.




Figure 3 Selection of key areas to examine further within Phase 2




Figure 4 An overview of intervention ideas after detailed consideration of how to implement them and taken to Phase 3




Figure 5 An overview of intervention ideas after final appraisal and modification with stakeholder groups

