

1 **Preventable deaths from SARS-CoV-2 in England and Wales: a systematic**
2 **case series of coroners' reports during the COVID-19 pandemic**

3
4
5 *Bethan Swift, MSc^{1,2*}, Carl Heneghan, DPhil^{3,4}, Jeffrey K. Aronson, DPhil³, David J. Howard,*
6 *DPhil^{4,5}, Georgia C. Richards, DPhil^{3,4}*

7
8 ¹Oxford Endometriosis CaRe Centre, Nuffield Department of Women's and Reproductive
9 Health, University of Oxford, Oxford OX3 9DU, UK.

10
11 ²Wellcome Centre for Human Genetics, University of Oxford, Oxford, OX3 7BN, UK.

12
13 ³Centre for Evidence-Based Medicine, Nuffield Department of Primary Care Health Sciences,
14 University of Oxford, Radcliffe Observatory Quarter, Woodstock Road, Oxford OX2 6GG, UK.

15
16 ⁴Global Centre on Healthcare and Urbanisation, Kellogg College, University of Oxford, 60-62
17 Banbury Road, Oxford OX2 6PN, UK.

18
19 ⁵Department for Continuing Education, University of Oxford, UK.

20
21 *Corresponding author:

22 Bethan Swift, MSc

23 Email: bethan@well.ox.ac.uk

24 Tel: +44 7921527208

25 Affiliations and address: Oxford Endometriosis CaRe Centre, Nuffield Department of Women's
26 and Reproductive Health, University of Oxford, Oxford OX3 9DU, UK. Wellcome Centre for
27 Human Genetics, University of Oxford, Oxford OX3 7BN, UK.

28
29
30 **Word count: 3110**

34 **Abstract**

35

36 **Objectives** To examine coroners' Prevention of Future Deaths reports (PFDs) to identify deaths
37 involving SARS-CoV-2 that coroners deemed preventable.

38

39 **Design** Consecutive case series.

40

41 **Setting** England and Wales.

42

43 **Participants** Patients reported in 510 PFDs dated between 01 January 2020 and 28 June 2021,
44 collected from the UK's Courts and Tribunals Judiciary website using web scraping to create an
45 openly available database, <https://preventabledeathstracker.net/>.

46

47 **Main outcome measures** Concerns reported by coroners.

48

49 **Public and Patient Involvement** Patients and members of the public were not involved in this
50 study.

51

52 **Results** SARS-CoV-2 was involved in 23 deaths reported by coroners in PFDs. Twelve deaths
53 were indirectly related to the COVID-19 pandemic, defined as those that were not medically
54 caused by SARS-CoV-2, but were associated with mitigation measures. In 11 cases the coroner
55 explicitly reported that COVID-19 had directly caused death. There was geographical variation
56 in the reporting of PFDs; most (39%) were written by coroners in the North-West of England.
57 The coroners raised 56 concerns, problems in communication being the most common (30%),
58 followed by failure to follow protocols (23%). Organizations in the National Health Service (NHS)
59 were sent the most PFDs (51%), followed by the Government (26%), but responses to PFDs by
60 these organizations were poor.

61

62 **Conclusions** PFDs contain a rich source of information on preventable deaths that has
63 previously been difficult to examine systematically. Our openly available tool
64 (<https://preventabledeathstracker.net/>) streamlines this process and has identified many
65 concerns raised by coroners that should be addressed during the Government's inquiry into the
66 handling of the COVID-19 pandemic, so that mistakes made are less likely to be repeated.

67

68 **Study protocol pre-registration:** <https://osf.io/bfypc/>

69

70

71 **Summary box**

72

73 **What is already known about this subject?**

- 74 • The UK Government has stated that there will be a public inquiry into the handling of the
75 COVID-19 pandemic, to learn lessons for future pandemics.
76 • Coroners in England and Wales have a duty to report and communicate information
77 about the deaths they investigate when the coroner believes that action should be taken
78 to prevent future deaths.
79 • These reports, called Prevention of Future Death reports (PFDs), had not yet been
80 systematically analysed to identify deaths that occurred during the COVID-19 pandemic.
81

82 **What are the new findings?**

- 83 • We created the Preventable Deaths Database (<https://preventabledeathstracker.net/>)
84 using web scraping to systematically assess PFDs published on the Courts and Tribunal
85 Judiciary website.
86 • Between 01 January 2020 and 28 June 2021, one in 20 (4.5%, n=23) PFDs that were
87 published by coroners involved SARS-CoV-2.
88 • Coroners raised many concerns about the care of patients in hospitals, care homes, and
89 people in the community during the COVID-19 pandemic, which require action to prevent
90 future deaths.
91

92 **How might it affect clinical practice in the foreseeable future?**

- 93 • Preventable deaths that occurred during the COVID-19 pandemic should be referred to
94 the coroner so that an inquest can be performed and a PFD issued, highlighting actions
95 that could be avoided in improving the handling of future pandemics in both the UK and
96 elsewhere.
97 • Our tool, <https://preventabledeathstracker.net/>, can be used by others to examine
98 preventable deaths in England and Wales, and to identify signals for quality
99 improvement to reduce avoidable harms in clinical practice.
100

101 **Introduction**

102

103 Over five million deaths worldwide have been attributed to the severe acute respiratory
104 syndrome due to coronavirus 2 (SARS-CoV-2)¹; some deaths may have been preventable.

105

106 In England and Wales, causes of deaths are investigated by coroners during an inquest, unless
107 the death is natural or referred to the criminal court. Under UK regulations, coroners have a duty
108 to report and communicate information about the deaths that they investigate when they believe
109 that actions should be taken to prevent similar deaths^{2,3,4}. Such reports, previously called Rule
110 43 reports, are named Prevention of Future Deaths reports or PFDs. Despite these regulations,
111 there is no formal system in place for auditing or systematically analysing PFDs, so concerns
112 raised by coroners in such reports may go unrecognized and unreported, reducing the impact of
113 the PFD system.

114

115 The PFD system has three processes: 1) coroners write PFDs after the inquest and send
116 reports to those with the power to act; 2) addressees respond to coroners regarding the
117 concerns raised in PFDs within 56 days; and 3) actions to prevent such deaths are proposed
118 and ideally implemented. In December 2020, a series was launched in BMJ Evidence Based
119 Medicine to disseminate PFDs that highlight lessons for clinical practice and policy⁵. Case
120 reports in the series have identified deaths from ingesting alcohol-based hand sanitizer,
121 misdiagnosed adverse drug reactions, problems with repeat opioid prescriptions, and fire
122 hazards from emollient products⁶⁻⁹. Case series of PFDs have also been conducted to
123 investigate preventable deaths from medicines and misused drugs, suicides, cardiovascular
124 disease, and cycling¹⁰⁻¹³.

125

126 During the COVID-19 pandemic, healthcare professionals in England and Wales called for the
127 deaths of their colleagues to be reported to coroners and for PFDs to be issued^{14,15}. However,
128 PFDs issued during the COVID-19 pandemic have not been analysed. We therefore aimed to
129 systematically analyse PFDs in which SARS-CoV-2 was directly or indirectly implicated in a
130 death.

131

132 **Methods**

133

134 We extracted a series of cases from the Courts and Tribunals Judiciary website and analysed
135 them according to a study protocol that was preregistered on an open repository¹⁶.

136

137 **Data collection**

138 PFDs are published on the Courts and Tribunals Judiciary website¹⁷. We used web scraping to
139 systematically collect all published PFDs and created a searchable database, the Preventable
140 Deaths Database¹⁸, which can be efficiently updated. The code for the web scraper is openly
141 available on GitHub and the methods have been previously described^{19,20}. The Preventable
142 Deaths Database contains the case reference number; the date of the report; the name of the

143 deceased; the coroner's name; the coroner's jurisdiction; the category of death (as assigned by
144 the Chief Coroner's office); to whom the report was sent; and the URL to the Judiciary website.

145 **Eligibility of cases**

146 We screened all 510 PFDs in the Preventable Deaths Database, dated from 01 January 2020 to
147 28 June 2021, for cases that mentioned COVID-19 as a direct or indirect cause of death. Direct
148 deaths were defined as those that the coroner explicitly attributed to COVID-19 as a cause of
149 death or were associated with a positive test for COVID-19 within 28 days of death. Indirect
150 deaths were defined as those that were not medically caused by COVID-19, but where coroners
151 mentioned that the associated mitigation measures during the COVID-19 pandemic (for
152 example, missed appointments due to lockdown), had contributed to the death. Cases that did
153 not mention SARS-CoV-2 or an effect of the COVID-19 pandemic were excluded.

154 **Data extraction**

155 For included cases, one study author (BS) manually extracted the following variables into a
156 predesigned Google Sheet, which was cross-examined by another study author (GCR): the
157 individuals or organizations to whom reports were sent and who responded; date of death; age;
158 sex; setting or location of death; medical cause(s) of death; the coroner's conclusion(s) of the
159 inquest; relevant medical, mental health and social history; substance(s) implicated in the death
160 and the type of substance(s); and the coroner's concerns. The data available for extraction were
161 limited by the information reported by coroners in the PFDs.

162 **Data analysis**

163 We used descriptive statistics to describe the numbers and types of cases that met the eligibility
164 criteria for inclusion. The numbers and types of individuals and organizations who received
165 PFDs were synthesized and response rates to coroners were calculated. A response rate of
166 100% meant that the individual or organization responded to all published PFDs issued by
167 coroners. Once all concerns were extracted, one study author (BS) read through all the
168 concerns and used content analysis to count and classify each concern inductively²¹.

169 **Software**

170 We used Tableau to present the coroners' concerns visually and Data Wrapper to
171 geographically map the number of PFDs reported in each region of England and Wales.

172 **Patient and public Involvement**

173 Patients and the public were not involved in this study.

174

175 **Results**

176 Figure 1 shows the results of the extraction process. Of 510 PFDs reported on the Judiciary
177 website between 01 January 2020 and 28 June 2021, 23 (4.5%) were attributed to COVID-19
178 and deemed preventable by coroners (Table 1).

179 **Table 1:** Summary of 23 deaths involving SARS-CoV-2 as reported in Prevention of Future Death reports in England and Wales between 1
 180 January 2020 and 28 June 2021, ordered by causes and date of death (created by the authors)

Age	Sex	Dates			Causes of death	Responses to PFDs	
		Death	Inquest	Report		Addressee(s)	Date of reply [‡]
Directly-related to COVID-19							
-	M	13/04/2020	14/04/2020	01/12/2020	1) Community-acquired pneumonia 2) COVID-19 3) Dementia, chronic obstructive pulmonary disease, asbestos-related pulmonary fibrosis, pleural plaques, type 2 diabetes	1) CQC 2) Vicarage Residential Care Home 3) Public Health England 4) NHS England 5) Greater Manchester Health and Social Care Partnership	1) 04/02/2021 2) Undated 3) 26/01/2021 4) 02/03/2021 5) 19/02/2021
86	F	17/04/2020	03/07/2020	11/12/2020	Natural causes and COVID-19	Whipps Cross Hospital	Received but not dated
74	M	21/04/2020	22/04/2020	01/12/2020	1) Hypovolemic shock 2) End-stage kidney disease 3) Polyneuropathy, frailty and COVID-19	1) Department of Health and Social Care 2) Royal London Hospital	Not yet received [^]
-	M	21/05/2020	26/05/2020	09/12/2020	1) COVID-19 pneumonia 2) Right sided neck of femur fracture, hypertension, atrial fibrillation	1) Public Health England 2) NHS England	1) 11/02/2021 2) 09/02/2021
18	M	31/07/2020	03/08/2020	30/03/2020	Drowning to which COVID-19 and asthma were contributory	1) Craven District Council 2) Yorkshire Dales National Park 3) Yorkshire Water	Not yet received [^]
-	M	07/09/2020	07/09/2020	24/04/2021	1) Bronchopneumonia in combination with COVID-19 2) Falls with vertebral fractures, type 2 diabetes mellitus, pulmonary fibrosis, heart failure and epilepsy	1) Greater Manchester Health and Social Care Partnership 2) NHS England	1) 07/09/21 2) 02/07/21

-	M	15/11/2020	16/11/2020	11/06/2021	1) Covid 19 Pneumonitis 2) Chronic obstructive pulmonary disease, ischaemic heart disease, previous right upper lobe resection for lung adenocarcinoma, type 2 diabetes mellitus	Tameside CCG	24/06/2021
90	M	28/01/2021	08/02/2021	23/04/2021	1) COVID-19 pneumonia 2) Dementia, heart failure, acute on chronic subdural haematoma, fall	Medway Maritime Hospital	07/06/2021
-	M	05/02/2021	08/02/2021	10/06/2021	1) COVID-19 on background of immunomodulatory treatment 2) Seborrheic eczema 3) Peripheral vascular disease	1) NHS England 2) Secretary of State of Health	Not yet received [^]
-	M	03/04/2021	17/11/2020	14/06/2021	1) Aspiration pneumonia on a background of a choking incident, COVID-19 pneumonitis 2) Alzheimers dementia	1) MHRA 2) NHS Stockport CCG	1) 27/07/21 2) 06/08/21
-	M	-	30/10/2020	19/02/2021	COVID-19 pneumonitis	1) Brighton Sussex University NHS Hospital Trust 2) West Sussex NHS Hospital Trust 3) Medico-Legal	1) 19/03/2021 2) Not yet received [^] 3) Not yet received [^]
Indirectly-related to COVID-19							
32	F	19/03/2020	30/03/2020	23/11/2020	1) Hanging 2) Bipolar affective disorder	1) Sussex Partnership Foundation NHS Trust 2) Brighton and Hove City Council	1) 10/02/2021 2) 10/02/2021
-	M	17/04/2020	24/04/2020	07/12/2020	Methadone toxicity	1) Public Health England 2) Haverhill Pharmacy	1) 13/01/2021 2) Undated
-	M	24/04/2020	05/05/2020	19/11/2020	Suicide	1) Woolwich Station Medical Centre 2) Ministry of Defence	1) Not received [^] 2) 16/02/2021
-	F	20/06/2020	23/06/2020	11/02/2021	1) Bronchopneumonia 2) Frailty 3) Dementia 4) Hypertension	1) CQC 2) Department of Health and Social Care	1) 04/06/2021 2) 03/06/2021

					5) Fractured neck of femur		
28	M	10/08/2020	14/08/2020	15/03/2021	Suicide	Sussex Partnership NHS Foundation Trust	Not yet received [^]
76	F	28/09/2020	03/12/2020	16/12/2020	Atherosclerosis and complete blockage of one artery	1) NHS Pathways* 2) COVID-19 Pandemic Response Service	10/02/2021
-	M	25/10/2020	26/10/2020	02/06/2021	Combined drug toxicology	Stockport CCG	07/07/21
77	M	-	08/10/2020	02/02/2021	1) Advanced dementia 2) Fractured neck of femur 3) Ischaemic heart disease	1) Adult Social Services, Norfolk County Council 2) Norfolk and Norwich University Hospital	1) 11/03/2021 2) 09/04/2021
88	M	-	20/08/2020	05/02/2021	1) Bronchopneumonia 2) Heat stroke 3) Dehydration	Care Outlook Ltd	18/04/2021
-	M	-	11/08/2020	07/05/2021	1) Small bowel obstruction and perforation 2) Ingestion of foreign body	Norfolk and Norwich University Hospital NHS Foundation Trust	23/07/21
68	F	-	20/09/2020	14/12/2020	1) Pneumothorax 2) Rib fractures 3) Fall 4) COPD and IHD	West Midlands Ambulance Service	08/01/2021
87	M	-	08/10/2020	07/05/2021	1) Urosepsis 2) Long term indwelling catheter not changed since October 2019 3) Alzheimer's dementia, cerebrovascular accident, chronic kidney disease, bladder cancer, prostate cancer	Lower Clapton Group Practice	24/06/21

181 *NHS Digital responded on behalf of NHS Pathways and the COVID-19 Pandemic Response Service. [‡]Recipients of PFDs have 56 days from the date of report to
182 respond to the coroner under Regulation 29 of The Coroners (Investigations) Regulations 2013. [^]replies were still overdue on 18/10/21; however not all replies are
183 posted on the Judiciary website.
184 CCG: Clinical Commissioning Group; CQC: Care Quality Commission; COPD: Chronic Obstructive Pulmonary Disease; MHRA: Medicines and Healthcare
185 products Regulatory Agency; NHS: National Health Service. Deaths directly related were defined as those that the coroner explicitly reported COVID-19 as a
186 cause of death or a positive test for COVID-19 within 28 days of death. Indirectly related deaths were defined as those that were attributed to mitigation measures
187 during the COVID-19 pandemic.

188 Most of the deaths (78%; n=18) occurred in men, and the median age at death was 76 years
189 (IQR: 50–87 years; range: 18–90 years; n=11). The deaths occurred between 13 April 2020
190 and 03 April 2021, but the date of death was unreported in six cases.

191 **Causes of death**

192 Eleven deaths (48%) were directly related to COVID-19, defined as those for which the
193 coroner explicitly reported COVID-19 as a cause of death or that were associated with a
194 positive test for COVID-19 within 28 days of death. Risk factors for death from COVID-19
195 included type 2 diabetes mellitus, kidney disease, hypertension, heart failure, and epilepsy
196 (Table 1). Pneumonitis co-occurred in 64% of deaths (n=7). One death was attributed to
197 drowning, to which COVID-19 and asthma contributed. In one-third (36%; n=4) of the direct
198 deaths, patients contracted COVID-19 as in-patients for other reasons, and another
199 contracted COVID-19 while a care-home resident. One man's symptoms of COVID-19 were
200 undiagnosed during a telephone appointment and untreated, resulting in death. In four cases
201 (36%) it was unclear how the deceased had contracted COVID-19.

202 Twelve deaths (52%) were indirectly related to COVID-19, defined as those that were
203 attributed to mitigation measures during the pandemic. There were three suicides and two
204 cases of inappropriate prescription of medications during lockdown. Two deaths occurred
205 because the deceased refused to go to hospital or a care home, against medical advice,
206 owing to fears of COVID-19. A further two deaths occurred from complications with medical
207 devices when medical appointments were cancelled owing to lockdown and inadequate
208 follow-up; a catheter was not changed in one case, causing urosepsis, and a salivary bypass
209 tube was accidentally left inside the patient, where it perforated the small bowel. In two
210 cases, deaths were attributed to undiagnosed medical problems after incorrect diagnoses of
211 COVID-19 via remote telehealth appointments. One death was due to natural causes,
212 contributed to by several falls and a fractured neck of femur in a patient with advanced
213 dementia.

214 The 23 deaths were classified into nine groups by the Chief Coroner's Office; hospital-
215 related (28%; n=11), community healthcare (18%; n=7), care homes (13%; n=5), other
216 (13%; n=5), emergency services (10%; n=4), alcohol, drugs of abuse, and medications (8%;
217 n=3), mental health related (5%; n=2), suicide (3%; n=1), and service personnel related (3%;
218 n=1).

219 **Geographical variation**

220 Twelve coroners across 12 jurisdictions wrote PFDs relating to COVID-19. Most were written
221 by coroners in the North West of England (39%; n=9), followed by the South East (22%;
222 n=5) and London (17%; n=4) (Figure 2; Supplementary Table 1). Coroners in the South
223 West, North East, West Midlands, and Wales did not report any deaths deemed preventable
224 from COVID-19.

225 **Coroners' concerns**

226 The coroners raised 56 individual concerns in the 23 PFDs. We categorized them into 28
227 groups and five higher-order categories (Figure 3; Supplementary Table 2). Poor
228 communication was reported in one-third of PFDs, followed by failure to follow protocols

229 (23%), lack of education and training (19%), lack of resources (16%), and safety concerns
230 (12%).

231 1) *Communication*

232

233 Most (42%, n=8) concerns involved poor communication, followed by failure to keep
234 accurate medical records/care plans (n=5) and failure to consider how the COVID-19
235 pandemic would affect appointments (n=3).

236 **Example 1:** *An elderly woman residing in a care home, at high risk of falls, became unwell.*
237 *Her GP remotely diagnosed suspected COVID-19, and she was isolated in her room, with*
238 *staff observation and sensor mats to ensure her wellbeing. She fell while unobserved and*
239 *was admitted to hospital with a fractured neck of femur, bronchopneumonia, and possible*
240 *COVID-19. A test for SARS-COV-2 was negative. She was unfit for surgery, deteriorated,*
241 *and later died in hospital from bronchopneumonia. The care home had a risk plan that*
242 *stipulated that she was to be observed during the day in communal areas. However, the*
243 *home was not staffed to provide one-to-one observation for residents in self-isolation.*

244 The coroner believed that it was unclear how care homes were being advised to safely
245 manage residents at risk of falls when isolation was required, and the home knew of no
246 guidance they could follow to manage the risk. Furthermore, restrictions on family visitors
247 when in hospital presented significant communication problems, which reduced appropriate
248 support and timely clinical decision making.

249 2) *Failure to follow protocols*

250 The failure to regularly monitor/observe the patient was the most common form of failure
251 (36%; n=4), followed by a general failure to follow protocols, omission of necessary
252 treatment, and delayed discharge (Figure 3).

253 **Example 2:** *A 28-year-old man with a lifelong history of low mood and depression with*
254 *suicidal ideation was socially isolated, which was exacerbated when he was furloughed from*
255 *his job. He had made five or six attempts to kill himself in 2020 alone and came to the*
256 *attention of an NHS Trust in May 2020. He registered with a GP at around the same time*
257 *and was assessed by a Mental Health practitioner, who referred him for treatment. The man*
258 *was aware of the referral and waited for treatment, but unfortunately the referral was lost. He*
259 *died by suicide at home 2 months later.*

260 The coroner concluded that the Care Programme Approach (CPA), as set out in national
261 guidance “Refocusing the CPA – Policy and Positive Practical Guidance”, had not been
262 followed. It was clear at the inquest that staff members were not aware of how matters
263 should be dealt with and that this depended heavily on geographical location.

264 3) *Education and training*

265 The most common concern was poor clinical decision making (30%, n=3), followed by
266 inappropriate dosage of medication (20%, n=2).

267 **Example 3:** *A man with drug dependency had been receiving support from a Recovery*
268 *Network and was generally fit, well, and in good spirits. He received a regular prescription*
269 *for methadone in daily dosage bottles three times a week. During the COVID-19 pandemic*
270 *this regimen was changed to once every 14 days, giving him access to a much larger*
271 *quantity of methadone. The deceased was found at home with a very high blood*
272 *concentration of methadone. There was no evidence that he intended to take his own life. At*
273 *the start of the pandemic, Public Health England (PHE) guidance was issued that patients*
274 *taking short-term methadone should be moved to long-term treatment. The doctor who*
275 *changed the prescription stipulated that the drug must be supplied in single daily dosage*
276 *bottles, and explanatory letters had been sent to all the pharmacies that supplied opiate*
277 *replacement therapies to his patients.*

278 The coroner saw evidence that the prescription was not in daily dose bottles and that there
279 was no measuring jug to enable accurate measurement of doses. The coroner believed that
280 it was probable that the man had guessed his first dose from the large methadone bottle. If
281 he had been given his daily dosage bottles, as prescribed, a measuring jug, and instructions
282 on how to use it, his death might have been prevented.

283 4) Resources

284 Lack of awareness of appropriate protocols and national guidelines was the most common
285 failure (40%, n=4), followed by understaffing (30%, n=3), a shortage or lack of availability of
286 medical equipment (20%, n=2), and unclear protocols (10%, n=1).

287 **Example 4:** *A man was admitted to hospital after an accidental fall at home. After surgery*
288 *for a fractured hip, he developed a chest infection. When fit for discharge, he was moved to*
289 *several different wards and eventually put in a bay where patients had been exposed to*
290 *COVID-19. He subsequently tested positive for COVID-19, deteriorated rapidly, and died.*

291 The coroner heard that the decision to move the deceased had been made on interpretation
292 of guidance from PHE. The Trust changed its policy, and such movements are reportedly no
293 longer taking place. However, when the PFD was written the guidance from PHE had not
294 been amended, and it was not known how other Trusts were choosing to interpret the
295 guidance, potentially putting other vulnerable in-patients at risk of COVID-19. Rapid and
296 national dissemination of coroner's concerns might have prevented similar errors in other
297 hospitals.

298 5) Safety

299 Half of the concerns about safety related to facilities (50%, n=3), followed by poor systems
300 (33%, n=2) and inadequate discharge processes (17%, n=1).

301

302 **Example 5:** *A resident in a care home, who had pulmonary fibrosis, had an unwitnessed fall.*
303 *He lay on the floor for over 4 hours awaiting an ambulance. Sepsis was diagnosed and it*
304 *was thought that he had symptoms consistent with COVID-19. He remained at the home*
305 *until 11 April, became unresponsive, deteriorated rapidly, was moved to palliative care, and*
306 *died on 13 April 2020.*

307

308 The coroner reported that the care had been of limited quality, notwithstanding the diagnosis
 309 of COVID-19 and his vulnerability. The care home was unclear if staff had brought COVID-
 310 19 into the home or if admission of residents from the community who had not being tested
 311 before admission had caused COVID-19 to enter the home. There was no risk assessment
 312 in place relating to the admission of new residents, creating an unsafe environment for
 313 vulnerable people.

314

315

316 **Responses to PFDs**

317 Forty-three unique individuals and organizations received PFDs from coroners (Table 2).
 318 Most reports were sent to the NHS (51%; n=22), followed by the Government and related
 319 bodies, other organizations, and professional bodies. Government organizations had the
 320 highest response rates (64%; n=7), followed by professional bodies (50%; n=2), other
 321 organizations (50%; n=3), and NHS organizations (45%; n=10). Of the 23 PFDs, nine had a
 322 100% response rate and ten had no responses on the Judiciary website.

323

324 **Table 2:** Recipients of Prevention of Future Deaths reports involving COVID-19 in England and Wales
 325 between 1 January 2020 and 28 June 2021 and their response rates (created by the authors)

Addressee	No. of PFDs sent	No. of responses [±]	Response rate (%)
NHS organizations	22	10	45%
Trusts	5	2	40%
NHS England	4	2	50%
NHS Hospitals	4	3	75%
CCGs	3	0	0%
Health and Social Care Partnerships	2	1	50%
NHS Pathways*	1	1	100%
Ambulance services	1	1	100%
GPs	2	0	0%
Government	11	7	64%
Public Health England	3	3	100%
Department of Health and Social Care	2	1	50%
Local authorities	3	2	67%
COVID-19 pandemic response service*	1	1	100%
Secretary of State of Health	1	0	0%
Ministry of Defence	1	0	0%
Professional bodies	4	2	50%
CQC	2	2	100%
General Pharmaceutical Council	1	0	0%
MHRA	1	0	0%
Other	6	3	50%
Care homes/providers	2	2	100%

Water board	1	0	0%
National Park	1	0	0%
Legal	1	0	0%
Pharmacy	1	1	100%

326 *NHS Digital responded on behalf of NHS Pathways and the COVID-19 Pandemic Response Service
327 ‡Recipients of PFDs have 56 days from the date of the report to respond to the coroner under
328 Regulation 29 of The Coroners (Investigations) Regulations 2013
329 CCG: Clinical Commissioning Group; CQC: Care Quality Commission; GPs: General Practitioners; MHRA:
330 Medicines and Healthcare products Regulatory Agency; NHS: National Health Service
331

332 Discussion

333 One in 20 PFDs published online involved COVID-19. Most of the deaths occurred in men
334 and older adults. There was wide geographical variation: no PFDs were reported by
335 coroners in Wales or in the North East, East Midlands, or South West of England. Coroners
336 raised several concerns, particularly regarding problems with communication and following
337 protocols. The largest numbers of PFDs were sent to NHS Trusts and the Government.

338 Eleven deaths were directly caused by SARS-CoV-2. In one-third of these cases, patients
339 acquired COVID-19 after admission to hospital for an unrelated reason, and another
340 acquired COVID-19 while in a care home. This suggests that measures to reduce the
341 transmission of SARS-COV-2 in healthcare settings were not adequate. Healthcare settings
342 should focus on evidence-based provisions, such as ventilation, personal protective
343 equipment (PPE), and regular testing to mitigate the risk to vulnerable patients²².

344 Twelve deaths were indirectly attributed to the COVID-19 pandemic. This finding highlights
345 the importance of considering the harms of measures and policies that were implemented to
346 reduce the transmission of SARS-COV-2 in the community. Reduced social interactions and
347 changed working conditions or loss of work and income have negatively affected adult
348 mental health in the UK²³. Six million patients in the UK did not seek treatment in 2020 (so-
349 called “missing patients”), owing to reprioritization of healthcare services²⁴. In some cases
350 reduced access to care because of lockdown, despite telemedicine, led to deaths.

351 Healthcare professionals in England and Wales have called for the deaths of their
352 colleagues to be reported to coroners and for PFDs to be issued^{14,15}. However, we did not
353 identify any PFDs that reported deaths of healthcare professionals.

354 The UK Government has stated that they will begin their public inquiry into the handling of
355 the COVID-19 pandemic to learn lessons for future pandemics²⁵. We have identified several
356 areas that the Government should address during this inquiry, including poor communication
357 and gaps in education and training. PFDs should be examined by the Government and
358 healthcare providers to inform quality improvement and patient safety initiatives.

359 The Office of the Chief Coroner, who is responsible for uploading PFDs to the Courts and
360 Tribunals Judiciary website, categorized the COVID-19-related PFDs into nine groups,
361 including hospital-related, community healthcare, and care homes. A new category
362 specifically for the effects of pandemics should be added to the Judiciary website to assist

363 the Government in examining these case reports so that policy measures can be
364 implemented for future pandemics.

365 Our study has several limitations. The 23 deaths do not represent all deaths during the
366 COVID-19 pandemic that could have been prevented in England and Wales. We are also
367 limited by the information reported by coroners in PFDs. In 52% of PFDs the age of the
368 deceased was not reported and 26% did not report the date of death. There were also
369 regions in England and Wales that did not report any PFDs.

370 Under-reporting of PFDs limits the capacity for actions to be taken to prevent future deaths.
371 However, it is likely that more PFDs relating to COVID-19 will be published, owing to the
372 backlog of inquests and the time it takes for inquests to conclude and PFDs to be written
373 and published.

374 There are no clear guidelines in England and Wales for referring deaths to coroners nor for
375 determining when a PFD should be issued and what information to include, hence the
376 missing data. There is also no auditing or quality control of PFDs and their responses. Thus,
377 whether action is taken to prevent such deaths, and the timeliness of such action, is
378 unknown and unmonitored. In the meantime, we encourage coroners across England and
379 Wales to continue writing PFDs when they believe that deaths could have been prevented.

380 PFDs contain a rich source of information that can be systematically analysed to share
381 information on preventing harms. The concerns identified in the 23 PFDs should be
382 considered during the UK Government's inquiry, including how communication, protocols,
383 education, training, resources, and patient safety can be improved. Guidelines on how and
384 when to report deaths to coroners, thresholds for issuing PFDs, and the
385 necessary mandatory types of information to report in PFDs, including age and date of death,
386 are needed. Coroners in England and Wales should be encouraged to continue writing
387 PFDs during the COVID-19 pandemic, particularly when deaths involve frontline healthcare
388 professionals.

389

390 **Declarations**

391

392 **Statement of ethical approval**

393 We are using publicly available information, for which ethics committee approval is not
394 required.

395

396 **Funding statement**

397 No grant or research funding was obtained to undertake this study.

398

399 **Competing interests statement**

400 BS receives funding from Mustafa Bahceci (Bahceci Health Group, Istanbul, Turkey) for her
401 Doctor of Philosophy studies at the University of Oxford (2019-2022) and has received
402 financial remuneration for consultancy work in women's health. CH is a National Institute for
403 Health Research (NIHR) Senior Investigator and has received expenses and fees for his
404 media work, received expenses from the WHO, FDA, and holds grant funding from the NIHR
405 School for Primary Care Research (SPCR) and the NIHR SPCR Evidence Synthesis
406 Working Group [Project 380], the NIHR BRC Oxford and the WHO. On occasion, CH

407 receives expenses for teaching EBM and is also paid for his GP work in NHS out of hours
408 (contract with Oxford Health NHS Foundation Trust). JKA has published articles and edited
409 textbooks on adverse drug reactions and interactions and has often given medicolegal
410 advice, including appearances as an expert witness in coroners' courts, often dealing with
411 the adverse effects of opioids and other medicines. DJH is the Director of Studies for
412 Sustainable Urban Development at the Department for Continuing Education, University of
413 Oxford. DJH has received financial remuneration for providing political and socioeconomic
414 country updates for Latin America and the Caribbean for IHS Global. GCR was financially
415 supported by the NIHR SPCR, the Naji Foundation, and the Rotary Foundation to study for a
416 Doctor of Philosophy (2017-2021), but no longer has any financial COIs. GCR is an
417 Associate Editor of BMJ Evidence Based Medicine and is developing
418 <https://preventabledeathstracker.net/>. The views expressed are those of the authors and not
419 necessarily those of the NHS, the NIHR, or the Department of Health and Social Care.

420

421 **Contributorship statement**

422 GCR developed the idea for this study, wrote the initial study protocol, ran the python code
423 to collect the most recent PFDs for screening, contributed to the first draft of the manuscript,
424 and provided supervisory support. BS contributed to the study protocol, screened the 510
425 PFDs for their eligibility, extracted the data from the 23 included cases, analysed the data,
426 and wrote the first draft of the manuscript. CH, JKA, and DJH contributed to the study
427 protocol, data analysis, and supervision of the project. All authors read, reviewed, and
428 approved the manuscript before submission.

429 **Data sharing**

430 The data, statistical code, and study materials are openly available via the Open Science
431 Framework (OSF) and GitHub^{16,19}.


432

433 **References**

- 434 1. WHO Coronavirus (COVID-19) Dashboard. *World Health Organization*, 2021.
435 <https://covid19.who.int>
- 436 2. The Coroners Rules 1984. *Rule 43*.
437 <http://www.legislation.gov.uk/ukxi/1984/552/article/43/made>.
- 438 3. Coroners and Justice Act 2009.
439 https://www.legislation.gov.uk/ukpga/2009/25/pdfs/ukpga_20090025_en.pdf.
- 440 4. The Coroners (Investigations) Regulations 2013.
441 http://www.legislation.gov.uk/ukxi/2013/1629/pdfs/ukxi_20131629_en.pdf.
- 442 5. Richards GC, Aronson JK, Heneghan C. Coroners' concerns to prevent harms: a series
443 of coroners' case reports to serve patient safety and educate the public, clinicians and
444 policy-makers. *BMJ Evid Based Med* 2021; 26(2): 37-8.

- 445 6. Richards GC. Alcohol-based hand sanitisers: a warning to mitigate future poisonings
446 and deaths. *BMJ Evid Based Med* 2021; 26(2): 65-8.
- 447 7. Cox AR, Ferner R. Tramadol: repeated prescriptions and repeated warnings. *BMJ Evid*
448 *Based Med* 2021 Mar 8; bmjebm-2020-111661.
- 449 8. Richards GC. Preventable opioid-related deaths in England and Wales. (2021)
450 doi:10.17605/OSF.IO/ECZ4R.
- 451 9. Bilip MK, Richards GC. Emollients and smoking: a fire hazard that could be prevented
452 to reduce future deaths. *BMJ Evid Based Med* 2021; 26(3): 131-4. .
- 453 10. Ferner RE, Easton C, Cox AR. Deaths from medicines: a systematic analysis of
454 coroners' Reports to Prevent Future Deaths. *Drug Saf* 2018; 41(1): 103-10.
- 455 11. Richards GC. *et al.* Preventable suicides: a systematic analysis of coroners' prevent
456 future death reports in England and Wales. (2020) doi:10.17605/OSF.IO/AD4UP
- 457 12. Ains A, Heneghan C, Aronson JK, DeVito NJ, Richards GC. Deaths from cardiovascular
458 disease involving anticoagulants: a systematic synthesis of coroners' case reports.
459 *BJGP Open* 2021; BJGPO.2021.0150.
- 460 13. Odgers L, Aronson JK, Richards GC. Preventable deaths involving cyclists in England
461 and Wales: solutions to improve their safety. *Global Centre on Healthcare and*
462 *Urbanisation*. [https://www.gchu.org.uk/2021/09/gchu-report-preventable-deaths-](https://www.gchu.org.uk/2021/09/gchu-report-preventable-deaths-involving-cyclists-in-england-and-wales)
463 *involving-cyclists-in-england-and-wales* (2021).
- 464 14. Kendrick D, Agius R, McKee M, Robertson J, Sewell H, Stewart M. COVID-19: Learning
465 lessons from the deaths of our colleagues. *BJGP Life* 2020;
466 <https://bjgplife.com/2020/05/18/gp-deaths/>.
- 467 15. Agius RM. Doctors' deaths from COVID-19 should be reported to the coroner. *BMJ*
468 2020; 369 m1622.
- 469 16. Swift B, Heneghan C, Aronson JK, Howard D, Richards. Preventable deaths from
470 SARS-CoV-2 in England and Wales: a systematic analysis of coroners' case reports.
471 *OSF Registries* 29 June 2021; <https://osf.io/ej6ks>
- 472 17. Judiciary. Prevention of Future Deaths. *Courts and Tribunals Judiciary* 2021;


- 473 <https://www.judiciary.uk/subject/prevention-of-future-deaths/>.
- 474 18. Richards, G.C.Preventable Deaths Tracker. 2021. <https://preventabledeathstracker.net>
- 475 19. [dataset] Richards, G.C. &. DeVito, N.J. Data from: Preventable Deaths Database.
- 476 2021, November 12 2021. georgiarichards.github.io DOI: 10.5281/zenodo.567586620.
- 477
- 478 20. DeVito NJ, Richards GC, Inglesby P. How we learnt to stop worrying and love web
- 479 scraping Nature 2020; 585(7826): 621-2.
- 480 21. Kleinheksel AJ, Rockich-Winston N, Tawfik H, Wyatt TR. Demystifying content analysis.
- 481 *Am J Pharm Educ* 2020; 84(1): 7113.
- 482 22. Richterman A, Meyerowitz EA, Cevik M. Hospital-acquired SARS CoV-2 infection
- 483 lessons for public health. *JAMA* 2020; 324(21): 2155-6.
- 484 23. Abbott A. COVID's mental-health toll: how scientists are tracking a surge in depression.
- 485 *Nature* 2021; 590(7845): 194-5.
- 486 24. The Health Foundation. Unequal pandemic, fairer recovery: The COVID-19 impact
- 487 inquiry report. 2021; <https://reader.health.org.uk/unequal-pandemic-fairer-recovery>.
- 488 25. Booth R. Johnson says public inquiry into Covid will begin this parliament. *The Guardian*
- 489 11 May 2021. [https://www.theguardian.com/world/2021/may/11/boris-johnson-public-](https://www.theguardian.com/world/2021/may/11/boris-johnson-public-inquiry-into-covid-begin-this-parliament)
- 490 [inquiry-into-covid-begin-this-parliament](https://www.theguardian.com/world/2021/may/11/boris-johnson-public-inquiry-into-covid-begin-this-parliament)





Absence/lack of awareness of appropriate protocols/national guidelines	40%
Understaffing	30%
Shortage/lack of availability of appropriate medical/safety equipment	20%
Protocols not clear and interpreted wrongly	10%

Safety of facilities	50%
Poor systems	33%
Inadequate discharge process	17%



Poor clinical decision making	30%
Inappropriate dosage for the patient	20%
Failure to appreciate the risk	10%
Inadequate training	10%
Lack of training in drug administration	10%
Poor awareness of symptoms	10%
The incorrect drug administered	10%

Poor communication	42%
Failure to keep accurate medical records/care plans	26%
Failure to consider how COVID-19 would affect appointments	16%
Failure to follow advice	5%
Failure to inform family on patient's progress	5%
Failure to take informed consent	5%

Failure to regularly monitor/observe the patient	36%
Delayed discharge	9%
Failure to arrange supervision	9%
Failure to follow a protocol	9%
Failure to review medicines	9%
Failure to triage patients appropriately	9%
Failure/delay in an appropriate assessment of the patient	9%
Omission of necessary treatment	9%