

1 Original Article

2 **Internet Use impact on Physical Health during COVID-19 pandemic in Bangladesh: A**
3 **Web-based Cross-sectional study**

4
5 **Authors:** Tanvir Abir¹, Uchechukwu Levi Osuagwu², Dewan Muhammad Nur–A Yazdani³,
6 Abdullah Al Mamun⁴, Kaniz Kakon⁵, Anas A. Salamah⁶, Noor Raihani Zainol⁷, Mansura
7 Khanam⁸, Kingsley Emwinyore Agho⁹

8 **Affiliation:**

9 ¹. Associate Professor, College of Business Administration—CBA, International University
10 of Business, Agriculture and Technology—IUBAT University, Dhaka 1230, Bangladesh;
11 tanvir.cba@iubat.edu;

12 ². Diabetes, Obesity and Metabolism Translational Research Unit, Western Sydney
13 University, Campbelltown, NSW 2560, Australia; African Vision Research Institute (AVRI),
14 University of KwaZulu-Natal, Westville Campus, Durban, 3629, South Africa;
15 losuagwu@westernsydney.edu.au;

16 ³. Assistant Professor, College of Business Administration—CBA, International University of
17 Business, Agriculture and Technology—IUBAT University, Dhaka 1230, Bangladesh;
18 dewanm@iubat.edu

19 ⁴. Faculty of Business and Management, UCSI University, Kuala Lumpur 56000, Malaysia;
20 abdullaham@ucsiuniversity.edu.my;

21
22 ⁵. Assistant Professor, Department of Philosophy, College of Arts and Sciences—CAAS,
23 International University of Business, Agriculture and Technology—IUBAT University,
24 Dhaka 1230, Bangladesh; kanizkakon@iubat.edu

25 ⁶. Department of Management Information Systems, College of Business Administration,
26 Prince Sattam Bin Aziz University, 165 Al-Kharj 11942, Saudi Arabia. Email:
27 a.salameh@psua.edu.sa

28 ⁷. Lecturer, Faculty of Entrepreneurship and Business, Universiti Malaysia Kelantan, 16100
29 Kota Bharu, Malaysia; raihani@umk.edu.my

30 ⁸. Maternal and Child Nutrition, Nutrition and Clinical Services Division, icddr,b
31 Bangladesh; mansura@icddr.org

32 ⁹. Associate Professor, School of Health Science, Western Sydney University, Campbelltown,
33 NSW 2560, Australia; African Vision Research Institute (AVRI), University of KwaZulu-
34 Natal, Westville Campus, Durban, 3629, South Africa; K.Agho@westernsydney.edu.au

35 * Correspondence: Tanvir Abir (PhD), Associate Professor, College of Business
36 Administration—CBA, International University of Business, Agriculture and Technology—
37 IUBAT University, Dhaka 1230, Bangladesh; tanvir.cba@iubat.edu

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

39 **Funding:** This research did not receive any funding.

40

41 **Declaration of Conflict of Interest:** The authors declare that they have no known competing
42 financial interests or personal relationships that could have appeared to influence the work
43 reported in this paper.

44

45 **Author Contribution:** Conceptualization, T.A. and D.M.N.-A. Y; Data curation, T.A. and
46 K.A; Formal analysis, U.L.O. and K.A; Investigation, D.M.N.-A.Y. and M.K.; Methodology,
47 T.A., A.A.M, A.A.S.; Project administration, K.A., D.M.N.-A.Y., K.K.; Resources, D.M.N.-
48 A.Y., A.A.M. and N.R.Z; Supervision, N.R.Z. and K.A.; Validation, T.A. and K.A.;
49 Visualization, M.K. and K.A.; Writing—original draft, T.A. and U.L.O.

50 All authors have read and agreed to the published version of the manuscript.

51

52 **Article Type:** Original Research.

53 **Keywords:** Internet use, Coronavirus, Survey, Headache, Back pain, Neck pain, Physical
54 health

55 **Running Head:** Internet use and Physical health of Bangladeshi Adults

56

57 **Ethics Statement:** Permission for this study (IUBAT/AR/2021/001) was obtained from the
58 Institutional Review Board of International University of Business, Agriculture and
59 Technology (IUBAT), Dhaka, Bangladesh. The study adhered to the tenets of the Declaration
60 of Helsinki as revised in Fortaleza. All the participants were informed about the specific
61 objective of this study before proceeding to the questionnaire. Consent was obtained from all
62 participants prior to completing the questionnaire through an online preamble. Participants
63 were able to complete the survey only once and could terminate the survey at any time they
64 desired. Anonymity and confidentiality of the data were ensured.

65 **Data Reference:** Our data are included in the manuscript and raw data can be released at
66 reasonable request.

67

68 **Abstract**

69 **Background:** Bangladesh ranked fifth largest internet user in Asia. Past studies on internet
70 use have focused on its impact on mental health, with little known about its impact on the
71 physical health of individuals during COVID-19 pandemic. Hence, this study examines the
72 impact of Internet use frequency on physical health during the Covid-19 lockdown in
73 Bangladesh.

74 **Methods:** A web-based cross-sectional study on 3242 individuals aged 18 and above was
75 conducted from 2nd August – 1st October 1, 2020, during the lockdown in Bangladesh. The
76 survey covers demographics, Internet use frequency and physical health questions. Multiple
77 linear regression analyses were used to examine the impact of internet use frequency on
78 physical health.

79 **Results:** The result indicated that 72.5%, 69.9%, 65.1% and 55.3% reported headache, back
80 pain, numbness of the fingers and neck pain, respectively. The multivariable analyses showed
81 increased physical health impact among regular (coefficient $\beta = 0.52$, 95% confidence interval
82 [CI]: 0.18-0.85, $P = 0.003$), frequent ($\beta = 1.21$, 95%CI: 0.88–1.54, $P < .001$) and intense (β
83 = 2.24, 95%CI: 1.91–2.57, $P < .001$) internet users. Other factors associated with physical
84 health scores were gender, income (in Taka), occupation, regions, and working status.

85 **Conclusion:** Frequent, intensive, and extensive use of the internet were strong predictors of
86 increased physical health problems, and the study suggests the need for raising awareness of
87 physical health problems triggered by high internet users among the high socioeconomic
88 group in Bangladesh.

89

90

91 INTRODUCTION

92 The coronavirus disease 2019 (COVID-19) pandemic and the associated public health
93 interventions, including social distancing, home quarantine measures, and ‘stay-at-home’
94 orders, put in place by the various governments, led to a surge in internet usage during the
95 pandemic ^[1]. The lockdown measures resulted in widespread and unprecedented social
96 disruption ^[2] as non-essential businesses were closed or some employees worked from home
97 and unemployment increased ^[3]. There were massive changes in internet usage patterns and
98 user behaviour with employees having to adjust to the new "normals" - with meetings going
99 completely online, office work shifting to the home, leading to new emerging patterns of
100 work. These changes have come across most organizations, businesses, societies, or
101 governments and suddenly, leaving barely any time for organizations and people to plan for,
102 prepare and implement new setups and arrangements. Society has had to adjust, try,
103 experiment, and find ways that did not exist before the pandemic ^[1], and these may have
104 implications on the physical health of individuals.

105 People are now spending even more time with technology while consuming news
106 media, watching television, using social media to connect with others, utilizing lifestyle apps
107 to shop for groceries and other consumer goods, and engaging in home workouts ^[2]. In
108 Bangladesh, this kept parts of the economy going ^[4] as the country witnessed a boom in
109 internet usage due to the fast-growing mobile internet and the government's push for
110 digitalization. Engaging in social media platforms is now a routine activity for children and
111 adults, enhancing communication, social connection, and even technical skills ^[5]. Platforms
112 such as Facebook and Twitter offer multiple daily opportunities for connecting with friends,
113 classmates, and people with shared interests. The number of people using such sites has also
114 increased dramatically over the last five years, as shown in a recent poll ^[6]. A past study
115 reported that about 22% of teenagers log on to their favorite social media site more than ten
116 times a day, and more than half of adolescents log on to a social media site more than once a

117 day ^[6,7]. Seventy-five percent of teenagers now own cell phones, 25% use them for social
118 media, 54% use them for texting, and 24% use them for instant messaging ^[7].

119 According to the Bangladesh Telecommunication Regulatory Commission, there were
120 99,428 million (approx. 60.9% of the population) internet users in February 2020 ^[8]. Though
121 there is anecdotal early evidence that this may lead to increased productivity, it has also led to
122 increased stress ^[9,10] where employees must learn new technologies, be available for work at
123 almost all times, stay with digital devices all the time, and cope with multi-tasking ^[1]. The
124 use of video-conferencing and content delivery services such as Zoom and Akamai also
125 increased ^[1]. Those working from home using this technology have found themselves under
126 intense scrutiny and tension ^[11].

127 That a large part of the peoples' social and emotional development is occurring while
128 on the Internet and cell phones may affect the physical health of individuals considering the
129 long hours spent on the internet during the mandatory stay at home orders which expose the
130 population to lots of misinformation ^[12] known to affect physical and mental health ^[13]. Since
131 this high use of information systems may become the new normal, this study examined the
132 impact of internet usage frequency during the COVID-19 pandemic on Bangladeshi residents'
133 physical and mental health and determined the level of addiction to social media platforms
134 used by the people.

135 **Ethical consideration**

136 Permission for this study (IUBAT/AR/2021/001) was obtained from the Institutional Review
137 Board of International University of Business, Agriculture and Technology (IUBAT), Dhaka,
138 Bangladesh. The study adhered to the tenets of the Declaration of Helsinki as revised in
139 Fortaleza. All the participants were informed about the specific objective of this study before
140 and consent was obtained from all participants before completing the questionnaire through
141 an online preamble. To avoid repeated responses and in order to ensure the validity of data to

142 some extent, participants were able to complete the survey only once as data was restricted to
143 their IP addresses and device. Participants could terminate the survey at any time they
144 desired. Anonymity and confidentiality of the data were ensured.

145 **METHODS**

146 **Study population and sampling**

147 By convenience sampling, 3242 questionnaires were collected from adults aged 18 and over
148 living in eight divisions across Bangladesh during the pandemic. All questionnaires were
149 web-based, participation was voluntary, and each participant was informed about the study's
150 background through an online preamble. No reward or incentive was offered for
151 participation. There may be more than one questionnaire filled by the same person that
152 affects the statistical analysis results of questionnaire data. In order to ensure the validity of
153 data to some extent, restricting the same IP address and device are applied to the system,
154 which means the participant of the same IP address or device only can fill once.

155 **Data Collection**

156 Data were collected using self-administered web-based questionnaires distributed to various
157 groups of people via the E-link using numerous mailing lists and social network sites
158 including Facebook, WhatsApp and Twitter. Due to the mandatory lockdown, no paper
159 questionnaires were distributed, making it difficult to reach the very remote regions.

160 **Dependent variable**

161 Ten items of physical complaints reported in *Table 2* below were used to determine the
162 impact of individual physical health on Internet use. The respondents used a close-end
163 response ['Yes' or 'No'] to indicate whether they suffered from the following ten problems
164 (Back and neck pain, numbness in fingers, headaches, inability to sleep, dry eyes or other
165 vision problems, poor nutrition, poor personal hygiene, weight gain/loss and loss of appetite]
166 during or after pro-longed Internet use. Binary scores for individual questions were summed

167 to give a physical health score which ranged from 0-10, and the Kuder Richardson coefficient
168 measuring internal consistency among the physical health scores ranged from 0.65 to 0.78,
169 indicating a satisfactory level of reliability.

170

171 **Main study factor**

172 The main study variable was the respondent's Internet usage, including the average daily
173 amount of time spent on the internet and dependence on networks. The frequency of internet
174 use was categorized as 'seldom' (<1 hour per day), 'casual' (≥ 1 and <3 hours per day),
175 'regular' (≥ 3 and <5 hours per day), 'frequent' (≥ 5 and <7 hours per day), 'intense' (≥ 7 hours
176 per day) [14].

177

178 **Independent variables**

179 Independent variables used for the study analyses were based on previous studies [14-16].
180 Independent variables included demographic characteristics (i.e., gender, age [categorized as
181 18-27, 28-37, 38 years and over], marital status [single, married, divorced/widowed], mothers
182 level of education [Higher education, Bachelor, Intermediate], place of residence,
183 employment status [Employed, unemployed/student], occupation [healthcare and non-
184 healthcare worker], and income in Taka [lower, middle and high-income earners].

185

186 **Statistical Analysis**

187 All statistical analyses were performed in Stata version 14.1 (Stata Corp. 2015, College
188 Station USA) mainly using descriptive statistics; t-test used compares the means of two
189 independent groups, One-way analysis of variance (one-way ANOVA) used compares the
190 means of more than two independent groups, correlation analyses, and regression analysis.
191 All confounding variables with a $P < .20$ were retained in the univariate linear regression
192 analysis and were used to build a multivariable model [17]. A manual elimination procedure

193 was applied for multivariate linear regression analyses to remove non-significant variables (P
194 $> .05$). The main study factor variable (Internet usage frequency) was added to all significant
195 confounding variables after elimination processes. The main study factor and independent
196 variables associated with physical health scores ($P < .05$) were reported. In the regression
197 analysis, we checked for homogeneity of variance and multicollinearity, including Variance
198 Inflation Factors (VIF) and the $VIF < 4$ was considered suitable [18].

199

200 RESULTS

201 Demographic Characteristics According to the Frequency of Internet Use among adults 202 in Bangladesh

203 *Table 1* depicts the main demographic characteristics, including gender, education and age
204 group on the five groups extracted. It is worth noting that the education in this paper refers to
205 that of the mothers' maximum educational level. The majority of participants were males
206 (1260, 61.1%), married (2632, 81.2%) and employed (2864, 88.3%) in non-healthcare sectors
207 (2567, 79.2%), and for many, their mothers had completed a bachelor degree or higher.
208 About half of the respondents earned more than 70,000 Taka at the time of data collection
209 (*Table 1*).

210 About two-thirds of the respondents that reported intense use of the internet were
211 males, while females were more likely to report regular and casual use of the internet. More
212 than half of intense and frequent internet users were aged 28-37years, and a similar
213 proportion of casual internet users were younger (18-27years). Nearly all intense internet
214 users were employed and working in the non-Health care sector during the study.

215 *Table 1* Demographic Characteristics According to the Frequency of Internet Use among
216 adults in Bangladesh. Values are expressed as numbers and percentages (n, %).

Variables	Total (N=3236)	Intense (n=1668)	Frequent (n=689)	Regular (n=559)	Casual (n=187)	Seldom (n=93)
Demography						

<i>Gender</i>						
Male	1976 (61.1)	1179 (70.7)	382 (55.4)	264 (44.1)	93 (44.7)	58 (62.4)
Female	1260 (38.9)	489 (29.3)	307 (44.6)	335 (55.9)	94 (50.3)	35 (37.6)
<i>Age in years</i>						
18-27	555 (17.2)	30 (1.8)	156 (22.6)	254 (42.1)	98 (52.4)	17 (17.9)
28-37	1657 (51.1)	956 (57.3)	366 (53.1)	238 (39.5)	60 (32.1)	37 (39.0)
38+	1030 (31.8)	682 (40.9)	167(24.2)	111 (18.4)	29 (15.5)	41 (43.2)
<i>Marital status</i>						
Single	508 (15.7)	29 (1.7)	150 (21.8)	223(37.0)	90 (48.1)	16 (16.8)
Married	2632 (81.2)	1632 (97.8)	505 (73.3)	339(56.2)	83 (44.40)	73 (76.8)
Divorced/widow	102 (3.1)	7 (0.4)	34 (4.9)	41 (6.8)	14(7.49)	6 (6.3)
<i>Place of residence (%)</i>						
Barisal Division	172 (5.31)	70 (4.20)	48 (7.0)	35 (5.0)	7 (3.7)	12 (12.6)
Chittagong Division	291 (9.0)	100 (6)	107 (15.5)	61 (10.1)	16(8.6)	7(7.4)
Dhaka Division	1574 (48.6)	1010 (60.6)	205 (29.8)	236 (39.1)	95 (50.8)	28 (29.5)
Khulna Division	352 (10.9)	115 (6.9)	109 (15.8)	92 (15.3)	20 (10.7)	16 (16.8)
Mymensingh Division	313 (9.7)	127 (7.6)	76 (11.0)	85 (14.1)	17 (9.1)	8 (8.4)
Rajshahi Division	213 (6.6)	99 (5.9)	47 (6.8)	41 (6.8)	18 (9.6)	8 (8.4)
Rangpur Division	179 (5.5)	93 (5.6)	45 (6.5)	22 (3.7)	9 (4.8)	10 (10.5)
Sylhet Division	148 (4.6)	54 (3.2)	52 (7.6)	31 (5.1)	5 (2.7)	6 (6.3)
<i>Mother's Level of Education</i>						
Higher education (above Bachelor	1287 (39.7)	684 (41.0)	341 (49.5)	178 (29.5)	46 (24.6)	38 (40.)
Intermediate (11–12)	441 (13.6)	23 (1.4)	122 (17.7)	192 (31.84)	74 (39.5)	30 (31.6)
<i>Working status</i>						
Employed	2864 (88.3)	1647 (98.7)	600 (87.1)	432 (71.6)	99 (52.9)	86 (91.0)
Not employed/student	378 (11.7)	21 (1.3)	89 (12.9)	171 (28.4)	88 (47.1)	9 (9.47)
<i>Income in Taka</i>						
lower-income (<30,000)	204 (6.3)	14 (0.9)	37 (5.4)	102 (16.9)	41 (21.9)	10 (10.4)
middle-income (30,000-70,000)	1496 (46.1)	510 (30.9)	433 (62.8)	389 (64.5)	108 (57.8)	56 (59.0)
High-income (>70,000)	1542 (47.7)	1144 (68.59)	219 (31.8)	112 (18.8)	38 (20.3)	29 (30.5)
<i>Occupation</i>						
Healthcare workers	675 (20.8)	31 (1.9)	212 (30.8)	304 (50.4)	89 (47.6)	39 (41.1)
Non-health care worker	2567 (79.2)	1637 (98.1)	477 (69.2)	299 (49.6)	98 (52.4)	56 (59.0)

217

218 **Prevalence and 95% confidence intervals of physical complaints by gender among**
 219 **internet users.**

220

221 *Table 2* depicts the prevalence and 95% CIs of the adverse effects on individual physical

222 health complaints examined in this study in males and females. Headache 72.50% [95%CI

223 70.93%, 74.01%] was the prevalent complaint of the participants, while 69.87% [68.27%,

224 71.43%] and 65.10% [63.42%, 66.71%] complained of Back pain and numbness of the


225 fingers, respectively. Males reported a higher prevalence of back pain, numbness of the
226 finger, headaches, and neck pains than females as the confidence intervals do not overlap (see
227 **Table 2**). By contrast, females reported a higher prevalence of poor nutrition, poor personal
228 hygiene, and loss of appetite compared with men (**Table 2**).

229 **Table 2** Gender differences in the prevalence of physical health scores among internet users
230 in Bangladesh. *Significant variables between males and females are bolded because the*
231 *confidence intervals did not overlap.*

Physical complaints	n	Prevalence (95%CI)		
		Overall	Male	Female
Back pain	2261	69.9 [68.27, 71.43]	78.14 [76.26, 79.91]	56.90 [54.15, 59.62]
Finger numbness	2106	65.10 [63.42, 66.71]	72.37 [70.35, 74.30]	53.65 [50.89, 56.39]
Headaches	2346	72.50 [70.93, 74.01]	78.80 [76.94, 80.54]	62.62 [59.91, 65.25]
Inability to sleep	1396	43.14 [41.44, 44.85]	41.95 [39.79, 44.14]	45.00 [42.27, 47.76]
Poor Nutrition	746	23.05 [21.63, 24.54]	16.55 [14.97, 18.25]	33.25 [30.70, 35.91]
Poor Personal Hygiene	605	18.70 [17.39, 20.08]	14.98 [13.47, 16.62]	24.52 [22.23, 26.98]
Neck pain	1789	55.28 [53.56, 56.99]	58.50 [56.31, 60.66]	50.24 [47.48, 53.00]
Dry eyes/other vision problems	1815	56.09 [54.37, 57.79]	57.54 [55.35, 59.70]	53.81 [51.05, 56.55]
Weight gain/loss	1653	51.08 [49.36, 52.80]	52.43 [50.22, 54.63]	48.97 [46.21, 51.73]
Loss of Appetite	499	15.42 [14.22, 16.71]	13.77 [12.31, 15.36]	18.02 [15.99, 20.24]

232 *CI=Confidence interval; n= number of respondents*

233 **Figure 1** shows the mean scores of physical health by internet use frequency in males
234 and females. From the figure, the mean scores of physical health by intensive internet use
235 was 5.6 for males and 5.4 for females. This physical health mean scores could be translated
236 as percentage mean scores of males (56%) and females (54%), respectively. Physical health
237 mean scores were lower among the casual user of internet (3.2 for females and 3.0 for males)
238 than seldom users of internet (3.1 for females and 3.4 for males)



239
240
241
242
243

Figure 1.

Mean scores by gender of the physical health by internet use frequency in Bangladesh

Unadjusted Analysis for the association between prolonged internet use and physical health complaints

Table 3 shows the mean scores of physical health complaints and the unadjusted coefficients for the associations with the sociodemographic variables. From the table, the mean scores for physical health complaints varied across variables. The unadjusted coefficients revealed significantly higher physical health problems due to prolonged internet use in those aged >28years, married, completed bachelor education, middle-high income earners and non-healthcare workers compared with other groups. In contrast, females, those who were divorced or widowed, those with intermediate education, non-workers/students and respondents who resided in Chittagong, Dhaka and Khulna during the study period had significantly lower mean scores of physical health complaints from prolonged internet usage compared with the other groups.

256
257
258

Table 3: Mean scores and Univariate linear regression of Physical Complaints among Internet Users.


Variables	Mean scores (SD)	Unadjusted Coefficient [95%CI]
Gender		
Male	4.85 (1.83)***	Reference
Female	4.47 (1.88)	-0.38 [-0.51, -0.25]
Age,		
18-27 yrs	3.52 (1.71)***	Reference
28-37 yrs	5.02 (1.81)	1.50 [1.33, 1.67]
38+ yrs	4.80 (1.77)	1.31 [1.12, 1.49]
Marital status		
Single	3.67 (1.76)***	Reference
married	4.96 (1.79)	1.30 [1.13, 1.47]
Divorced/widow	3.00 (1.19)	-0.67 [-1.04, -0.29]
Place of residence		
Barisal Division	5.35 (1.83)***	Reference
Chittagong Division	4.66 (1.87)	-0.69 [-1.04, -0.35]
Dhaka Division	4.36 (1.77)	-0.99 [-1.28, -0.70]
Khulna Division	4.83 (1.83)	-0.52 [-0.85, -0.19]
Mymensingh Division	5.03 (1.92)	-0.32 [-0.66, 0.02]
Rajshahi Division	5.21 (1.91)	-0.14 [-0.51, 0.22]
Rangpur Division	5.47 (1.88)	0.12 [-0.26, 0.50]
Sylhet Division	4.99 (1.83)	-0.36 [-0.76, 0.04]
Mother's Level of Education		
Higher education	4.71 (1.82)***	Reference
Bachelor	5.00 (1.87)	0.30 [0.16, 0.43]
Intermediate (11–12)	3.65 (1.50)	-1.06 [-1.26, -0.86]
Working status		
Employed	4.89 (1.79)***	Reference
Not employed/student	3.24 (1.72)	-1.65 [-1.84, -1.46]
Income in Taka		
lower-income (<30,000)	3.05 (1.50)***	Reference
middle-income (30,000)	4.82 (1.88)	1.77 [1.51, 2.04]
High-income (>70,000)	4.80 (1.77)	1.75 [1.48, 2.01]
Occupation		
Healthcare workers	3.62 (1.26)***	Reference
Non-health care workers	4.98 (1.89)	1.36 [1.21, 1.51]

259 ^{SD=standard deviation; CI=confidence interval. ***= $P < .001$ based on t-test and Anova. Bolded are significant coefficients.}

260

261 **The impact of the frequency of internet use on physical health scores**

262 *Figure 2* presents the unadjusted and adjusted odds ratio and their confidence intervals for
 263 the association between frequency of internet use and physical health symptoms during the
 264 COVID-19. In the unadjusted analysis, frequent and intensive internet users during the
 265 COVID-19 significantly reported higher physical health problems than seldom users of the
 266 internet. After adjusting for the independent variables used in this study, regular, frequent,
 267 and intensive internet users during the COVID-19 had significantly higher physical health
 268 problems than those who seldom used the internet (see *Figure 2*).



269

270 *Independent variables adjusted for are Gender, Age group, Division of residence, Mothers Education, marital status, income, working status, occupation.*

271 **Figure 2:** The impact of frequency of internet use on physical health scores: Unadjusted and
 272 adjusted coefficient and their 95% confidence intervals.

273

274 Marital status, age, gender, place of residence, level of education, working status,
 275 occupation, and income level were all associated with the physical health complaints among
 276 internet users in Bangladesh (*Supplementary Table S1 for details*).

277

278 **DISCUSSION**

279 A considerable body of literature has emerged over the past two decades assessing the
 280 relationship between problematic or addictive (excessive) use of the internet and individuals'
 281 psychological/mental well-being. Comparatively, very little research has evaluated the
 282 relationship between various internet use frequencies and the individual's physical health.
 283 The COVID-19 pandemic caused a drastic surge in the use of the internet globally and
 284 Bangladesh ^[1], and this cross-sectional study investigated the relationship between the
 285 frequency of internet use and the physical health of Bangladeshi respondents. The findings
 286 demonstrated that more than one in every two persons in this study used the internet for seven
 287 or more hours each day. This was associated with 2.2 folds increase in physical health
 288 problems compared with seldom users (<1 hour per day). Headache, back pain and numbness
 289 of the fingers (reported by over seventy percent of internet users) were the predominant
 290 complaints. There was a positive association between physical health problems and the
 291 number of daily hours spent on the internet. After adjusting for the frequency of internet use,

292 we found that middle to high-income earners and those working in non-healthcare sectors
293 experienced more significant physical health problems due to internet use. In comparison,
294 students and widowed or divorced persons reported fewer physical health problems compared
295 with other respondents.

296 This study found that headache was the most prevalent symptom reported by internet
297 users in Bangladesh, followed by back pain and numbness of the fingers. These complaints
298 were significantly increased among frequent internet users and intense users, which agrees
299 with the findings of significant association between extensive use of the internet and physical
300 health problems in previous studies ^[19-22]. A study in Spain reported that headache was the
301 most common complaint among internet users during the COVID-19 pandemic ^[22]. Similarly,
302 Cao et al.s study (2009), after examining adolescents, found that those who used the internet
303 excessively were more likely to report psychosomatic symptoms including lack of physical
304 energy, physiological dysfunction, weakened immunity, emotional symptoms, behavioural
305 symptoms and social adaptation problems and had lower life satisfaction scores compared
306 with normal internet users ^[20].

307 There are also reports that school children and other graduate students in China ^[23]
308 and Italy ^[24] experienced increased psychological distress following excessive internet use
309 during the COVID-19 pandemic. Such addiction to the internet can result in the problem of
310 self-care, difficulty in performing daily routine, and mental health effects of anxiety and
311 depression ^[25]. The negative effects of internet addiction on health may be related to the
312 alterations in the individuals neuroanatomical and neurochemical mechanisms, including
313 cortical thinning of various components of the brain and dopaminergic reward circuitry,
314 during excessive use of the internet ^[25]. While it is imperative that such behaviours,
315 especially during a pandemic, remain at a moderate and regulated level, identifying the
316 population at risk of the problematic use of the internet is necessary in order to prevent the
317 associated mental health problems ^[26] and improve work or school functions ^[27].

318 The present study further revealed that, compared with women, a higher proportion of
319 men used the internet excessively during the pandemic and thus experienced more significant
320 physical health problems. However, this association was significant only when it interacted
321 with the frequency of internet use. Previous studies among Taiwanese residents ^[28] and those
322 in mainland China ^[29] found that male gender was a potential risk factor for internet addiction
323 among adolescents. This association of gender and physical health complaints found in the
324 present study could be explained by the patriarchal nature of the Bangladesh society, where
325 women and girls rarely go outside of their homes even before the pandemic ^[30] but preferred
326 to stay home and spend more time on the Internet. Whereas these women may have
327 developed adaptive strategies, it was a different experience for the male counterpart, which
328 were used to staying outside/working prior to the lockdown. For these men, the lockdown
329 measures and transition to work online and/or staying at home required them to adjust
330 quickly, try, experiment, and find ways that did not exist before the pandemic ^[1] as
331 educational institutions, businesses and workplaces were shut down for such a long duration
332 ^[31]. These adjustments may have affected their physical health.

333 Another finding of this study was the significant relationship between income class
334 and occupational status (non-healthcare worker) with the physical health problems
335 experienced by internet users. Earning above 30,000 Taka was associated with higher
336 physical health problems due to internet usage after adjusting for the frequency of internet
337 use. In line with this finding, Islam et al ^[32] on the correlation between problematic
338 (excessive) internet use (PIU) and lifestyle during the COVID-19 pandemic using data from a
339 large sample of Bangladeshi youth and adults (n=13,525) found that younger age, higher
340 education, cigarette smoking, more sleep, less physical activity, gaming and social media use
341 were associated with PIU ^[32]. This research has depicted the physical health concerns related
342 to prolonged internet usage during the COVID-19 pandemic. A similar association was found
343 in this research between educational status, age and physical health complaints among

344 internet users in Bangladesh; however, these associations were dependent on the frequency of
345 internet use, which was not assessed in the previous study [32].

346 The finding that non-healthcare workers had higher physical health problems
347 compared with health care workers in this study is consistent with the finding of poorer
348 psychological well-being among non-health-care workers who used the Internet to obtain
349 information on COVID-19 compared with those who received COVID-19 information from
350 medical staff in health care settings [33]. Although this research were unable to identify
351 whether the non-healthcare workers in our study used the internet for COVID-19 information
352 retrieval, it has been shown that the influx of misinformation around COVID-19 via online
353 platforms had adverse effects on people's health [33]. Such problematic internet use (PIU) can
354 lead to clinical impairment, distress [34] and/or deterioration in financial, familial, social,
355 educational and/or occupational domains [35].

356 This study has some limitations. First, the cross-sectional design of this study does not
357 allow for causality or the direction of relationships to be determined. However, appropriate
358 analysis of cross-sectional data represents a useful initial step in identifying associations
359 between extensive internet usage and the physical health condition of the participants.
360 Second, as the responses were self-reported, it was not possible to verify the participants'
361 physical health complaints, which may lead to a response bias. Multiple assessments,
362 interviews, and informants may have provided a richer and more thorough understanding of
363 this topic. Third, since the survey was administered during COVID-19 lockdown using an
364 online platform, it is possible that some respondents, especially those living in rural areas
365 with limited internet access or people addicted to other online platforms such as gaming or
366 social media, may not have participated in this survey. Although this findings may not
367 represent the opinion of all internet users, a face-to-face survey was not possible at the time
368 due to the lockdown measures. Despite these limitations, this was the first study to investigate
369 the associations between extensive internet use and physical health symptoms in a sample of

370 adolescents in mainland Bangladesh. Future studies could address nesting follow-ups in
371 existing random samples that could provide timely access to longitudinal population-
372 representative data.

373

374 **Conclusions**

375 In summary, the study demonstrated that uncontrolled use of the internet is prevalent among
376 the respondents in this study as more than two-third reported excessive use of the internet
377 during the lockdown, and this doubled the adverse effects on their physical health compared
378 with seldom users. Headaches, back pain, numbness of the fingers and neck pain were
379 predominant among internet users as spending 3 hours or more each day on the internet led to
380 a significant increase in the level of adverse physical health problems. The positive
381 associations between physical complaints and non-healthcare workers, middle to high-income
382 earners, indicate the target group for interventions to minimize this growing epidemic. The
383 health department of the Bangladesh government has now recognized extensive internet
384 usage as a serious public health problem. Arguably, it is time for the World Health
385 Organization and health departments worldwide to develop effective health policies to
386 increase public awareness of extensive internet use, especially during the pandemic. This has
387 the potential to affect millions of children and adults and, eventually the societies and
388 economies.

389

390

391 **References**

- 392 1. De' R, Pandey N, Pal A. Impact of digital surge during Covid-19 pandemic: A
393 viewpoint on research and practice. *Int J Inf Manage.* 2020;55(102171):102171.
394 2. Garfin DR. *Technology as a Coping Tool during the COVID-19 Pandemic:*
395 *Implications and Recommendations.* Stress and Health; 2020.
396 3. Kawohl W, Nordt C. COVID-19, unemployment, and suicide. *Lancet Psychiatry.*
397 2020;7(5):389-390.
398 4. Király O, Potenza MN, Stein DJ, King DL, Hodgins DC, Saunders JB, Griffiths MD,
399 Gjoneska B, Billieux J, Brand M, Abbott MW, Chamberlain SR, Corazza O,
400 Burkauskas J, Sales CMD, Montag C, Lochner C, Grünblatt E, Wegmann E,
401 Martinotti G, Lee HK, Rumpf H-J, Castro-Calvo J, Rahimi-Movaghar A, Higuchi S,
402 Menchon JM, Zohar J, Pellegrini L, Walitza S, Fineberg NA, Demetrovics Z.
403 Preventing problematic internet use during the COVID-19 pandemic: Consensus
404 guidance. *Compr Psychiatry.* 2020;100(152180):152180.
405 5. Ito M, Horst HA, Bittanti M, Herr Stephenson B, Lange PG, Pascoe CJ, Robinson L.
406 Living and learning with new media. In: The MIT Press; 2009:128.
407 6. O'Keeffe GS, Clarke-Pearson K, Council on Communications and Media. The impact
408 of social media on children, adolescents, and families. *Pediatrics.* 2011;127(4):800-
409 804.
410 7. Hinduja S, Patchin JW. Offline consequences of online victimization: School violence
411 and delinquency. *Journal of school violence.* 2007;6(3):89–112.
412 8. Internet. Gov.bd. Accessed August 6, 2021. <http://www.btrc.gov.bd/telco/internet>
413 9. Ayyagari, Grover, Purvis. Technostress: Technological Antecedents and
414 Implications. *MIS Q.* 2011;35(4):831.
415 10. Tarafdar M, Tu Q, Ragu-Nathan BS, Ragu-Nathan TS. The impact of technostress on
416 role stress and productivity. *J Manag Inf Syst.* 2007;24(1):301-328.
417 11. Kalia A. *The Zoom Boom: How Video Calling Became a Blessing—and a Curse.* The
418 Guardian; 2020.
419 12. Osuagwu UL, Miner CA, Bhattarai D, Mashige KP, Oloruntoba R, Abu EK,
420 Ekpenyong B, Chikasirimobi TG, Goson PC, Oveneri-Ogbomo GO, Langsi R,
421 Charwe DD, Ishaya T, Nwaeze O, Agho KE. Misinformation about COVID-19 in
422 sub-Saharan Africa: Evidence from a cross-sectional survey. *Health Secur.*
423 2021;19(1):44-56.
424 13. Lee JJ, Kang K-A, Wang MP, Zhao SZ, Wong JYH, O'Connor S, Yang SC, Shin S.
425 Associations between COVID-19 misinformation exposure and belief with COVID-
426 19 knowledge and preventive behaviors: Cross-sectional online study. *J Med Internet*
427 *Res.* 2020;22(11):e22205.
428 14. Zheng Y, Wei D, Li J, Zhu T, Ning H. Internet use and its impact on individual
429 physical health. *IEEE Access.* 2016;4:5135-5142.
430 15. Usgaonkar U, Shet Parkar SR, Shetty A. Impact of the use of digital devices on eyes
431 during the lockdown period of COVID-19 pandemic. *Indian J Ophthalmol.*
432 2021;69(7):1901-1906.
433 16. Lam LT, Peng ZW, Mai JC, Jing J. Faktorer forbundet med internettavhengighet blant
434 ungdom. *Cyberpsychol Behav.* 2009;12:551–555.

- 435 17. Heinze G, Dunkler D. Five myths about variable selection. *Transpl Int*. 2017;30(1):6-
436 10.
- 437 18. Vatcheva KP, Lee M, McCormick JB, Rahbar MH. Multicollinearity in regression
438 analyses conducted in epidemiologic studies. *Epidemiology (Sunnyvale)*. 2016;6(2).
439 doi:10.4172/2161-1165.1000227
- 440 19. Morrison CM, Gore H. The relationship between excessive Internet use and
441 depression: a questionnaire-based study of 1,319 young people and
442 adults. *Psychopathology*. 2010;43(2):121-126.
- 443 20. Cao H, Sun Y, Wan Y, Hao J, Tao F. Problematic Internet use in Chinese adolescents
444 and its relation to psychosomatic symptoms and life satisfaction. *BMC Public Health*.
445 2011;11:802.
- 446 21. Tsai HF, Cheng SH, Yeh TL, Shih CC, Chen KC, Yang YC, Yang YK. The risk
447 factors of Internet addiction—a survey of university freshmen. *Psychiatry research*.
448 2009;167(3):294–299.
- 449 22. López-Bravo A, García-Azorín D, Belvís R, González-Oria C, Latorre G, Santos-
450 Lasaosa S, Guerrero-Peral AL. Impact of the COVID-19 pandemic on headache
451 management in Spain: an analysis of the current situation and future
452 perspectives. *Neurol (Engl Ed)*. 2020;35(6):372-380.
- 453 23. Chen C-Y, Chen I-H, Pakpour AH, Lin C-Y, Griffiths MD. Internet-related behaviors
454 and psychological distress among schoolchildren during the COVID-19 school
455 hiatus. *Cyberpsychol Behav Soc Netw*. 2021;(cyber.2020.0497).
456 doi:10.1089/cyber.2020.0497
- 457 24. Servidio R, Bartolo MG, Palermi AL, Costabile A. Fear of COVID-19, depression,
458 anxiety, and their association with Internet addiction disorder in a sample of Italian
459 students. *Journal of Affective Disorders Reports*. 2021;4(100097):100097.
- 460 25. Tripathi A. Impact of internet addiction on mental health: An integrative therapy is
461 needed. *Integr Med Int*. 2018;4(3-4):215-222.
- 462 26. Wieckiewicz M, Danel D, Pondel M, Smardz J, Martynowicz H, Wieczorek T, Mazur
463 G, Pudlo R, Wieckiewicz G. Identification of risk groups for mental disorders,
464 headache and oral behaviors in adults during the COVID-19 pandemic. *Sci Rep*.
465 2021;11(1):10964.
- 466 27. Kelley KJ, Gruber EM. Psychometric properties of the Problematic Internet Use
467 Questionnaire. *Comput Human Behav*. 2010;26(6):1838-1845.
- 468 28. Yen C-F, Ko C-H, Yen J-Y, Chang Y-P, Cheng C-P. Multi-dimensional
469 discriminative factors for Internet addiction among adolescents regarding gender and
470 age: Internet addiction in adolescence. *Psychiatry Clin Neurosci*. 2009;63(3):357-364.
- 471 29. Lam LT, Peng Z-W, Mai J-C, Jing J. Factors associated with Internet addiction
472 among adolescents. *Cyberpsychol Behav*. 2009;12(5):551-555.
- 473 30. Lata LN, Walters P, Roitman S. The politics of gendered space: Social norms and
474 purdah affecting female informal work in Dhaka, Bangladesh. *Gend Work Organ*.
475 2021;28(1):318-336.
- 476 31. Husain T, Uddin MM, Chowdhury SA, Kalimullah NA. Mental Health Status of
477 Adolescents During the COVID-19 pandemic: A cross-sectional survey among the
478 Bangladeshi graduate students at Dhaka city. *bioRxiv*. Published online 2020.
479 doi:10.1101/2020.11.12.379487

- 480 32. Islam MS, Sujan MSH, Tasnim R, Ferdous MZ, Masud JHB, Kundu S, Mosaddek
481 ASM, Choudhuri MSK, Kircaburun K, Griffiths MD. Problematic internet use among
482 young and adult population in Bangladesh: Correlates with lifestyle and online
483 activities during the COVID-19 pandemic. *Addict Behav Rep.*
484 2020;12(100311):100311.
- 485 33. Ko N-Y, Lu W-H, Chen Y-L, Li D-J, Wang P-W, Hsu S-T, Chen C-C, Lin Y-H,
486 Chang Y-P, Yen C-F. COVID-19-related information sources and psychological well-
487 being: An online survey study in Taiwan. *Brain Behav Immun.* 2020;87:153-154.
- 488 34. Shaw M, Black DW. Internet addiction: Definition, assessment, epidemiology
489 and??Clinical management. *CNS Drugs.* 2008;22(5):353-365.
- 490 35. ICD-11 for mortality and morbidity statistics. Who.int. Accessed August 6, 2021.
491 <https://icd.who.int/browse11/l-m/en>

492

493

494

495

496 **Supplementary Table 1** Results of Multiple Regression Analysis in Prediction of Level of
497 Physical Complaints Among Internet Users. *Bold are significant variables.*

Variables	Adjusted Coefficients [95%CI]	P-value
Gender		
Male	Reference	
Female	-0.08 [-0.20, 0.04]	0.179
Age groups		
18-27 yrs	Reference	
28-37 yrs	0.09 [-0.13, 0.30]	0.440
38+ yrs	-0.03 [-0.29, 0.23]	0.818
Marital status		
Single	Reference	
married	-0.08[-0.29, 0.12]	0.431
Divorced/widow	-1.09[-1.44, -0.74]	<0.0005
Place of residence		
Barisal Division	Reference	
Chittagong Division	-0.70 [-0.99, -0.41]	<0.0005
Dhaka Division	-1.27 [-1.51, -1.02]	<0.0005
Khulna Division	-0.42 [-0.70, -0.14]	0.003
Mymensingh Division	-0.31 [-0.59, -0.03]	0.031
Rajshahi Division	-0.29 [-0.60, 0.01]	0.060
Rangpur Division	-0.11[-0.43, 0.21]	0.487
Sylhet Division	-0.40[-0.74, -0.07]	0.018
Mother's Level of Education		
Higher education	Reference	
Bachelor	-0.00 [-0.16, 0.16]	0.976
Intermediate (11–12)	0.01 [-0.19, 0.21]	0.924
Working status		
Employed	Reference	
Not employed/student	-0.49 [-0.72, -0.26]	<0.0005
Income in Taka		
lower-income (<30,000)	Reference	
middle-income (30,000-70,000)	0.72 [0.48, 0.96]	<0.0005
High-income (>70,000)	0.26 [-0.00, 0.53]	0.052
Occupation		
Healthcare workers	Reference	
Non-health care worker	0.62 [0.47, 0.78]	<0.0005
Internet use Frequency		
Seldom	Reference	-
Casual	0.36 [-0.02, 0.74]	0.067
Regular	0.52 [0.18, 0.85]	0.003
Frequent	1.21 [0.88, 1.54]	<0.0005
Intense	2.24 [1.91, 2.57]	<0.0005
Constant	3.18 [2.69, 3.67]	<0.0005

498 [†]The final model included Gender, Age group, Division, Mother's education, Income status, Employment status, Occupational status, Marital
499 status and Internet use frequency.