1

Original Article 1

Internet Use impact on Physical Health during COVID-19 pandemic in Bangladesh: A 2 Web-based Cross-sectional study 3

- 4
- Authors: Tanvir Abir¹, Uchechukwu Levi Osuagwu², Dewan Muhammad Nur-A Yazdani³, 5
- Abdullah Al Mamun⁴, Kaniz Kakon⁵, Anas A. Salamah⁶, Noor Raihani Zainol⁷, Mansura 6
- Khanam⁸, Kingsley Emwinyore Agho⁹ 7

Affiliation: 8

9 ¹. Associate Professor, College of Business Administration—CBA, International University of Business, Agriculture and Technology-IUBAT University, Dhaka 1230, Bangladesh; 10 tanvir.cba@iubat.edu: 11

². Diabetes, Obesity and Metabolism Translational Research Unit, Western Sydney 12 13 University, Campbelltown, NSW 2560, Australia; African Vision Research Institute (AVRI), University of KwaZulu-Natal, Westville Campus, Durban, 3629, South 14 Africa: l.osuagwu@westernsvdnev.edu.au: 15

³. Assistant Professor, College of Business Administration—CBA, International University of 16 Business, Agriculture and Technology-IUBAT University, Dhaka 1230, Bangladesh; 17 dewanm@iubat.edu 18

19 ⁴. Faculty of Business and Management, UCSI University, Kuala Lumpur 56000, Malaysia; abdullaham@ucsiuniversitv.edu.mv: 20

21

⁵. Assistant Professor, Department of Philosophy, College of Arts and Sciences-CAAS, 22 International University of Business, Agriculture and Technology-IUBAT University, 23 24 Dhaka 1230, Bangladesh; kanizkakon@iubat.edu

- 25 ⁶. Department of Management Information Systems, College of Business Administration,
- Prince Sattam Bin Aziz University, 165 Al-Kharj 11942, Saudi Arabia. Email: 26
- a.salameh@psua.edu.sa 27
- ⁷. Lecturer, Faculty of Entrepreneurship and Business, Universiti Malaysia Kelantan, 16100 28 Kota Bharu, Malaysia; raihani@umk.edu.my 29
- ⁸. Maternal and Child Nutrition, Nutrition and Clinical Services Division, icddr,b 30
- Bangladesh; mansura@icddrb.org 31
- ⁹. Associate Professor, School of Health Science, Western Sydney University, Campbelltown, 32
- NSW 2560, Australia; African Vision Research Institute (AVRI), University of KwaZulu-33
- Natal, Westville Campus, Durban, 3629, South Africa; K.Agho@westernsydney.edu.au 34
- * Correspondence: Tanvir Abir (PhD), Associate Professor, College of Business 35
- 36 Administration—CBA, International University of Business, Agriculture and Technology—
- INJREATHISUTATISTY IS Debate and and the stange ball scherification of the ball of the stand of 37

2

39	Funding: This research did not receive any funding.
40 41	Declaration of Conflict of Interest: The authors declare that they have no known competing
42	financial interests or personal relationships that could have appeared to influence the work
43	reported in this paper.
44 45	Author Contribution: Conceptualization, T.A. and D.M.NA. Y; Data curation, T.A. and
46	K.A; Formal analysis, U.L.O. and K.A; Investigation, D.M.NA.Y. and M.K.; Methodology,
47	T.A., A.A.M, A.A.S.; Project administration, K.A., D.M.NA.Y., K.K.; Resources, D.M.N
48	A.Y., A.A.M. and N.R.Z; Supervision, N.R.Z. and K.A.; Validation, T.A. and K.A.;
49	Visualization, M.K. and K.A.; Writing-original draft, T.A. and U.L.O.
50	All authors have read and agreed to the published version of the manuscript.
51 52	Article Type: Original Research.
53	Keywords: Internet use, Coronavirus, Survey, Headache, Back pain, Neck pain, Physical
54	health
55	Running Head: Internet use and Physical health of Bangladeshi Adults
56 57	Ethics Statement: Permission for this study (IUBAT/AR/2021/001) was obtained from the
58	Institutional Review Board of International University of Business, Agriculture and
59	Technology (IUBAT), Dhaka, Bangladesh. The study adhered to the tenets of the Declaration
60	of Helsinki as revised in Fortaleza. All the participants were informed about the specific
61	objective of this study before proceeding to the questionnaire. Consent was obtained from all
62	participants prior to completing the questionnaire through an online preamble. Participants
63	were able to complete the survey only once and could terminate the survey at any time they
64	desired. Anonymity and confidentiality of the data were ensured.
65	Data Reference: Our data are included in the manuscript and raw data can be released at

66 reasonable request.

67

3

68 Abstract

Background: Bangladesh ranked fifth largest internet user in Asia. Past studies on internet use have focused on its impact on mental health, with little known about its impact on the physical health of individuals during COVID-19 pandemic. Hence, this study examines the impact of Internet use frequency on physical health during the Covid-19 lockdown in Bangladesh.

Methods: A web-based cross-sectional study on 3242 individuals aged 18 and above was
conducted from 2nd August – 1st October 1, 2020, during the lockdown in Bangladesh. The
survey covers demographics, Internet use frequency and physical health questions. Multiple
linear regression analyses were used to examine the impact of internet use frequency on
physical health.

Results: The result indicated that 72.5%, 69.9%, 65.1% and 55.3% reported headache, back pain, numbness of the fingers and neck pain, respectively. The multivariable analyses showed increased physical health impact among regular (coefficient $\beta = 0.52$, 95% confidence interval [CI]: 0.18-0.85, *P*=0.003), frequent ($\beta = 1.21$, 95% CI: 0.88–1.54, *P* < .001) and intense (β = 2.24, 95% CI: 1.91–2.57, *P* < .001) internet users. Other factors associated with physical health scores were gender, income (in Taka), occupation, regions, and working status.

85 Conclusion: Frequent, intensive, and extensive use of the internet were strong predictors of 86 increased physical health problems, and the study suggests the need for raising awareness of 87 physical health problems triggered by high internet users among the high socioeconomic 88 group in Bangladesh.

89

90

4

91 INTRODUCTION

The coronavirus disease 2019 (COVID-19) pandemic and the associated public health 92 interventions, including social distancing, home quarantine measures, and 'stay-at-home' 93 orders, put in place by the various governments, led to a surge in internet usage during the 94 pandemic^[1]. The lockdown measures resulted in widespread and unprecedented social 95 disruption^[2] as non-essential businesses were closed or some employees worked from home 96 and unemployment increased ^[3]. There were massive changes in internet usage patterns and 97 user behaviour with employees having to adjust to the new "normals" - with meetings going 98 completely online, office work shifting to the home, leading to new emerging patterns of 99 work. These changes have come across most organizations, businesses, societies, or 100 governments and suddenly, leaving barely any time for organizations and people to plan for, 101 prepare and implement new setups and arrangements. Society has had to adjust, try, 102 experiment, and find ways that did not exist before the pandemic ^[1], and these may have 103 implications on the physical health of individuals. 104

People are now spending even more time with technology while consuming news 105 media, watching television, using social media to connect with others, utilizing lifestyle apps 106 to shop for groceries and other consumer goods, and engaging in home workouts ^[2]. In 107 Bangladesh, this kept parts of the economy going ^[4] as the country witnessed a boom in 108 109 internet usage due to the fast-growing mobile internet and the government's push for digitalization. Engaging in social media platforms is now a routine activity for children and 110 adults, enhancing communication, social connection, and even technical skills ^[5]. Platforms 111 such as Facebook and Twitter offer multiple daily opportunities for connecting with friends, 112 classmates, and people with shared interests. The number of people using such sites has also 113 increased dramatically over the last five years, as shown in a recent poll ^[6]. A past study 114 reported that about 22% of teenagers log on to their favorite social media site more than ten 115 times a day, and more than half of adolescents log on to a social media site more than once a 116

5

day ^[6,7]. Seventy-five percent of teenagers now own cell phones, 25% use them for social
media, 54% use them for texting, and 24% use them for instant messaging ^[7].

According to the Bangladesh Telecommunication Regulatory Commission, there were 119 99,428 million (approx. 60.9% of the population) internet users in February 2020^[8]. Though 120 there is anecdotal early evidence that this may lead to increased productivity, it has also led to 121 increased stress ^[9,10] where employees must learn new technologies, be available for work at 122 almost all times, stay with digital devices all the time, and cope with multi-tasking ^[1]. The 123 use of video-conferencing and content delivery services such as Zoom and Akamai also 124 increased ^[1]. Those working from home using this technology have found themselves under 125 intense scrutiny and tension^[11]. 126

127 That a large part of the peoples' social and emotional development is occurring while on the Internet and cell phones may affect the physical health of individuals considering the 128 long hours spent on the internet during the mandatory stay at home orders which expose the 129 population to lots of misinformation ^[12] known to affect physical and mental health ^[13]. Since 130 this high use of information systems may become the new normal, this study examined the 131 impact of internet usage frequency during the COVID-19 pandemic on Bangladeshi residents' 132 physical and mental health and determined the level of addiction to social media platforms 133 used by the people. 134

135 **Ethical consideration**

Permission for this study (IUBAT/AR/2021/001) was obtained from the Institutional Review Board of International University of Business, Agriculture and Technology (IUBAT), Dhaka, Bangladesh. The study adhered to the tenets of the Declaration of Helsinki as revised in Fortaleza. All the participants were informed about the specific objective of this study before and consent was obtained from all participants before completing the questionnaire through an online preamble. To avoid repeated responses and in order to ensure the validity of data to

6

some extent, participants were able to complete the survey only once as data was restricted to
their IP addresses and device. Participants could terminate the survey at any time they
desired. Anonymity and confidentiality of the data were ensured.

145 METHODS

146 Study population and sampling

By convenience sampling, 3242 questionnaires were collected from adults aged 18 and over 147 living in eight divisions across Bangladesh during the pandemic. All questionnaires were 148 web-based, participation was voluntary, and each participant was informed about the study's 149 background through an online preamble. No reward or incentive was offered for 150 participation. There may be more than one questionnaire filled by the same person that 151 affects the statistical analysis results of questionnaire data. In order to ensure the validity of 152 data to some extent, restricting the same IP address and device are applied to the system, 153 which means the participant of the same IP address or device only can fill once. 154

155 Data Collection

Data were collected using self-administered web-based questionnaires distributed to various groups of people via the E-link using numerous mailing lists and social network sites including Facebook, WhatsApp and Twitter. Due to the mandatory lockdown, no paper questionnaires were distributed, making it difficult to reach the very remote regions.

160 **Dependent variable**

161 Ten items of physical complaints reported in *Table 2* below were used to determine the 162 impact of individual physical health on Internet use. The respondents used a close-end 163 response ['Yes' or 'No'] to indicate whether they suffered from the following ten problems 164 (Back and neck pain, numbress in fingers, headaches, inability to sleep, dry eyes or other 165 vision problems, poor nutrition, poor personal hygiene, weight gain/loss and loss of appetite] 166 during or after pro-longed Internet use. Binary scores for individual questions were summed

7

to give a physical health score which ranged from 0-10, and the Kuder Richardson coefficient
measuring internal consistency among the physical health scores ranged from 0.65 to 0.78,
indicating a satisfactory level of reliability.

170

171 Main study factor

The main study variable was the respondent's Internet usage, including the average daily amount of time spent on the internet and dependence on networks. The frequency of internet use was categorized as 'seldom' (<1 hour per day), 'casual' (≥ 1 and <3 hours per day), 'regular' (≥ 3 and <5 hours per day), 'frequent' (≥ 5 and <7 hours per day), 'intense' (≥ 7 hours per day) ^[14].

177

178 Independent variables

Independent variables used for the study analyses were based on previous studies ^[14-16].
Independent variables included demographic characteristics (i.e., gender, age [categorized as
181 18-27, 28-37, 38 years and over], marital status [single, married, divorced/widowed], mothers
182 level of education [Higher education, Bachelor, Intermediate], place of residence,
183 employment status [Employed, unemployed/student], occupation [healthcare and non184 healthcare worker], and income in Taka [lower, middle and high-income earners].

185

186 Statistical Analysis

All statistical analyses were performed in Stata version 14.1 (Stata Corp. 2015, College Station USA) mainly using descriptive statistics; t-test used compares the means of two independent groups, One-way analysis of variance (one-way ANOVA) used compares the means of more than two independent groups, correlation analyses, and regression analysis. All confounding variables with a P < .20 were retained in the univariate linear regression analysis and were used to build a multivariable model ^[17]. A manual elimination procedure

8

was applied for multivariate linear regression analyses to remove non-significant variables (P> .05). The main study factor variable (Internet usage frequency) was added to all significant confounding variables after elimination processes. The main study factor and independent variables associated with physical health scores (P < .05) were reported. In the regression analysis, we checked for homogeneity of variance and multicollinearity, including Variance Inflation Factors (VIF) and the VIF < 4 was considered suitable ^[18]:

199

200 **RESULTS**

Demographic Characteristics According to the Frequency of Internet Use among adults in Bangladesh

Table 1 depicts the main demographic characteristics, including gender, education and age
group on the five groups extracted. It is worth noting that the education in this paper refers to
that of the mothers' maximum educational level. The majority of participants were males
(1260, 61.1%), married (2632, 81.2%) and employed (2864, 88.3%) in non-healthcare sectors
(2567, 79.2%), and for many, their mothers had completed a bachelor degree or higher.
About half of the respondents earned more than 70,000 Taka at the time of data collection
(*Table 1*).

About two-thirds of the respondents that reported intense use of the internet were males, while females were more likely to report regular and casual use of the internet. More than half of intense and frequent internet users were aged 28-37years, and a similar proportion of casual internet users were younger (18-27years). Nearly all intense internet users were employed and working in the non-Health care sector during the study.

Table 1 Demographic Characteristics According to the Frequency of Internet Use among
 adults in Bangladesh. Values are expressed as numbers and percentages (n, %).

Variables	Total	Intense	Frequent	Regular	Casual	Seldom
	(N=3236)	(n=1668)	(n=689)	(n=559)	(n=187)	(n=93)
Demography						

All rights reserved. No reuse allowed without permission.

9

Gender Male 1976 (61.1) 1179 (70.7) 382 (55.4) 264 (44.1) 93 (44.7) 58 (62.4) Female 1260 (38.9) 489 (29.3) 335 (55.9) 35 (37.6) 307 (44.6) 94 (50.3) Age in years 18-27 555 (17.2) 30 (1.8) 156 (22.6) 254 (42.1) 98 (52.4) 17 (17.9) 28-37 238 (39.5) 1657 (51.1) 956 (57.3) 366 (53.1) 60 (32.1) 37 (39.0) 38 +1030 (31.8) 682 (40.9) 167(24.2) 111 (18.4) 29 (15.5) 41 (43.2) Marital status Single 508 (15.7) 150 (21.8) 223(37.0) 90 (48.1) 16 (16.8) 29 (1.7) Married 2632 (81.2) 1632 (97.8) 505 (73.3) 339(56.2) 83 (44.40) 73 (76.8) Divorced/widow 102 (3.1) 7 (0.4) 34 (4.9) 41 (6.8) 14(7.49) 6 (6.3) *Place of residence* (%) 70 (4.20) **Barisal Division** 172 (5.31) 48 (7.0) 35 (5.0) 7 (3.7) 12 (12.6) Chittagong Division 291 (9.0) 100 (6) 107 (15.5) 61 (10.1) 16(8.6) 7(7.4) Dhaka Division 1574 (48.6) 1010 (60.6) 205 (29.8) 236 (39.1) 95 (50.8) 28 (29.5) Khulna Division 352 (10.9) 115 (6.9) 109 (15.8) 92 (15.3) 20 (10.7) 16 (16.8) 313 (9.7) 17 (9.1) Mymensingh Division 127 (7.6) 76 (11.0) 85 (14.1) 8 (8.4) Rajshahi Division 213 (6.6) 99 (5.9) 47 (6.8) 41 (6.8) 18 (9.6) 8 (8.4) **Rangpur Division** 179 (5.5) 93 (5.6) 45 (6.5) 22 (3.7) 9 (4.8) 10 (10.5) Sylhet Division 148 (4.6) 54 (3.2) 52 (7.6) 31 (5.1) 5 (2.7) 6 (6.3) Mother's Level of Education Higher education 1287 (39.7) 684 (41.0) 341 (49.5) 178 (29.5) 46 (24.6) 38 (40.) (above Bachelor 1514 (46.7) 961 (57.6) 226 (32.8) 233 (38.6) 27(28.4)67 (35.8) Intermediate (11–12) 192 (31.84) 74 (39.5) 30 (31.6) 441 (13.6) 23 (1.4) 122 (17.7) Working status Employed 2864 (88.3) 1647 (98.7) 600 (87.1) 432 (71.6) 99 (52.9) 86 (91.0) Not employed/student 9 (9.47) 378 (11.7) 21 (1.3) 89 (12.9) 171 (28.4) 88 (47.1) Income in Taka lower-income 204 (6.3) 14 (0.9) 37 (5.4) 102 (16.9) 41 (21.9) 10 (10.4) (<30,000) 510 (30.9) 389 (64.5) middle-income 1496 (46.1) 433 (62.8) 108 (57.8) 56 (59.0) (30,000-70,000)High-income 1542 (47.7) 1144 219 (31.8) 112 (18.8) 38 (20.3) 29 (30.5) (>70,000) (68.59) **Occupation** Healthcare workers 675 (20.8) 212 (30.8) 304 (50.4) 89 (47.6) 39 (41.1) 31 (1.9) Non-health care 2567 (79.2) 1637 (98.1) 477 (69.2) 299 (49.6) 98 (52.4) 56 (59.0) worker

217

Prevalence and 95% confidence intervals of physical complaints by gender among internet users.

220

Table 2 depicts the prevalence and 95%CIs of the adverse effects on individual physical
health complaints examined in this study in males and females. Headache 72.50% [95%CI
70.93%, 74.01%] was the prevalent complaint of the participants, while 69.87% [68.27%,
71.43%] and 65.10% [63.42%, 66.71%] complained of Back pain and numbness of the

10

fingers, respectively. Males reported a higher prevalence of back pain, numbness of the 225

finger, headaches, and neck pains than females as the confidence intervals do not overlap (see 226

Table 2). By contrast, females reported a higher prevalence of poor nutrition, poor personal 227

hygiene, and loss of appetite compared with men (Table 2). 228

Table 2 Gender differences in the prevalence of physical health scores among internet users 229

in Bangladesh. Significant variables between males and females are bolded because the 230

231 confidence intervals did not overlap.

	п	Prevalence (95%CI)		
Physical complaints		Overall	Male	Female
Back pain	2261	69.9 [68.27, 71.43]	78.14 [76.26, 79.91]	56.90 [54.15, 59.62]
Finger numbness	2106	65.10 [63.42, 66.71]	72.37 [70.35, 74.30]	53.65 [50.89, 56.39]
Headaches	2346	72.50 [70.93, 74.01]	78.80 [76.94, 80.54]	62.62 [59.91, 65.25]
Inability to sleep	1396	43.14 [41.44, 44.85]	41.95 [39.79, 44.14]	45.00 [42.27, 47.76]
Poor Nutrition	746	23.05 [21.63, 24.54]	16.55 [14.97, 18.25]	33.25 [30.70, 35.91]
Poor Personal Hygiene	605	18.70 [17.39, 20.08]	14.98 [13.47, 16.62]	24.52 [22.23, 26.98]
Neck pain	1789	55.28 [53.56, 56.99]	58.50 [56.31, 60.66]	50.24 [47.48, 53.00]
Dry eyes/other vision	1815	56.09 [54.37, 57.79]	57.54 [55.35, 59.70]	53.81 [51.05, 56.55]
problems				
Weight gain/loss	1653	51.08 [49.36, 52.80]	52.43 [50.22, 54.63]	48.97 [46.21, 51.73]
Loss of Appetite	499	15.42 [14.22, 16.71]	13.77 [12.31, 15.36]	18.02 [15.99, 20.24]

232

CI=Confidence interval; n= number of respondents

Figure 1 shows the mean scores of physical health by internet use frequency in males 233 and females. From the figure, the mean scores of physical health by intensive internet use 234 was 5.6 for males and 5.4 for females. This physical health mean scores could be translated 235 as percentage mean scores of males (56%) and females (54%), respectively. Physical health 236 237 mean scores were lower among the casual user of internet (3.2 for females and 3.0 for males)

than seldom users of internet (3.1 for females and 3.4 for males) 238

11



239 Figure 1. 240

Mean scores by gender of the physical health by internet use frequency in Bangladesh 241 242

243

Unadjusted Analysis for the association between prolonged internet use and physical 244 health complaints 245

Table 3 shows the mean scores of physical health complaints and the unadjusted coefficients 246 for the associations with the sociodemographic variables. From the table, the mean scores for 247 248 physical health complaints varied across variables. The unadjusted coefficients revealed 249 significantly higher physical health problems due to prolonged internet use in those aged >28 years, married, completed bachelor education, middle-high income earners and non-250 251 healthcare workers compared with other groups. In contrast, females, those who were 252 divorced or widowed, those with intermediate education, non-workers/students and respondents who resided in Chittagong, Dhaka and Khulna during the study period had 253 significantly lower mean scores of physical health complaints from prolonged internet usage 254 255 compared with the other groups.

256

Table 3: Mean scores and Univariate linear regression of Physical Complaints among 257 258 Internet Users.

perpetuity. All rights reserved. No reuse allowed without permission.

12

Variables	Mean scores (SD)	Unadjusted Coefficient [95%CI]			
Gender					
Male	4.85 (1.83)***	Reference			
Female	4.47 (1.88)	-0.38 [-0.51, -0.25]			
Age,					
18-27 yrs	3.52 (1.71)***	Reference			
28-37 yrs	5.02 (1.81)	1.50 [1.33, 1.67]			
38+ yrs	4.80 (1.77)	1.31 [1.12, 1.49]			
Marital status					
Single	3.67 (1.76)***	Reference			
married	4.96 (1.79)	1.30 [1.13, 1.47]			
Divorced/widow	3.00 (1.19)	-0.67 [-1.04, -0.29]			
Place of residence					
Barisal Division	5.35 (1.83)***	Reference			
Chittagong Division	4.66 (1.87)	-0.69 [-1.04, -0.35]			
Dhaka Division	4.36 (1.77)	-0.99 [-1.28, -0.70]			
Khulna Division	4.83 (1.83)	-0.52 [-0.85, -0.19]			
Mymensingh Division	5.03 (1.92)	-0.32 [-0.66, 0.02]			
Rajshahi Division	5.21 (1.91)	-0.14 [-0.51, 0.22]			
Rangpur Division	5.47 (1.88)	0.12 [-0.26, 0.50]			
Sylhet Division	4.99 (1.83)	-0.36 [-0.76, 0.04]			
Mother's Level of Education					
Higher education	4.71 (1.82)***	Reference			
Bachelor	5.00 (1.87)	0.30 [0.16, 0.43]			
Intermediate (11–12)	3.65 (1.50)	-1.06 [-1.26, -0.86]			
Working status					
Employed	4.89 (1.79)***	Reference			
Not employed/student	3.24 (1.72)	-1.65 [-1.84, -1.46]			
Income in Taka					
lower-income (<30,000	3.05 (1.50)***	Reference			
middle-income (30,000	4.82 (1.88)	1.77 [1.51, 2.04]			
High-income (>70,000)	4.80 (1.77)	1.75 [1.48, 2.01]			
Occupation					
Healthcare workers	3.62 (1.26)***	Reference			
-Non-health care workers SD=standard deviation; CI=confidence interval. **	$\frac{4.98 (1.89)}{P} < .001$ based on t-test and	1.36 [1.21, 1.51] Anova. Bolded are significant coefficients.			

260

259

261 The impact of the frequency of internet use on physical health scores

Figure 2 presents the unadjusted and adjusted odds ratio and their confidence intervals for the association between frequency of internet use and physical health symptoms during the COVID-19. In the unadjusted analysis, frequent and intensive internet users during the COVID-19 significantly reported higher physical health problems than seldom users of the internet. After adjusting for the independent variables used in this study, regular, frequent, and intensive internet users during the COVID-19 had significantly higher physical health problems than those who seldom used the internet (see *Figure 2*).

13



²⁷⁰ Independent variables adjusted for are Gender, Age group, Division of residence, Mothers Education, marital status, income, working status, occupation.

Figure 2: The impact of frequency of internet use on physical health scores: Unadjusted and 271 adjusted coefficient and their 95% confidence intervals. 272

273

Marital status, age, gender, place of residence, level of education, working status, 274 occupation, and income level were all associated with the physical health complaints among 275 276 internet users in Bangladesh (Supplementary Table S1 for details).

277

DISCUSSION 278

279 A considerable body of literature has emerged over the past two decades assessing the relationship between problematic or addictive (excessive) use of the internet and individuals' 280 psychological/mental well-being. Comparatively, very little research has evaluated the 281 relationship between various internet use frequencies and the individual's physical health. 282 The COVID-19 pandemic caused a drastic surge in the use of the internet globally and 283 Bangladesh^[1], and this cross-sectional study investigated the relationship between the 284 frequency of internet use and the physical health of Bangladeshi respondents. The findings 285 demonstrated that more than one in every two persons in this study used the internet for seven 286 or more hours each day. This was associated with 2.2 folds increase in physical health 287 problems compared with seldom users (<1 hour per day). Headache, back pain and numbness 288 of the fingers (reported by over seventy percent of internet users) were the predominant 289 complaints. There was a positive association between physical health problems and the 290 291 number of daily hours spent on the internet. After adjusting for the frequency of internet use,

14

we found that middle to high-income earners and those working in non-healthcare sectors 292 experienced more significant physical health problems due to internet use. In comparison, 293 students and widowed or divorced persons reported fewer physical health problems compared 294 with other respondents. 295

This study found that headache was the most prevalent symptom reported by internet 296 users in Bangladesh, followed by back pain and numbness of the fingers. These complaints 297 298 were significantly increased among frequent internet users and intense users, which agrees with the findings of significant association between extensive use of the internet and physical 299 health problems in previous studies ^[19-22]. A study in Spain reported that headache was the 300 most common complaint among internet users during the COVID-19 pandemic ^[22]. Similarly, 301 Cao et al.s study (2009), after examining adolescents, found that those who used the internet 302 303 excessively were more likely to report psychosomatic symptoms including lack of physical 304 energy, physiological dysfunction, weakened immunity, emotional symptoms, behavioural symptoms and social adaptation problems and had lower life satisfaction scores compared 305 with normal internet users ^[20]. 306

There are also reports that school children and other graduate students in China^[23] 307 and Italy ^[24] experienced increased psychological distress following excessive internet use 308 309 during the COVID-19 pandemic. Such addiction to the internet can result in the problem of self-care, difficulty in performing daily routine, and mental health effects of anxiety and 310 depression ^[25]. The negative effects of internet addiction on health may be related to the 311 alterations in the individuals neuroanatomical and neurochemical mechanisms, including 312 cortical thinning of various components of the brain and dopaminergic reward circuitry, 313 during excessive use of the internet ^[25]. While it is imperative that such behaviours, 314 especially during a pandemic, remain at a moderate and regulated level, identifying the 315 population at risk of the problematic use of the internet is necessary in order to prevent the 316 associated mental health problems ^[26] and improve work or school functions ^[27]. 317

15

The present study further revealed that, compared with women, a higher proportion of 318 men used the internet excessively during the pandemic and thus experienced more significant 319 physical health problems. However, this association was significant only when it interacted 320 with the frequency of internet use. Previous studies among Taiwanese residents ^[28] and those 321 in mainland China^[29] found that male gender was a potential risk factor for internet addiction 322 among adolescents. This association of gender and physical health complaints found in the 323 present study could be explained by the patriarchal nature of the Bangladesh society, where 324 women and girls rarely go outside of their homes even before the pandemic ^[30] but preferred 325 326 to stay home and spend more time on the Internet. Whereas these women may have developed adaptive strategies, it was a different experience for the male counterpart, which 327 were used to staying outside/working prior to the lockdown. For these men, the lockdown 328 329 measures and transition to work online and/or staying at home required them to adjust quickly, try, experiment, and find ways that did not exist before the pandemic^[1] as 330 educational institutions, businesses and workplaces were shut down for such a long duration 331 ^[31]. These adjustments may have affected their physical health. 332

Another finding of this study was the significant relationship between income class 333 and occupational status (non-healthcare worker) with the physical health problems 334 experienced by internet users. Earning above 30,000 Taka was associated with higher 335 physical health problems due to internet usage after adjusting for the frequency of internet 336 use. In line with this finding, Islam et al ^[32] on the correlation between problematic 337 (excessive) internet use (PIU) and lifestyle during the COVID-19 pandemic using data from a 338 large sample of Bangladeshi youth and adults (n=13,525) found that younger age, higher 339 education, cigarette smoking, more sleep, less physical activity, gaming and social media use 340 were associated with PIU^[32]. This research has depicted the physical health concerns related 341 to prolonged internet usage during the COVID-19 pandemic. A similar association was found 342 in this research between educational status, age and physical health complaints among 343

16

internet users in Bangladesh; however, these associations were dependent on the frequency of
internet use, which was not assessed in the previous study ^[32].

The finding that non-healthcare workers had higher physical health problems 346 compared with health care workers in this study is consistent with the finding of poorer 347 psychological well-being among non-health-care workers who used the Internet to obtain 348 information on COVID-19 compared with those who received COVID-19 information from 349 medical staff in health care settings ^[33]. Although this research were unable to identify 350 whether the non-healthcare workers in our study used the internet for COVID-19 information 351 352 retrieval, it has been shown that the influx of misinformation around COVID-19 via online platforms had adverse effects on people's health ^[33]. Such problematic internet use (PIU) can 353 lead to clinical impairment, distress ^[34] and/or deterioration in financial, familial, social, 354 educational and/or occupational domains ^[35]. 355

This study has some limitations. First, the cross-sectional design of this study does not 356 allow for causality or the direction of relationships to be determined. However, appropriate 357 analysis of cross-sectional data represents a useful initial step in identifying associations 358 between extensive internet usage and the physical health condition of the participants. 359 Second, as the responses were self-reported, it was not possible to verify the participants' 360 physical health complaints, which may lead to a response bias. Multiple assessments, 361 interviews, and informants may have provided a richer and more thorough understanding of 362 this topic. Third, since the survey was administered during COVID-19 lockdown using an 363 online platform, it is possible that some respondents, especially those living in rural areas 364 with limited internet access or people addicted to other online platforms such as gaming or 365 social media, may not have participated in this survey. Although this findings may not 366 represent the opinion of all internet users, a face-to-face survey was not possible at the time 367 due to the lockdown measures. Despite these limitations, this was the first study to investigate 368 369 the associations between extensive internet use and physical health symptoms in a sample of

17

adolescents in mainland Bangladesh. Future studies could address nesting follow-ups in
existing random samples that could provide timely access to longitudinal populationrepresentative data.

373

374 Conclusions

In summary, the study demonstrated that uncontrolled use of the internet is prevalent among 375 the respondents in this study as more than two-third reported excessive use of the internet 376 during the lockdown, and this doubled the adverse effects on their physical health compared 377 378 with seldom users. Headaches, back pain, numbness of the fingers and neck pain were predominant among internet users as spending 3 hours or more each day on the internet led to 379 a significant increase in the level of adverse physical health problems. The positive 380 associations between physical complaints and non-healthcare workers, middle to high-income 381 earners, indicate the target group for interventions to minimize this growing epidemic. The 382 health department of the Bangladesh government has now recognized extensive internet 383 usage as a serious public health problem. Arguably, it is time for the World Health 384 Organization and health departments worldwide to develop effective health policies to 385 increase public awareness of extensive internet use, especially during the pandemic. This has 386 the potential to affect millions of children and adults and, eventually the societies and 387 economies. 388

389

390

391 References

392	1.	De' R, Pandey N, Pal A. Impact of digital surge during Covid-19 pandemic: A
393		viewpoint on research and practice. Int J Inf Manage. 2020;55(102171):102171.
394	2.	Garfin DR. Technology as a Coping Tool during the COVID-19 Pandemic:
395		Implications and Recommendations. Stress and Health; 2020.
396	3.	Kawohl W, Nordt C. COVID-19, unemployment, and suicide. <i>Lancet Psychiatry</i> .
397		2020;7(5):389-390.
398	4.	Király O, Potenza MN, Stein DJ, King DL, Hodgins DC, Saunders JB, Griffiths MD,
399		Gjoneska B, Billieux J, Brand M, Abbott MW, Chamberlain SR, Corazza O,
400		Burkauskas J, Sales CMD, Montag C, Lochner C, Grünblatt E, Wegmann E,
401		Martinotti G, Lee HK, Rumpf H-J, Castro-Calvo J, Rahimi-Movaghar A, Higuchi S,
402		Menchon JM, Zohar J, Pellegrini L, Walitza S, Fineberg NA, Demetrovics Z.
403		Preventing problematic internet use during the COVID-19 pandemic: Consensus
404		guidance. Compr Psychiatry. 2020;100(152180):152180.
405	5.	Ito M, Horst HA, Bittanti M, Herr Stephenson B, Lange PG, Pascoe CJ, Robinson L.
406		Living and learning with new media. In: The MIT Press; 2009:128.
407	6.	O'Keeffe GS, Clarke-Pearson K, Council on Communications and Media. The impact
408		of social media on children, adolescents, and families. Pediatrics. 2011;127(4):800-
409		804.
410	7.	Hinduja S, Patchin JW. Offline consequences of online victimization: School violence
411		and delinquency. Journal of school violence. 2007;6(3):89-112.
412	8.	Internet. Gov.bd. Accessed August 6, 2021. http://www.btrc.gov.bd/telco/internet
413	9.	Ayyagari, Grover, Purvis. Technostress: Technological Antecedents and
414		Implications. <i>MIS Q</i> . 2011;35(4):831.
415	10.	. Tarafdar M, Tu Q, Ragu-Nathan BS, Ragu-Nathan TS. The impact of technostress on
416		role stress and productivity. J Manag Inf Syst. 2007;24(1):301-328.
417	11.	Kalia A. The Zoom Boom: How Video Calling Became a Blessing-and a Curse. The
418		Guardian; 2020.
419	12.	. Osuagwu UL, Miner CA, Bhattarai D, Mashige KP, Oloruntoba R, Abu EK,
420		Ekpenyong B, Chikasirimobi TG, Goson PC, Ovenseri-Ogbomo GO, Langsi R,
421		Charwe DD, Ishaya T, Nwaeze O, Agho KE. Misinformation about COVID-19 in
422		sub-Saharan Africa: Evidence from a cross-sectional survey. <i>Health Secur</i> .
423		2021;19(1):44-56.
424	13	Lee JJ. Kang K-A. Wang MP. Zhao SZ. Wong JYH. O'Connor S. Yang SC. Shin S.
425		Associations between COVID-19 misinformation exposure and belief with COVID-
426		19 knowledge and preventive behaviors: Cross-sectional online study. J Med Internet
427		<i>Res.</i> 2020:22(11):e22205.
428	14	Zheng Y. Wei D. Li J. Zhu T. Ning H. Internet use and its impact on individual
429		physical health. <i>IEEE Access</i> . 2016;4:5135-5142.
430	15	Usgaonkar U. Shet Parkar SR. Shetty A. Impact of the use of digital devices on eves
431	10	during the lockdown period of COVID-19 pandemic <i>Indian I Ophthalmol</i>
432		2021:69(7):1901-1906
433	16	Lam I.T. Peng ZW Mai IC Jing I. Faktorer forbundet med internettayhengighet blant
434	10.	ungdom Cybernsychol Rehay 2009:12:551-555
		anguonn cycerpsychol Denuv. 2007,12.331 333.

435	17.	Heinze G, Dunkler D. Five myths about variable selection. Transpl Int. 2017;30(1):6-
436		10.
437	18.	Vatcheva KP, Lee M, McCormick JB, Rahbar MH. Multicollinearity in regression
438		analyses conducted in epidemiologic studies. <i>Epidemiology (Sunnyvale)</i> . 2016;6(2).
439		doi:10.4172/2161-1165.1000227
440	19.	Morrison CM, Gore H. The relationship between excessive Internet use and
441		depression: a questionnaire-based study of 1,319 young people and
442		adults. Psychopathology. 2010;43(2):121-126.
443	20.	Cao H, Sun Y, Wan Y, Hao J, Tao F. Problematic Internet use in Chinese adolescents
444		and its relation to psychosomatic symptoms and life satisfaction. BMC Public Health.
445		2011;11:802.
446	21.	Tsai HF, Cheng SH, Yeh TL, Shih CC, Chen KC, Yang YC, Yang YK. The risk
447		factors of Internet addiction—a survey of university freshmen. Psychiatry research.
448		2009;167(3):294–299.
449	22.	López-Bravo A, García-Azorín D, Belvís R, González-Oria C, Latorre G, Santos-
450		Lasaosa S, Guerrero-Peral ÁL. Impact of the COVID-19 pandemic on headache
451		management in Spain: an analysis of the current situation and future
452		perspectives. Neurol (Engl Ed). 2020;35(6):372-380.
453	23.	Chen C-Y, Chen I-H, Pakpour AH, Lin C-Y, Griffiths MD. Internet-related behaviors
454		and psychological distress among schoolchildren during the COVID-19 school
455		hiatus. Cyberpsychol Behav Soc Netw. 2021;(cyber.2020.0497).
456		doi:10.1089/cyber.2020.0497
457	24.	Servidio R, Bartolo MG, Palermiti AL, Costabile A. Fear of COVID-19, depression,
458		anxiety, and their association with Internet addiction disorder in a sample of Italian
459		students. Journal of Affective Disorders Reports. 2021;4(100097):100097.
460	25.	Tripathi A. Impact of internet addiction on mental health: An integrative therapy is
461		needed. Integr Med Int. 2018;4(3-4):215-222.
462	26.	Wieckiewicz M, Danel D, Pondel M, Smardz J, Martynowicz H, Wieczorek T, Mazur
463		G, Pudlo R, Wieckiewicz G. Identification of risk groups for mental disorders,
464		headache and oral behaviors in adults during the COVID-19 pandemic. Sci Rep.
465		2021;11(1):10964.
466	27.	Kelley KJ, Gruber EM. Psychometric properties of the Problematic Internet Use
467		Questionnaire. Comput Human Behav. 2010;26(6):1838-1845.
468	28.	Yen C-F, Ko C-H, Yen J-Y, Chang Y-P, Cheng C-P. Multi-dimensional
469		discriminative factors for Internet addiction among adolescents regarding gender and
470		age: Internet addiction in adolescence. Psychiatry Clin Neurosci. 2009;63(3):357-364.
471	29.	Lam LT, Peng Z-W, Mai J-C, Jing J. Factors associated with Internet addiction
472		among adolescents. Cyberpsychol Behav. 2009;12(5):551-555.
473	30.	Lata LN, Walters P, Roitman S. The politics of gendered space: Social norms and
474		purdah affecting female informal work in Dhaka, Bangladesh. Gend Work Organ.
475		2021;28(1):318-336.
476	31.	Husain T, Uddin MM, Chowdhury SA, Kalimullah NA. Mental Health Status of
477		Adolescents During the COVID-19 pandemic: A cross-sectional survey among the
478		Bangladeshi graduate students at Dhaka city. <i>bioRxiv</i> . Published online 2020.
479		doi:10.1101/2020.11.12.379487

480	32. Islam MS, Sujan MSH, Tasnim R, Ferdous MZ, Masud JHB, Kundu S, Mosaddek
481	ASM, Choudhuri MSK, Kircaburun K, Griffiths MD. Problematic internet use among
482	young and adult population in Bangladesh: Correlates with lifestyle and online
483	activities during the COVID-19 pandemic. Addict Behav Rep.
484	2020;12(100311):100311.
485	33. Ko N-Y, Lu W-H, Chen Y-L, Li D-J, Wang P-W, Hsu S-T, Chen C-C, Lin Y-H,
486	Chang Y-P, Yen C-F. COVID-19-related information sources and psychological well-
487	being: An online survey study in Taiwan. Brain Behav Immun. 2020;87:153-154.
488	34. Shaw M, Black DW. Internet addiction: Definition, assessment, epidemiology
489	and??Clinical management. CNS Drugs. 2008;22(5):353-365.
490	35. ICD-11 for mortality and morbidity statistics. Who.int. Accessed August 6, 2021.
491	https://icd.who.int/browse11/l-m/en
492	
493	
494	
495	

21

Supplementary Table 1 Results of Multiple Regression Analysis in Prediction of Level of 496 Physical Complaints Among Internet Users. Bold are significant variables. 497

Variables	Adjusted Coefficients [95%CI]	P-value
Gender		
Male	Reference	
Female	-0.08 [-0.20, 0.04]	0.179
A go groups		
18-27 vrs	Reference	
28_{-37} yrs	0.09[-0.13, 0.30]	0.440
38+ yrs	-0.03 [-0.29, 0.23]	0.818
Marital status		
Single	Reference	0.421
married	-0.08[-0.29, 0.12]	0.431
Divorced/widow	-1.09[-1.44, -0.74]	<0.0005
Place of residence		
Barisal Division	Reference	
Chittagong Division	-0.70 [-0.99, -0.41]	< 0.0005
Dhaka Division	-1.27 [-1.51, -1.02]	< 0.0005
Khulna Division	-0.42 [-0.70, -0.14]	0.003
Mymensingh Division	-0.31 [-0.59, -0.03]	0.031
Rajshahi Division	-0.29 [-0.60, 0.01]	0.060
Rangpur Division	-0.11[-0.43, 0.21]	0.487
Sylhet Division	-0.40[-0.74, -0.07]	0.018
Mother's Level of Education		
Higher education	Reference	
Bachelor	-0.00 [-0.16, 0.16]	0.976
Intermediate (11–12)	0.01 [-0.19, 0.21]	0.924
Working status		
Employed	Reference	
Not employed/student	-0.49 [-0.72, -0.26]	< 0.0005
Income in Take		
lower income $(<30,000)$	Pafaranca	
(< 50,000)		<0.0005
High-income (>70,000)	0.26 [-0.00, 0.53]	< 0.0003
	- / -	
Occupation		
Healthcare workers	Reference	0 000 -
Non-health care worker	0.62 [0.47, 0.78]	< 0.0005
Internet use Frequency		
Seldom	Reference	-
Casual	0.36 [-0.02, 0.74]	0.067
Regular	0.52 [0.18, 0.85]	0.003
Frequent	1.21 [0.88, 1.54]	< 0.0005
Intense	2.24 [1.91, 2.57]	< 0.0005
_Constant The final model included Gender, Age group, Divis	3.18 [2.69, 3.67] sion, Mother's education, Income status, Employment	<0.0005 status, Occupational status, Marita

498 status and Internet use frequency. 499