

1 **Title:**

2 Treatment interruption is a risk factor for sickness presenteeism: a large-scale cross-sectional study
3 during the COVID-19 pandemic

4

5 **Authors:**

6 Makoto Okawara¹, M.D., Tomohiro Ishimaru¹, M.D., M.P.H., Ph.D., Seiichiro Tateishi², M.D., Ph.D.,
7 Ayako Hino³, M.D., Ph.D., Mayumi Tsuji⁴, M.D., Ph.D., Akira Ogami⁵, M.D., Ph.D., Tomohisa
8 Nagata⁶, M.D., Ph.D., Shinya Matsuda⁷, M.D., Ph.D., Yoshihisa Fujino¹, M.D., M.P.H., Ph.D., for
9 the CORoNaWork project

10

11 **Affiliations:**

12 1 Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,
13 University of Occupational and Environmental Health, Japan

14 2 Department of Occupational Medicine, School of Medicine, University of Occupational and
15 Environmental Health, Japan

16 3 Department of Mental Health, Institute of Industrial Ecological Sciences, University of
17 Occupational and Environmental Health, Japan

18 4 Department of Environmental Health, School of Medicine, University of Occupational and

19 Environmental Health, Japan

20 5 Department of Work Systems and Health, Institute of Industrial Ecological Sciences, University of

21 Occupational and Environmental Health, Japan

22 6 Department of Occupational Health Practice and Management, Institute of Industrial Ecological

23 Sciences, University of Occupational and Environmental Health, Japan

24 7 Department of Preventive Medicine and Community Health, School of Medicine, University of

25 Occupational and Environmental Health, Japan

26

27 Corresponding author:

28 Yoshihisa Fujino, M.D., M.P.H., Ph.D.

29 Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences, University

30 of Occupational and Environmental Health, Japan

31 1-1, Iseigaoka, Yahatanishiku, Kitakyushu, 807-8555, Japan

32 Tel: +81-93-691-7401

33 E-mail: zenq@med.uoeh-u.ac.jp

34

35 **Abstract**

36 Objective: This study examined the relationship between interruption to routine medical care during

37 the coronavirus disease 2019 pandemic and sickness presenteeism in Japan.

38 Methods: An internet monitor questionnaire was conducted. Data from 27,036 people were analyzed.

39 Interruption to medical care was defined based on the response “I have not been able to go to the

40 hospital or receive treatment as scheduled.” The number of sickness presenteeism days in the past 30

41 days was employed as the primary outcome. A zero-inflated negative binomial model was used for

42 analysis.

43 Results: The incidence rate ratio was significantly higher among workers who experienced

44 interrupted medical care (2.26; 95% confidence interval: 2.03–2.52) than those who did not require

45 routine medical care.

46 Conclusions: This study suggests the importance of continuing necessary treatment during a

47 pandemic to prevent presenteeism.

48

49 Keywords: occupational health; patient dropouts; presenteeism; COVID-19; regression analysis

50

51 **Introduction**

52 Sickness presenteeism is an increasingly important issue in occupational health. Aronsson
53 defined sickness presenteeism as “people, despite complaints and ill health that should prompt rest
54 and absence from work, still turning up at their jobs”¹. Sickness presenteeism is the result of a choice
55 made by a worker with ill-health, disease, or capacity loss between sickness presenteeism and
56 sickness absence². This decision is influenced by the individual's personality, values, economic
57 status, workplace “demands for presence” and support for adaptation, and national culture and
58 employment customs². Evidence suggests that sickness presenteeism can lead to sickness absence
59 and future worsening of physical and mental health conditions³⁻⁹. In addition, the impact of working
60 while ill on productivity is also gaining attention, especially in the US^{10,11}. A variety of diseases and
61 health conditions have been found to be associated with sickness presenteeism, suggesting the
62 importance of managing disease and maintaining good condition^{4,12}.

63 Under the coronavirus disease 2019 (COVID-19) pandemic, there is concern that both
64 organizational and individual factors will increase sickness presenteeism above that observed under
65 normal conditions¹³⁻¹⁵. Organizational factors that may lead to more sickness presenteeism include
66 increased workload and working hours per person, increased work pressure due to a manpower
67 shortage in the organization and changes in work demands on short notice during the pandemic. This

68 increased workload on workers can lead to a negative work culture around taking sick leave, such as
69 where workers who choose to work while ill are valued for their loyalty to the company and
70 motivation to work, thus promoting sickness presenteeism¹⁶. Examples of individual factors that may
71 increase sickness presenteeism include worsening economic situations and job insecurity; increased
72 telecommuting, which can make it easier for workers to work even while sick; the impact of the
73 pandemic on anxiety and mental health; worsening of health conditions and diseases due to lifestyle
74 changes; and worsening of chronic diseases due to the inability to access medical resources. All of
75 these factors are expected to lead to an increase in sickness presenteeism.

76 Interruption to medical care is an important problem in the COVID-19 pandemic. Access to
77 necessary routine medical care and medical resources is reportedly being affected in many countries
78 around the world¹⁷⁻¹⁹. In Japan, there is data showing that the number of prescriptions issued has
79 decreased²⁰. There are multiple reasons for such interruptions to medical care, including fear of
80 being infected with COVID-19 when leaving the house to visit a hospital, worsening personal
81 economic situations, and shortages in medical personnel, all of which affect the treatment schedule
82 for chronic diseases²¹. Interruption to medical care can adversely affect management of chronic
83 diseases and delay the detection and treatment of new diseases²². In fact, excess deaths unrelated to
84 COVID-19 have been reported²³. Thus, interruption to medical care during the COVID-19 pandemic
85 may lead to worsening of non-COVID-19 diseases and health conditions.

86 During the COVID-19 pandemic, the number of people working while ill may increase as a
87 result of worsening health conditions arising from treatment interruptions and delays. This may
88 result in an increase in sickness presenteeism. However, few studies have examined the effect of
89 medical care interruption on sickness presenteeism during the COVID-19 pandemic. We
90 hypothesized that sickness presenteeism has increased among workers who experienced
91 interruptions to their medical care during the COVID-19 pandemic. The purpose of this study was to
92 examine the association between medical care interruption and sickness presenteeism in Japanese
93 workers during the COVID-19 pandemic.

94

95 **Materials and Methods**

96 We performed a cross-sectional study based on baseline survey data obtained in the
97 Collaborative Online Research on the Novel- Coronavirus and Work (CORoNaWork) project, a
98 prospective cohort study that performed a questionnaire-based survey of Internet monitors to
99 determine the effect of the COVID-19 pandemic on workers' health. Before completing the online
100 survey, participants read a description of the survey's aims and details about the handling of their
101 information. Only participants who agreed with the contents of the description were allowed to

102 participate. Participation was anonymized. This study was approved by the Ethics Committee of the
103 University of Occupational and Environmental Health, Japan (Approval No. R2-079 and R3-006).

104 The baseline survey was conducted from December 22 to 26, 2020. A total of 33,302
105 participants aged from 20 and 65 years who indicated they were working when completing the
106 survey were included. Participants were selected such that sex and occupation (office and non-office
107 workers) were approximately equal among the regions of residence. Regions of residence were
108 categorized according to the cumulative COVID-19 infection rate. Participants who provided
109 fraudulent responses (n=6,266) according to the survey company or a predefined definition of a
110 fraudulent response were excluded. Fraudulent responses included an unusually short response time
111 (below 6 minutes), unusually short height (below 140 cm), unusually low weight (below 30 kg),
112 varying answers to similar questions in the survey (e.g., varying answers to questions about marital
113 status or area of residence), and incorrect answers to tiered questions used to identify inappropriate
114 responses (e.g., choose the third highest number from the following five numbers). After exclusion,
115 responses from 27,036 participants were included in the analysis.

116

117 Assessment of treatment status

118 We assessed the presence of disease and participants' need for medical care using the
119 following question: "Do you have a disease that requires regular visits to the hospital or treatment?"
120 Responses were "I do not have any such disease"; "I am able to go to the hospital or receive
121 treatment as scheduled"; "I have not been able to go to the hospital or receive treatment as
122 scheduled."

123 Those who answered "I do not have any such disease" were defined as workers who did not
124 require routine medical care, and thus did not have any disease that requires hospital visits or
125 treatment. Those who answered "I am able to go to the hospital or receive treatment as scheduled"
126 were defined as workers who used medical care. Those who answered "I have not been able to go to
127 the hospital or receive treatment as scheduled" were defined as workers who experienced interrupted
128 medical care.

129

130 Assessment of sickness presenteeism and other covariates

131 Respondents' number of sickness presenteeism days was ascertained based on the following
132 question and used as the primary outcome: "In the last 30 days, how many days have you worked
133 (including work from home) despite feeling that you really should have taken sick leave due to your
134 state of health?"

135 Socioeconomic and work-related factors included sex, age, job type (mainly desk work,
136 mainly interpersonal communication, mainly physical work), marital status (married,
137 divorced/deceased, never married), annual household income, education (junior high school, high
138 school, vocational school/junior college/college of technology, university/graduate school), company
139 size (total number of employees in the respondent's main place of work [1 for self-employed]), and
140 the number of days worked per week.

141 The cumulative infection rate of COVID-19 in the province of residence was employed as a
142 community-level variable.

143 To control for potential confounders, we also asked participants to indicate their main
144 symptoms using the following question: "Which of the following conditions or body parts give you
145 the most trouble during your work?" The options were "No problem"; "pain"; "movement";
146 "tightness, loss of energy, appetite, fever, dizziness, or feeling poor"; "toileting or elimination";
147 "mental health"; "skin, hair, or beauty"; "sleep"; "eyes"; "nose"; "ears"; and "other."

148

149 Statistical analysis

150 Age and the number of sickness presenteeism days were expressed as continuous variables,
151 with median and interquartile range (IQR). Other covariates were expressed as categorical variables
152 using percentages.

153 We compared the results of linear regression, Poisson regression, Zero-inflated Poisson
154 regression (ZIP), negative binomial regression, and Zero-inflated Negative Binomial regression
155 (ZINB) as statistical models, as they treat the number of sickness presenteeism days as continuous
156 count data. Negative binomial regression can handle over-dispersed data, where the variance is much
157 higher than the mean, which cannot be assumed in Poisson distribution. Further, to handle data with
158 excess zeros, which indicates a population at low risk of sickness presenteeism, we used a
159 zero-inflated model. In addition to dealing with the excess zeros that often occur in count data, a
160 zero-inflated model has also been proposed as a way to handle the difficulty of defining sickness
161 presenteeism cutoffs^{24,25}. As a measure of model fitness, we compared the Akaike's Information
162 Criterion (AIC), and ultimately adopted the ZINB model.

163 ZINB regression analysis was conducted with the number of continuous sickness
164 presenteeism days as the dependent variable, the respondents' category of treatment status as an
165 independent variable, and the number of days worked per week as an offset variable.

166 We adjusted for the following potential confounders: sex, age, job type, marital status,
167 household income, education, company size, cumulative infection rate by prefecture, and main
168 symptoms.

169 In further analysis, we estimated the margins of sickness presenteeism days for each
170 treatment status and symptom. First, we used the same statistical model as that in the main analysis.
171 Second, we calculated the predictive margins of sickness presenteeism days, substituting measured
172 values for other covariates, dividing the data into 36 groups (3 treatment statuses and 12
173 symptoms)²⁶. Preliminarily, we confirmed the simple main effects for each treatment status
174 compared to workers who did not require routine medical care by adding the interaction term
175 between treatment status and symptoms to the model used for the main analysis. For all analyses, the
176 Bonferroni method was used to adjust for multiple comparisons.

177 All comparisons were performed in Stata (Stata Statistical Software: Release 16; StataCorp
178 LLC, TX, USA), with $p < 0.05$ indicating statistical significance.

179

180 **Results**

181 Of the 33,302 responses, 6,266 were excluded (215 were deemed fraudulent according to
182 the survey company and 6,051 satisfied the exclusion criteria during data cleaning), leaving a total of
183 27,036 responses for analysis. Because all responses were mandatory, there were no missing data in
184 this study.

185 The demographic and sociological characteristics of the analyzed population are shown in
186 Table 1. A total of 13,814 (51%) were men, with a median age of 48 years (IQR: 39–55). Of the total
187 population, 17,526 (65%) were workers who did not require routine medical care, 8,451 (31%) were
188 using medical care as scheduled, and 1,059 (4%) experienced interrupted medical care. The overall
189 median number of sickness presenteeism days was 0.0 (IQR: 0–2). The distribution of sickness
190 presenteeism is shown for the three treatment statuses in a histogram in Figure 1.

191 The association between the number of sickness presenteeism days and treatment status is
192 shown in Table 2. There was a significant association between the number of sickness presenteeism
193 days and treatment status based on the count model part of ZINB. After adjusting for other
194 covariates in the multivariate model, the incidence rate ratio (IRR) of sickness presenteeism days
195 was significantly higher among workers who used medical care (IRR: 1.15, 95% CI: 1.08–1.22,
196 $p<0.001$) and workers who experienced interrupted medical care (IRR: 2.26, 95% CI: 2.03–2.52,
197 $p<0.001$) compared to workers who did not require routine medical care.

198 The association between the number of sickness presenteeism days and participants' main
199 symptoms is shown in Table 3. There were significant associations between the number of sickness
200 presenteeism days and some symptoms using the model presented in Table 2. The highest IRR of
201 sickness presenteeism days was observed for mental health symptoms (adjusted IRR: 1.67, 95% CI:
202 1.52–1.83, $p<0.001$).

203 The predictive margins of sickness presenteeism days for each treatment status and
204 symptom are shown in Table 4 and Figure 2. When the analysis was performed based on the three
205 treatment statuses, irrespective of symptom, the predictive margin of sickness presenteeism days
206 among workers who experienced interrupted medical care was 7.06 days (standard error [SE]=0.34),
207 while that among workers who did not require routine medical care was 1.43 days (SE=0.03). When
208 the analysis was performed based on the 36 treatment-symptom groups (3 treatment statuses and 12
209 symptoms), the largest predictive margin of sickness presenteeism days was observed for mental
210 health symptoms AND interrupted medical care (PM: 10.05 days, SE=0.57). The simple effect
211 comparisons test, which included the interaction term between treatment status and symptoms,
212 showed that there were significant differences between workers with the same symptoms who did
213 and did not require routine medical care, and between workers with the same symptoms who
214 experienced interruption to medical care and who did not require routine medical care. For example,
215 the number of sickness presenteeism days significantly differed between those with mental health

216 symptoms who used medical care and those with mental health symptoms who did not require

217 routine medical care ($p < 0.001$).

218

219 **Discussion**

220 This study demonstrated an association between treatment interruption and sickness

221 presenteeism among Japanese workers in the COVID-19 pandemic. Compared to workers who did

222 not require routine medical care, workers who had diseases that required routine medical care

223 reported more days of sickness presenteeism, and those who experienced interrupted medical care

224 reported even more such days. Furthermore, our findings reveal differences in the occurrence of

225 sickness presenteeism depending on workers' symptoms.

226 We found that workers who experienced interrupted medical care had increased sickness

227 presenteeism. This is because appropriate treatment can improve work function and productivity by

228 improving workers' health and subjective symptoms^{10,27}. This is supported by the fact that workers

229 who used medical care reported fewer sickness presenteeism days than those who experienced

230 interrupted medical care, although workers who used medical care had a higher incidence of

231 sickness presenteeism than workers who did not require routine medical care. Employees who

232 experience interrupted treatment for chronic diseases may be forced to return to work due to fear of

233 being laid off, depending on the financial situation of their workplace during the pandemic. It is thus
234 important to continue regular treatment during the COVID-19 pandemic to manage disease and
235 maintain good condition²⁸.

236 We found that the occurrence of sickness presenteeism depends on the type of symptoms
237 experienced by workers. Workers with symptoms related to mental health problems, loss of
238 energy/fever, and body movements reported more sickness presenteeism than those who reported
239 having “no problem.” In contrast, workers with symptoms related to sleep, pain, and elimination
240 reported comparable or fewer days of sickness presenteeism to those who reported having “no
241 problem.” Sickness presenteeism is the result of a worker's choice to be absent from work or to
242 attend work despite being unwell. Many previous studies have evaluated sickness presenteeism
243 based on whether or not workers “worked one or more days in a certain period of time with a health
244 condition for which they think they really should be absent”^{2,29}. However, workers experiencing
245 symptoms not typically associated with sickness presenteeism may not consider their symptoms
246 suitable for an absence from work, and thus may not have indicated that they experienced sickness
247 presenteeism. For example, symptoms related to beauty are not directly related to an individual's
248 ability to work; sleep may be considered something that the individual simply needs to get more of
249 on holidays; and chronic pain may be considered an instruction to move rather than rest. However,
250 some symptoms that can pose a health risk are also unlikely to be recognized as contributing to

251 sickness presenteeism. These include symptoms that can lead to delayed detection or worsening of a
252 disease if left untreated when rest or treatment is in fact required. For individuals with insomnia or
253 elimination symptoms, for example, resting or visiting a hospital when feeling unwell can lead to
254 prevention or early diagnosis and treatment of mental health problems or inflammatory bowel
255 disease, respectively. While forcing oneself to work with such symptoms can pose a health risk,
256 workers may not consider this sickness presenteeism due to differences in interpretation of “health
257 conditions that require absence from work.” This is an important point when evaluating sickness
258 presenteeism.

259 We also found that the impact of continuing treatment on the prevention of sickness
260 presenteeism varied by symptom. Sickness presenteeism was more frequent in workers who
261 experienced interrupted medical care with symptoms related to tightness and loss of energy, toileting
262 and elimination, sleep, and eyes than those who did not require routine medical care. In contrast, no
263 difference in sickness presenteeism was observed between workers who used medical care and those
264 who did not require routine medical care, suggesting the importance of continuing necessary routine
265 medical care for preventing sickness presenteeism due to these symptoms. Workers who experienced
266 interrupted medical care with these symptoms may be able to reduce the incidence of sickness
267 presenteeism by continuing appropriate treatment to maintain and improve their health condition.
268 Symptoms that led to more sickness presenteeism in both workers who used medical care and

269 workers who experienced interrupted medical care compared to workers who did not require routine
270 medical care were pain and mental health-related symptoms. For these symptoms, sickness
271 presenteeism remained high even with continued treatment, indicating the need to identify
272 appropriate treatment and manage one's daily health condition in addition to continuing treatment. In
273 contrast, symptoms that led to comparable sickness presenteeism in workers who used medical care
274 and workers who experienced interrupted medical care compared to workers who did not require
275 routine medical care were related to movement and mobility; skin, hair, or beauty; nose; ears; and
276 other symptoms, for which treatment is ineffective to prevent sickness presenteeism. The lack of a
277 difference in sickness presenteeism for these symptoms may be due to the fact that individuals do
278 not consider these symptoms sufficiently adverse to require an absence from work. Alternatively,
279 some individuals, such as those with physical movement symptoms, may experience chronic
280 symptoms for which support and adaptive behaviors have already been put into place; thus, whether
281 or not these individuals experience sickness presenteeism may be unrelated to their treatment status.
282 Thus, the impact of continuing treatment on sickness presenteeism may be related to whether an
283 individual considers their symptoms to be sufficiently adverse to require an absence from work, or
284 whether or not the symptoms can be improved with treatment.

285 There are several limitations to this study. First, we did not obtain detailed information
286 related to treatment interruptions, including the type of disease, duration, and reasons for interruption.

287 We were thus unable to determine whether the reason for interruption to treatment was due to
288 patient-related reasons (e.g., economic situation and anxiety) or hospital-related reasons (e.g.,
289 schedule adjustment). Second, interruptions to treatment may be the result of better disease control
290 and improved health. It is unclear how these factors would affect the occurrence of sickness
291 presenteeism. Finally, we did not consider all possible confounders affecting sickness presenteeism
292 because we did not obtain information on some confounders, such as job insecurity, annual leave
293 rights, and the culture around employment and sick leave in each company.

294

295 **Conclusion**

296 Interruption to medical care during the COVID-19 pandemic was associated with the
297 occurrence of sickness presenteeism. This study demonstrates the importance of maintaining one's
298 health condition and continuing necessary treatment even during an infectious disease pandemic.

299

300 References

- 301 1. Aronsson G, Gustafsson K, Dallner M. Sick but yet at work. An empirical study of sickness
302 presenteeism. *J Epidemiol Community Health*. 2000;54: 502–509.
- 303 2. Aronsson G, Gustafsson K. Sickness presenteeism: prevalence, attendance-pressure factors, and
304 an outline of a model for research. *J Occup Environ Med*. 2005;47: 958–966.
- 305 3. Skagen K, Collins AM. The consequences of sickness presenteeism on health and wellbeing
306 over time: A systematic review. *Soc Sci Med*. 2016;161: 169–177.
- 307 4. Johns G. Presenteeism in the workplace: A review and research agenda. *J Organ Behav*.
308 2010;31: 519–542.
- 309 5. Kivimäki M, Head J, Ferrie JE, Hemingway H, Shipley MJ, Vahtera J, et al. Working while ill
310 as a risk factor for serious coronary events: the Whitehall II study. *Am J Public Health*.
311 2005;95: 98–102.
- 312 6. Janssens H, Clays E, De Clercq B, De Bacquer D, Braeckman L. The relation between
313 presenteeism and different types of future sickness absence. *J Occup Health*. 2013;55: 132–141.
- 314 7. Bergström G, Bodin L, Hagberg J, Aronsson G, Josephson M. Sickness presenteeism today,
315 sickness absenteeism tomorrow? A prospective study on sickness presenteeism and future
316 sickness absenteeism. *J Occup Environ Med*. 2009;51: 629–638.
- 317 8. Gustafsson K, Marklund S. Associations between health and combinations of sickness presence
318 and absence. *Occup Med* . 2014;64: 49–55.
- 319 9. Gustafsson K, Marklund S. Consequences of sickness presence and sickness absence on health
320 and work ability: a Swedish prospective cohort study. *Int J Occup Med Environ Health*.
321 2011;24: 153–165.
- 322 10. Schultz AB, Chen C-Y, Edington DW. The cost and impact of health conditions on
323 presenteeism to employers: a review of the literature. *Pharmacoeconomics*. 2009;27: 365–378.
- 324 11. Nagata T, Mori K, Ohtani M, Nagata M, Kajiki S, Fujino Y, et al. Total Health-Related Costs
325 Due to Absenteeism, Presenteeism, and Medical and Pharmaceutical Expenses in Japanese
326 Employers. *J Occup Environ Med*. 2018;60: e273–e280.
- 327 12. Burton WN, Pransky G, Conti DJ, Chen C-Y, Edington DW. The association of medical
328 conditions and presenteeism. *J Occup Environ Med*. 2004;46: S38-45.

- 329 13. Reuter M, Dragano N, Wahrendorf M. Working while sick in context of regional
330 unemployment: a Europe-wide cross-sectional study. *J Epidemiol Community Health*. 2021;75:
331 574–580.
- 332 14. Tilchin C, Dayton L, Latkin CA. Socioeconomic Factors Associated With an Intention to Work
333 While Sick From COVID-19. *J Occup Environ Med*. 2021;63: 363–368.
- 334 15. Kinman G, Grant C. Presenteeism during the COVID-19 pandemic: risks and solutions. *Occup*
335 *Med* . 2020. doi:10.1093/occmed/kqaa193
- 336 16. Kinman G. Sickness presenteeism at work: prevalence, costs and management. *Br Med Bull*.
337 2019;129: 69–78.
- 338 17. Das AK, Mishra DK, Gopalan SS. Reduced access to care among older American adults during
339 CoVID-19 pandemic: results from a prospective cohort study. *bioRxiv*. medRxiv; 2020.
340 doi:10.1101/2020.11.29.20240317
- 341 18. Czeisler MÉ, Mpp KM, Clarke KEN, Salah Z, Shakya I, Thierry JAM, et al. Delay or
342 avoidance of medical care because of COVID-19–related concerns — .
343 doi:10.1101/2020.04.22.20076141v1
- 344 19. van Ballegooijen H, Goossens L, Bruin RH, Michels R, Krol M. Concerns, quality of life,
345 access to care and productivity of the general population during the first 8 weeks of the
346 coronavirus lockdown in Belgium and the Netherlands. *BMC Health Serv Res*. 2021;21: 227.
- 347 20. Lopresti M Seo T Sato. Pandemics and access to care: use of real-world DATA to examine the
348 IMPACT of COVID-19 on pharmacy visits in Japan. *Value Health*. [cited 28 May 2021].
349 doi:10.1016/j.jval.2020.08.1707
- 350 21. Bong C-L, Brasher C, Chikumba E, McDougall R, Mellin-Olsen J, Enright A. The COVID-19
351 Pandemic: Effects on Low- and Middle-Income Countries. *Anesth Analg*. 2020;131: 86–92.
- 352 22. Erol MK, Kayıkçıoğlu M, Kılıçkap M, Güler A, Yıldırım A, Kahraman F, et al. Treatment
353 delays and in-hospital outcomes in acute myocardial infarction during the COVID-19
354 pandemic: A nationwide study. *Anatol J Cardiol*. 2020;24: 334–342.
- 355 23. Woolf SH, Chapman DA, Sabo RT, Weinberger DM, Hill L. Excess Deaths From COVID-19

- 356 and Other Causes, March-April 2020. *JAMA*. 2020;324: 510–513.
- 357 24. Navarro A, Salas-Nicás S, Llorens C, Moncada S, Molinero-Ruíz E, Moriña D. Sickness
358 presenteeism: Are we sure about what we are studying? A research based on a literature review
359 and an empirical illustration. *Am J Ind Med*. 2019;62: 580–589.
- 360 25. Tashiro Y, Nakamura K, Seino K, Ochi S, Ishii H, Hasegawa M, et al. The impact of a
361 school-based tooth-brushing program on dental caries: a cross-sectional study. *Environ Health
362 Prev Med*. 2019;24: 83.
- 363 26. Williams R. Using the Margins Command to Estimate and Interpret Adjusted Predictions and
364 Marginal Effects. *Stata J*. 2012;12: 308–331.
- 365 27. Burton WN, Morrison A, Wertheimer AI. Pharmaceuticals and worker productivity loss: a
366 critical review of the literature. *J Occup Environ Med*. 2003;45: 610–621.
- 367 28. Chudasama YV, Gillies CL, Zaccardi F, Coles B, Davies MJ, Seidu S, et al. Impact of
368 COVID-19 on routine care for chronic diseases: A global survey of views from healthcare
369 professionals. *Diabetes Metab Syndr*. 2020;14: 965–967.
- 370 29. Hansen CD, Andersen JH. Going ill to work--what personal circumstances, attitudes and
371 work-related factors are associated with sickness presenteeism? *Soc Sci Med*. 2008;67:
372 956–964.


Table 1. Basic characteristics of the study subjects

	Total (n=27,036)	Workers who did not require routine medical care (n= 17,526)	Workers who used medical care (n=8,451)	Workers who experienced interrupted medical care (n=1,059)
	n (%)	n (%)	n (%)	n (%)
Age, median (IQR)	48 (39-55)	46 (38-53)	52 (45-58)	47 (39-54)
Sex, men	13814 (51.1%)	8422 (48.1%)	4885 (57.8%)	507 (47.9%)
Marital status, married	15029 (55.6%)	9627 (54.9%)	4894 (57.9%)	508 (48.0%)
Job type				
Mainly desk work	13468 (49.8%)	8545 (48.8%)	4440 (52.5%)	483 (45.6%)
Mainly interpersonal communication	6927 (25.6%)	4621 (26.4%)	2032 (24.0%)	274 (25.9%)
Mainly physical work	6641 (24.6%)	4360 (24.9%)	1979 (23.4%)	302 (28.5%)
Equivalent income (Japanese Yen)				
550000-2500000	5710 (21.1%)	3636 (20.7%)	1800 (21.3%)	274 (25.9%)
2500000-4250000	7550 (27.9%)	5036 (28.7%)	2227 (26.4%)	287 (27.1%)
4250000-5500000	6625 (24.5%)	4353 (24.8%)	2039 (24.1%)	233 (22.0%)
>5500000	7151 (26.4%)	4501 (25.7%)	2385 (28.2%)	265 (25.0%)
Education				
Junior high school	368 (1.4%)	244 (1.4%)	107 (1.3%)	17 (1.6%)
High school	6953 (25.7%)	4381 (25.0%)	2284 (27.0%)	288 (27.2%)
Vocational school, junior college, or technical college	6544 (24.2%)	4378 (25.0%)	1913 (22.6%)	253 (23.9%)

University or graduate school	13171 (48.7%)	8523 (48.6%)	4147 (49.1%)	501 (47.3%)
Number of employees				
1	2556 (9.5%)	1625 (9.3%)	848 (10.0%)	83 (7.8%)
2-49	7999 (29.6%)	5378 (30.7%)	2320 (27.5%)	301 (28.4%)
50-999	9703 (35.9%)	6262 (35.7%)	3036 (35.9%)	405 (38.2%)
1000-9999	4719 (17.5%)	2986 (17.0%)	1552 (18.4%)	181 (17.1%)
≥10000	2059 (7.6%)	1275 (7.3%)	695 (8.2%)	89 (8.4%)
Sickness presenteeism days, median (IQR)	0.0 (0-2)	0.0 (0-1)	0.0 (0-2)	4.0 (1-10)

IQR: interquartile range

1 Figure 1. Sickness presenteeism days under each treatment status



2

Table 2. Association between treatment status and sickness presenteeism

Treatment status	Univariate			Multivariate*		
	IRR	95% CI	<i>p</i>	IRR	95% CI	<i>p</i>
Workers who did not require routine medical care	Reference			Reference		
Workers who used medical care	1.13	1.06 1.21	<0.001	1.15	1.08 1.22	<0.001
Workers who experienced interrupted medical care	2.30	2.06 2.57	<0.001	2.26	2.03 2.52	<0.001

*Adjusted for sex, age, marital status, household income, education, company size, job type, cumulative infection rate for COVID-19, and main symptoms

Offset variable: number of days worked per week

IRR: incidence rate ratio, CI: confidence interval

Table 3. Association between main symptoms and sickness presenteeism

Main symptom	IRR	95% CI		<i>p</i>
No problem	Reference			
Pain	1.03	0.92	1.15	0.587
Movement	1.28	1.11	1.48	0.001
Tightness, loss of energy, appetite, fever, dizziness, or feeling poor	1.35	1.21	1.50	<0.001
Toileting or elimination	0.97	0.81	1.17	0.780
Mental health	1.67	1.52	1.83	<0.001
Skin, hair, or beauty	0.90	0.75	1.09	0.284
Sleep	1.05	0.95	1.17	0.352
Eyes	0.86	0.75	0.98	0.024
Nose	0.68	0.49	0.95	0.023
Ears	0.85	0.63	1.15	0.299
Other	1.35	1.18	1.54	<0.001

Adjusted for sex, age, marital status, household income, education, company size, job type, cumulative infection rate for COVID-19, and treatment status

Offset variable: number of days worked per week

IRR: incidence rate ratio, CI: confidence interval

Table 4. Predictive margins of sickness presenteeism days for each symptom and comparison between each treatment status

	Workers who did not require routine medical care (n=17,526)				Workers who used medical care (n=8,451)					Workers who experienced interrupted medical care (n=1,059)					
	n	%	PM	SE	n	%	PM	SE	<i>p</i>	n	%	PM	SE	<i>p</i>	
Total	17526	100	1.43	0.03	8451	100	2.16	0.05	<0.001	1059	100	7.06	0.34	<0.001	
Main symptom															
No problem	10938	62.4	0.76	0.02	3642	43.1	0.97	0.04	0.001	160	15.1	4.85	0.35	0.006	
Pain	849	4.8	2.11	0.10	842	10.0	2.53	0.11	0.017	144	13.6	6.19	0.38	<0.001	
Movement	481	2.7	2.61	0.17	344	4.1	3.09	0.20	1.000	54	5.1	7.11	0.57	1.000	
Tightness, loss of energy, appetite, fever, dizziness, or feeling poor	715	4.1	3.67	0.16	583	6.9	3.96	0.17	0.056	133	12.6	8.32	0.53	<0.001	
Toileting or elimination	263	1.5	1.91	0.16	201	2.4	2.35	0.20	1.000	40	3.8	5.87	0.56	<0.001	
Mental health	1143	6.5	4.09	0.15	909	10.8	4.69	0.17	<0.001	219	20.7	10.05	0.57	<0.001	
Skin, hair, or beauty	389	2.2	1.62	0.14	160	1.9	1.99	0.17	0.069	29	2.7	5.45	0.54	1.000	
Sleep	997	5.7	2.34	0.10	637	7.5	2.74	0.12	1.000	116	11.0	6.20	0.39	<0.001	
Eyes	795	4.5	1.46	0.08	512	6.1	1.76	0.10	1.000	65	6.1	4.57	0.33	<0.001	
Nose	98	0.6	1.29	0.19	71	0.8	1.54	0.23	1.000	9	0.9	4.03	0.67	1.000	
Ears	80	0.5	1.76	0.24	84	1.0	2.14	0.29	1.000	12	1.1	5.25	0.80	0.343	
Other	778	4.4	1.96	0.12	466	5.5	2.51	0.16	1.000	78	7.4	7.86	0.60	0.351	

PM: predictive margins; mean predicted number of sickness presenteeism days for each symptom and treatment status using the model in Table 2 and 3 with substitution of measured values for other covariates (adjusted using the Bonferroni method)

Offset variable: number of days worked per week


IRR: incidence rate ratio, SE: standard error, %: proportion of the total number of respondents for each treatment status

p : p-value for simple main effects for each treatment status compared to workers who did not require routine medical care using a model that included the interaction term for treatment status and main symptoms (adjusted using the Bonferroni method)

1

2

1 Figure 2. Predictive margins of each treatment status and symptom



2

3 *others include loss of energy, appetite, fever, dizziness, or feeling poor