

Evaluation of the relationship between quantitative PCR results and cell culturing of SARS2-CoV with respect to symptoms onset and Viral load – a systematic review

Rozenberg Gilad (1), Erster Oran (2), Ghersin Itai (1), , Mandelboim Michal (2), Neuberger Ami (1,5), Schwartz Eli (3,4).

1. Internal Medicine B, Rambam Medical Center, Haifa, Israel
2. Central Virology Laboratory, Ministry of Health, Sheba Medical Center, Tel Hashomer, Ramat-Gan, Israel.
3. The Center for Geographic Medicine and Tropical Diseases, Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel.
4. Sackler Faculty of Medicine, Tel-Aviv University, Tel-Aviv, Israel.
5. Bruce Rappaport Faculty of Medicine, Technion, Haifa, Israel

Word count: 3582

Corresponding Author:

Rozenberg Gilad

Rambam medical center, Haifa, Israel

rozenberg.gilad@gmail.com

Tel: +972547626424

Abstract

Background: Viral culture is currently the most accurate method to demonstrate viability and infectivity of Severe acute respiratory syndrome Coronavirus (SARS-2 CoV). Routine clinical diagnosis, however, is mostly performed by PCR – based assays that do not discriminate between infectious and non-virus. Herein, we aimed to determine the correlation between positive viral cultures and either PCR positivity, the Cycle Threshold (Ct) or the number of viral copies.

Methods: A systematic electronic literature search was performed and studies that reported both viral SARS-CoV-2 culture and PCR–based assays were included. A separate search for samples from blood, urine, stool, breast milk and tears were performed. To convert Ct values reported in the reviewed studies were to viral genomic copies, calibration experiments with four different reaction performed, using quantified RNA molecules.

Results: A total 540 articles were reviewed, and 38 studies were included in this review. Out of 276 positive-culture of non-severe patients, 272 (98.55%) were negative ten days after symptoms onset, while PCR assays remained positive for up to 67 days. In severely ill or immunocompromised patients positive-culture was obtained up to 32 days and out of 168 cultures, 31 (18.45%) stayed positive after day 10. In non-severe patients, in Ct value greater than 30 only 10.8% were still culture-positive while in Ct >35 it was nearly universally negative. The minimal calculated number of viral genome copies in culture-

positive sample was 2.5×10^3 copies / mL. These findings were similar in immunocompromised patients. Recovering positive culture from non-respiratory samples was sporadically obtained in stool or urine samples. Conversion of Ct values to viral genome copies showed variability between different PCR assays and highlighted the need to standardize reports to correctly compare results obtained in different laboratories.

Conclusion: During the pandemic phase, non-severe COVID-19 patients who are recovering and are not immuno-suppressed, can be regarded as non-infectious, within 10 days from symptom onset, or with Ct value greater than 35 (or a calculated viral load lower than 1.2×10^3 copies / mL). These findings have important implications for recovering patients and asymptomatic patients, with respect to isolation criteria. The conversion of Cq values to viral genome copies described herein may be useful in future work, enabling a more standardized comparison between results reported in different studies from different laboratories.

Key words:

SARS-COV-2, quantitative RT-PCR, Cell culture, immunosuppressed patients, viral infectivity

Introduction:

By the end of February 2021, coronavirus disease 2019 (COVID-19) has infected around 108 million people worldwide and caused more than 2.3 million deaths¹.

In addition to its tragic medical implications, the COVID-19 outbreak also has considerable social and economic consequences^{2,3,4}. The Organization for Economic Co-operation and Development (OECD) has forecast that the global economy will shrink by 6% in 2020⁵, the deepest recession since the Second World War⁶. Isolation and quarantine of infected people is one of the contributors to these consequences.

Viral culture is currently the most accurate method to demonstrate COVID-19 viability and infectivity⁷. However, it is not routinely performed due to its time-consuming, its labor-intensive nature, and the required safety measures for such viral pathogen propagation. Therefore, diagnosis of COVID-19 has largely relied on real-time reverse transcriptase polymerase chain reaction (rRT-PCR)⁸. This method is both rapid, highly sensitive and specific⁷. It can also provide quantitative data, under suitable conditions. However, RT-qPCR has a major limitation, as it might detect viral RNA of non-viable, and thus non-infective, viruses⁹. This phenomenon has been demonstrated in several viruses, such as Ebola virus⁹, respiratory syncytial virus¹⁰ and influenza¹¹, as well as for SARS-CoV-2^{12,13}. Recent studies using viral cultures deemed COVID-19 patients as non-infective after 5 to 7 days from onset of symptoms^{12,14}. In contrast, the median duration of RNA shedding, as assessed by rRT-PCR, was 31 days in a cohort of Chinese patients¹⁵.

The RT-qPCR assay is based on a specific fluorescent signal that is generated when the target nucleic acid (viral RNA in the case of SARS-2) is present in the examined sample. The target sequence is amplified by PCR. The quantity of the target RNA is inverse correlated with the number of amplification cycles required to generate the positive signal. Thus, lower Ct levels represent a larger number of target nucleic acid copies in the examined sample¹⁶.

Despite the above-mentioned significant limitation, RT-qPCR results are usually reported in medical systems as a binary result (positive or negative). A positive result is frequently used to determine how long COVID19 patients need to remain isolated. Determining the isolation period of patients who have recovered from COVID-19 has immense medical, economical, psychological and social implications.

In this systematic review, we aimed to determine the correlation between COVID-19 infectivity, as expressed by positive viral cultures, in relation to the Ct thresholds and to number of viral copies which is a more accurate measurement, and in relation to time from symptoms onset.

Methods

Type of studies

We screened all clinical studies that included viral SARS-CoV-2 cultures taken from symptomatic and asymptomatic COVID19 patients of any age and described their

positivity as a function of symptoms duration, cycle threshold (Ct) or number of viral copies determined by PCR. We have included randomized controlled trials (RCTs), non-randomized comparative cohorts and case series studies until August 15, 2020. We included viral cultures taken from the upper respiratory tract for the main analysis of outcomes, but also included studies that described cultures of materials from other sites (stool, urine, tears, breast milk) for further considerations. We also extracted the gene probes used in measuring the cycle threshold for *qPCR* positivity and methods of viral growth in culture.

Outcome measures

1. Duration of viral culture positivity in comparison to *qPCR* positivity as measured from the day of symptoms onset.
2. Duration of viral culture positivity in comparison to either Ct of a *qPCR* assay or to viral genome copies.
3. Ability to isolate SARS-CoV-2 from non-respiratory tract samples.

Search methods for identification of studies

We conducted a systematic electronic literature search with the PubMed and MedRxiv search engines. We used the following search terms: "COVID-19"\"coronavirus\" SARS-CoV-2\" combined with "culture". We also used references of retrieved papers, including reviews or systematic reviews, to identify further studies. Two reviewers independently screened all studies published or appeared online as non-peer-reviewed pre-prints

before August 1, 2020. We excluded all initially identified and retrieved articles that did not fulfill the inclusion criteria. In case of disagreement, a third reviewer acted as arbitrator.

Laboratory criteria for included studies.

Nucleic acid testing: To be able to analyze the data presented in the included studies, they were selected so that either the PCR assays described by Corman et al¹⁷, or the SARS2 test developed by the US CDC¹⁸ were used. Additionally, a description of the sampling and RNA extraction procedures needed to be included, so that the conversion from Ct values to target copies could be performed.

Virus isolation

SARS-CoV-2 isolation was determined by the presence of a typical cytopathic effect and increased viral RNA in the supernatant.

Conversion of Ct values to copy number and vice versa

Many of the studied reported Ct values without converting them to viral target copy number. In order to estimate the viral load in the samples described in these studies, we converted the Ct values reported, to viral genomic copies, based on the procedure performed in each study, as described herein.

RNA segments containing the amplified targets of E-Sarbeco, RdRp, N-Sarbeco and CDC N1 reactions were generated by *In vitro* transcription in the Israel Central Virology

Laboratory (CVL). The RNA molecules were purified, quantified, and served as standards for the standard curve calibration. The number of target copies in 1 ng of purified RNA was calculated based on the amplified sequence length, thereby enabling the conversion from copy number to Ct values. Serial dilutions of each amplicon were examined by RT-qPCR to generate the standard curve for each reaction. The Ct values were then plotted against the calculated concentration using a semi-logarithmic scale graph, and the regression formula was deduced. The copy number per ml was then calculated for each report, based on its specific sampling and extraction method. Detailed description of the procedure is included in the Supplementary material of this review.

Results

We reviewed 540 articles (figure 1). Finally, 14 studies were included in the comparisons between the duration of PCR positivity and viral culture positivity (Table 1A+1B). Twelve studies were included in the comparison between viral culture positivity and the cycle threshold and / or viral load of the molecular analyses (Table 2A+2B). Ten studies described viral cultures performed in specimens other than respiratory tract secretions (Table 3).

Examination of the duration of PCR and viral culture positivity from symptoms onset showed that out of 1668 positive PCR samples reported in the included studies, 444 (26.6%) cultures were positive, and 1224 (73.4%) were negative. Viral cultures were positive from between 6 days before the onset of symptoms, until 32 days after the onset of symptoms¹⁹. However, upon separating the immunocompetent-non-severe patients 272 out of 276 cultures (98.5%) were negative within 10 days of symptoms onset (Table 1A), while PCR assays remained positive for up to 67 days after the onset of symptoms²⁰. In the seven studies that reported the duration of culture positivity in detail, the medians ranged between 1.5 and 8 days.¹²¹⁴²¹²²²³²⁴²⁵

From the group of studies that included immunocompetent patients with non-severe disease, cases positive after day 10 from symptoms onset included a case from Taiwan of a 50-year-old woman without any comorbidities and a clinically mild covid-19 had a positive viral culture by day 18²⁶, another 2 samples from a series from England²⁵, and one sample from second study from Taiwan¹⁴. Thus altogether, only 4 out of 276 (1.45%) samples stayed positive after day 10 of symptoms.

In case-series of severely ill-patients or immunocompromised patients, culture-positive samples were obtained up to 32 days of symptoms onset. For example, a recipient of a heart transplant, despite having mild disease, was reported to have positive viral cultures for 21 days²⁷. In a work from the Netherlands that included immunocompromised patients, and more severely ill patients they had positive cultures up to 20 days from symptoms onset²⁴. Finally, in a study from Spain, a severely ill

patient had positive PCR assays for up to 32 days¹⁹. Altogether, for this group, 31 out of 168 positive samples (18.45%) stayed positive after day 10 of symptoms (Table 4).

The summary of the 12 studies that examined the correlation between culture positivity and either cycle threshold (Ct) or viral load (VL) as expressed in genomic copies is presented in table 2.

The majority of samples had Ct level of <30, and only 59 samples out of 547 (10.8%) were culture-positive with Ct above 30. While positive samples with viable culture in Ct value above 35 were found only in 9 samples out of 547 (1.6%)¹⁴¹⁹²⁵. In the five studies that reported the median Ct or VL in samples with positive cultures, Ct values range from 23, to the maximal median Ct was 31²⁵⁷²⁸. Median viral loads were 1.7×10^8 and 8.14×10^6 .²²²⁴ (Table 2)

In this regard, there were no major differences between immunocompetent and immunosuppressed or severely ill patients (Table 2 B). One exceptional case described a positive viral culture from a sample with a Ct of 41. No clinical data were provided for that specific patient regarding either immune status or illness severity.²⁵

In addition to the information obtained from the literature, we added calculation of the viral genome copy number based on the Ct values. This was determined by generating standard curves for the E-sarbeco, N-sarbeco, RdRp and CDC-N1 reactions using diluted *In-vitro* transcribed RNA targets, as detailed in the Supplementary material file. The minimal calculated number of viral genome copies that were found in samples that

were culture positive were 2.5×10^3 and 1.2×10^3 copies /mL in immunocompetent and immunosuppressed patients, respectively¹⁹. However, when studies where the median viral load is reported, there was a difference between the two population: 1.7×10^8 copies /mL in immunocompetent patients and 8.14×10^6 in immunosuppressed patients [Table 2]. A comparison of data among immunocompetent vs immunocompromised patients presented in Table 4.

Attempts to isolate the SARS-CoV-2 from non-respiratory tract samples are detailed in Table 3. No viral blood culture was positive in any patient with PCR-positive blood samples²¹²⁹. In studies of PCR-positive stool samples, stool viral cultures were positive in 4 out of 21 (19%) samples¹²²¹³⁰³¹³². In one of the above-mentioned studies, no patient with culture –positive stools had diarrhea or any other GI symptoms¹². A single case report describing a patient with PCR-positive and culture-positive urine was published³³, in another study that tested urine specimens, none of the 5 samples were culture positive³². In a cohort of women with COVID-19, only one out of 64 samples of breast milk were PCR – positive, and none were culture positive³⁴. Two studies tested tears in patients with or without conjunctivitis, in one study PCR and viral culture were negative for all samples³⁵. In the second study two samples out of 60 yielded positive PCR results from a patient with conjunctivitis, non-yielded positive culture results³⁶.

Discussion

In this systematic review we analyzed data on COVID-19 viral culture positivity among patients with positive molecular tests for the virus. Among patients the were reported to be immunocompetent with mild clinical disease, nearly all patients (~99%) are culture negative by day 10, even if PCR tests still were positive. In patients with severe COVID-19, or among immunocompromised hosts, viral cultures may remain positive for longer periods up to 32 days [table 1b].

In many viral infections including the coronaviruses (SARS-CoV, Middle East respiratory syndrome coronavirus) as well as in other RNA viruses such as influenza, Ebola and measles, it is well known that viral RNA can be detected long after infectious virus is cleared. This phenomenon is due to neutralization of the virus by the immune system. As viral RNA undergoes physical interaction with host cells and is protected by viral proteins, it can still be detected when the patient is no longer infectious. With measles virus for example, viral RNA can still be detected 6–8 weeks after the clearance of infectious virus³⁷³⁸³⁹.

This *In-vitro* determination of the duration of infectivity correlates well with epidemiological studies. In a study from Taiwan, the authors describe a higher rate of infection in people exposed to index patients within 5 days of symptoms onset and lower rates of transmission after this time. In this study, the actual attack rate among 1818 patients whose exposure to index cases started within 5 days of symptom onset was much higher (1.0% [95% CI, 0.6%-1.6%]) when compared with those who were

exposed later (0 cases from 852 contacts; 95% CI, 0%-0.4%)⁴⁰. These epidemiological findings, supported by other similar observations, are consistent with our summary of *In-vitro* studies that showed nearly no viral infectivity beyond 10 days of symptoms onset¹³.

The other factor that may affect culture positivity is the viral load reflected either by a high number of viral genome copy number, or by a relatively low Ct count. In this study we found that practically all patients with a positive PCR test, but with Ct > 35 or with a calculated viral copies less than 1.2×10^3 copies / mL, were culture negative. The reported result of a positive culture derived from a single clinical sample with a Ct value of 41²⁵ is exceptional and warrants further investigation, in light of the accumulating data suggesting that samples with Ct values above 30-35 are non-infectious.

In this respect, it is important to consider the specific properties of the test performed. Different studies use different qPCR assays to evaluate the viral load in the human sample or in the cell culture. Although the vast majority of these tests have sufficient sensitivity to detect down to 50-100 copies in the reaction tube, they do vary, and the same sample can give a Ct result of 35 in one test, and Ct of 40 in another.

Furthermore, since the sensitivity of different PCR assays differs, and different laboratories use different extraction procedures (not including direct PCR tests, without RNA extraction), it is very challenging to accurately determine the actual viral load reported, based on Ct values. Ideally, results from quantitative PCR should be reported

in viral RNA copies, rather than Ct values. However, accurate calculation of the viral load, as expressed by the number of target viral RNA copies detected in the sample, requires using calibrated standards. This is depended on availability of RNA standards in an established concentration and on performing a calibration test in parallel to the diagnostic test. This is not practical for many diagnostic laboratories, currently struggling with an overload of work. To accurately evaluate the viral load in the studies reviewed here, we undertook to experimentally determine the sensitivity of some of the qPCR assays described, as detailed in the Supplementary data. We show that while the E-Sarbeco and CDC N1 reactions have a similar sensitivity, the N-Sarbeco and RdRP reactions are approximately 5 to 10-fold less sensitive.

Adult patients with severe COVID-19 have prolonged viral shedding as reflected by a continuously positive PCR. A study from China showed that patients with severe disease had a median duration of viral shedding of 31.0 (IQR, 24.0–40.0) days from illness onset¹⁵. In a comparative observational study the median duration of viral shedding, reflected by positive PCR assays, was significantly longer among patients with severe disease (21 days, 14-30 days) than among patients with mild disease (14 days, 10-21 days; $P=0.04$)⁴¹. However, in none of these studies the presence of viable virus was studied. In this review, we present reports of samples from patients with either a severe disease or immunosuppression who were shedding infectious virus for longer than 10 days, up to 32 days after symptoms onset. Interestingly, the Ct levels and number of viral copies were similar in these cases to immunocompetent patients without severe disease. A possible explanation stems from the fact that the PCR test does not

necessarily reflect infectivity; it provides information on the number of target copies present in the sample. Therefore, it is conceivable that in some immunocompromised patients, most or all of the detectable viral RNA is from infectious virus, while in other patients, most of the RNA is from partially degraded, non-infectious virions. In such cases, the PCR results are not consistent with actual infectivity and can be misleading. These assumptions are theoretical at this point and definitely warrant further investigation to allow comprehensive understanding of the infectivity state of such cases. Furthermore, persistent infection in immunocompromised patients can accelerate the development of new variants with increased virulence or infectivity, as suspected of the development of the recent British VOC⁴².

Interestingly in children, the rate of infectious virus isolation was largely comparable to that of adults, although two specimens yielded an isolate at a lower VL (1.2×10^4 and 1.4×10^5 copies/ml, respectively) than what was observed in adults²². Since SARS-CoV-2 shedding patterns in symptomatic children resemble those observed in adults, transmission of SARS-CoV-2 from children is plausible. The notion that small children are less infectious should therefore be attributed to other factors rather than the viral load per se⁴³.

PCR-positive samples that were cultured successfully from non-respiratory tract such as: urine and stool, were reported in very few cases, thereby leading to the conclusion that

infection via these secretion routes is probably negligible compared with respiratory route transmission.

No positive viral cultures of blood, tears and breast milk samples were reported thus far. Fecal-oral transmission, or transmission via contact with urine is therefore likely to be either non-existent or extremely rare. Similarly, as viable virus could not be detected in breast milk, breast feeding should be recommended for COVID-19 infected mothers who can take the necessary precautions⁴⁴.

As the economic, social, and psychological toll of the COVID-19 pandemic becomes ever more evident, and second waves of infections inflicting many countries, it is vital to keep to a necessary minimum the number of days in which recovering COVID-19 patients remain isolated. From the data analyzed in this systematic review, it seems that for nearly all immuno-competent recovering patients with a non – severe infection, no more than ten days from the onset of symptoms would suffice. Such a recommendation would negate the need for additional PCR testing in such patients, as accepted now by several countries. The logistical, economic, social, and psychological benefits of such change are evident.

The maximal Ct values in a PCR – based assay that should be considered truly positive (rather than representing degrading genetic material) should be < 35 in nearly all cases. It is likely that for immunocompetent patients recovering clinically, a PCR test with Ct> 35 should be considered negative, leading to shortening of the recommended isolation periods. In the context of a local outbreak that strain laboratory resources and with

limited isolation capabilities, a Ct value of 30 may also be considered, as only 10% of samples remained culture positive above this value.

A more standardized assessment of the infection state is calculating the viral genome copies, rather than relying on Ct values that are affected by the assay used. Reporting the test result using viral copy number, may minimize bias due to differences in assay performance, and allow comparison between results from different laboratories and different methods. This, however, requires the diagnostic laboratory to determine the reaction sensitivity and run routine calibration curves while performing clinical tests. In this study, we provide a schematic workflow for converting Cq values into target copies, allowing standardization of the method and comparison of results from different laboratories which use different qPCR tests.

The limitations of this review include lack of larger cohorts of COVID-19 with serial viral cultures, lack of standardization of laboratory techniques, and a reporting bias that probably leads to overestimation of cases with prolonged viral shedding. Another fundamental limitation is the need to draw practical conclusions from cell culture data. Due to obvious safety limitations, it is very complicated to perform SARS-CoV-2 infection experiments with animal model. Therefore, cell tissue culture (T/C) is the best practical alternative. However, inferring human viral infectivity based on T/C experiments needs to be done carefully, due to multiple stages, which the sample undergoes, before the test is performed. Sampling method, sampling media, transport timeline and storage conditions, may all affect the test results. Additionally, factors such as different susceptibility and presence/absence of effective immune response may also affect the

quantity of infectious virus in the sample. A second limitation concerning T/C results is the different practices between different laboratories, different conditions of the cells used and different sample processing methods. Each of these factors separately can have a profound effect on the outcome. Combined, they render the comparison of different studies performed under different conditions very complicated. The reasons for unsuccessful culturing may stem from technical issues, and should therefore be interpreted carefully, when trying to correlate qPCR results with T/C infectivity. Nevertheless, as described above, the available epidemiological studies support the timeline delineated by culture positivity, suggesting that the infectious state is almost entirely during the early post-symptoms state.

In summary, the insights of this systematic review are important for policy makers worldwide. During the pandemic phase, symptomatic COVID-19 patients who are recovering from a mild infection and are not immuno-suppressed, can be regarded as non-infectious, within 10 days from symptom onset, with no additional testing needed. Based on the described findings, calculated viral load below 1.2×10^3 copies per mL can be regarded as non-infectious. Recovering COVID-19 patients or asymptomatic patients whose tests yield Ct values greater than 35 may be regarded as non-infectious and may be freed from isolation. When the time of near elimination would (hopefully) come, more stringent criteria for defining non-infectiousness may be used. Urine, feces, blood, and breast milk may be regarded as largely non – infectious, based on the described results.




Figure 1:

Table 1 – Comparison of the duration of SARS-CoV-2 PCR and culture positivity

Country	study design	Positive cultures/total samples	population; median age – years (range)	PCR positive ²	Med. culture pos. ³	Max. culture pos. ⁴	Target	Cell type
Taiwan ¹⁴	retrospective case series	23/60	inpatients and outpatients; NA	26	3	11 (1 case)	E, N, nsp12	Vero-E6, MK-2
Canada ⁷	retrospective cross-sectional	26/90	N\A; 45 (30-59)	21	N\A	8	E	Vero
USA ²⁸	Retrospective cohort	31/47	asymptomatic and symptomatic elderly patients; N\A	21	N\A	9	N1 and N2	Vero CCL-81
Taiwan ²⁶	case report	1/1	50 years old inpatient	63	N\A	18	E, N, RdRp	Vero E6, LLC-MK2
Germany ¹²	Retrospective case series	9/16	inpatients; N\A	28	5	8	E, RdRp	Vero E6
USA ²¹	Retrospective case series	9/12	53 (21-68) ¹	36	4	9	N\A	Vero CCL-81
Switzerland ²²	Retrospective case series	12/23	neonates, children, adolescents; 12 (0-15)	N\A	1.5	3	E	Vero E6
Hong Kong ²⁰	retrospective cohort	16/68	symptomatic and asymptomatic; 39 (21-73)	67	N\A	8	N	N\A
France ⁴⁵	retrospective case series	46/80	52.5 (20-88)	14	N\A	9	E	Vero E6
England ²⁵	retrospective study	103/246	symptomatic and asymptomatic; N\A	N\A	4	12	RdRp	Vero E6

Studies that included immunocompromised and severely ill patients

Country	study design	Positive cultures /total samples	population; median age – years (range)	population; median age – years (range)	PCR positive ²	Med. culture pos. ³	Max. culture pos. ⁴	Target	Cell type
Germany ²⁷	Case report	1 (1)	62 years old immunocompromised patient	N\A	35	N\A	21	N\A	N\A
Netherlands ²⁴	retrospective case series	690 (62)	immunocompetent, immunocompromised; 65 (57-72)	N\A	36	8	20	E	Vero clone 118
Spain ¹⁹	retrospective case series	106 (49)	inpatients and outpatients; 42 (32-50)	16.5, 19.6	N\A	N\A	32	E	Vero E6

Australia ²³	Retrospective case series	228 (56)	ICU, hospitalized, outpatients; 40 (8-78)	11	29	3.5	18	E, RdRp, N, M, ORF1ab	Vero E6
-------------------------	---------------------------	----------	---	----	----	-----	----	-----------------------	----------------

¹ Median duration of PCR positivity was 19 days

² Maximal duration of PCR positivity (days)

³ Median duration of culture positivity (days)

⁴ Maximal duration of culture positivity (days)

Table 2 – Minimal viral loads and maximal cycle threshold for positive SARS-CoV-2 cultures

Country	study design	Cultured/Total ¹	Pop., Med. age ²	Max. Ct ³	Med. Ct ⁴	Min. copies / mL ⁵	Med. copies / mL ⁶	Target	Cell type
Taiwan ¹⁴	retrospective case series	23/60	inpatients and outpatients; N\A	35.2	N\A	5.4x10 ⁶	N\A	E, N, nsp12	Vero-E6, MK-2
France ⁴⁶	retrospective case series	129/183	N\A	34		2.5X10 ³		E	Vero E6
Canada ⁷	retrospective cross-sectional case series	26/90	45 (30-59)	24	23 (17-32)	3.7X10 ⁶	N\A	E	Vero
USA ²⁸	retrospective cohort	31/47	asymptomatic and symptomatic elderly patients; N\A	34	24.8	4.1X10 ³	N\A	N1 and N2	Vero, CCL-81
Germany ¹²	retrospective case series	9/16	inpatients; N\A	26.5	N\A	E – 7X10 ⁵ RdRp-4x10 ⁶	N\A	E, RdRp	Vero E6
England ⁴⁷	prospective, cross-sectional case series	1/19	symptomatic and asymptomatic; 39 (19–68)	26.2	N\A	2-4 X10 ⁶	N\A	RdRp, E, N	Vero E6
Switzerland ²²	retrospective case series	12/23	neonates, children, adolescents; 12 (0 -15)	N\A	N\A	1.2x10 ⁴	1.7x10 ⁸	E	Vero E6
Hong Kong ²⁰	retrospective cohort	16/68	symptomatic and asymptomatic; 39 (21-73)	N\A		1X10 ⁵	N\A	N	N\A
England ²⁵	retrospective study	133/324	symptomatic and asymptomatic; N\A	41.78	31.15 (27.5-33.86)	N\A	N\A	RdRp	Vero E6

Studies that included immunocompromised and severely ill patients

Country	study design	Cultured/Total ¹	Pop., Med. age ²	Max. Ct ³	Med. Ct ⁴	Min. copies / mL ⁵	Med. copies / mL ⁶	Target	Cell type
Australia ²³	retrospective		ICU, inpatients, outpatients; 40			E: 1.1X10 ⁴			
	case series	228 (56)	(8-78)	32		RdRp: 1.2X10 ⁵ N - ~5.6 X10 ⁵		E, RdRp, N, M, ORF1ab	Vero E6
Spain ¹⁹	retrospective		inpatients and outpatients; 42						
	case series	106 (49)	(32-50)	35		1.2X10 ³	N\A	E	Vero E6
Netherlands ²⁴	retrospective		immunocompetent,				8.14x10 ⁶		Vero clone
	case series	690 (62)	immunocompromised; 65 (57-72)	N\A		1x10 ^{5.5}		E	118

¹ No. of positive culture / Total No. of samples

² Population; Median age – years (range)

³ Maximal Ct With Culture Positivity

⁴ Median Ct of positive culture (IQR)

⁵ Minimal copies / mL with culture positivity

⁶ Median copies / mL with positive culture

Table 3 – Comparison of PCR positivity and culture positivity in non-respiratory body fluids

Body Fluid	Country	Study Design	Pos. culture/Pos. PCR ¹	Pop.; Med. age ²	Target	Culture type
Blood	United Kingdom ²⁹	Retrospective cohort	0/16	Acute and convalescent adult patients; 57 (46-76)	RdRp, ORF1ab	Vero E6
Blood	USA ³⁰	Retrospective case series	0/1	30-39 years old inpatient	N\A	Vero CCL-81
Stool	USA ²¹	Retrospective case series	0/7	53 (21-68)	N\A	Vero CCL-81
Stool	China ³⁰	Retrospective cohort	2/4	Inpatients; 44 (5-67)	ORF1ab	N\A
Stool	China ³¹	Retrospective case series	2/3	Adult inpatients; N\A	ORF1ab	Vero E6
Stool	Germany ¹²	Retrospective case series	0/4	Adult inpatients; N\A	E, RdRp	Vero E6
Stool	South Korea ³²	Retrospective case series	0/3	63 (51-64); mild to critical disease	N\A	Vero
Urine	China ³³	Case report	1/1	72 year old male inpatient with Severe COVID-19	N\A	Vero E6
Urine	South Korea ³²	Retrospective case series	0/5	63 (51-64); mild to critical disease	N\A	Vero
Breast Milk	USA ³⁴	Prospective cohort	0/1	25-29 years old woman	N\A	Vero E6
Tears	Singapore ³⁵	Prospective study	0/0	Adult inpatients; N\A	N\A	Vero E6
Tears	China ³⁶	Prospective case series	0/2	54 (13-83)	N\A	N\A

¹ No. of positive culture / Total No. of samples² Population; Median age – years (range)

Table 4 - Positive culture Characteristics

	Immunocompetent –mild disease	Immunocompromised or severe disease
Positive culture samples	276	168
Positive culture>10 days	4 (1.4%)	31 (18%)
Positive culture>20 days	0	3 (1.8%)
Maximal reported days post-symptoms onset	18 days	32 days
Maximal Ct level with positive culture	41.48	35
Minimal Viral Load (copies) with positive culture	2.5×10^3	1.2×10^3
Median viral load (copies)	1.7×10^8	8.1×10^6

References

1. Who. *COVID-19 Weekly Epidemiological Update Global Overview*.
2. Bayham J, Fenichel EP. Impact of school closures for COVID-19 on the US health-care workforce and net mortality: a modelling study. *Lancet Public Heal*. 2020;5(5):e271-e278. doi:10.1016/S2468-2667(20)30082-7
3. Tandon P. COVID-19: Impact on health of people & wealth of nations. *Indian J Med Res*. 2020;0(0):0. doi:10.4103/ijmr.IJMR_664_20
4. Ayittey FK, Ayittey MK, Chiwero NB, Kamasah JS, Dzuvor C. Economic impacts of Wuhan 2019-nCoV on China and the world. *J Med Virol*. 2020;92(5):473-475. doi:10.1002/jmv.25706
5. Economy News: OECD Sees Virus Damage and Warns of Second Wave - Bloomberg. <https://www.bloomberg.com/news/articles/2020-06-10/virus-splinters-global-economy-exposing-inequality-faultlines?cmpid=BBD061020`CORONAVIRUS&utm`medium=email&utm`source=newsletter&utm`term=200610&utm`campaign=coronavirus>. Accessed January 28, 2021.
6. COVID-19 to Plunge Global Economy into Worst Recession since World War II. <https://www.worldbank.org/en/news/press-release/2020/06/08/covid-19-to-plunge-global-economy-into-worst-recession-since-world-war-ii>. Accessed January 28, 2021.
7. Bullard J, Dust K, Funk D, et al. Predicting Infectious Severe Acute Respiratory Syndrome Coronavirus 2 From Diagnostic Samples. *Clin Infect Dis*. 2020;71(10):2663-2666. doi:10.1093/cid/ciaa638
8. Bedford J, Enria D, Giesecke J, et al. COVID-19: towards controlling of a pandemic. *Lancet*. 2020;395(10229):1015-1018. doi:10.1016/S0140-6736(20)30673-5
9. Strong JE, Feldmann H. The Crux of Ebola Diagnostics. *J Infect Dis*. 2017;216(11):1340-1342. doi:10.1093/infdis/jix490
10. Falsey AR, Formica MA, Treanor JJ, Walsh EE. Comparison of quantitative reverse transcription-PCR to viral culture for assessment of respiratory syncytial virus shedding. *J Clin Microbiol*. 2003;41(9):4160-4165. doi:10.1128/JCM.41.9.4160-4165.2003
11. Ip DKM, Lau LLH, Chan KH, et al. The Dynamic Relationship between Clinical Symptomatology and Viral Shedding in Naturally Acquired Seasonal and Pandemic Influenza Virus Infections. *Clin Infect Dis*. 2015;62(4):431-437. doi:10.1093/cid/civ909
12. Wölfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature*. 2020;581(7809):465-469. doi:10.1038/s41586-020-2196-x
13. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*. 2020;26(5):672-675. doi:10.1038/s41591-020-0869-5
14. Huang CG, Lee KM, Hsiao MJ, et al. Culture-based virus isolation to evaluate potential infectivity of clinical specimens tested for COVID-19. *J Clin Microbiol*. 2020;58(8). doi:10.1128/JCM.01068-20

15. Zhou B, She J, Wang Y, Ma X. Duration of Viral Shedding of Discharged Patients With Severe COVID-19. *Clin Infect Dis*. 2020;71(16):2240-2242. doi:10.1093/cid/ciaa451
16. Zou J, Zhi S, Chen M, et al. Heat inactivation decreases the qualitative real-time RT-PCR detection rates of clinical samples with high cycle threshold values in COVID-19. *Diagn Microbiol Infect Dis*. 2020;98(1):115109. doi:10.1016/j.diagmicrobio.2020.115109
17. Corman VM, Landt O, Kaiser M, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Eurosurveillance*. 2020;25(3). doi:10.2807/1560-7917.ES.2020.25.3.2000045
18. Real-time RT-PCR Primers and Probes for COVID-19 | CDC. <https://www.cdc.gov/coronavirus/2019-ncov/lab/rt-pcr-panel-primer-probes.html>. Accessed February 14, 2021.
19. Folgueira MD, Luczkowiak J, Lasala F, Pérez-Rivilla A, Delgado R. Persistent SARS-CoV-2 replication in severe COVID-19. *medRxiv*. June 2020:2020.06.10.20127837. doi:10.1101/2020.06.10.20127837
20. Perera RAPM, Tso E, Tsang OTY, et al. SARS-CoV-2 virus culture from the upper respiratory tract: Correlation with viral load, subgenomic viral RNA and duration of illness. *medRxiv*. July 2020:2020.07.08.20148783. doi:10.1101/2020.07.08.20148783
21. Kujawski SA, Wong KK, Collins JP, et al. Clinical and virologic characteristics of the first 12 patients with coronavirus disease 2019 (COVID-19) in the United States. *Nat Med*. 2020;26(6):861-868. doi:10.1038/s41591-020-0877-5
22. Huillier AGL, Torriani G, Pigny F, Kaiser L, Eckerle I. Culture-Competent SARS-CoV-2 in Nasopharynx of Symptomatic Neonates, Children, and Adolescents. 2020;26(10):3-6.
23. Basile K, Mcphie K, Carter I, et al. (which was not certified by peer review) is the author / funder , who has granted medRxiv a license to display the preprint in perpetuity . All rights reserved . No reuse allowed without permission . Cell-based culture of SARS-CoV-2 informs infectivity . 2020.
24. van Kampen JJA, van de Vijver DAMC, Fraaij PLA, et al. Duration and key determinants of infectious virus shedding in hospitalized patients with coronavirus disease-2019 (COVID-19). *Nat Commun*. 2021;12(1):1-6. doi:10.1038/s41467-020-20568-4
25. Singanayagam A, Patel M, Charlett A, et al. Duration of infectiousness and correlation with RT-PCR cycle threshold values in cases of COVID-19, England, January to May 2020. *Eurosurveillance*. 2020;25(32). doi:10.2807/1560-7917.ES.2020.25.32.2001483
26. Liu W Da, Chang SY, Wang JT, et al. Prolonged virus shedding even after seroconversion in a patient with COVID-19. *J Infect*. 2020;81(2):318-356. doi:10.1016/j.jinf.2020.03.063
27. Decker A, Welzel M, Laubner K, et al. Prolonged SARS-CoV-2 shedding and mild course of COVID-19 in a patient after recent heart transplantation. *Am J Transpl*. 2020;20:3239-3245. doi:10.1111/ajt.16133
28. Arons MM, Hatfield KM, Reddy SC, et al. Presymptomatic SARS-CoV-2 Infections and Transmission in a Skilled Nursing Facility. *N Engl J Med*. 2020;382(22):2081-2090. doi:10.1056/nejmoa2008457

29. Andersson MI, Arancibia-Cárcamo C V., Auckland K, et al. SARS-CoV-2 RNA detected in blood samples from patients with COVID-19 is not associated with infectious virus. *medRxiv*. June 2020:2020.05.21.20105486. doi:10.1101/2020.05.21.20105486
30. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA - J Am Med Assoc*. 2020;323(18):1843-1844. doi:10.1001/jama.2020.3786
31. Xiao F, Sun J, Xu Y, et al. Infectious SARS-CoV-2 in feces of patient with severe COVID-19. *Emerg Infect Dis*. 2020;26(8):1920-1922. doi:10.3201/eid2608.200681
32. Jeong HW, Kim SM, Kim HS, et al. Viable SARS-CoV-2 in various specimens from COVID-19 patients. *Clin Microbiol Infect*. 2020;26(11):1520-1524. doi:10.1016/j.cmi.2020.07.020
33. Sun J, Zhu A, Li H, et al. Isolation of infectious SARS-CoV-2 from urine of a COVID-19 patient. *Emerg Microbes Infect*. 2020;9(1):991-993. doi:10.1080/22221751.2020.1760144
34. Chambers C, Krogstad P, Bertrand K, et al. Evaluation for SARS-CoV-2 in Breast Milk from 18 Infected Women. *JAMA - J Am Med Assoc*. 2020;324(13):1347-1348. doi:10.1001/jama.2020.15580
35. Seah IYJ, Anderson DE, Kang AEZ, et al. Assessing Viral Shedding and Infectivity of Tears in Coronavirus Disease 2019 (COVID-19) Patients. *Ophthalmology*. 2020;127(7):977-979. doi:10.1016/j.optha.2020.03.026
36. Xia J, Tong J, Liu M, Shen Y, Guo D. Evaluation of coronavirus in tears and conjunctival secretions of patients with SARS-CoV-2 infection. *J Med Virol*. 2020;92(6):589-594. doi:10.1002/jmv.25725
37. Peiris JSM, Chu CM, Cheng VCC, et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: A prospective study. *Lancet*. 2003;361(9371):1767-1772. doi:10.1016/S0140-6736(03)13412-5
38. Chan KH, Poon LLLM, Cheng VCC, et al. Detection of SARS Coronavirus in Patients with Suspected SARS. *Emerg Infect Dis*. 2004;10(2):294-299. doi:10.3201/eid1002.030610
39. Lin WHW, Kouyos RD, Adams RJ, Grenfell BT, Griffin DE. Prolonged persistence of measles virus RNA is characteristic of primary infection dynamics. *Proc Natl Acad Sci U S A*. 2012;109(37):14989-14994. doi:10.1073/pnas.1211138109
40. Cheng HY, Jian SW, Liu DP, Ng TC, Huang WT, Lin HH. Contact Tracing Assessment of COVID-19 Transmission Dynamics in Taiwan and Risk at Different Exposure Periods before and after Symptom Onset. *JAMA Intern Med*. 2020;180(9):1156-1163. doi:10.1001/jamainternmed.2020.2020
41. Zheng S, Fan J, Yu F, et al. Viral load dynamics and disease severity in patients infected with SARS-CoV-2 in Zhejiang province, China, January-March 2020: Retrospective cohort study. *BMJ*. 2020;369. doi:10.1136/bmj.m1443
42. Ecdc. *Suggested Citation: European Centre for Disease Prevention and Control. Rapid Increase of a SARS-CoV-2 Variant with Multiple Spike Protein Mutations Observed in the Rapid Increase of a SARS-CoV-2 Variant with Multiple Spike Protein Mutations Observed in the United Kingdom.*; 2020. <http://covid19-country->

overviews.ecdc.europa.eu/#34_United_Kingdom. Accessed February 14, 2021.

43. Lee B, Raszka W V. COVID-19 transmission and children: The child is not to blame. *Pediatrics*. 2020;146(2). doi:10.1542/peds.2020-004879
44. Coronavirus Disease (COVID-19) and Breastfeeding | Breastfeeding | CDC. <https://www.cdc.gov/breastfeeding/breastfeeding-special-circumstances/maternal-or-infant-illnesses/covid-19-and-breastfeeding.html>. Accessed February 14, 2021.
45. Gautret P, Lagier JC, Parola P, et al. Clinical and microbiological effect of a combination of hydroxychloroquine and azithromycin in 80 COVID-19 patients with at least a six-day follow up: A pilot observational study. *Travel Med Infect Dis*. 2020;34. doi:10.1016/j.tmaid.2020.101663
46. La Scola B, Le Bideau M, Andreani J, et al. Viral RNA load as determined by cell culture as a management tool for discharge of SARS-CoV-2 patients from infectious disease wards. *Eur J Clin Microbiol Infect Dis*. 2020;39(6):1059-1061. doi:10.1007/s10096-020-03913-9
47. Brown CS, Clare K, Chand M, et al. Snapshot PCR surveillance for SARS-CoV-2 in hospital staff in England. *J Infect*. 2020;81(3):427-434. doi:10.1016/j.jinf.2020.06.069