

1 **Assessing the impact of community-based interventions on hypertension and**
2 **diabetes management in three Minnesota communities: findings from the**
3 **prospective evaluation of US HealthRise programs**

4
5 Nancy Fullman MPH^{1†}, Krycia Cowling PhD², Luisa S Flor PhD¹, Shelley Wilson BA¹, Paurvi Bhatt MPH³,
6 Miranda F Bryant MPH¹, Joseph N Camarda BA¹, Danny V Colombara PhD⁴, Jessica Daly MPH³, Rose K
7 Gabert MPH MD⁵, Katie Panhorst Harris MPA¹, Casey K Johanns MPH¹, Charlie Mandile MPH⁶, Susan
8 Marshall BA MAESL⁷, Claire R McNellan MPH⁸, Vasudha Mulakaluri MS¹, Bryan K Phillips MPH⁹, Marissa B
9 Reitsma ScB¹, Naomi Sadighi MPH¹⁰, Tsega Tamene AB¹¹, Blake Thomson DPhil¹², Alexandra Wollum
10 MPH¹³, Emmanuela Gakidou PhD¹

11
12
13 ¹Institute for Health Metrics and Evaluation, Department of Health Metrics Sciences, University of
14 Washington, Seattle, Washington, USA

15 ²US Department of Health and Human Services, Washington DC, USA

16 ³Medtronic Foundation, Minneapolis, Minnesota, USA

17 ⁴Public Health Seattle and King County, Seattle, Washington, USA

18 ⁵University of Washington School of Medicine, University of Washington, Seattle, Washington, USA

19 ⁶HealthFinders Collaborative, Northfield, Minnesota, USA

20 ⁷International School of Djibouti – Ministere de l'Education Nationale, Djibouti City, Djibouti

21 ⁸School of Social Work, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

22 ⁹The Permanente Medical Group, Oakland, California, USA

23 ¹⁰Small Sums, St Paul, Minnesota, USA

24 ¹¹Pillsbury United Communities, Minneapolis, Minnesota, USA

25 ¹²American Cancer Society, Atlanta, Georgia, USA

26 ¹³Ibis Reproductive Health, Cambridge, Massachusetts, USA

27
28
29 †Corresponding author:

30 Nancy Fullman

31 Institute for Health Metrics and Evaluation

32 Department of Health Metrics Sciences

33 University of Washington

34 Seattle, WA 98195

35 USA

36 Email: nf4@uw.edu

37

38

39

40

41

42 **Abstract**

43 **Background**

44 Community-based health interventions are increasingly viewed as models of care that can bridge
45 healthcare gaps experienced by underserved communities in the United States (US). With this study, we
46 sought to assess the impact of such interventions, as implemented through the US HealthRise program,
47 on hypertension and diabetes among underserved communities in Hennepin, Ramsey, and Rice
48 Counties, Minnesota.

49
50 **Methods and findings**

51 HealthRise patient data from June 2016 to October 2018 were assessed relative to comparison patients
52 in a difference-in-difference analysis, quantifying program impact on reducing systolic blood pressure
53 (SBP) and hemoglobin A1c, as well as meeting clinical targets (< 140 mmHg for hypertension, < 8% A1c
54 for diabetes), beyond routine care. For hypertension, HealthRise participation was associated with SBP
55 reductions in Rice (6.9 mmHg [95% confidence interval: 0.9–12.9]) and higher clinical target
56 achievement in Hennepin (27.3 percentage-points [9.8–44.9]) and Rice (17.1 percentage-points [0.9 to
57 33.3]). For diabetes, HealthRise was associated with A1c decreases in Ramsey (1.3 [0.4–2.2]). Qualitative
58 data showed the value of home visits alongside clinic-based services; however, challenges remained,
59 including community health worker retention and program sustainability.

60
61 **Conclusions**

62 HealthRise participation had positive effects on improving hypertension and diabetes outcomes at some
63 sites. While community-based health programs can help bridge healthcare gaps, they alone cannot fully
64 address structural inequalities experienced by many underserved communities.

65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82

83 Introduction

84 Longstanding health disparities occur throughout the US,[1] with differences often manifesting across
85 multiple dimensions (e.g., geography, gender, race or ethnicity, socioeconomic status).[2] Non-
86 communicable diseases (NCDs) and NCD-related risks like hypertension can uniquely affect underserved
87 communities, as a constellation of factors – from service affordability and inadequate insurance to more
88 entrenched socioeconomic obstacles like low access to nutritional food – can quickly give way to high
89 rates of chronic, debilitating conditions without good access to care or appropriate services.
90 Community-based interventions with integrated care have emerged as approaches to bridge gaps in
91 NCD care, particularly for underserved areas in the US.[3–9] Nonetheless, the effectiveness of these
92 interventions vary across settings; interventions implemented; roles of community health workers
93 (CHWs)(e.g., direct involvement in patient care[10] *versus* assisting health providers who are then
94 responsible for service provision[11]); and NCDs targeted. While CHWs and community-based
95 interventions appear to be promising models of care for hypertension and diabetes in underserved
96 communities,[5–7] more rigorous evaluations across more local contexts and populations are needed.

97
98 To strengthen this evidence base, the HealthRise program was developed to implement and pilot locally-
99 tailored programs for improving screening, diagnosis, management and control of hypertension and
100 diabetes among underserved communities.[12–14] Funded by the Medtronic Foundation, HealthRise
101 took place in nine communities in four countries – Brazil, India, South Africa, and the US – from 2014 to
102 2018. In the US, Minnesota was selected, a state which generally surpasses national averages and ranks
103 among the healthiest across many health measures.[1,15] Nonetheless, county- and sub-county level
104 health disparities remain in Minnesota, particularly among populations that face compounding barriers
105 to care and improved health outcomes.[15,16] To maximize the potential impact of HealthRise
106 programs, especially within a relatively short time span (i.e., the earliest US program began in 2016),
107 HealthRise targeted geographic areas with the greatest need and highest disease burden. A 2014-2015
108 needs assessment identified communities in three counties – Hennepin, Ramsey, and Rice – as
109 candidates for HealthRise programs based on their high NCD burden and risk profiles, challenges in
110 healthcare access, and sociodemographic characteristics (e.g., large minority and/or immigrant
111 populations).[15,17] Three main recommendations emerged from this assessment: (1) focus on people
112 with the highest need and poorest clinical outcomes (2) use multi-faceted interventions to address
113 multiple risks and comorbid conditions; and (3) identify opportunities to integrate CHWs within formal
114 health system functions. Based on this needs assessment and grantee applications, three implementing
115 partners were selected (i.e., one per county), community-based interventions were developed, and US
116 HealthRise program implementation took place from June 2016 to October 2018.

117
118 With this study, we provide key findings from HealthRise programs in Hennepin, Ramsey, and Rice
119 Counties, Minnesota. Based on quantitative and qualitative data collected over the course of program
120 implementation, we evaluated the potential impact of these community-based interventions on
121 improving clinical and health outcomes for hypertension and diabetes patients. We conducted
122 difference-in-difference analyses in relation to comparison patients to quantify this impact above and
123 beyond what might be expected for demographically similar patients under routine care in the same

124 communities. This study contributes to the science supporting the role of community-based programs in
125 elevating the health of underserved communities, in the US and elsewhere.

126
127

128 **Materials and Methods**

129 **Study overview, design, and interventions**

130 This analysis follows the global HealthRise prospective evaluation framework, which was established in
131 2014 and agreed upon by all partners; greater detail on the global team structure, interventions, and
132 analyses are provided elsewhere.[12–14] In sum, HealthRise was funded by the Medtronic Foundation,
133 with global and US implementation partners coordinated by Abt Associates and evaluation activities
134 overseen by the Institute for Health Metrics and Evaluation (IHME). Though ongoing coordination and
135 collaboration occurred across implementation and evaluation organizations, these grant streams were
136 purposefully structured and funded separately to support an independent assessment of the HealthRise
137 programs.

138

139 For US HealthRise, patient-level monitoring data were routinely collected and collated by grantees
140 during program implementation (June 2016 to October 2018). Using a mixed-methods quasi-
141 experimental design, we synthesized qualitative and quantitative data from HealthRise and comparison
142 patients and stakeholders (e.g., service providers, administrators, and policymakers) to inform its
143 endline evaluation. Table 1 summarizes key information on each US HealthRise site and interventions
144 implemented by grantee, as interventions were tailored to address gaps or challenges identified for
145 each site during the 2014-2015 needs assessment;[15,17] more detail is available elsewhere.[14]

146

147 **Definitions and data**

148 ***Definitions***

149 We used following case definitions for hypertension and diabetes at each time point: (1) prevalent cases
150 were patients with documented diagnoses, or patients without prior diagnoses but with clinical readings
151 that would qualify for diagnosis (i.e., systolic blood pressure [SBP] \geq 140 mmHg or diastolic blood
152 pressure [DBP] \geq 90 mmHg for hypertension; hemoglobin A1c \geq 6.5% for diabetes); (2) diagnosed cases
153 were patients with documented diagnoses; and (3) patients meeting treatment targets were prevalent
154 cases with SBP $<$ 140 mmHg and DBP $<$ 90 mmHg for hypertension, and A1c $<$ 8% for diabetes. If DBP
155 measures were not available for a given patient, then only SBP readings were used.

156

157 ***Endline evaluation data collection***

158 **HealthRise patient data.** Each US grantee collected patient-level data from existing sources and
159 provided de-identified data over time. To best capture potential program impact, analyses were limited
160 to HealthRise patients who (1) remained enrolled in HealthRise at endline (i.e., never withdrew from
161 programs); and (2) had at least two separate biometric data points for blood pressure (i.e., ideally both
162 SBP and DBP, but at minimum, SBP) or A1c (Table 2). Subsequently, evaluation results reflected
163 potential effects from HealthRise participation, and not “intention to treat,” which would have included
164 patients who enrolled but then withdrew from the program at some point. Rates of any program
165 withdrawal varied by site, ranging from 16.7% (n=19) for Hennepin to 32.5% (n=25) for Ramsey.

166 **Table 1. Overview of interventions by US HealthRise site.** More detailed descriptions of HealthRise interventions, as provided by grantees and compiled by
 167 Abt Associates, are available elsewhere.[14]
 168

	Hennepin County	Ramsey County	Rice County
HealthRise grantee and/or local partners	Pillsbury United Communities (PUC) North Rising partnership comprised of PUC and North Memorial Health, a network of hospitals and clinics	Regions Hospital Foundation Minnesota Community Care (MCC) (formerly named West Side Community Health Services, abbreviated WSCHS)	HealthFinders Collaborative, Inc (HFC)
HealthRise implementation location	North Minneapolis, Minnesota	Saint Paul, Minnesota	Northfield, Minnesota
HealthRise program period	July 2016 to September 2018	June 2016 to September 2018	September 2016 to October 2018
Key characteristics and/or challenges for communities served by HealthRise	Shared across sites <ul style="list-style-type: none"> • High levels of poverty and/or unemployment (e.g., 40% at or below 200% poverty line in Hennepin; 97% of MCC clinic patients in Saint Paul are below 200% poverty line) • High proportion of population are non-white and/or immigrants or refugees (e.g., past HFC patients were 60% Latino immigrants and 25% Somali refugees in Rice County; 30-65% of MCC clinic patients do not speak English as their primary language; North Minneapolis population is about 50% African American, 20% Asian, 15% Caucasian, and 15% Hispanic/Other) • Insufficient support of disease management and behavioral changes to improve health within formal health system structures or communities (e.g., poor access to high-quality education, healthcare, and nutritious foods in Hennepin County; HFC specifically targeted uninsured or individuals with public insurance plans in Rice County) • Inadequate or poor integration of community health care systems, including data systems (e.g., minimal integration of electronic health records across community health care sites in Hennepin; data systems between MCC clinics and hospitals in Ramsey were not integrated) 		
	Site-specific	<ul style="list-style-type: none"> • Urban setting (Twin Cities area) • Reported low trust in local health systems 	<ul style="list-style-type: none"> • Urban setting (Twin Cities area) • Reported high levels of emergency department repeat users
Key HealthRise interventions and activities	Shared across sites <ul style="list-style-type: none"> • Community-based programs and training: hired and trained community health workers (CHWs) and community paramedics (CPs) to provide home-based care and linkages to clinic-based provider teams (e.g., doctors, nurses, pharmacists, clinical care coordinators, and diabetes education in Hennepin County); for Rice County, where HFC already had pre-existing CHW/CP care teams and networks, additional training and expanded services occurred (e.g., mental health, on-site lab for easier access to diagnostic tests) • Home-based care: CHWs and CPs visited patients for disease management (i.e., monitor health status, medications), health education (e.g., health education, healthy food cooking demonstrations), and support for social needs or social determinants of health (e.g., insurance, housing, transportation); often tailored frequency of in-home visits to patient care plans and based on trends in clinical targets • Technologies for care coordination: implemented tools to better coordinate care between CHW/CP teams and clinic-based teams (e.g., Pathways from Care Coordination Systems for Ramsey County) or incorporated home visit information into electronic medical record (EMR) systems (e.g., HFC designed EMRs to include in-home information into patient medical records at clinics) • Community activities and wellness programs: led via CHWs or supported via community centers to support nutrition education and resource connection (e.g., healthy eating demonstrations in Hennepin County and Opportunity grant nutrition-focused program in Ramsey County); disease management tailored to cultural and linguistic needs (e.g., monthly and quarterly diabetes management classes and Somali Health series in Rice County); and exercise/wellness programs (e.g., Pura Vida which included exercise classes, cooking and nutrition classes, etc. in Rice County). 		
	Site-specific	<ul style="list-style-type: none"> • Established community-care teams of CHWs and CPs linked to clinic-based care teams: this model of care was relatively new to PUC and Hennepin County partners, so recruitment of CHWs/CPs and training with care teams occurred alongside other HealthRise-supported activities • Implemented interdisciplinary approaches to improving health: established a full-service grocery store (North Market) with linkages to an interdisciplinary wellness team (e.g., CHWs, nutritionist, pharmacy liaison, coordinator) and a Wellness Resource Center with North Memorial Health 	<ul style="list-style-type: none"> • Established community-care teams of CHWs and CPs linked to clinic-based care teams: this model of care was relatively new to Regions and Ramsey County partners, so recruitment of CHWs/CPs and training with care teams occurred alongside other HealthRise-supported activities • Focused on community-based nutrition programs: used Opportunity Grant to develop and implement a nutrition-focused program, in both English and Spanish, wherein sessions focused on nutrition education, effects of non-nutrition factors on blood sugar (e.g., physical activity, stress management), and grocery store tours highlighting ways to shop for healthy and affordable foods

170 **Table 2. Endline data availability and patient sample sizes by HealthRise site and for intervention and comparison patients.** “Total patients” reflect the
 171 number of patients included in original samples, irrespective of additional inclusion or exclusion criteria. “Patients with baseline data and meeting inclusion
 172 criteria” reflect the number of patients who met inclusion criteria at baseline (i.e, prevalent case of hypertension or diabetes and aged 30-89 years old); for
 173 HealthRise patients, this also reflects the number of patients who formally enrolled in the program, whereas comparison patients were limited to individuals
 174 who could have met eligibility requirements at each site within the program implementation time periods (2016-2018). “Total patients enrolled at endline”
 175 reflect the number of patients who did not withdraw from HealthRise during program duration. At each site and group, some proportion of patients have both
 176 hypertension and diabetes.
 177

Data collection	Hennepin County		Ramsey County		Rice County	
	HealthRise	Comparison	HealthRise	Comparison	HealthRise	Comparison
Quantitative						
Total patients	121	135	78	104	217	311
Patients with baseline data and meeting inclusion criteria	114	113	77	95	157	311
Patients enrolled at endline	95	113	52	95	152	311
Patients with hypertension	85	83	32	66	84	182
Patients with hypertension and ≥ 2 biometric readings	80	83	32	66	80	170
Patients with diabetes	76	75	48	78	125	303
Patients with diabetes and ≥ 2 biometric readings	37	39	43	72	96	296
Qualitative						
Total interviews and focus groups	5	-	9	-	6	-
Community health workers and frontline health workers	2	-	3	-	2	-
Facility- or clinic-based providers	1	-	3	-	2	-
Facility or clinic managers and administrators	2	-	3	-	2	-
Policymakers				3		

178
 179
 180
 181
 182
 183
 184

185 For Rice, 3.2% of patients (n=5) lacked a clinic visit or biometric data since baseline, and thus were considered
186 withdrawn. In Ramsey, most patients who withdrew did so after a few months of enrollment and within the first
187 year of HealthRise implementation.

188

189 For baseline measures, we used biometric data collected at HealthRise program enrollment. If such data were
190 not available at the precise enrollment date, then biometric data were used from the data closest to that of
191 enrollment. For endline measures, we used patients' most recent biometric measurements.

192

193 **Comparison patient data.** Grantees provided comparison data drawn from patient populations similar to those
194 enrolled in HealthRise. Upon receiving each site's dataset, we sought to reconstruct samples of comparison
195 patients that were similar demographically and in terms of baseline health conditions to HealthRise patients
196 (i.e., excluding comparison patients younger than 30 years and 90 years or older, and those without a diagnosis
197 of hypertension or diabetes and had baseline biometric data that fell within disease control categories). As
198 necessary, comparison patient data were censored to correspond with each site's HealthRise program
199 implementation period (Table 1), and thus better approximate similar follow-up times for comparison patients.
200 After this censoring step, we then excluded any comparison patients who lacked more than one measurement
201 of A1c and systolic blood pressure and therefore could not contribute to baseline *versus* endline comparisons.
202 Included comparison patients, by site, are provided in Table 2.

203

204 Comparison patient data selection occurred between October 2018 and January 2019, with criteria determined
205 by site. For Hennepin, data for patients who formed the comparison group were extracted from clinics
206 associated with North Memorial but had not enrolled in HealthRise. Selection criteria included having at least
207 two biometric readings for A1c or SBP – one in 2016 and one in 2018 – to approximate baseline and endline
208 measures for HealthRise; and being between the ages of 30 and 89 years at "baseline." For Ramsey, comparison
209 patient data were extracted through Minnesota Community Care (formerly named West Side Community Health
210 Services); eligible individuals were patients who had not enrolled in HealthRise and had similar baseline levels of
211 A1c or SBP as HealthRise patients. For Rice, data were extracted from a partner clinic where HealthRise
212 interventions were not implemented. Unlike other comparison patient datasets, International Classification of
213 Disease codes for diabetes and hypertension were not available for patient diagnosis; instead, the diagnosis
214 variable for Rice comparison patients listed active diagnoses. Consequently, a text-matching algorithm was
215 applied to assign diabetes and/or hypertension diagnosis based on the text data in this variable.

216

217 **Qualitative data.** Twenty-three key informant interviews (KIIs) were conducted with local policymakers (non-site
218 specific) and with different types of staff at each site (Table 2). Interviews were not conducted with patients or
219 with staff at clinics from which comparison patient data were selected.

220

221 Initial potential interviewees were identified via leadership from HealthRise grantees and partner organizations,
222 and then additional staff (e.g., clinic-based providers, community paramedics [CPs], CHWs) were contacted via
223 snowball sampling. Of the original individuals identified, 79% completed one-hour interviews via telephone with
224 an IHME evaluation team member.

224

225 All interviews were audio-recorded and listened to multiple times by a single researcher. Key components were
226 transcribed in an Excel template, with thematic coding applied to identify both site-specific and overarching
227 themes across sites.

228 **Endline evaluation analyses**

229 To quantify potential effects of HealthRise participation, we used two outcome indicators to measure patient-
230 level changes from baseline to endline: (1) the proportion of patients meeting treatment targets (i.e., SBP < 140
231 mmHg and DBP < 90 mmHg for hypertension; < 8% A1c for diabetes); and (2) patient biometric measures (i.e.,
232 SBP for hypertension, A1c for diabetes). All analyses were limited to patients who were prevalent cases at
233 baseline and had corresponding biometric data for each time point.

234
235 We conducted difference-in-difference analyses in two steps for each site and by condition. First, we ran an
236 unadjusted model, only including binary variables for HealthRise status and timing (i.e., baseline or endline) and
237 an interaction term for HealthRise at endline to capture the effect of HealthRise participation over time. We
238 then ran an adjusted model, including the following covariates to account for potential systematic differences in
239 HealthRise and comparison patients: sex (female, male); age (< 50 years, ≥ 50 years); time elapsed from baseline
240 to endline (< 12 months, ≥ 12 months); and comorbidities at baseline (prevalent case of only hypertension or
241 diabetes; prevalent case of both hypertension and diabetes). We specified robust standard errors for each
242 model, and used Welch's t-tests (i.e., assuming unequal variance between each group) to evaluate statistically
243 significant differences between HealthRise and comparison patients. All analyses were conducted in Stata
244 version 15 and R version 3.6.2.[18,19]

245

246 **Ethical approval**

247 Ethical approval for this study was obtained from the University of Washington's institutional review board, as
248 well as the local data collection agencies and government entities for each site. All personal identifiers were
249 removed prior to data sharing with IHME; only de-identified data were analyzed.

250

251 **Role of the funding source**

252 Funding for HealthRise came from the Medtronic Foundation. The corresponding author had full access to all
253 the data in the study and had final responsibility for the decision to submit for publication.

254

255

256 **Results**

257 **Quantitative results**


258 Overall, hypertension and diabetes indicators generally improved for HealthRise patients compared with
259 baseline measures (Figure 1, Table 3). At the same time, clinical improvements were heterogeneous since
260 program enrollment (Figure 1), emphasizing potential challenges in effective case management among
261 underserved populations.

262

263 Across sites, a considerable percentage of hypertension patients with baseline SBP measures exceeding 140
264 mmHg improved endline levels to below 140 mmHg (Figure 1A); this trend was particularly pronounced for
265 Hennepin and Ramsey. In Hennepin, 76.8% (95% confidence interval: 65.6-84.4%) of hypertension patients
266 enrolled in HealthRise recorded endline SBP measures below 140 mmHg and 50.0% (32.3-67.3%) of HealthRise
267 patients with hypertension in Ramsey met this threshold. Nonetheless, some percentage of hypertension
268 patients shifted into worse SBP categories by endline: 17.5% in Hennepin, 9.4% in Ramsey, and 13.8% in Rice.


269 **Figure 1. HealthRise patient shifts in disease severity categories between baseline and endline based on biometric readings for hypertension (A) and**
 270 **diabetes (B).** The height of each column reflects 100% of patients at each time point (baseline and endline), while the categories within each column
 271 represents the percentage of patients in each category at baseline and endline. Patient groups are color-coded by their categorization at endline (right column
 272 per site) and flow from their categorization at baseline (left column per site)
 273

A) Hypertension



274

B) Diabetes



275
276
277

278 **Table 3. Baseline and endline measures by HealthRise site and for intervention and comparison patients for hypertension (A) and diabetes (B).** For
 279 HealthRise, samples reflect patients who were prevalent cases of hypertension or diabetes at baseline; remained enrolled throughout the program; and had at
 280 least two biometric readings during program participation.
 281

A) Hypertension

HealthRise program site	Sample size	Percent of patients meeting treatment targets (SBP < 140 and DBP < 90) (%)		Average SBP reading (mmHg)		Average changes in SBP from baseline to endline	
		Baseline	Endline	Baseline	Endline	Absolute change (mmHg)	Percent change (%)
Hennepin County							
HealthRise patients	80	53.8 (42.7 to 64.5)	76.3 (65.6 to 84.4)	136.2 (131.8 to 140.6)	128.5 (125.6 to 131.5)	-7.7 (-12.1 to -3.2)	-4.2 (-7.2 to -1.1)
Comparison patients	83	83.1 (73.3 to 89.8)	78.3 (68.0 to 86.0)	127.3 (124.3 to 130.3)	125.3 (121.9 to 128.7)	-2.0 (-6.0 to 2.0)	-0.7 (-3.9 to 2.5)
Ramsey County							
HealthRise patients	32	12.5 (4.6 to 29.8)	50.0 (32.7 to 67.3)	150.4 (144.9 to 156.0)	136.0 (126.6 to 145.5)	-14.4 (-23.0 to -5.8)	-9.4 (-15.0 to -3.9)
Comparison patients	66	30.3 (20.3 to 42.6)	45.5 (33.7 to 57.7)	146.5 (141.5 to 151.6)	142.9 (137.4 to 148.3)	-3.7 (-10.2 to 2.9)	-0.9 (-5.5 to 3.6)
Rice County							
HealthRise patients	80	42.5 (32.0 to 53.7)	63.8 (52.5 to 73.6)	142.5 (137.5 to 147.6)	134.6 (129.9 to 139.3)	-7.9 (-13.4 to -2.5)	-4.0 (-7.7 to -0.4)
Comparison patients	170	78.2 (71.4 to 83.8)	82.4 (75.8 to 87.4)	128.3 (126.0 to 130.6)	127.3 (124.9 to 129.7)	-1.0 (-3.6 to 1.5)	0.0 (-2.0 to 2.1)

B) Diabetes

HealthRise program site	Sample size	Percent of patients meeting treatment targets (A1c < 8%)		Average A1c reading (%)		Average changes in A1c from baseline to endline	
		Baseline	Endline	Baseline	Endline	Absolute change (%)	Percent change (%)
Hennepin County							
HealthRise patients	37	24.3 (12.9 to 41.1)	24.3 (12.9 to 41.1)	9.9 (9.1 to 10.7)	9.4 (8.6 to 10.3)	-0.5 (-1.4 to 0.5)	-0.8 (-12.6 to 11.1)
Comparison patients	39	71.8 (55.8 to 84.0)	76.9 (60.7 to 87.8)	7.6 (7.1 to 8.2)	7.5 (6.9 to 8.1)	-0.1 (-0.7 to 0.5)	-0.2 (-6.3 to 5.9)
Ramsey County							
HealthRise patients	43	4.7 (1.1 to 17.4)	25.6 (14.5 to 41.0)	10.4 (9.7 to 11.2)	9.0 (8.4 to 9.5)	-1.5 (-2.1 to -0.8)	-11.4 (-16.9 to -5.9)
Comparison patients	72	26.4 (17.4 to 37.9)	29.2 (19.7 to 40.8)	10.0 (9.4 to 10.6)	9.8 (9.2 to 10.4)	-0.2 (-0.8 to 0.5)	2.7 (-4.9 to 10.3)
Rice County							
HealthRise patients	96	43.8 (34.1 to 53.9)	51.0 (41.0 to 61.0)	8.8 (8.3 to 9.2)	8.5 (8.1 to 9.0)	-0.3 (-0.6 to 0.1)	-1.3 (-4.9 to 2.4)
Comparison patients	296	50.3 (44.6 to 56.0)	54.7 (49.0 to 60.3)	8.6 (8.3 to 8.8)	8.3 (8.1 to 8.6)	-0.2 (-0.5 to 0.0)	0.1 (-2.4 to 2.6)

282

283 Sizeable improvements occurred for diabetes patients meeting clinical targets since enrollment (Figure 1B),
284 especially for Ramsey. Compared with baseline, where fewer than 5% of patients with diabetes were meeting
285 treatment targets, 25.6% (14.5-41.0%) of Ramsey HealthRise patients with diabetes had A1c levels lower than
286 8%. Yet many HealthRise patients with diabetes still had A1c levels of 8% or higher by endline across sites: 75.7%
287 in Hennepin, 74.4% in Ramsey, and 49.0% in Rice. A1c category shifts between baseline and endline were
288 especially varied for Hennepin and Rice; for nearly every A1c category at baseline (i.e., < 7%, 7-7.9%, 8-9.9%,
289 $\geq 10\%$), some portion of patients moved to one of the other three A1c categories by endline.

290

291 Unadjusted and adjusted difference-in-difference model results were nearly identical for the effect of HealthRise
292 (Table 4); accordingly, we report on the adjusted model results here. Overall, HealthRise patients trended
293 toward greater progress in reducing biometric measures and meeting treatment targets than comparison
294 patients; however, these differences were not consistently significant across indicators and sites. For
295 hypertension patients, HealthRise participation was associated with statistically significant SBP reductions
296 relative to comparison patients in Rice (6.9 mmHg decrease [0.9-13.0; $p < 0.05$]). Relative to comparison
297 patients, HealthRise was also associated with a statistically significant increase in the percentage of
298 hypertension patients meeting treatment targets in Hennepin (27.3 percentage-point rise [9.7-45.0; $p < 0.01$])
299 and Rice (17.1 percentage-point increase [0.9 to 33.4; $p < 0.05$]). In Ramsey, changes in hypertension indicators
300 were not statistically different between HealthRise and comparison group, though program participation
301 trended toward improvement: a 10.7 mmHg decrease (-0.3-21.8; $p=0.057$) in SBP and 22.3 percentage-point
302 increase (-0.01 to 44.4=5; $p=0.054$) in meeting treatment targets since baseline.

303

304 Among diabetes patients, HealthRise participation was associated with statistically significant reductions in A1c
305 in Ramsey (1.3 decrease in A1c [0.4-3.2; $p < 0.01$) relative to comparison patients. While the percentage of
306 HealthRise patients meeting treatment targets for diabetes did not statistically differ from that of comparison
307 patients in Ramsey, this indicator trended toward improvement as well (a 18.2 percentage-point increase [-0.4-
308 36.7; $p=0.054$]).

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324 **Table 4. Unadjusted and adjusted difference-in-difference regression results, by HealthRise grantee, for hypertension (A)**
 325 **and diabetes (B) patients.** Patient samples included in this analysis are those who met all inclusion criteria, remained
 326 enrolled throughout the program (for HealthRise patients), and had at least two biometric readings to reflect potential
 327 changes between baseline and endline. Bolded values reflect statistically significant estimates at $p < 0.05$.
 328

A) Hypertension

	Hennepin County		Ramsey County		Rice County	
	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value
Change in systolic blood pressure (mmHg)						
Unadjusted model						
HealthRise-endline interaction	-5.6 (-11.7 to 0.4)	0.068	-10.7 (-21.7 to 0.2)	0.054	-6.9 (-12.9 to -0.9)	0.024
Adjusted model						
HealthRise-endline interaction	-5.6 (-11.7 to 0.5)	0.070	-10.7 (-21.8 to 0.3)	0.057	-6.9 (-13.0 to -0.9)	0.025
Age						
< 50 years	-	-	-	-	-	-
≥ 50 years	1.1 (-4.2 to 6.4)	0.677	3.6 (-2.8 to 10.1)	0.266	1.4 (-2.9 to 5.7)	0.531
Sex						
Male	-	-	-	-	-	-
Female	-0.3 (-4.6 to 4.1)	0.909	2.6 (-4.3 to 9.6)	0.455	0.5 (-3.2 to 4.2)	0.780
Duration from baseline to endline						
< 12 months	-	-	-	-	-	-
≥ 12 months	-0.8 (-8.0 to 6.4)	0.829	-2.0 (-11.3 to 7.2)	0.663	-9.1 (-14.1 to -4.2)	< 0.001
Comorbid						
No (hypertension only)	-	-	-	-	-	-
Yes (hypertension and diabetes)	-5.6 (-11.7 to 0.5)	0.266	-7.4 (-15.1 to 0.3)	0.061	8.5 (2.5 to 14.5)	0.006
Change in patients meeting treatment targets (% points)						
Unadjusted model						
HealthRise-endline interaction	27.3 (9.8 to 44.9)	0.002	22.3 (-0.0 to 44.8)	0.051	17.1 (0.9 to 33.3)	0.038
Adjusted model						
HealthRise-endline interaction	27.3 (9.7 to 45.0)	0.003	22.3 (-0.0 to 44.5)	0.054	17.1 (0.9 to 33.4)	0.039
Age						
< 50 years	-	-	-	-	-	-
≥ 50 years	2.9 (-12.7 to 18.6)	0.714	-2.4 (-18.8 to 14.1)	0.777	7.8 (-1.6 to 17.2)	0.102
Sex						
Male	-	-	-	-	-	-
Female	0.1 (-9.0 to 11.0)	0.844	-6.5 (-21.4 to 8.4)	0.389	-1.6 (-9.8 to 6.5)	0.697
Duration from baseline to endline						
< 12 months	-	-	-	-	-	-
≥ 12 months	6.0 (-12.1 to 24.1)	0.511	5.2 (11.9 to 22.3)	0.546	21.2 (11.1 to 31.2)	< 0.001
Comorbid						
No (hypertension only)	-	-	-	-	-	-
Yes (hypertension and diabetes)	11.0 (-0.4 to 22.4)	0.058	10.3 (-6.9 to 27.4)	0.238	-8.2 (-21.8 to 5.4)	0.236

329
 330
 331

B) Diabetes

	Hennepin County		Ramsey County		Rice County	
	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value
Change in A1c (%)						
Unadjusted model						
HealthRise-endline interaction	-0.4 (-1.5 to 0.8)	0.551	-1.3 (-2.2 to -0.4)	0.007	-0.03 (-0.5 to 0.4)	0.904
Adjusted model						
HealthRise-endline interaction	-0.4 (-1.5 to 0.8)	0.556	-1.3 (-2.3 to -0.4)	0.007	-0.03 (-0.5 to 0.4)	0.905
Age						
< 50 years	-	-	-	-	-	-
≥ 50 years	-1.8 (-3.0 to -0.6)	0.006	0.1 (-0.8 to 1.0)	0.828	-0.1 (-0.6 to 0.4)	0.369
Sex						
Male	-	-	-	-	-	-
Female	0.1 (-0.7 to 0.9)	0.833	-0.1 (-0.9 to 0.7)	0.791	-0.1 (-0.5 to 0.3)	0.622
Duration from baseline to endline						
< 12 months	-	-	-	-	-	-
≥ 12 months	-1.5 (-2.8 to -0.1)	0.031	0.2 (-0.8 to 1.1)	0.807	0.2 (-0.2 to -0.6)	0.369
Comorbid						
No (diabetes only)	-	-	-	-	-	-
Yes (hypertension and diabetes)	-0.5 (-1.7 to 0.7)	0.417	-0.5 (-1.3 to 0.4)	0.260	0.1 (-0.3 to -0.6)	0.510
Change in patients meeting treatment targets (% points)						
Unadjusted model						
HealthRise-endline interaction	-5.1 (-30.2 to 19.9)	0.684	18.2 (-0.2 to 36.5)	0.052	3.0 (-8.3 to 14.1)	0.612
Adjusted model						
HealthRise-endline interaction	-5.1 (-30.5 to 20.2)	0.688	18.2 (-0.4 to 36.7)	0.055	3.0 (-8.4 to 14.2)	0.613
Age						
< 50 years	-	-	-	-	-	-
≥ 50 years	9.0 (-10.1 to 28.2)	0.351	4.5 (-8.6 to 17.6)	0.498	-2.6 (-12.5 to 7.2)	0.596
Sex						
Male	-	-	-	-	-	-
Female	-7.8 (-24.0 to 8.3)	0.336	-2.1 (-14.5 to 10.3)	0.739	0.8 (-8.2 to 9.8)	0.867
Duration from baseline to endline						
< 12 months	-	-	-	-	-	-
≥ 12 months	13.8 (-11.2 to 38.8)	0.274	-0.3 (-14.8 to 14.1)	0.963	-2.7 (-12.4 to 7.0)	0.582
Comorbid						
No (diabetes only)	-	-	-	-	-	-
Yes (hypertension and diabetes)	0.9 (-19.6 to 21.5)	0.929	1.9 (-10.5 to 14.3)	0.760	2.9 (-8.4 to 14.2)	0.648

332

333 **Qualitative findings**

334 Across HealthRise sites, six main themes emerged for the qualitative data synthesis (Table 5). First, respondents
335 viewed home-based providers as critical to bridging barriers experienced by patients (e.g., linguistic and cultural
336 divides), and clinic-based providers indicated high value in meeting patients beyond clinical settings.
337 Coordination of care and a focus on social determinants of health, such as access to healthier food and nutrition,
338 were highlighted as key program features. Second, program strengths involved learning from HealthRise sites in
339 other countries and enabling many clinical staff to work with in-home providers for the first time. Clinical
340 providers reported improved quality and efficiency in clinical appointments due to having additional details
341 about patient needs from in-home providers. Home visits also enabled providers to connect patients with non-
342 clinical resources (e.g., housing) to support improved outcomes. The theme of perceived program impacts
343 extended program strengths, with providers reporting positive changes in the health and lives of patients and
344 their families. Further, several interviewees emphasized the synergistic effects of pairing CHWs and community
345 paramedics (CPs) within care teams, and reported efforts to adopt home-based provider models by other local
346 organizations because of HealthRise experiences.

347
348 Common challenges emerged across sites, often relating to new program establishment and incorporation of in-
349 home providers within care teams. For example, some clinical providers showed initial skepticism about the
350 added value of in-home providers, and most administrators did not have prior experience managing CHWs and
351 CPs. Site-specific challenges also occurred; for instance, patient consent for enrollment took a long time during
352 the initial phase of program implementation in Rice, while CHW turnover was an ongoing obstacle for both
353 Hennepin and Ramsey. Data sharing was another pervasive challenge, mainly from poor interoperability and
354 coordination between clinic data systems and electronic medical record (EMR) systems. Providers also
355 expressed frustration with expectations for rapidly improving clinical outcomes, especially given the
356 longstanding challenges in healthcare access and social determinants of health most patients faced.

357
358 The fifth theme pertained to recommendations for improvement, many of which stemmed from acknowledged
359 challenges. Such suggestions included prioritizing better communication and coordination among care teams as
360 well as EMR systems that could more seamlessly accommodate patient updates and provider notes from
361 multiple care-team members. Interviewees reported having inadequate clarity on the ideal frequency and length
362 of home visits, an area where efficiencies in resource deployment could be improved. The sixth theme, future
363 program directions, involved many ideas about adapting and expanding HealthRise programming to new
364 locations and conditions (e.g., mental health). Another common thread concerned program sustainability,
365 namely longer-term financing and retaining home-based providers for the HealthRise model.

Table 5. Summary of key themes, components, and quotes from qualitative data synthesized across US HealthRise sites.

HealthRise thematic area components and contexts	Sample thematic quotes
Theme 1: Key program features	
<ul style="list-style-type: none"> • Home-based providers as the cornerstone to HealthRise model, bridging linguistic and cultural divides and gaining valuable new information on home visits 	<p>“...through care coordination, communication, and use of frontline health workers...in ways we’ve never been able to before, connect[ing] with families and follow[ing] up with specific patients to help them really understand and manage their chronic disease.” – <i>Clinic-based provider</i></p>
<ul style="list-style-type: none"> • Emphasis on care coordination and extending care outside the clinic to address social determinants of health 	<p>“There are so many hard things about managing diabetes...it takes so much time for any patient to fully understand how to put the different parts of diabetes treatment together...giving people the time they need to really understand all the components of diabetes control...that’s where our CHWs have really been massive assets.” – <i>Clinic-based provider</i></p> <p>“If 80% of health happens outside the clinical setting, what are the ways we can foster healthy environments that allow individuals more capacity and agency to focus on these chronic diseases?” – <i>Administrator</i></p>
Theme 2: Program strengths	
<ul style="list-style-type: none"> • Global learning from other HealthRise sites to inform intervention design 	<p>“The global aspect is quite unique...utilizing similar strategies in different countries with very different health systems but with a similar population focus and similar workforce approaches.... I’m not aware of other projects that have attempted that across a set of different jurisdictions and landscapes.” – <i>Policymaker</i></p>
<ul style="list-style-type: none"> • Introduction to the value of home-based providers for many clinic staff 	
<ul style="list-style-type: none"> • Strong relationships built between different types of providers over course of program 	<p>“The home visits contributed to more rational use of clinic time...and improved care on my end. From listening to CHWs, I have a better understanding of what’s going on in people’s lives.” – <i>Clinic-based provider</i></p>
<ul style="list-style-type: none"> • Patients’ receipt of extra support beyond what was typically possible in limited time of clinical appointment 	
Theme 3: Program challenges	
<ul style="list-style-type: none"> • Clinical providers lacked familiarity with home-based providers 	<p>“We’ve learned that a lot of the hurdle we have to get past is educating other health care providers on what we do...what is a CP and how can we be part of their team and help to better serve their patients...the ones who do now understand our role...they are our champions, they get so excited...we definitely see resistance in the beginning.” – <i>CP</i></p>
<ul style="list-style-type: none"> • Facility administrators lacked experience managing home-based providers 	
<ul style="list-style-type: none"> • High turnover of staff, both community health workers and management 	<p>“It was too short of a time...it took forever to get these communities up and running, get people hired...people quit, etc...need a longer lifespan than three years...to show enough impact to indicate policy change.” - <i>Policymaker</i></p>
<ul style="list-style-type: none"> • Expectations to improve clinical outcomes in short period of time 	
<ul style="list-style-type: none"> • Times lags between patient recruitment/consent and actual program enrollment 	<p>“One of the largest hurdles and barrier to successful implementation was the lack of cohesive patient data systems...you need layers of permission, use agreements, consent, you can’t compare across systems, you can’t look at anybody’s system but your own...if somebody could fix that, we could do a lot more good.” - <i>Administrator</i></p>
<ul style="list-style-type: none"> • Patients’ barriers to accessing healthcare (e.g., social determinants of health), as well as provider barriers to effectively accessing patients (e.g., language) 	
<ul style="list-style-type: none"> • Interoperability and data sharing between health systems, across platforms, and among care team (e.g., electronic medical record access for all team members) 	

HealthRise thematic area components and contexts (<i>continued</i>)	Sample thematic quotes
Theme 4: Perceived program impacts	
<ul style="list-style-type: none"> • Information gained during home visits improved the quality and efficiency of clinical appointments • Pairing of CP and CHW brought together complementary skill sets • Home visits helped connect patients with non-clinical resources to improve health • Positive changes in health and lives of patients and their families • Efforts to institutionalize home-based providers as part of care model 	<p>“There’s just a synergy when you combine the two...the CP and the medical side and the CHW looking at the social issues...it was kind of exponential how much benefit we were able to provide versus just one or the other.” - CP</p> <p>“We have nutritionists and a diabetic educator in the clinic, but patients for a variety of reasons are not always open to meeting with them....to have somebody go to their house and figure out what their specific food interests are and come up with recipes, that was great.” – <i>Clinic-based provider</i></p>
Theme 5: Recommendations for improvement	
<ul style="list-style-type: none"> • Tailor EMR software for care teams that include home-based providers • Determine ideal duration and/or frequency of home visits • Offer additional mental health resources to support patients with chronic disease • Provide more trainings for home-based providers (e.g. motivational interviewing) 	<p>“EMRs are designed around providers and reimbursements, our model is around holistic care coordination across contexts. The tools we have been using are imperfect, we’re looking for other tools that might be able to plug in to our model better...We haven’t yet found the silver bullet.” – <i>Administrator</i></p> <p>“We’re aggressively moving into providing more mental health care, plugging it into our model that we’ve perfected during HealthRise...Behavioral health, mental health, and chemical dependency are NCDs, similar to diabetes in that it’s all about what happens in the meantime.” – <i>Administrator</i></p>
Theme 6: Future program directions	
<ul style="list-style-type: none"> • Expand use of home-based providers to other patient populations, particularly for other chronic conditions • Generate additional evidence establishing the cost-effectiveness of home-based care for chronic conditions • Identify sustainable funding for home-based providers 	<p>“I think it should be implemented everywhere...that’s the response we’re getting from our partners, from others in the health care setting...everyone is saying ‘CHW! I need 7 of you in my facility!’...everyone has a million questions about how to get it started, set up...I definitely think it’s a model to follow.” - CHW</p>

Discussion

Increasingly more evidence shows that community-based programs can help underserved communities in the US better access health services, alleviate barriers to care, and improve at least some health behaviors and outcomes. The present study contributes to this evidence base through its prospective evaluation of HealthRise programs in three Minnesota communities. Relative to comparison patients, HealthRise participation was significantly associated with SBP reductions in Rice, while the percentage of hypertension patients meeting treatment targets increased in Hennepin and Rice. For diabetes, HealthRise patients saw larger A1c declines in Ramsey than comparison patients. Heterogeneous patterns in patient improvements since baseline highlight potential case management challenges among underserved individuals and communities, especially under short program implementation periods. As emphasized by HealthRise care teams, community-based programs show promise for improving NCD care and outcomes for underserved populations; nonetheless, more work is needed to better understand how such programs can be further brought to scale and sustained long-term.

HealthRise participation was related to SBP or A1c decreases and a higher percentage of patients meeting treatment targets at some sites relative to comparison patients. Variations across sites may be related to different demographic characteristics or relative health status of each population at baseline. For instance, Ramsey HealthRise patients with diabetes averaged 11.4% reductions in A1c and 9.4% declines in SBP by endline; yet Ramsey patients also began HealthRise with highest risk profiles, averaging 150 mmHg SBP and 10% A1c at baseline. As found in past studies,[5–7] several factors may have contributed to the observable effects of HealthRise. These included focusing on specific barriers patients faced in each community (e.g., home-based care provided by CHW and CP teams); maintaining small patient loads, enabling more individualized attention and tailoring of visit frequency to patient need; and explicitly providing non-medical support, such as health education and community resources like transportation. Further, the overall positive views of HealthRise by grantees and care teams alike may have contributed to the program's effects. Despite challenges during earlier stages of program implementation (e.g., recruiting and retaining CHWs, ensuring adequate access to EMRs), providers voiced valuing home-based health workers and were eager to expand this model of care. In combination, these factors may have set the foundation for HealthRise's impact for underserved patients with hypertension and diabetes in the US.

Amid such promising findings, however, important challenges remained for each site and for broader applications of the HealthRise model elsewhere. Across sites, some proportion of HealthRise patients failed to meet clinical targets for hypertension or diabetes at both time points – and concerningly, some percentage moved from being below biometric thresholds at baseline to exceeding them at endline. These patterns underscore the complexity of effectively managing chronic conditions like hypertension and diabetes, especially in environments where patients face compounding obstacles to medical care and health-promoting behavior (e.g., limited budgets for nutritious food, minimal time for exercise amid job and family demands, inferior access to adequate transportation and housing). CHWs and integrated care teams may be able to mitigate some of these barriers and better support patients' medical needs, a critical step in addressing deep-seeded health disparities; however, in the absence of more macro-level

socioeconomic policies and health system investments to support underserved patients in the US, many community-based programs will continue facing need and demand that far exceeds their limited capacities and resources. As laid bare by COVID-19, the health challenges underserved communities experience do not begin and end at clinics: rather, they stem from and are exacerbated by structural inequalities that require intervention and engagement well beyond the formal health system.[20,21] CHWs can provide a vital role in at least overcoming some of these access and sociocultural barriers, ranging from house-based visits made by CHWs fluent in patients' native languages[3] to connecting patients with services that can facilitate better overall care. Nevertheless, the potential benefits of encouraging health programs can be easily blunted if actors – and actions – beyond the immediate health system are not also actively addressing fundamental drivers of health disparities.

Limitations

This study is subject to several limitations. First, small study samples at each site likely affected the degree to which potential program impact could be detected and conclusively attributed to HealthRise participation. This is particularly true for site-condition combinations in which very few patients were prevalent cases at baseline and had at least two biometric readings within the program implementation period (e.g., 37 HealthRise patients with diabetes in Hennepin, 32 HealthRise patients with hypertension in Ramsey). The inclusion or exclusion of even a few patients for several site-condition groupings could shift effect sizes and statistical significance estimated by the difference-in-difference models.

Second, comparison groups were constructed retrospectively based on available patient record information and were not selected by random assignment. While efforts were made to ensure that comparison patient data were chosen to generally represent individuals who would have been eligible for HealthRise enrollment, they may have differed from individuals who enrolled.

Third, only patients who remained enrolled at endline were included in the present study; by taking this 'as treated' analytic approach, which provides insights into program effects closer to full adherence, these patients may not represent all potential target populations for HealthRise interventions and results may be positively biased. For instance, relatively high rates of program withdrawal at some sites could have led to a bias toward healthier patients remaining in the program (i.e., sicker patients may not go into the clinic). However, due to the home-based care model espoused by HealthRise, it is equally possible that patients who did not withdraw were less healthy and stayed enrolled because HealthRise offered important access to services, like home visits, otherwise unavailable to them.

Fourth, single biometric readings comprised baseline and endline measures, as well as patients meeting clinical targets at each time period; subsequently, analyses could be sensitive to outliers in patient records, particularly given the relatively small sample sizes for each site. If more readings could have informed baseline and endline indicators, it is possible patient-level patterns could have differed from what observed on the basis of single readings.

Fifth, medication data were not available and thus we were unable to assess the full cascade of care for hypertension and diabetes case management. Medication adherence may have been important factor

for patients who either failed to see improvements in clinical indicators or experienced worsening outcomes.

Sixth, due to the small sample sizes for each site, we could not further analyze the potential effects of visit frequency (and thus approximate dose-response) or intensity of intervention exposure on patient outcomes. This is further complicated by issues related to endogeneity, such that patients with worse clinical profiles and thus greater need are likely to receive more frequent visits by CHWs or care teams.

Seventh, small sample sizes also limited our ability to conduct more disaggregated analyses by sex, race or ethnicity, income, and other important characteristics for understanding how community-based health interventions can promote greater equity for underserved communities.

Conclusions

With its focus on community-based health programs and improving NCD care for underserved populations, HealthRise showed some positive effects for hypertension and diabetes patients in Hennepin, Ramsey, and Rice counties. Provider experiences indicated enthusiasm for expanding the home-based model of care for NCDs, though the resource requirements – as well feasibility – to sustain impact at a larger scale remain unknown. While community-based NCD interventions show promise for overcoming barriers to effective care for hypertension and diabetes among underserved populations, continued monitoring and robust evaluations of local impact are vital to ensuring maximum benefit for individuals with the greatest need.

Acknowledgments

We would like to acknowledge the support and insights from the US HealthRise program implementers at Pillsbury United Communities, Regions Hospital Foundation, and HealthFinders Collaborative Inc., of whom also provided the de-identified data for this study. Without them, this work would not be possible. We also acknowledge the state and local governments in HealthRise areas for their cooperation and support with HealthRise intervention and evaluation activities. We thank Kelsey Bannon, Erin Palmisano, and Laura Di Giorgio for assisting in the management and execution of the HealthRise evaluation. Last, we thank all individuals who participated in this study.

References

1. Mokdad AH, Ballestros K, Echko M, Glenn S, Olsen HE, Mullany E, et al. The State of US Health, 1990–2016: Burden of Diseases, Injuries, and Risk Factors Among US States. *JAMA*. 2018;319:1444–1472. doi:10.1001/jama.2018.0158
2. National Academies of Sciences E, Division H and M, Practice B on PH and PH, States C on C-BS to PHE in the U, Baciu A, Negussie Y, et al. The State of Health Disparities in the United States.

- Communities in Action: Pathways to Health Equity. National Academies Press (US); 2017. Available: <https://www.ncbi.nlm.nih.gov/books/NBK425844/>
3. Witmer A, Seifer SD, Finocchio L, Leslie J, O’Neil EH. Community health workers: integral members of the health care work force. *Am J Public Health*. 1995;85: 1055–1058. doi:10.2105/AJPH.85.8_Pt_1.1055
 4. Rosenthal EL, Brownstein JN, Rush CH, Hirsch GR, Willaert AM, Scott JR, et al. Community Health Workers: Part Of The Solution. *Health Affairs*. 2010;29: 1338–1342. doi:10.1377/hlthaff.2010.0081
 5. Norris SL, Chowdhury FM, Le KV, Horsley T, Brownstein JN, Zhang X, et al. Effectiveness of community health workers in the care of persons with diabetes. *Diabetic Medicine*. 2006;23: 544–556. doi:10.1111/j.1464-5491.2006.01845.x
 6. Brownstein JN, Chowdhury FM, Norris SL, Horsley T, Jack L, Zhang X, et al. Effectiveness of Community Health Workers in the Care of People with Hypertension. *American Journal of Preventive Medicine*. 2007;32: 435–447. doi:10.1016/j.amepre.2007.01.011
 7. Kim K, Choi JS, Choi E, Nieman CL, Joo JH, Lin FR, et al. Effects of Community-Based Health Worker Interventions to Improve Chronic Disease Management and Care Among Vulnerable Populations: A Systematic Review. *Am J Public Health*. 2016;106: e3–e28. doi:10.2105/AJPH.2015.302987
 8. Kangovi S, Mitra N, Grande D, Huo H, Smith RA, Long JA. Community Health Worker Support for Disadvantaged Patients With Multiple Chronic Diseases: A Randomized Clinical Trial. *Am J Public Health*. 2017;107: 1660–1667. doi:10.2105/AJPH.2017.303985
 9. Kangovi S, Mitra N, Norton L, Harte R, Zhao X, Carter T, et al. Effect of Community Health Worker Support on Clinical Outcomes of Low-Income Patients Across Primary Care Facilities: A Randomized Clinical Trial. *JAMA Intern Med*. 2018;178: 1635–1643. doi:10.1001/jamainternmed.2018.4630
 10. Gary TL, Bone LR, Hill MN, Levine DM, McGuire M, Saudek C, et al. Randomized controlled trial of the effects of nurse case manager and community health worker interventions on risk factors for diabetes-related complications in urban African Americans. *Prev Med*. 2003;37: 23–32. doi:10.1016/s0091-7435(03)00040-9
 11. Brown SA, Garcia AA, Kouzekanani K, Hanis CL. Culturally competent diabetes self-management education for Mexican Americans: the Starr County border health initiative. *Diabetes Care*. 2002;25: 259–268. doi:10.2337/diacare.25.2.259
 12. Flor LS, Wilson S, Bhatt P, Bryant M, Burnett A, Camarda JN, et al. Community-based interventions for detection and management of diabetes and hypertension in underserved communities: a mixed-methods evaluation in Brazil, India, South Africa and the USA. *BMJ Global Health*. 2020;5: e001959. doi:10.1136/bmjgh-2019-001959
 13. Institute for Health Metrics and Evaluation (IHME). HealthRise Evaluation Final Report. Seattle, WA, USA: IHME; Forthcoming.

14. Abt Associates. HealthRise Final Report: Expanding Access to Chronic Disease Care through Community Approaches in Four Countries. Forthcoming.
15. Roth G, Gabert R, Thomson B, Gakidou E. HealthRise Preliminary Needs Assessment Report US Sites: Hennepin, Ramsey, Rice Counties. Seattle, WA: Institute for Health Metrics and Evaluation; 2015.
16. MN Community Measurement. 2017 Health Equity of Care Report: Stratification of Health Care Performance Results in Minnesota by Race, Hispanic Ethnicity, Preferred Language, & Country of Origin. Minneapolis, MN: MN Community Measurement; 2018. Available: https://mncm.org/wp-content/uploads/2018/01/2017-Health-Equity-of-Care-Report_unencrypted-1.pdf
17. Gabert R, Thomson B, Gakidou E, Roth G. Identifying High-Risk Neighborhoods Using Electronic Medical Records: A Population-Based Approach for Targeting Diabetes Prevention and Treatment Interventions. PLOS ONE. 2016;11: e0159227. doi:10.1371/journal.pone.0159227
18. StataCorp. Stata Statistical Software: Release 15. College Station, TX: StataCorp LLC; 2017.
19. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2018. Available: <https://www.R-project.org>
20. Haynes Norrisa, Cooper Lisa A., Albert Michelle A. At the Heart of the Matter: Unmasking and Addressing COVID-19's Toll on Diverse Populations. Circulation. 0. doi:10.1161/CIRCULATIONAHA.120.048126
21. Dorn A van, Cooney RE, Sabin ML. COVID-19 exacerbating inequalities in the US. Lancet. 2020;395: 1243–1244. doi:10.1016/S0140-6736(20)30893-X