- 1 Title
- 2 Morphological alternations of intracerebral arterial during middle age: a community
- 3 cohort study
- 4 Authors
- 5 Boyu Zhang^{1,2}, Zidong Yang^{1,2}, Jing Li³, Bei Wang^{1,2}, Huazheng Shi⁵, He Wang^{1,2,4*},
- 6 Yuehua Li^{3*}
- ⁷ ¹Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University,
- 8 Shanghai, China
- 9 ²Key Laboratory of Computational Neuroscience and Brain-Inspired Intelligence
- 10 (Fudan University), Ministry of Education, China
- ¹¹ ³Department of Radiology, Shanghai Jiao Tong University Affiliated Sixth People's
- 12 Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China
- ¹³ ⁴Human Phenome Institute, Fudan University, Shanghai, China
- 14 ⁵Shanghai Universal Medical Imaging Diagnostic Center, Shanghai, China
- 15 **Correspondence to:**
- 16 Prof He Wang,
- 17 Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University,
- 18 Shanghai 200433, China
- 19 Email: hewang@fudan.edu.cn
- 20 Prof Yuehua Li,
- 21 Department of Radiology, Shanghai Jiao Tong University Affiliated Sixth People's
- 22 Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200233, China
- 23 Email: liyuehua77@sjtu.edu.cn
- 24

25 Morphological alternations of intracerebral arterial during middle age: a community cohort study 26

Abstract 27

28 Background

Ample evidence has suggested that vascular modifications are associated with aging. 29

30 To expand previous understanding of age-related vascular changes, we examined the

association between aging and cerebrovascular morphologies. 31

32 Methods

1176 participants aged 35 to 75 years recruited from Shanghai, China were included in 33 this study. Cerebrovascular morphological features comprising arterial branch density, 34 radius, and tortuosity were quantified using three-dimensional magnetic resonance 35 angiography. Linear regression was used to examine the association between age and 36 vasculature features. 37

Results 38

Age was found to be a significant predictor for cerebrovascular morphological 39 40 alterations after adjusting for vascular risk factors. However, after dividing subjects into subgroups based on their age, aging was found to be significantly correlated with all 41 three morphometric features only in the 45-54 subgroup after adjusting for the other 42 vascular risk factors. Smoking gives rise to a more rapid age-related changes in vascular 43 morphologies, while alcohol consumption could decelerate those age-related alterations. 44 Conclusions 45

Rapid alternations in all three morphological features assessed have been noticed to be 46

47 associated with aging in the 45-54 subgroup, suggesting the potential importance of the

5th decade for early preservation method of vascular aging. 48

49 Funding

National Natural Science Foundation of China. 50

51

Introduction 52

The power-hungry nature of the human brain requires a consistent supply of oxygen 53 and nutrients, which is dependent on the structural and functional integrity of the 54

intracranial vasculature, to maintain its normal functioning (Zhang and Raichle 2010). 55 Indeed, the cerebral vasculature has been implicated in many dementing diseases, such 56 57 as vascular dementia and Alzheimer's disease (AD) (Leeuwis et al. 2017, Alber et al. 2019). Several previous studies have suggested a synergistic relationship between 58 cerebrovascular dysfunctions and the neurological symptoms (Silvestrini et al. 2006, 59 60 Kalaria 2002). For instance, reduced cerebral blood flow (CBF) has been implicated in cognitive decline (Swan et al. 1998, Leeuwis et al. 2017). Exposure to cerebrovascular 61 62 risk (CVR) could cause cerebral vascular system impairments. In particular, elevated blood pressure is associated with the remodeling of both small and large arteries, which 63 paves the way for the impairment of the brain functions (Laurent and Boutouyrie 2015). 64 Among the common CVR factors, aging has been recognized to pose a significant risk 65 for the development of cardiovascular diseases, and its effect on the cerebral vascular 66 67 system deserves more attention (D'Agostino et al. 2008).

Association between advanced age and a general reduction in CBF in normal subjects 68 has been reported earlier (Melamed et al. 1980). During the past few years, clinical 69 70 studies have revealed that major vascular modifications, including generalized 71 endothelial dysfunction and central arterial stiffness, in the cerebral arteriolar system also transpire with aging (Xu et al. 2017, Donato, Machin and Lesniewski 2018). With 72 the advancements in time-of-flight magnetic resonance angiography (TOF MRA), age-73 related intracranial arterial morphological alterations, including reduced vessel branch 74 number and increased mean tortuosity have been reported(Chen et al. 2019, Bullitt et 75 al. 2010). Such alterations have recently been implicated in cerebral small vessel 76 77 disease (CSVD), a brain disease that is believed to be associated with a wide range of 78 cognitive disorders including nearly a half of dementias, making those promising 79 markers for brain damage and cognitive decline (Banerjee et al. 2016, Zhang et al. 2021). 80

81 Moreover, extensive research has focused on midlife since exposure to CVR factors 82 during midlife was reported to influence cognitive outcomes later on significantly (Walker et al. 2019, Kloppenborg et al. 2008, Swan et al. 1998). Namely, midlife 83 exposure to CVRs, including obesity, hypertension, and smoking, has been reported to 84

85 markedly accelerate the deterioration of the executive functions and was believed to be a better predictor than exposure in later life (Tolppanen et al. 2012, Debette et al. 2011). 86 87 Furthermore, a subtle decline in cognitive functions including cognitive reaction time and episodic memory, has been identified in middle-aged healthy subjects (younger 88 than 50 years old), adding to the importance of midlife for the prevention of the 89 pathological aspects of aging (Zimprich and Mascherek 2010, Ferreira et al. 2015). 90 Findings as such have raised the question as to whether the observed declines in 91 92 cognitive or executive abilities were mediated by the structural or functional alterations associated with aging or the exposure to other CVR factors in the cerebral vasculature 93 during midlife. However, for the elucidation of mechanisms underlying those 94 neurological disorders and the preservation of brain functions, a better and more 95 comprehensive understanding of the structural modifications of the intracranial 96 vasculature with aging is required. Nevertheless, previous studies investigating the 97 effect of aging or general CVR factors on the intracranial vessel morphologies usually 98 suffered from a relatively smaller sample size with a restricted age range, minimizing 99 100 their power to detect earlier changes (Bullitt et al. 2010, Chen et al. 2019).

Therefore, in this study, we set out to address two questions. First, how does the 101 characteristics of cerebral vasculature precipitated by MRA alter with aging. Thanks to 102 improved MRA resolution and automatic vascular quantification, in vivo studies 103 104 investigating the age-related morphological changes of the cerebral vasculature are possible. Second, whether aging affects the cerebral vascular structures differently in 105 106 different age groups, especially in the middle-aged group (45-54 years), given changes in the cerebrovascular system during this period seemed to have marked predicting 107 108 power for the cognitive outcomes in later life. In the current study, a large natural population in China aged from 35 to 75 was recruited. A more detailed division of 109 subjects into subgroups based on age has been made to study the effect of aging as well 110 as the conventional CVR factors on vascular morphologies in different age groups. 111

112

Results 113

The participant characteristics are presented in Table 1. Figure 1 summarizes the 114 recruitment in the current study. A total of 1240 participants were included; of these, 115 116 33 participants with poor image quality that is not adequate for vessel segmentation and 31 participants with incomplete cerebrovascular risk information were excluded. 117 Therefore, 1176 participants were included in the subsequent analysis. 118

Association between cerebrovascular morphology and aging 119

We first investigated the effect of age on the morphometric features of the cerebral 120 vasculature in general. Age was found to be significantly associated with decreased 121 vessel density (β =-5.7×10⁻⁴, 95% CI -0.001 to -4.3×10⁻⁴, p<0.001), increased vessel 122 radius, (B=0.002, 95% CI 0.002 to 0.002, p<0.001), and increased vessel tortuosity, 123 $(\beta = 8.5 \times 10^{-4}, 95\% \text{ CI } 6.7 \times 10^{-4} \text{ to } 0.001, \text{ p} < 0.001)$ after adjusting for CVR factors, as 124 shown in Table 2. The participants were further divided into four groups based on their 125 age to further examine whether the observed effect of aging differed in different age 126 groups (Figure 2a-c). Surprisingly, vessel density was only found to be negatively 127 associated with aging in participants aged 45-54, (β =-0.002, 95% CI -0.002 to -0.001, 128 p<0.001), and 65-75, (β =-0.001, 95% CI -0.002 to -7.6×10⁻⁵, p=0.036), and only the 129 association found in people aged 45-54 remained significant after adjusting for CVR 130 factors, (β =-0.001, 95% CI -0.002 to -0.001, p=0.001), suggesting the effects of CVR 131 factors are not negligible for people in the 65-75 age group, whereas the effect of aging 132 dominates the changes of vascular features in the 45-54 age group. Similarly, increased 133 vessel radius was only found to be associated with aging in participants aged 45-54, 134 (β=0.005, 95% CI 0.002 to 0.008, p<0.001), and 65-75, (β=0.003, 95% CI 2.9×10⁻⁴ to 135 0.006, p=0.03), and, after adjusting for CVR factors, vessel radius was found to be 136 137 significantly associated with advanced age only in participants aged 45-54, (β =0.005, 95% CI 0.003 to 0.008, p<0.001). Increased vessel tortuosity was found to be 138 significantly associated with age in subjects aged 35-44 (β =0.001, 95% CI 0.001 to 139 0.003, p=0.03), and 45-54, (β =0.002, 95% CI 0.001 to 0.003, p=0.003). The 140 associations remained significant after adjusting for CVR factors for participants aged 141 35-44. (β =0.002, 95% CI 3.4×10⁻⁴ to 0.003, p=0.01), and for participants aged 45-54, 142 $(\beta=0.001, 95\% \text{ CI } 4.4 \times 10^{-4} \text{ to } 0.003, p=0.006).$ 143

To better delineate the underlying mechanisms for dilated arteries, we further assessed 144 the proportion of the smaller vessels visualized by MRA. As shown in Table 3, a general 145 decrease in the proportion of the smaller vessels with advancing age was identified, (β = 146 -3.7×10^{-4} , 95% CI: -6.1×10^{-4} , -1.3×10^{-4} , p=0.03). The decrease in mean vessel radius 147 reported before, however, was found not to be due to the decreased proportion of the 148 smaller vessels, since the association between aging and vessel radius was not much 149 affected after adjusting for the proportion of the smaller vessels, ($\beta = 2.7 \times 10^{-4}$, 95% CI: 150 6.6×10^{-5} , 4.8×10^{-4} , p=0.01), hence suggesting compensatory vasodilatation might be 151 the cause here. 152

153 Association between cerebrovascular morphologies and CVR factors

The relationships between cerebrovascular morphologies and CVR factors were further 154 examined. Vessel radius and mean tortuosity were found not to be associated with any 155 of the individual risk factors listed in Table 4. A decrease in vessel density was found 156 in participants with diabetes after adjusting for age and other risk factors (β =-0.005, 157 95% CI -0.01 to -0.001, p<0.046). Interestingly, the interaction between age and 158 smoking was found to significantly impact vessel density (β = -6.7×10⁻⁴, 95% CI: -159 8.8×10^{-4} , 4.6×10^{-4} , p=0.002) and vessel radius (β = 0.002, 95% CI: 9.6 \times 10^{-4}, 0.002, 160 p=0.01). However, the effect of the interaction between age and alcohol use seems to 161 counteract the effect of aging alone on vascular density and vessel radius. While 162 decreased vascular density and increased vessel radius were found in older adults in 163 general, the interaction of age and alcohol consumption seems to result in a higher 164 vascular density (β = 5.8×10⁻⁴, 95% CI: 8.6×10⁻⁴, 3.1×10⁻⁴, p=0.03) and a lower vessel 165 radius (β = -0.002, 95% CI: -0.003, -8.6×10⁻⁴, p=0.04). In general, smoking was found 166 167 to facilitate the deterioration of the vascular structures, while alcohol use seemed to decelerate the changes (Figure 2d-i). No significant interaction between age and other 168 CVR factors was found. 169

170 Association between cerebrovascular morphologies and CVR score

171 A cumulative CVR score of 0 to 6 was assigned accordingly, with a higher score indicating exposure

172 to fewer CVR factors. The associations between CVR score and vascular morphologies have been

173 reported in Table 5. Overall, a significant association was identified between CVR score and vessel

174 radius after adjusting for age and sex (β = -0.004 95% CI: -0.008, -2.0×10⁻⁴, p=0.039). However,

175 when participants were subdivided into age groups, CVR score was only found to be correlated with

an increase in vascular density adjusted for age and sex (β = 0.002, 95% CI: 1.0×10⁻⁴, 0.004, p=0.041)

- 177 in the 55-64 group, and a decrease in vessel radius after adjusting for age and sex (β = -0.010, 95%
- 178 CI: -0.010, -1.6×10⁻⁴, p=0.048) in participants aged 35-44.
- 179

180 **Discussion**

In this cross-sectional study, the effect of aging and other conventionally defined CVR 181 factors on the intracerebral arterial morphologies acquired by 3D TOF MRA were 182 investigated. As per previous studies, advanced age was found in our research to be 183 associated with increased vessel radius, increased vessel tortuosity and decreased 184 branch density (Bullitt et al. 2010, Chen et al. 2019). Thanks to the relatively large 185 sample size of this study, assessments of the age- or CVR-related morphological 186 alternations of the cerebral arteries through adulthood were carried out in four 187 subgroups. Indeed, our study might be the first to report the changes in arterial 188 189 morphologies in relation to aging in different age groups.

In subgroups, a significant age-related increase in tortuosity was first identified in the 190 35-44 age subgroup, appearing earlier than the age-related increase in mean arterial 191 radius and decrease in branch density, both of which were first evident in the 45-54 age 192 subgroup. In fact, aging was found to significantly influence all three vascular 193 morphologies only in subjects aged 45-54, and such associations remained significant 194 after adjusting for the vascular risk factors. On the contrary, although significant 195 associations of aging with branch density and mean vascular radius were identified in 196 197 the 65-75 subgroup initially, those ceased to be significant after adjusting for the vascular risk factors. Our results combined have at least two implications. First, 198 tortuosity may be a more sensitive metrics for vascular aging since it appears to be 199 changing rapidly with aging, even in the youngest subjects. Second, while aging 200 dominates the changes of the morphological features reported above in the 45-54 201 subgroup, common vascular risk factors may have played a more crucial part in the 65-202 75 subgroup. However, since a full history of chronic conditions from the patients was 203

unavailable, and the vascular risk factors were dichotomously coded (1 for the exposure
of risk and 0 for the absence), the investigation of the effect of the chronic conditions
duration on vessel structures was not feasible. Further studies may benefit from having
such information. For example, it is possible that the effects of the common vascular
risks, gradually re-shaping the cerebral vascular system, accrue through a lifetime until
significant changes can be observed.

Overall, the observed age-related morphological alternations may reflect impaired 210 211 vascular functions. First, since tortuosity increases the total vessel length, leading to increased flow resistance or, in advanced cases, occlusion of blood flow, it may 212 underpin the previously reported age-related reduction in cerebral perfusion (Han 2012, 213 Melamed et al. 1980). Indeed, tortuosity has been considered to increase the risk of 214 stroke (Alazzaz et al. 2000, Diedrich et al. 2011). Moreover, a recent study found that 215 the tortuosity index of the intracranial vessels can be used to differentiate healthy 216 individuals from subjects with mild cognitive impairment or AD, suggesting that 217 tortuosity may also reflect some pathological changes in the brain (Barbara-Morales et 218 219 al. 2020). Second, due to the failure of the current MRA to detect under-perfused vessels, the age-related reduction in the branch density may also be the result of impaired 220 cerebral perfusion. Nevertheless, as suggested previously (Bullitt et al. 2010), such 221 reduction in branch density with age may also result from a physical loss of smaller 222 vessels. In view of the restrictions posed by the current imaging method, it is not 223 plausible to attribute the observed reduction in branch density to either one of the causes. 224 Nonetheless, the fact that lower branch density was identified in the older population 225 suggests decrements in vascular functions in the aging population. Third, our results 226 227 also showed that advanced age was significantly associated with an increase in the mean vessel radius, in agreement with the results from previous MRA studies (Bullitt et al. 228 2010, Chen et al. 2019). Such results may arise from the rarefaction of smaller 229 branches, resulting in a relative increase in the proportion of larger branches, 230 compensatory vasodilation, or both. Our results, however, suggest that compensatory 231 vessel dilation may be a more viable explanation, since aging was still significantly 232 associated with increased vessel density after adjusting for the proportion of smaller 233

vessels. As mentioned before, tortuous vessels increase the flow resistance, resulting in 234 a more prominent drop in the affected sections of vessels and eventually leading to 235 236 decreased cerebral flow in the downstream vessels. Although very tempting, age-related compensatory vasodilation, which, as indicated by our results, appears later than age-237 related alternation in the mean vessel tortuosity, may work to counteract the pressure 238 drop brought about by increased tortuosity, and thus maintaining a constant blood flow 239 in the downstream parts. It is even possible that early intervention methods that either 240 241 prevent or slow further increase of tortuosity may be effective for the preservation of normal vascular functions. Taken together, our results may imply an age-related 242 disturbance of the cerebral hemostasis, which may underlie age-related brain 243 pathologies and would have important implications for early clinical prevention or 244 intervention methods targeting the middle-aged population. 245

In the current study, vascular risk factors were less related to cerebrovascular 246 morphologies after adjusting for age. Whereas having less vascular risks (indicated by 247 a higher CVR score) might mitigate the aging-related morphological alterations, as we 248 249 found that vascular score was positively correlated with vascular density at the age of 55-75. Our results combined emphasized that the common vascular risk factors may 250 play a more prominent role in aged groups. The research in young adults has shown 251 similar findings that lower modifiable cardiovascular score (more cardiovascular risk 252 factors) is associated with lower cerebral vascular density, raising the potential that 253 some individuals may be starting to diverge to different risk trajectories for brain 254 vascular health in early adulthood(Williamson et al. 2018). These results further 255 underline the importance of managing CVR to protect the health of cerebrovascular 256 257 system. Since reducing vascular risk factors and sustained lifestyle intervention can reduce cardiovascular burden (Spring et al. 2014, Shah et al. 2016), and improve 258 cerebral blood flow (Espeland et al. 2018). 259

Interaction of aging and risk factors on vascular morphology was presented. Greater effects of aging on cerebrovascular morphologies in smokers were observed in this study. A previous report has indicated that cigarette smoking could induce vascular smooth muscle cell senescence (Centner, Bhide and Salazar 2020). The accumulation

of senescent VSMCs with aging may be the underlying cause for our observation. The interaction of aging with alcohol consumption also significantly affects the vascular morphologies was the. Although the relationship between aging and vascular morphology was steeper in non-drinking participants, vascular density was lower in drinking participants for almost the entire age range. The possible explanation is that few participants are drinkers in our study, leading to the minimal power of our results. Future studies with more participants would help verify these interactions.

271 This study suffered from several limitations: 1) only one MRI modality, TOF MRA, was available in the study, which makes further investigation of the potential causes of 272 the discussed vascular alterations implausible. For example, to extract the general trend 273 of the age-related cerebral perfusion changes in different age subgroups, arterial spin-274 labeling would be necessary. 2) Risk factors were coded in our analysis as only having 275 dichotomous values indicating whether the subject had the chronic condition at the time 276 when their data was collected, and their medical history was not available. However, as 277 implied by our results, the effect of those risk factors may accumulate throughout time 278 279 until the consequence is noticeable. Future studies would benefit from having access to the medical history and more detailed information of the subjects' chronic conditions 280 such as the duration. 3) The cerebral vessel structural information obtained from TOF 281 MRA was averaged for each individual, and thus might be unable to identify subtle 282 correlations between aging and local vessels. Existing studies have also primarily 283 focused on the cerebral vasculature in terms of whole-brain averaging or local 284 averaging (Williamson et al. 2018, Mouches et al. 2021), since it is laborious to 285 distinguish vessels by anatomical structure on images manually. More intelligent 286 287 techniques, such as machine learning-based approaches, are needed to automate the distinction between different cerebral vessels. 288

In the current study, we present an analysis of the 3D TOF MRA images of 1176 subjects recruited from 2 communities in Shanghai. Significant differences in the cerebrovascular morphologies were identified between older subjects and younger subjects even after adjusting for vascular risk factors. In subgroups based on age, age was a significant predictor for the alternations of all three vessel morphologies extracted

by MRA only in the 45-54 subgroup. In the 65-75 subgroup, although significant 294 associations of aging with branch density and vessel radius were initially identified, the 295 296 significance has ceased to exist after adjusting for vascular risk factors. Furthermore, the age-related increase in tortuosity was the only significant age-related change 297 identified in the youngest subgroup. These results provide two key insights. First, 298 morphometric changes visualizable by MRA may underlie different physiological 299 processes. Second, early intervention methods that either prevent or slow further 300 301 increases of tortuosity may be effective for preserving normal vascular functions.

302

303 Materials and Methods

304 Participants

Participants in the present study also completed the survey of unruptured cerebral 305 aneurysms in Chinese adults (Li et al. 2013). This cross-sectional study initially 306 enrolled 4813 participants (aged 35 to 75 years) who completed the survey between 307 June 2007 and June 2011 from two different Shanghai communities in China. Ten years 308 309 later (2017 to 2021), participants were invited to call back and reported on their health status. As shown in Figure 1, 3198 participants at the first survey who were free of 310 cardiac disease, stroke, arteriostenosis or a history of cancer were first selected for the 311 call back. Participants with identified intracranial aneurysms were not included to 312 prevent potential influence of cerebral aneurysms on vascular morphology. Among the 313 3198 subjects, 1751 participants that either withdrew or refused to respond to the call 314 315 back and 207 participants who had severe cardiovascular disease (CVD) events, 316 including myocardial infarction, unstable angina, heart failure, coronary heart disease 317 or stroke over the 10 years were excluded. 1240 participants were included in final the 318 analysis. In these participants, although some have suffered chronic conditions, such as hypertension and diabetes, these conditions were well managed, and no severe CVD 319 events were reported during the past ten years. The study was approved by the 320 321 institutional review board of the Sixth People's Hospital, Shanghai, China. All participants were voluntarily enrolled and provided written informed consent before 322 323 data collection.

324 Assessment of Risk Factors

Initially, all participants were asked to complete a standard questionnaire to provide 325 326 demographic characteristics, personal and family medical history, and lifestyle risk factors. Information about smoking, alcohol consumption, hypertension, diabetes, 327 hyperlipidemia, stroke, coronary heart disease, myocardial infarction, and arrhythmia 328 was obtained. In addition, participants completed physical examinations including 329 measurement of height, weight, and blood pressure. Venous blood samples were 330 331 collected by certified nurses between 7 AM and 8 AM after overnight fasting. The presence of hypertension was defined as systolic blood pressure ≥ 140 mm Hg or 332 diastolic blood pressure \geq 90 mm Hg, a previous diagnosis of hypertension, or use of 333 anti-hypertensive medications. Smokers were defined as smoked>100 cigarettes in a 334 335 lifetime, and drinker was defined as consuming >30 g of alcohol per week for 1 or more year. The presence of diabetes was defined by a fasting level of plasma glucose 336 \geq 7.0 mmol/L, a previous diagnosis of diabetes, or treatment with antidiabetic drugs. 337 Hyperlipidemia was defined by total plasma cholesterol (TC) > 5.2 mmol/L, 338 339 triglycerides (TG) > 1.7 mmol/L, a previous diagnosis of hyperlipidemia, or use of lipid-lowering drugs. During the call back, a new questionnaire was administered by 340 trained staff to obtain information on medical history within the past 10 years. Due to 341 the limitations of the call back, accurate assessments of the vascular risk factors were 342 not possible. Hence the vascular risk information from baseline was used in the 343 subsequent analysis. 344

345 MRI Scanning and Processing

All MRI examinations were performed on one 3.0T MRI system (Philips Healthcare,
Amsterdam, The Netherlands). 3D-TOF MRA was completed using 3-dimensional T1weighted fast-field sequences with the following settings: repetition time/echo time,
35/7; flip angle, 20 degrees; field of view, 250×190×108 mm3; 4 slabs (180 slices);
slice thickness, 0.8 mm; matrix, 732×1024; acquisition time, 8 minutes and 56 seconds.
Any adverse events during the MRI examination, including panic attack,
claustrophobia, dizziness, or falling, were queried and monitored by the investigators.

Cerebral vessel segmentation and morphologies extraction were completed on TOF 353 MRA with the following steps: 1) vascular structures were enhanced using the Jerman 354 355 Enhancement Filter (Jerman et al. 2016); 2) the level-set based method was used to obtain vessel segmentation; 3) segmentation results were firstly checked by B.Y.Z and 356 B.W. Segmentations with too few vascular branches, nonvascular tissue remnants, or 357 358 incorrect vascular branch adhesions were re-segmented by adjusting the algorithm parameters. Then all segmentation results were visually checked twice by two 359 360 radiologist J.L and Y.H.L who have over 3 years' experience with neuroimaging interpretation. The segmented vessels that were still showing too few branches, 361 remnants of nonvascular tissue, or severe vascular branch adhesions were defined as 362 poor vascular segmentation and excluded from the analysis; 4) vessel centerlines were 363 extracted using the Skeleton 3D toolbox(Kollmannsberger et al. 2017); 5) 364 cerebrovascular morphologic measurements, including vessel density, radius and 365 tortuosity were calculated. To avoid the influence of extracerebral vessels, the brain 366 extraction tool of FSL (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/) was used to segment the 367 brain regions on MRA images. Only vessels involved in the brain region were included 368 in the subsequent analysis. The vessel density was defined as the ratio of intracranial 369 370 artery branch numbers to the brain volume. The vessel tortuosity represents the averaged tortuosity for each segment and was calculated as the ratio of the shortest path 371 and the actual length of a given vessel segment. The vessel radius was determined in 372 each centerline points and then averaged over each individual participant. The smaller 373 374 vessels were defined as vessels with a mean radius less than a certain threshold (global 375 thresholding based on the radius distribution of the whole intracranial arteries) (Zhang 376 et al. 2021). All data processing processes were accomplished on the Matlab2019a

377 (MathWorks, Natick, Massachusetts, USA).

378 Statistical Analysis

Bivariable and multivariable analyses were completed to investigate association between cerebrovascular morphologies (cerebral vessel density, radius and tortuosity) and vascular risk makers (age, body mass index, smoking, alcohol consumption, diabetes, hypertension and hyperlipidemia). Linear regression models were used with

the risk makers as the determinant and cerebrovascular morphologies as outcome variables. The differential influence of aging on cerebrovascular morphologies by other risk makers was tested using interaction terms in the multivariable model. Participants were divided into different age groups according to their age: middle-youth (35-54 years), middle-aged (55-75 years), 35 to 44 years, 45 to 54 years, 55 to 64 years and 65 to 75 years.

The individuals' combined CVR score from 6 risk factors was also analyzed, as existing 389 390 evidence indicated combined risk factor score was correlated with the brain health in young adults (Williamson et al. 2018). The score was cumulative based on each of the 391 following factors: body mass index ≤ 24 ; no smoking; not a drinker; no hyperlipidemia; 392 no hypertension; no diabetes, with a higher number indicating less exposure to 393 cerebrovascular risks. The relationships between the CVR score and MRA findings 394 were studied using the linear regression model. In addition, linear analysis was 395 completed to investigate the correlations between aging and cerebrovascular 396 morphologies in different age subgroups. 397

All statistical analysis was completed using Matlab2019a (MathWorks, Natick, Massachusetts, USA). Shapiro-Wilk test was performed to confirm the normality of continuous variables. Results of normally distributed continuous variables are presented as mean and standard deviation (SD) or as median and interquartile range (IQR). For categorical variables, frequencies, and proportions are presented. The twosample t-test was used to evaluate group differences. All tests were 2-sided and a probability value <0.05 was considered significant.

405

Funding: This work was supported by the National Natural Science Foundation of 406 China (No. 81971583, No. 81871329), National Key R&D Program of China (No. 407 2018YFC1312900), New developing and Frontier Technologies of Shanghai Shen 408 Kang Hospital Development Center (No. SHDC12018117), Shanghai Municipal 409 Education Commission-Gaofeng Clinical Medicine Grant (No. 2016427), Shanghai 410 Municipal Science and Technology Major Project (No.2018SHZDZX01, 411 No.2017SHZDZX01) and ZJLab. 412

413

Competing interests statement 414

415 There are no competing interests.

416

Reference 417

- Alazzaz, A., J. Thornton, V. A. Aletich, G. M. Debrun, J. I. Ausman & F. Charbel (2000) Intracranial 418 419 Percutaneous Transluminal Angioplasty for Arteriosclerotic Stenosis. Archives of Neurology, 57, 420 1625-1630.
- 421 Alber, J., S. Alladi, H.-J. Bae, D. A. Barton, L. A. Beckett, J. M. Bell, S. E. Berman, G. J. Biessels, S. E. 422 Black, I. Bos, G. L. Bowman, E. Brai, A. M. Brickman, B. L. Callahan, R. A. Corriveau, S. 423 Fossati, R. F. Gottesman, D. R. Gustafson, V. Hachinski, K. M. Hayden, A. M. Helman, T. M. 424 Hughes, J. D. Isaacs, A. L. Jefferson, S. C. Johnson, A. Kapasi, S. Kern, J. C. Kwon, J. Kukolja, 425 A. Lee, S. N. Lockhart, A. Murray, K. E. Osborn, M. C. Power, B. R. Price, H. F. M. Rhodius-426 Meester, J. A. Rondeau, A. C. Rosen, D. L. Rosene, J. A. Schneider, H. Scholtzova, C. E. 427 Shaaban, N. C. B. S. Silva, H. M. Snyder, W. Swardfager, A. M. Troen, S. J. van Veluw, P. 428 Vemuri, A. Wallin, C. Wellington, D. M. Wilcock, S. X. Xie & A. H. Hainsworth (2019) White matter hyperintensities in vascular contributions to cognitive impairment and dementia (VCID): 429 430 Knowledge gaps and opportunities. Alzheimer's & dementia (New York, N. Y.), 5, 107-117.
- 431 Banerjee, G., D. Wilson, H. R. Jäger & D. J. Werring (2016) Novel imaging techniques in cerebral small 432 vessel diseases and vascular cognitive impairment. Biochim Biophys Acta, 1862, 926-38.
- 433 Barbara-Morales, E., J. Perez-Gonzalez, K. C. Rojas-Saavedra & V. Medina-Banuelos (2020) Evaluation 434 of Brain Tortuosity Measurement for the Automatic Multimodal Classification of Subjects with 435 Alzheimer's Disease. Computational Intelligence and Neuroscience, 2020.
- 436 Bullitt, E., D. Zeng, B. Mortamet, A. Ghosh, S. R. Aylward, W. Lin, B. L. Marks & K. Smith (2010) The effects of healthy aging on intracerebral blood vessels visualized by magnetic resonance 437 438 angiography. Neurobiology of Aging, 31, 290-300.
- 439 Centner, A. M., P. G. Bhide & G. Salazar (2020) Nicotine in Senescence and Atherosclerosis. Cells, 9.
- 440 Chen, L., J. Sun, D. S. Hippe, N. Balu, Q. Yuan, I. Yuan, X. Zhao, R. Li, L. He, T. S. Hatsukami, J.-N. 441 Hwang & C. Yuan (2019) Quantitative assessment of the intracranial vasculature in an older 442 adult population using iCafe. Neurobiology of Aging, 79, 59-65.
- 443 D'Agostino, R. B., R. S. Vasan, M. J. Pencina, P. A. Wolf, M. Cobain, J. M. Massaro & W. B. Kannel 444 (2008) General Cardiovascular Risk Profile for Use in Primary Care. Circulation, 117, 743-753.
- 445 Debette, S., S. Seshadri, A. Beiser, R. Au, J. J. Himali, C. Palumbo, P. A. Wolf & C. DeCarli (2011) 446 Midlife vascular risk factor exposure accelerates structural brain aging and cognitive decline. 447 Neurology, 77, 461-468.
- 448 Diedrich, K. T., J. A. Roberts, R. H. Schmidt, C. K. Kang, Z. H. Cho & D. L. Parker (2011) Validation 449 of an arterial tortuosity measure with application to hypertension collection of clinical 450 hypertensive patients. BMC Bioinformatics, 12 Suppl 10, S15.
- Donato, A. J., D. R. Machin & L. A. Lesniewski (2018) Mechanisms of Dysfunction in the Aging 451 452 Vasculature and Role in Age-Related Disease. Circulation Research, 123, 825-848.
- 453 Espeland, M. A., J. A. Luchsinger, R. H. Neiberg, O. Carmichael, P. J. Laurienti, X. Pi-Sunyer, R. R. 454 Wing, D. Cook, E. Horton, R. Casanova, K. Erickson, R. N. Bryan & M. Action Hlth Diabet

P01	
It is made available under a	CC-BY 4.0 International license

455	Brain (2018) Long Term Effect of Intensive Lifestyle Intervention on Cerebral Blood Flow.
456	Journal of the American Geriatrics Society, 66, 120-126.
457 459	Ferreira, D., R. Correia, A. Nieto, A. Machado, Y. Molina & J. Barroso (2015) Cognitive decline before the age of 50 can be detected with sensitive cognitive measures. <i>Psicothama</i> , 27, 216, 22
450	Han H. C. (2012) Truisted blood vessels, symptoms, stiele sy and biomechanical mechanisms. Journal
459	Han, HC. (2012) I wisted blood vessels: symptoms, ettology and biomechanical mechanisms. <i>Journal</i>
460	of vascular research, 49, 185-197.
401	Jerman, I., F. Pernus, B. Likar & Z. Spicin (2016) Ennancement of vascular Structures in 3D and 2D
462	Angiographic images. <i>Ieee Transactions on Medical Imaging</i> , 35, 2107-2118.
463	Kalaria, R. N. (2002) Small vessel Disease and Alzneimer's Dementia: Pathological Considerations.
404	Cerebrovascular Diseases, 15(suppl 2), 48-52.
405	Kloppenborg, R. P., E. van den Berg, L. J. Kappelle & G. J. Blessels (2008) Diabetes and other vascular
400	fisk factors for dementia: which factor matters most? A systematic review. Eur J Pharmacol,
467	585, 97-108.
468	Kolimannsberger, P., M. Kerschnitzki, F. Repp, W. Wagermaier, R. Weinkamer & P. Fratzi (2017) The
409	small world of osteocytes: connectomics of the lacuno-canalicular network in bone. New
470	Journal of Physics, 19.
471	Laureni, S. & P. Boulouyne (2015) The Structural Factor of Hypertension Large and Small Artery
472	Anterations. Circulation Research, 116, 1007-1021.
473	Leeuwis, A. E., M. R. Benedicius, J. P. A. Kuljer, M. A. A. Binnewijzend, A. M. Hoogmemstra, S. C. J.
4/4 475	vertainie, I. Koene, P. Schenens, F. Barkhol, N. D. Prins & W. M. van der Filer (2017) Lower
475	disease Alabeiru and R. Dementin, 12, 521, 540
470	Li M H S W Chen V D Li V C Chen V S Cheng D L Hu H O Ten O Wu W Weng 7 K
477	LI, MH., SW. Chen, ID. LI, IC. Chen, IS. Cheng, DJ. Hu, HQ. Tan, Q. Wu, W. Wang, ZK.
470	D. P. Dong, S. Y. Ding, Y. F. Yu, Y. H. Hou, & W. D. Lie (2012) Drevelonce of Unruntured
479	DR. Feng, SA. Ding, AF. Au, AH. Hou & WF. Jia (2015) Flevalence of Unituptuted
400	S14
401 400	JI4-T. Malamad E. S. Laury, S. Bantin, G. Caapar, & V. Binat (1980), DEDUCTION IN DECIONAL
40Z 102	CEDEDDAL DLOOD ELOW DUDING NODMAL AGING IN MAN. Stroke 11, 21, 25
403	Mouches P S Language M Domin M D Hill & N D Forkert (2021) Influence of cardiovascular risk
404 105	fostors on morphological changes of carebral arteries in healthy adults across the life span
405	Scientific Paparts 11
400	Sheh P. V. V. I. Murthy I. A. Colongolo, I. Pais, P. A. Venkatash P. Sharma, S. A. Abhasi, D. C. Coff
407	LI Carr I S Pana I G Terry C Bouchard M A Sarzynski A Eisman T Neilan S Das
400	M. Jerosch Herold, C. E. Lewis, M. Carnethon, G. D. Lewis & I. A. C. Lima (2016) Association
409	of Fitness in Young Adulthood With Survival and Cardiovascular Risk The Coronary Artery
430 /01	Risk Development in Young Adults (CARDIA) Study Jama Internal Medicine, 176, 87-95
491	Silvestrini M P. Pasqualetti P. Baruffaldi M. Bartolini V. Handouk, M. Matteis, F. Moffa I.
492	Provinciali & F. Vernieri (2006) Cerebrovascular Reactivity and Cognitive Decline in Patients
493	With Alzheimer Disease Stroke 37, 1010-1015
495	Spring B A C Moller L A Colangelo I Siddique M Roehrig M I Daviglus I F Polak I P Reis
496	S. Sidney & K. Liu (2014) Healthy Lifestyle Change and Subclinical Atherosclerosis in Voung
497	Adults Coronary Artery Risk Development in Young Adults (CARDIA) Study Circulation 130
498	10-+
100	

- Swan, G. E., C. DeCarli, B. L. Miller, T. Reed, P. A. Wolf, L. M. Jack & D. Carmelli (1998) Association
 of midlife blood pressure to late-life cognitive decline and brain morphology. *Neurology*, 51,
 986-993.
- Tolppanen, A. M., A. Solomon, H. Soininen & M. Kivipelto (2012) Midlife vascular risk factors and
 Alzheimer's disease: evidence from epidemiological studies. *J Alzheimers Dis*, 32, 531-40.
- Walker, K. A., A. R. Sharrett, A. Wu, A. L. C. Schneider, M. Albert, P. L. Lutsey, K. Bandeen-Roche, J.
 Coresh, A. L. Gross, B. G. Windham, D. S. Knopman, M. C. Power, A. M. Rawlings, T. H.
 Mosley & R. F. Gottesman (2019) Association of Midlife to Late-Life Blood Pressure Patterns
 With Incident Dementia. *Jama-Journal of the American Medical Association*, 322, 535-545.
- Williamson, W., A. J. Lewandowski, N. D. Forkert, L. Griffanti, T. W. Okell, J. Betts, H. Boardman, T.
 Siepmann, D. McKean, O. Huckstep, J. M. Francis, S. Neubauer, R. Phellan, M. Jenkinson, A.
 Doherty, H. Dawes, E. Frangou, C. Malamateniou, C. Foster & P. Leeson (2018) Association of
 Cardiovascular Risk Factors With MRI Indices of Cerebrovascular Structure and Function and
 White Matter Hyperintensities in Young Adults. *Jama-Journal of the American Medical Association*, 320, 665-673.
- Xu, X., B. Wang, C. Ren, J. Hu, D. A. Greenberg, T. Chen, L. Xie & K. Jin (2017) Age-related Impairment
 of Vascular Structure and Functions. *Aging and Disease*, 8, 590-610.
- Zhang, B., Y. Wang, B. Wang, Y.-H. Chu, Y. Jiang, M. Cui, H. Wang & X. Chen (2021) MRI-Based
 Investigation of Association Between Cerebrovascular Structural Alteration and White Matter
 Hyperintensity Induced by High Blood Pressure. *Journal of Magnetic Resonance Imaging*.
- 519 Zhang, D. & M. E. Raichle (2010) Disease and the brain's dark energy. *Nature Reviews Neurology*, 6, 15520 28.
- 521 Zimprich, D. & A. Mascherek (2010) Five views of a secret: does cognition change during middle
 522 adulthood? *European Journal of Ageing*, 7, 135-146.

523

525 Table 1. Characteristics of the Study Participants					
Age Range	35-44 y	45-54 y	55-64 y	65-75 y	Overall
Demographic					
Participants, No.	201	421	342	212	1176
Age, mean (SD), years	40.4 (3.3)	51.1 (2.7)	59.3 (2.8)	70.3 (3.2)	55.1 (9.9)
Female, No. (%)	121 (0.60)	279 (0.66)	189 (0.55)	113 (0.53)	702 (0.60)
Body mass index, mean (SD)	23.3 (3.0)	23.5 (3.0)	23.8 (2.9)	23.2 (2.6)	23.5 (2.9)
Smoking, No. (%)	30 (0.15)	87 (0.21)	70 (0.20)	39 (0.18)	226 (0.19)
Alcohol consumption, No. (%)	16 (0.08)	42 (0.10)	55 (0.16)	24 (0.11)	137 (0.12)
Diabetes diagnosis, No. (%)	1 (0)	29 (0.07)	34 (0.10)	30 (0.14)	94 (0.08)
Hypertension, No. (%)	22 (0.11)	78 (0.19)	117 (0.34)	84 (0.40)	301 (0.26)
Hyperlipidemia, No. (%)	7 (0.03)	14 (0.03)	30.0 (0.09)	15.0 (0.07)	66 (0.06)
Cerebrovascular morphology					
Vessel density, mean (SD)	0.09 (0.02)	0.08 (0.02)	0.07 (0.02)	0.07 (0.02)	0.08 (0.02)
Vessel radius, mean (SD), mm	0.78 (0.07)	0.80 (0.08)	0.82 (0.07)	0.83 (0.07)	0.81 (0.07)
Vessel tortuosity, mean (SD)	1.19 (0.03)	1.20 (0.03)	1.20 (0.03)	1.21 (0.03)	1.2 (0.03)

Table 2. Association between age and cerebrovascular morphologies.

529 In the multivariable adjusted models, sex and cerebrovascular risk factors were adjusted, including body mass index, smoking status, alcohol consumption, diabetes status, hypertension 530 status and hyperlipidemia status. Middle-youth represented participants aged from 35 to 54 531 years. Middle-aged represented participants aged from 55 to 75 years. 532

	Bivariable		Multivariable adjusted	
	Estimation. β (95%)	p value	Estimation. β (95%)	p value
Vessel density				
Middle-youth	-8.1×10 ⁻⁴ (-0.001 to -4.9×10 ⁻⁴)	p<0.001	-8.0×10 ⁻⁴ (-0.001 to -4.8×10 ⁻⁴)	p<0.001
Middle-aged	-4.3×10 ⁻⁴ (-0.001 to -1.2×10 ⁻⁴)	0.008	-4.0×10 ⁻⁴ (-0.001 to -7.5×10 ⁻⁵)	0.02
35-44 years	-8.8×10^{-4} (-0.002 to 1.4×10^{-4})	0.09	-9.2×10 ⁻⁴ (-0.002 to 1.1×10 ⁻⁴)	0.08
45-54 years	-0.002 (-0.002 to -0.001)	p<0.001	-0.001 (-0.002 to -0.001)	0.001
55-64 years	-3.1×10 ⁻⁴ (-0.001 to 0.001)	0.48	-2.6×10 ⁻⁴ (-0.001 to 0.001)	0.56
65-75 years	-0.001 (-0.002 to -7.6×10 ⁻⁵)	0.036	-8.8×10^{-4} (-0.002 to 1.7×10^{-4})	0.10
Overall	-5.9×10 ⁻⁴ (-0.001 to -4.5×10 ⁻⁴)	p<0.001	-5.7×10 ⁻⁴ (-0.001 to -4.3×10 ⁻⁴)	p<0.001
Vessel radius				
Middle-youth	0.003 (0.002 to 0.003)	p<0.001	0.003 (0.002 to 0.004)	p<0.001
Middle-aged	0.002 (0.001 to 0.003)	p<0.001	0.002 (0.001 to 0.003)	0.001
35-44 years	0.001 (-0.002 to 0.004)	0.45	0.001 (-0.001 to 0.004)	0.33
45-54 years	0.005 (0.002 to 0.008)	p<0.001	0.005 (0.003 to 0.008)	p<0.001
55-64 years	0.003 (-5.3×10 ⁻⁵ to 0.006)	0.06	$0.003 (-2.3 \times 10^{-4} \text{ to } 0.005)$	0.07
65-75 years	0.003 (2.9×10 ⁻⁴ to 0.006)	0.03	0.002 (-0.001 to 0.005)	0.13
Overall	0.002 (0.002 to 0.002)	p<0.001	0.002 (0.002 to 0.002)	p<0.001
Vessel tortuosity				
Middle-youth	0.001 (0.001 to 0.002)	p<0.001	0.001 (0.001 to 0.002)	p<0.001
Middle-aged	$7.8 \times 10^{-4} (3.6 \times 10^{-4} \text{ to } 0.001)$	p<0.001	7.7×10^{-4} (3.5×10 ⁻⁴ to 0.001)	p<0.001
35-44 years	0.001 (1.6×10 ⁻⁴ to 0.003)	0.03	0.002 (3.4×10 ⁻⁴ to 0.003)	0.01
45-54 years	0.002 (0.001 to 0.003)	0.003	0.001 (4.4×10 ⁻⁴ to 0.003)	0.006
55-64 years	2.6×10 ⁻⁴ (-0.001 to 0.001)	0.65	3.2×10 ⁻⁴ (-0.001 to 0.001)	0.58
65-75 years	7.2×10 ⁻⁴ (-0.001 to 0.002)	0.31	7.5×10 ⁻⁴ (-0.001 to 0.002)	0.32
Overall	8.4×10^{-4} (0.001 to 0.001)	p<0.001	8.5×10 ⁻⁴ (0.001 to 0.001)	p<0.001

533

528

In the multivariable adjusted models, sex and cerebrovascular risk factors were adjusted, including 536

537 body mass index, smoking status, alcohol consumption, diabetes status, hypertension status and

hyperlipidemia status. 538

	Estimation, β (95%CI)	p value
Proportion of smaller vessels (PSV)		
Bivariable	-4.0×10 ⁻⁴ (-6.3×10 ⁻⁴ to -1.7×10 ⁻⁴)	0.001
Multivariable adjusted	-3.7×10 ⁻⁴ (-6.1×10 ⁻⁴ to -1.3×10 ⁻⁴)	0.003
Vessel Radius adjusted for PSV	2.7×10 ⁻⁴ (6.6×10 ⁻⁵ to 4.8×10 ⁻⁴)	0.01

Table 4. Association between cerebrovascular morphologies and cerebrovascular risk

542	factor	rs			
543 Only signif	icant interaction terms were presented	. Middle-youth	n represented participants aged		
544 from 35 to 54 years. Middle-aged represented participants aged from 55 to 75 years.					
	Bivariable		Multivariable		
	Estimation, β (95%CI)	p value	Estimation, β (95%CI)	p value	
Vessel density					
Age	-5.9×10 ⁻⁴ (-7.2×10 ⁻⁴ to -4.5×10 ⁻⁴)	p<0.001	-5.7×10 ⁻⁴ (-7.1×10 ⁻⁴ to -4.3×10 ⁻⁴)	p<0.001	
Body mass index	-2.4×10 ⁻⁶ (-4.8×10 ⁻⁴ to 4.7×10 ⁻⁴)	0.99	1.4×10^{-5} (-4.6×10 ⁻⁴ to 4.9×10 ⁻⁴)	0.95	
Smoking	-0.002 (-0.006 to 0.001)	0.25	-2.7×10 ⁻⁴ (-0.004 to 0.004)	0.89	
Alcohol consumption	-0.004 (-0.009 to -1.5×10 ⁻⁴)	0.043	-0.003 (-0.008 to 0.001)	0.16	
Diabetes diagnosis	-0.008 (-0.01 to -0.003)	0.001	-0.005 (-0.01 to -9.1×10 ⁻⁵)	0.046	
Hypertension	-0.003 (-0.006 to 3.4×10 ⁻⁴)	0.08	5.8×10 ⁻⁴ (-0.003 to 0.004)	0.72	
Hyperlipidemia	0.003 (-0.003 to 0.009)	0.31	$0.005 (-5.4 \times 10^{-4} \text{ to } 0.01)$	0.07	
Age*Smoking			-6.7×10 ⁻⁴ (-8.8×10 ⁻⁴ to -4.6×10 ⁻⁴)	0.002	
Age*Alcohol consumption			5.8×10^{-4} (8.6×10 ⁻⁴ to 3.1×10 ⁻⁴)	0.03	
Vessel radius					
Age	0.002 (0.002 to 0.002)	p<0.001	0.002 (0.002 to 0.002)	p<0.001	
Body mass index	3.6×10 ⁻⁴ (-0.001 to 0.002)	0.63	4.2×10 ⁻⁴ (-0.001 to 0.002)	0.56	
Smoking	0.01 (8.0×10 ⁻⁴ to 0.02)	0.035	0.01 (-0.001 to 0.02)	0.08	
Alcohol consumption	0.009 (-0.004 to 0.02)	0.19	-7.7×10 ⁻⁴ (-0.02 to 0.01)	0.92	
Diabetes diagnosis	0.02 (0.008 to 0.04)	0.003	0.01 (-0.004 to 0.03)	0.14	
Hypertension	0.009 (-3.5×10 ⁻⁴ to 0.02)	0.06	-0.002 (-0.01 to 0.008)	0.68	
Hyperlipidemia	-0.007 (-0.03 to 0.01)	0.42	-0.02 (-0.03 to 0.003)	0.10	
Age*Smoking			$0.002 (9.6 \times 10^{-4} \text{ to } 0.002)$	0.01	
Age*Alcohol consumption			-0.002 (-0.003 to -8.6×10 ⁻⁴)	0.036	
Vessel tortuosity					
Age	8.4×10^{-4} (6.7×10 ⁻⁴ to 0.001)	p<0.001	8.5×10^{-4} (6.7×10 ⁻⁴ to 0.001)	p<0.001	
Body mass index	2.7×10^{-4} (-3.5×10 ⁻⁴ to 8.8×10 ⁻⁴)	0.39	3.0×10^{-4} (-3.1×10 ⁻⁴ to 9.1×10 ⁻⁴)	0.33	
Smoking	9.3×10 ⁻⁴ (-0.004 to 0.005)	0.69	0.002 (-0.003 to 0.007)	0.52	
Alcohol consumption	-2.8×10 ⁻⁴ (-0.006 to 0.005)	0.92	-0.003 (-0.009 to 0.004)	0.40	
Diabetes diagnosis	0.005 (-0.002 to 0.01)	0.17	-6.2×10 ⁻⁴ (-0.007 to 0.006)	0.85	
Hypertension	0.005 (5.8×10 ⁻⁴ to 0.009)	0.03	-1.5×10 ⁻⁴ (-0.004 to 0.004)	0.95	
Hyperlipidemia	0.002 (-0.005 to 0.01)	0.57	-4.0×10 ⁻⁴ (-0.008 to 0.007)	0.92	

547 Table 5. Association between cerebrovascular score and cerebrovascular morphologies.

Middle-youth represented participants aged from 35 to 54 years. Middle-aged represented 548

549 participants aged from 55 to 75 years.

			Adjusted for age and sex	
	Estimation. β (95%)	p value	Estimation. β (95%)	p value
Vessel density				
Middle-youth	-1.9×10 ⁻⁴ (-0.002 to 0.002)	0.83	-6.3×10 ⁻⁴ (-0.002 to 0.001)	0.50
Middle-aged	0.002 (0.001 to 0.004)	0.007	0.002 (0.001 to 0.004)	0.01
35-44 years	0.001 (-0.002 to 0.005)	0.45	0.001 (-0.002 to 0.005)	0.44
45-54 years	-0.001 (-0.004 to 0.001)	0.18	-0.001 (-0.003 to 0.001)	0.28
55-64 years	0.002 (2.8×10 ⁻⁴ to 0.004)	0.03	$0.002 (1.0 \times 10^{-4} \text{ to } 0.004)$	0.041
65-75 years	0.003 (-4.7×10 ⁻⁴ to 0.006)	0.10	0.002 (-0.001 to 0.005)	0.17
Overall	0.002 (0.001 to 0.003)	0.005	8.8×10 ⁻⁴ (-3.8×10 ⁻⁴ to 0.002)	0.17
Vessel radius				
Middle-youth	-0.004 (-0.010 to 0.001)	0.14	-0.003 (-0.009 to 0.002)	0.22
Middle-aged	-0.004 (-0.010 to 0.001)	0.10	-0.004 (-0.010 to 0.001)	0.09
35-44 years	-0.009 (-0.02 to 5.4×10 ⁻⁵)	0.053	-0.010 (-0.02 to -1.6×10 ⁻⁴)	0.048
45-54 years	-2.5×10 ⁻⁴ (-0.007 to 0.007)	0.94	-0.001 (-0.008 to 0.005)	0.69
55-64 years	-0.003 (-0.010 to 0.004)	0.40	-0.003 (-0.009 to 0.004)	0.45
65-75 years	-0.008 (-0.02 to 2.8×10 ⁻⁴)	0.06	-0.007 (-0.02 to 0.001)	0.09
Overall	-0.007 (-0.01 to -0.003)	0.001	-0.004 (-0.008 to -2.0×10 ⁻⁴)	0.039
Vessel tortuosity				
Middle-youth	-4.4×10 ⁻⁴ (-0.003 to 0.002)	0.71	5.1×10^{-4} (-0.002 to 0.003)	0.66
Middle-aged	-0.001 (-0.003 to 0.001)	0.38	-9.6×10 ⁻⁴ (-0.003 to 0.001)	0.42
35-44 years	-0.003 (-0.008 to 0.001)	0.12	-0.004 (-0.008 to 3.4×10 ⁻⁴)	0.07
45-54 years	0.002 (-0.001 to 0.005)	0.16	$0.002 (-3.6 \times 10^{-4} \text{ to } 0.005)$	0.09
55-64 years	-6.2×10 ⁻⁴ (-0.003 to 0.002)	0.65	-5.2×10 ⁻⁴ (-0.003 to 0.002)	0.71
65-75 years	-0.002 (-0.007 to 0.002)	0.29	-0.002 (-0.006 to 0.002)	0.36
Overall	-0.002 (-0.003 to -1.4×10 ⁻⁵)	0.048	-2.6×10 ⁻⁴ (-0.002 to 0.001)	0.76
550				



Figure 1. Flow chart of recruitment in the current study



perpetuity. It is made available under a CC-BY 4.0 International license .

Figure 2. Association of cerebrovascular morphology with age and cerebrovascular risk factors.

557 (a)~(c) Regression results between age and cerebrovascular morphologies, including vessel

density, vessel radius and vessel tortuosity, in different age groups. (d)~(f) Association between

vessel density and age in different groups (smoking and non-smoking, drinker and non-drinker,

smoker or drinker and both not). Shaded area represents the 95% confidence interval. (g)~(i)
 Association between vessel radius and age in different groups.



