- 1 Depressive symptoms are associated with cigarette smoking among tuberculosis patients in
- 2 **Botswana**
- 3
- 4 Alexandria Jones-Patten MSN, RN. Sue & Bill Gross School of Nursing, University of
- 5 California Irvine, Irvine, CA, USA;
- 6 Qiao Wang MPH: Fielding School of Public Health, University of California Los Angeles, Los
- 7 Angeles, CA, USA;
- 8 Keneilwe Molebatsi. Department of Psychiatry, Faculty of Medicine, University of Botswana,
- 9 Gaborone, Botswana;
- 10 Thomas E. Novotny MD, MPH: School of Public Health, San Diego State University, San
- 11 Diego, CA, USA;
- 12 Kamran Siddiqi MBBS, MRCP, MPH, FFPH, PhD. Department of Health Sciences, University
- 13 of York, York, England;
- 14 Chawangwa Modongo: Victus Global Botswana Organisation, Gaborone, Botswana;
- 15 Nicola M. Zetola MD, MPH: Victus Global Botswana Organisation, Gaborone, Botswana;
- 16 Bontle Mbongwe, PhD Department of Environmental Health, Faculty of Health Sciences,
- 17 University of Botswana, Gaborone, Botswana;
- 18 Sanghyuk S. Shin PhD. Sue & Bill Gross School of Nursing, University of California Irvine,
- 19 Irvine, CA, USA.
- 20
- 21
- 22
- 23

It is made available under a CC-BY-NC-ND 4.0 International license .

24 Corresponding Author:

- 25 Sanghyuk S. Shin, PhD
- 26 Associate Professor, Sue & Bill Gross School of Nursing
- 27 Director, UCI Infectious Disease Science Initiative
- 28 University of California Irvine
- 29 106F Berk Hall
- 30 Irvine, CA 92697
- 31 ssshin2@uci.edu
- 32 Tel: 949-576-8675

33

34

35 Acknowledgements

- 36 We would like to thank study participants for their time and efforts. We would also like to thank
- 37 University of California, Irvine and Botswana Ministry of Health and Wellness Human Research
- 38 Development Committee. Dr. Siddiqi was funded by Medical Research Council (UK) from TB
- 39 Multimorbidity Grant (Ref No: MC_PC_MR/T037806/1).

40

41 **Conflict of Interest Declaration**

42 All authors declare we have no conflict of interest.

43

It is made available under a CC-BY-NC-ND 4.0 International license .

45 Abstract

46 Background: Researchers have increasingly recognized the adverse effects of smoking on 47 tuberculosis (TB) outcomes. Smoking may be a maladaptive coping mechanism for depression 48 and anxiety among TB patients; however, this association has not yet been investigated. 49 50 Design/Methods: We conducted a cross-sectional study among newly diagnosed TB patients 51 between January and December 2019 in Gaborone, Botswana, and evaluated factors associated 52 with cigarette smoking. Using the Patient Health Questionnaire-9 and the Zung Self-Rating 53 Anxiety scale, we collected depression and anxiety scores, respectively; scores of ≥ 10 indicate 54 depression and scores of \geq 36 indicate anxiety. We performed Poisson regression analyses with 55 robust variance to examine whether depression and anxiety were associated with smoking. 56 57 Results: One hundred and eighty participants with TB were enrolled from primary health clinics. 58 Among all enrollees, depression was reported in 46 (27.1%) participants, while anxiety was 59 reported in 60 (44.4%) participants. Overall, 45 (25.0%) participants reported current smoking, 60 and the median number of cigarettes per day was 10. Depressive symptoms were associated with 61 a higher prevalence of smoking (aPR: 1.82; 95% CI = 1.11, 3.01) after adjusting for sex, HIV 62 status, food insecurity, anxiety, and income. The association between anxiety symptoms and 63 cigarette smoking did not reach statistical significance (aPR 1.26; 95% CI: 0.78-2.05). 64 65 Conclusions: We found the association between depressive symptoms and smoking among TB 66 patients in Botswana to be significant and the association between anxiety symptoms and

67	cigarette smoking insignificant. Future studies should further investigate these associations when
68	addressing TB care.
69	
70	Key words (4-6): Depression, cigarette smoking cessation, tuberculosis, Botswana
71 72 73	Association between depressive symptoms and cigarette smoking among tuberculosis
74	patients in Botswana.
75	
76	Introduction
77	An estimated 1.4 million people died from tuberculosis (TB) globally in 2019; it is one of
78	the top ten global causes of death worldwide [1]. Tobacco smoking is associated with an
79	increased risk of TB infection and TB disease [2]. Research suggests smoking is associated with
80	more extensive lung disease at time of TB diagnosis [3], higher rates of TB infection [4], and
81	higher rates of TB treatment failures [5]. Despite evidence showing these harmful links between
82	smoking and TB, the World Health Organization [6] estimates that 1.3 billion people use tobacco
83	globally, and more than 80% of smokers live in low- and middle-income countries [7]. In
84	Botswana, an upper middle-income country, 18% of adults use tobacco and 14.2% of adults
85	smoke tobacco [8]. Further, depression is reportedly higher in TB patients than the general
86	population [9]. Studies suggest an association between depression and smoking [10-12] as well
87	as between anxiety and smoking [13, 14]; however, to our knowledge, these associations have
88	not been studied in TB-infected populations. With both smoking and depression being more
89	common in people with TB than in the general population [15], their association may be even
90	stronger with potential adverse effects on patient outcomes. This study explores the association

It is made available under a CC-BY-NC-ND 4.0 International license .

91 between smoking and common mental health issues among TB patients, including depression
92 and anxiety.

93

94 Methods

95 Study Design, Population and Site Description

96 The methods for this study of TB patients have been described in detail elsewhere [16, 97 17]. Briefly, in an effort to explore the association between depression, anxiety and cigarette 98 smoking in TB patients, this cross-sectional study was conducted from January to December, 99 2019 at 12 primary health care facilities in Gaborone, Botswana. Participants were eligible for 100 inclusion if they were at least 18 years of age and older, with a microbiologically confirmed TB 101 diagnosis, and were recruited by our research staff at the time of receiving TB diagnosis. All 102 patients diagnosed with TB at the participating facilities were consecutively enrolled for this 103 study. All TB patients regardless of disease site and drug susceptibility status were eligible.

104

105 Measures

106 The primary outcome variable was smoking status. For analysis of smoking habits, 107 participants were asked, "Are you a current smoker?" Current smoking was self-reported by 108 participants, in which case participants were categorized into two groups: smokers and non-109 smokers. The independent variables were depression and anxiety. Depressive symptoms were 110 assessed using the Patient Health Questionnaire-9. This nine-item instrument analyzes frequency 111 of depressive symptoms up to two weeks prior to taking the questionnaire, using a 4-point Likert 112 scale. Scores of 10 or higher indicate depressive symptoms [18]. The internal reliability of the 113 Patient Health Questionnaire-9 in the current study was acceptable, with a Cronbach's alpha of

It is made available under a CC-BY-NC-ND 4.0 International license .

114	0.74. This instrument has been validated in primary care clinic settings [18]. The Zung Self-
115	Rating Anxiety Scale is a 20-item self-rating instrument which assesses cognitive, autonomic,
116	motor and central nervous system symptoms of anxiety using a 4-point Likert-type scale ranging
117	from 'a little of the time,' 'some of the time,' 'good part of the time,' and 'most of the time' [19].
118	Scores of 36 or higher on the Zung scale was indicative of anxiety symptoms. The Zung scale
119	demonstrated acceptable internal reliability with a Cronbach's alpha of 0.80 in this study.
120	
121	We also administered a questionnaire to collect data on age, sex, income, education,
122	employment status, history of alcohol use, and smoking status. Clinical information collected
123	included TB diagnosis, TB symptoms and their duration, HIV status and antiretroviral treatment
124	history, and CD4 counts (from the national HIV database for HIV-infected participants).
125	Rifampin-resistant results from Xpert MTB/RIF were available for some participants.
126	
127	Food insecurity was evaluated by the Household Food Insecurity Access Scale. This
128	nine-item questionnaire distinguishes food-insecure households and determines the degree of
129	food insecurity using a Likert-type scale that ranged from 'never', 'rarely (1 or 2 times)',
130	'sometimes (3–10 times)', and 'often (more than 10 times)'.
131	
132	Data Collection
133	Potential participants were identified first from the lab and clinic register and were
134	approached for recruitment by Research Assistants, who then screened them for eligibility and
135	obtained informed consent. The Research Assistants read consent materials in English and/or
136	Setswana (official languages in Botswana) to patients who were illiterate; all participants verified

It is made available under a CC-BY-NC-ND 4.0 International license .

137 their consent to participate by signature. Data on socioeconomic, demographic characteristics

138 and clinical information were collected in one-time interviews using the standardized

139 questionnaires described above, prepared in both English and Setswana.

140

Participants whose total depressive symptom score corresponded with mild depression
(score 5-9) were given a contact sheet for psychiatric services to seek help if symptoms
worsened or persisted. Research Assistants helped make appointments for participants scoring a
10 or higher on the Patient Health Questionnaire-9 or those presenting with suicidal ideation.
This study was approved by the Institutional Review Board of the University of California,
Irvine and Botswana Ministry of Health Human Research Development Committee.

147

148 Statistical Analysis

149 We investigated whether depression or anxiety was associated with the primary outcome 150 of interest, smoking status, in people with TB. Participants were categorized into two exposure 151 groups: current smokers and non-smokers. Participants were identified as experiencing 152 depressive and anxiety symptoms if they scored ≥ 10 on Patient Health Questionnaire-9 and ≥ 36 153 on the Zung scale, respectively. To control for potential confounding, participants were 154 categorized as experiencing 'food security or mild insecurity' or 'moderate to severe insecurity 155 [20]. We also dichotomized participants into having "no monthly income" and "some income," 156 and having "no education" and "some education," as the variation between these categories was 157 minimal. We examined the bivariate associations of current smoking with age, education, sex, 158 monthly income, HIV status, food insecurity, anxiety, and depression. Poisson regression 159 modeling with robust variance was used to estimate prevalence ratios (PRs) and 95% confidence

It is made available under a CC-BY-NC-ND 4.0 International license .

160	intervals (95% CI) for bivariate and multivariable analysis with current smoking status specified
161	as the outcome variable [21, 22]. The multivariate Poisson regression model assessed the
162	association between mental health disorders and current smoking status adjusting for sex,
163	income, HIV status, and food insecurity. A sensitivity analysis was also conducted; the analysis
164	was restricted to men only as there were very few women identifying as smokers. We also
165	conducted a sensitivity analysis to assess whether alcohol consumption would confound our
166	results. Data were analyzed using SAS version 9.4 (SAS Institute, Cary, North Carolina, U.S.).
167	
168	Results
169	A total of 180 participants were included in the study; the majority were male (64.4%,
169 170	A total of 180 participants were included in the study; the majority were male (64.4%, n=116), and the mean age was 37 years. Among all enrollees, depression was reported in 46
169 170 171	A total of 180 participants were included in the study; the majority were male (64.4%, n=116), and the mean age was 37 years. Among all enrollees, depression was reported in 46 (27.1%) participants, while anxiety was reported in 60 (44.4%) participants. One quarter (n=45)
169 170 171 172	A total of 180 participants were included in the study; the majority were male (64.4%, n=116), and the mean age was 37 years. Among all enrollees, depression was reported in 46 (27.1%) participants, while anxiety was reported in 60 (44.4%) participants. One quarter (n=45) of participants identified as smokers. Smokers were more likely to be male (91.1%, n=41), and
169 170 171 172 173	A total of 180 participants were included in the study; the majority were male (64.4%, n=116), and the mean age was 37 years. Among all enrollees, depression was reported in 46 (27.1%) participants, while anxiety was reported in 60 (44.4%) participants. One quarter (n=45) of participants identified as smokers. Smokers were more likely to be male (91.1%, n=41), and smoked a median of ten cigarettes per day (67%, n=30). Compared to non-smokers, current
169 170 171 172 173 174	A total of 180 participants were included in the study; the majority were male (64.4%, n=116), and the mean age was 37 years. Among all enrollees, depression was reported in 46 (27.1%) participants, while anxiety was reported in 60 (44.4%) participants. One quarter (n=45) of participants identified as smokers. Smokers were more likely to be male (91.1%, n=41), and smoked a median of ten cigarettes per day (67%, n=30). Compared to non-smokers, current smokers were more likely to report depressive symptoms (PHQ-9 ≥10: 40.0% vs. 21.5%) and
 169 170 171 172 173 174 175 	A total of 180 participants were included in the study; the majority were male (64.4%, n=116), and the mean age was 37 years. Among all enrollees, depression was reported in 46 (27.1%) participants, while anxiety was reported in 60 (44.4%) participants. One quarter (n=45) of participants identified as smokers. Smokers were more likely to be male (91.1%, n=41), and smoked a median of ten cigarettes per day (67%, n=30). Compared to non-smokers, current smokers were more likely to report depressive symptoms (PHQ-9 ≥10: 40.0% vs. 21.5%) and anxiety symptoms (Zung ≥ 36: 55.6% vs. 44.4%). Moderate to severe food insecurity was also
 169 170 171 172 173 174 175 176 	A total of 180 participants were included in the study; the majority were male (64.4%, n=116), and the mean age was 37 years. Among all enrollees, depression was reported in 46 (27.1%) participants, while anxiety was reported in 60 (44.4%) participants. One quarter (n=45) of participants identified as smokers. Smokers were more likely to be male (91.1%, n=41), and smoked a median of ten cigarettes per day (67%, n=30). Compared to non-smokers, current smokers were more likely to report depressive symptoms (PHQ-9 \geq 10: 40.0% vs. 21.5%) and anxiety symptoms (Zung \geq 36: 55.6% vs. 44.4%). Moderate to severe food insecurity was also reported more in smokers than non-smokers (48.9% vs. 34.8%), as was alcohol use (100% vs.

178

In the unadjusted model, we found female participants were less likely than male
participants to be current smokers (cPR: 0.18; 95% CI: 0.07-0.47). Having income (cPR: 1.98;
95% CI: 1.05-3.73) and depressive symptoms (cPR:1.89; 95% CI: 1.15-3.10) were more likely to
be associated with current smoking (Table 2). After adjusting for HIV co-infection, income,

183	anxiety symptoms, and food insecurity, depressive symptoms (aPR: 1.82; 95% CI: 1.11-3.01)
184	and male sex (aPR: 0.18; 95% CI: 0.07-0.48) remained to be significantly associated with current
185	smoking (Table 2). Depression (aPR: 1.09; 95% CI: 1.03-1.15) and income (aPR: 2.15; 95% CI:
186	1.02-4.53), were each positively associated with smoking intensity. Compared to men, women
187	demonstrated lower smoking intensity (aPR: 0.18; 95% CI: 0.07-0.48). In our sensitivity
188	analyses, we found similar results for male participants only, (aPR: 2.22, 95%CI: 1.41-3.48), and
189	a reduced level of association among alcohol consumers only (aPR: 1.44, 95% CI: 1.00-2.07) for
190	the association between depression and smoking. The association between anxiety symptoms and
191	cigarette smoking did not reach statistical significance (aPR 1.26; 95% CI: 0.78-2.05).
192	
193	Discussion
194	In this study, depression was associated with current smoking among TB patients after
195	adjusting for potential confounders. Additionally, we found that men were more likely to engage
196	in smoking than women in this study. Patients with TB have reported both depression [23, 24]
197	and smoking during treatment of TB [3, 25], with both potentially leading to reduced treatment
198	success. In addition to the hypothesis that this relationship could be bidirectional, another
199	hypothesis for the association between depression and smoking is the desire to self-medicate
200	[26]. Nicotine causes the release of dopamine; when smoking stops, depressed mood can occur
201	as a result of nicotine withdrawal [27]. The negative effect of depression may be enhanced
202	among people with TB, as treatment for TB can take a minimum of six to nine months. Our
203	findings suggest that current efforts to help TB patients quit smoking may benefit from the
204	management of multi-morbidity, including mental health and smoking cessation counseling.

It is made available under a CC-BY-NC-ND 4.0 International license .

205	The findings of this study should be considered with several limitations. First, these data
206	are from a cross-sectional study, leaving the direction of causality between smoking and
207	depression unclear. However, other studies show consistent associations between depression and
208	smoking [28, 29]. Given the high proportion of smoking and depression among TB patients, a
209	more consistent approach would be addressing the multi-morbidities of smoking and depression
210	in TB treatment programs. Additionally, data about recent quit attempts was not collected. It is
211	possible people recently diagnosed with TB could have quit smoking upon learning of their TB
212	diagnosis and therefore classified as non-smokers. Further, because we used a convenience
213	sample in Botswana for this study, our results may not be generalizable to other TB populations.
214	Finally, some data collected in our study were based on self-report (anxiety scores, smoking
215	status), which may lead to misclassification of variables included in our model. However,
216	research staff were trained to minimize this bias via objective interview techniques; for questions
217	related to mental health disorders, participants were able to directly record their answers on the
218	questionnaire.

219

220 Conclusion

This study expands our understanding of the association between depression and smoking status among TB patients in Botswana. Additionally, this study adds evidence to the need for specific public health efforts including smoking cessation interventions as part of TB treatment programs. Previous programs have evaluated smoking cessation as part of the core treatment program for people with TB [30], and should be considered in future TB treatment programs. Mental health screenings should also be included in treatment of TB to provide person-centered care among these higher risk patients and further develop treatment plans for patients with TB

228	multimorbidity. Future studies should employ larger sample size in different settings to validate
229	our findings. Additionally, future research should review the course of depression and anxiety
230	and its association with smoking cessation during TB treatment [31].
231	
232 233	Statements
234	Informed Consent
235	Informed consent was obtained from all individual participants included in the study.
236	
237	Ethical Approval
238	All procedures performed in studies involving human participants were in accordance with the
239	ethical standards of the institutional and national research committee and with the 1964 Helsinki
240	declaration and its later amendments or comparable ethical standards. This study was approved
241	by the UCI IRB and the Botswana Ministry of Health Human Research Development Program.
242	
243	Welfare of Animals
244	This article does not contain any studies with animals performed by any of the authors.
245	
246	
247	
248	
249	
250	
251	

It is made available under a CC-BY-NC-ND 4.0 International license .

252 Table 1. Characteristics of study participants in Gaborone, Botswana, 2019 (n=180)

253

Characteristic		Non smoker	Smoker
		n/N (%) or median	n/N (%) or median
		(IQR)	(IQR)
Overall		135/180 (75.0)	45/180 (25.0)
Age (in years)		36.0 (27.0-46.0)	39.0 (30.0-43.0)
Number of cigarettes		NA	10.0 (5.0-20.0)
smoked per day			
HIV	Negative	60/135 (44.4)	21/45 (46.7)
	Positive	75/135 (55.6)	24/45 (53.3)
Food insecurity	None or mildly insecure	88/135 (65.2)	23/45 (51.1)
	Moderate to severe	47/135 (34.8)	22/45 (48.9)
Depression	No	106/135 (78.5)	27/45 (60.0)
	Yes	29/135 (21.5)	18/45 (40.0)
Anxiety	No	75/135 (55.6)	20/45 (44.4)
	Yes	60/135 (44.4)	25/45 (55.6)
Education	No	12/135 (8.9)	5/45 (11.1)
	Yes	123/135 (91.1)	40/45 (88.9)
Income	No	55/135 (40.7)	10/45 (22.2)
	Yes	80/135 (59.3)	35/45 (77.8)
Gender	Male	75/135 (55.6)	41/45 (91.1)
	Female	60/135 (44.4)	4/45 (8.9)
Alcohol	No	101/135 (74.8)	0
	Yes	34/135 (25.2)	45/45 (100.0)
CD4* (cells/mm3)	≤ 200	23/75 (30.7)	7/24 (29.2)
	> 200	40/75 (53.3)	15/24 (62.5)
	Missing	12/75 (16.0)	2/24 (8.3)
ART*	Never taken ART	25/75 (33.3)	6/24 (25.0)
	Taking ART	45/75 (60.0)	14/24 (58.3)
	Took ART but stopped	5/75 (6.7)	4/24 (16.7)

254 Abbreviations: IQR=Interquartile range; ART=Antiretroviral therapy

255 *Among HIV-positive patients only (n = 99)

256

It is made available under a CC-BY-NC-ND 4.0 International license .

258 Table 2. Correlates of current smoking (n=180)

HIVNegative 1.00 1.00 Positive $0.94 (0.56-1.55)$ $0.87 (0.54-1.39)$ Food insecurityNone or mildly 1.00 1.00 insecureModerate to $1.54 (0.93-2.54)$ $1.43 (0.85-2.40)$ SevereNo 1.00 1.00 DepressionNo 1.00 1.00 AnxietyNo 1.00 1.00 Yes $1.40 (0.84-2.33)$ $1.26 (0.78-2.05)$ EducationNo 1.00 Not includedYes $0.83 (0.38-1.83)$ $1.67 (0.92-3.04)$ IncomeNo 1.00 1.00 Yes $1.98 (1.05-3.73)$ $1.67 (0.92-3.04)$ GenderMale 1.00 1.00 Age $1.00 (0.99-1.02)$ Not includedAbbreviations: PR=Prevalence ratio; CI=Confidence intervalNot included	HIV I Food insecurity I Depression I Anxiety I Education I Income I Gender I Age Abbreviations: PR=Prev	Negative Positive None or mildly nsecure Moderate to severe No Yes No Yes No Yes No Yes No Yes Male Female	1.00 0.94 (0.56-1.55) 1.00 1.54 (0.93-2.54) 1.00 1.89 (1.15-3.10) 1.00 1.40 (0.84-2.33) 1.00 0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.00 0.87 (0.54-1.39) 1.00 1.43 (0.85-2.40) 1.00 1.82 (1.11-3.01) 1.00 1.26 (0.78-2.05) Not included 1.00 1.67 (0.92-3.04) 1.00
Positive $0.94 (0.56-1.55)$ $0.87 (0.54-1.39)$ Food insecurityNone or mildly 1.00 1.00 insecureModerate to $1.54 (0.93-2.54)$ $1.43 (0.85-2.40)$ severeNo 1.00 1.00 DepressionNo 1.00 1.00 Yes $1.89 (1.15-3.10)$ $1.82 (1.11-3.01)$ AnxietyNo 1.00 1.00 Yes $1.40 (0.84-2.33)$ $1.26 (0.78-2.05)$ EducationNo 1.00 Not includedYes $0.83 (0.38-1.83)$ $1.67 (0.92-3.04)$ IncomeNo 1.00 1.00 Yes $1.98 (1.05-3.73)$ $1.67 (0.92-3.04)$ GenderMale 1.00 1.00 Kes $1.00 (0.99-1.02)$ Not includedAge $1.00 (0.99-1.02)$ Not included	Food insecurity	Positive None or mildly nsecure Moderate to severe No Yes No Yes No Yes No Yes Male Female	0.94 (0.56-1.55) 1.00 1.54 (0.93-2.54) 1.00 1.89 (1.15-3.10) 1.00 1.40 (0.84-2.33) 1.00 0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	0.87 (0.54-1.39) 1.00 1.43 (0.85-2.40) 1.00 1.82 (1.11-3.01) 1.00 1.26 (0.78-2.05) Not included 1.00 1.67 (0.92-3.04) 1.00
Food insecurity None or mildly insecure 1.00 1.00 Moderate to severe 1.54 (0.93-2.54) 1.43 (0.85-2.40) Depression No 1.00 1.00 Yes 1.89 (1.15-3.10) 1.82 (1.11-3.01) Anxiety No 1.00 1.00 Yes 1.40 (0.84-2.33) 1.26 (0.78-2.05) Education No 1.00 Not included Yes 0.83 (0.38-1.83) 1.26 (0.79-2.05) Income No 1.00 1.00 Yes 0.83 (0.38-1.83) 1.67 (0.92-3.04) Gender Male 1.00 1.00 Female 0.18 (0.07-0.47) 0.18 (0.07-0.48) Age 1.00 (0.99-1.02) Not included	Food insecurity	None or mildly nsecure Moderate to severe No Yes No Yes No Yes No Yes Male Female	1.00 1.54 (0.93-2.54) 1.00 1.89 (1.15-3.10) 1.00 1.40 (0.84-2.33) 1.00 0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.00 1.43 (0.85-2.40) 1.00 1.82 (1.11-3.01) 1.00 1.26 (0.78-2.05) Not included 1.00 1.67 (0.92-3.04) 1.00
Moderate to severe $1.54 (0.93-2.54)$ $1.43 (0.85-2.40)$ 1.00 DepressionNo 1.00 1.00 Yes $1.89 (1.15-3.10)$ $1.82 (1.11-3.01)$ AnxietyNo 1.00 1.00 Yes $1.40 (0.84-2.33)$ $1.26 (0.78-2.05)$ EducationNo 1.00 Not includedYes $0.83 (0.38-1.83)$ $1.67 (0.92-3.04)$ IncomeNo 1.00 1.00 Yes $1.98 (1.05-3.73)$ $1.67 (0.92-3.04)$ GenderMale 1.00 1.00 Female $0.18 (0.07-0.47)$ $0.18 (0.07-0.48)$ Age $1.00 (0.99-1.02)$ Not included	Depression	Moderate to severe No Yes No Yes No Yes Male Female	1.54 (0.93-2.54) 1.00 1.89 (1.15-3.10) 1.00 1.40 (0.84-2.33) 1.00 0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.43 (0.85-2.40) 1.00 1.82 (1.11-3.01) 1.00 1.26 (0.78-2.05) Not included 1.00 1.67 (0.92-3.04) 1.00
Depression No 1.00 1.00 Yes 1.89 (1.15-3.10) 1.82 (1.11-3.01) Anxiety No 1.00 1.00 Anxiety No 1.00 1.00 Yes 1.40 (0.84-2.33) 1.26 (0.78-2.05) Education No 1.00 Not included Yes 0.83 (0.38-1.83) Not included Income No 1.00 1.00 Gender Male 1.00 1.00 Gender Male 1.00 1.00 Age 1.00 (0.99-1.02) Not included	Depression	No Yes No Yes No Yes Male Female	1.00 1.89 (1.15-3.10) 1.00 1.40 (0.84-2.33) 1.00 0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.00 1.82 (1.11-3.01) 1.00 1.26 (0.78-2.05) Not included 1.00 1.67 (0.92-3.04) 1.00
Yes $1.89 (1.15-3.10)$ $1.82 (1.11-3.01)$ AnxietyNo 1.00 1.00 Yes $1.40 (0.84-2.33)$ $1.26 (0.78-2.05)$ EducationNo 1.00 Not includedYes $0.83 (0.38-1.83)$ 1.00 IncomeNo 1.00 1.00 Yes $1.98 (1.05-3.73)$ $1.67 (0.92-3.04)$ GenderMale 1.00 1.00 Female $0.18 (0.07-0.47)$ $0.18 (0.07-0.48)$ Age $1.00 (0.99-1.02)$ Not included	Anxiety	Yes No Yes No Yes Male Female	1.89 (1.15-3.10) 1.00 1.40 (0.84-2.33) 1.00 0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.82 (1.11-3.01) 1.00 1.26 (0.78-2.05) Not included 1.00 1.67 (0.92-3.04) 1.00
Anxiety No 1.00 1.00 Yes 1.40 (0.84-2.33) 1.26 (0.78-2.05) Education No 1.00 Not included Yes 0.83 (0.38-1.83)	Anxiety	No Yes No Yes Yes Male Female	1.00 1.40 (0.84-2.33) 1.00 0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.00 1.26 (0.78-2.05) Not included 1.00 1.67 (0.92-3.04) 1.00
Yes 1.40 (0.84-2.33) 1.26 (0.78-2.05) Education No 1.00 Not included Yes 0.83 (0.38-1.83) 1.00 1.00 Income No 1.00 1.00 1.00 Yes 1.98 (1.05-3.73) 1.67 (0.92-3.04) 1.00 Gender Male 1.00 1.00 1.00 Female 0.18 (0.07-0.47) 0.18 (0.07-0.48) Not included Age 1.00 (0.99-1.02) Not included	Education	Yes No Yes No Yes Male Female	1.40 (0.84-2.33) 1.00 0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.26 (0.78-2.05) Not included 1.00 1.67 (0.92-3.04) 1.00
Education No 1.00 Not included Yes 0.83 (0.38-1.83) 1.00 1.00 Income No 1.00 1.00 1.00 Yes 1.98 (1.05-3.73) 1.67 (0.92-3.04) 1.00 Gender Male 1.00 1.00 1.00 Female 0.18 (0.07-0.47) 0.18 (0.07-0.48) Not included Age 1.00 (0.99-1.02) Not included	Education	No Yes No Yes Male Female	1.00 0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	Not included 1.00 1.67 (0.92-3.04) 1.00
Yes 0.83 (0.38-1.83) Income No 1.00 1.00 Yes 1.98 (1.05-3.73) 1.67 (0.92-3.04) Gender Male 1.00 1.00 Female 0.18 (0.07-0.47) 0.18 (0.07-0.48) Age 1.00 (0.99-1.02) Not included	Income	Yes No Yes Male Female	0.83 (0.38-1.83) 1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.00 1.67 (0.92-3.04) 1.00
Income No 1.00 1.00 Yes 1.98 (1.05-3.73) 1.67 (0.92-3.04) Gender Male 1.00 1.00 Female 0.18 (0.07-0.47) 0.18 (0.07-0.48) Age 1.00 (0.99-1.02) Not included	Income	No Yes Male Female	1.00 1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.00 1.67 (0.92-3.04) 1.00
Yes 1.98 (1.05-3.73) 1.67 (0.92-3.04) Gender Male 1.00 1.00 Female 0.18 (0.07-0.47) 0.18 (0.07-0.48) Age 1.00 (0.99-1.02) Not included	Gender I Age Abbreviations: PR=Prev	Yes Male Female	1.98 (1.05-3.73) 1.00 0 18 (0 07-0 47)	1.67 (0.92-3.04) 1.00
GenderMale1.001.00Female0.18 (0.07-0.47)0.18 (0.07-0.48)Age1.00 (0.99-1.02)Not includedAbbreviations: PR=Prevalence ratio; CI=Confidence interval	Gender I Age Abbreviations: PR=Prev	Male Female	1.00 0 18 (0 07-0 47)	1.00
Female 0.18 (0.07-0.47) 0.18 (0.07-0.48) Age 1.00 (0.99-1.02) Not included Abbreviations: PR=Prevalence ratio; CI=Confidence interval Vertice Vertice	Age Abbreviations: PR=Prev	Female	0 18 (0 07-0 47)	
Age1.00 (0.99-1.02)Not includedAbbreviations: PR=Prevalence ratio; CI=Confidence interval	Age Abbreviations: PR=Prev			0.18 (0.07-0.48)
Abbreviations: PR=Prevalence ratio; CI=Confidence interval	Abbreviations: PR=Prev		1.00 (0.99-1.02)	Not included

It is made available under a CC-BY-NC-ND 4.0 International license .

281 **References**

- 282 1. World Health Organization. Tuberculosis 2020 [Available from: <u>https://www.who.int/news-</u>
 283 room/fact-sheets/detail/tuberculosis.
- 284
 2. Lin HH, Ezzati M, Murray M. Tobacco smoke, indoor air pollution and tuberculosis: a
 285 systematic review and meta-analysis. *PLoS Med.* 2007;4(1):e20.
- 286 3. Leung CC, Yew WW, Chan CK, Chang KC, Law WS, Lee SN, et al. Smoking adversely
 287 affects treatment response, outcome and relapse in tuberculosis. *Eur Respir J*.
 288 2015;45(3):738-45.
- 4. Alavi-Naini R, Sharifi-Mood B, Metanat M. Association between tuberculosis and smoking.
 Int J High Risk Behav Addict. 2012;1(2):71-4.
- 5. Mahishale V, Patil B, Lolly M, Eti A, Khan S. Prevalence of Smoking and Its Impact on
 Treatment Outcomes in Newly Diagnosed Pulmonary Tuberculosis Patients: A Hospital Based Prospective Study. *Chonnam Med J.* 2015;51(2):86-90.
- 6. World. Health Organization. WHO Report on the global tobacco epidemic 2019: Offer help to
 quit tobacco use 2019 [Available from: <u>https://www.who.int/teams/health-</u>
 promotion/tobacco-control/who-report-on-the-global-tobacco-epidemic-2019.
- 7. World Health Organization. Tobacco 2020 [Available from: <u>https://www.who.int/health-</u>
 topics/tobacco#tab=tab 2.
- 8. Republic of Botswana WHOCfDCP. GLOBAL ADULT TOBACCO SURVEY. 2018.
- 300 9. Duko B, Bedaso A, Ayano G. The prevalence of depression among patients with tuberculosis:
 301 a systematic review and meta-analysis. *Annals of general psychiatry*. 2020;19(1):30.
- 302 10. Ojo-Fati O, Thomas JL, Vogel RI, Ogedegbe O, Jean-Louis G, Okuyemi KS. Predictors of
 303 Adherence to Nicotine Replacement Therapy (Nicotine Patch) Among Homeless Persons
 304 Enrolled in a Randomized Controlled Trial Targeting Smoking Cessation. *J Fam Med.* 305 2016;3(7).
- 11. Harris T, Winetrobe H, Rhoades H, Wenzel S. The Role of Mental Health and Substance Use
 in Homeless Adults' Tobacco Use and Cessation Attempts. *J Dual Diagn*. 2019;15(2):76 87.
- 309 12. Weinberger AH, Kashan RS, Shpigel DM, Esan H, Taha F, Lee CJ, et al. Depression and
 310 cigarette smoking behavior: A critical review of population-based studies. *Am J Drug* 311 *Alcohol Abuse*. 2017;43(4):416-31.
- 312 13. Stanton R, To QG, Khalesi S, Williams SL, Alley SJ, Thwaite TL, et al. Depression, Anxiety
 313 and Stress during COVID-19: Associations with Changes in Physical Activity, Sleep,
 314 Tobacco and Alcohol Use in Australian Adults. *Int J Environ Res Public Health*.
 315 2020;17(11).
- 316 14. Zvolensky MJ, Bakhshaie J, Norton PJ, Smits JAJ, Buckner JD, Garey L, et al. Visceral
 317 sensitivity, anxiety, and smoking among treatment-seeking smokers. *Addict Behav.*318 2017;75:1-6.
- 319 15. Marshall AM, Barua D, Mitchell A, Keding A, Huque R, Khan A, et al. Smoking prevalence
 320 among tuberculosis patients: A crosssectional study in Bangladesh and Pakistan. *Tob* 321 *Induc Dis.* 2020;18:70.

322 16. Molebatsi K, Wang Q, Dima M, Ho-Foster A, Modongo C, Zetola N, et al. Depression and 323 delayed tuberculosis treatment initiation among newly diagnosed patients in Botswana. 324 *Glob Public Health*. 2020:1-11.

325 17. Wang Q, Dima M, Ho-Foster A, Molebatsi K, Modongo C, Zetola NM, et al. The association 326 of household food insecurity and HIV infection with common mental disorders among 327 newly diagnosed tuberculosis patients in Botswana. Public Health Nutr. 2020:1-9. 328 18. Kroenke K, Spitzer RL, Williams JB. The PHQ 9: validity of a brief depression severity 329 measure. J Gen Intern Med. 2001;16(9):606-13. 330 19. Zung WW. A rating instrument for anxiety disorders. Psychosomatics. 1971;12(6):371. 331 20. Maxwell D, Coates J, Vaitla B, Maxwell D. How Do Different Indicators of Household Food 332 Security Compare? Empirical Evidence from Tigray2013. 333 21. Greenland S. Model-based Estimation of Relative Risks and Other Epidemiologic Measures 334 in Studies of Common Outcomes and in Case-Control Studies. Am J Epidemiol. 335 2004;160(4):301-5. 336 22. Zou G. A modified poisson regression approach to prospective studies with binary data. Am J 337 Epidemiol. 2004;159(7):702-6. 338 23. Amreen, Nbsp, Rizvi N. Frequency of Depression and Anxiety among Tuberculosis Patients. 339 Journal of Tuberculosis Research. 2016;04(04):183-90. 340 24. Shrestha P, Subba UK, Brouwer M, Sweetland AC. Depression among TB patients and 341 associated factors in Kathmandu Valley, Nepal. Glob Ment Health (Camb). 2020;7:e4. 342 25. Altet N, Latorre I, Jiménez - Fuentes MÁ, Maldonado J, Molina I, González - Díaz Y, et al. 343 Assessment of the influence of direct tobacco smoke on infection and active TB 344 management. info:eu-repo/grantAgreement/ISCIII/PI 13/01546. 345 26. Fluharty M, Taylor AE, Grabski M, Munafò MR. The Association of Cigarette Smoking 346 With Depression and Anxiety: A Systematic Review. Nicotine Tob Res. 2017;19(1):3-13. 347 27. Benowitz NL. Pharmacology of nicotine: addiction, smoking-induced disease, and 348 therapeutics. Annu Rev Pharmacol Toxicol. 2009;49:57-71. 349 28. Ellis EM, Orom H, Giovino GA, Kiviniemi MT. Relations between negative affect and 350 health behaviors by race/ethnicity: Differential effects for symptoms of depression and 351 anxiety. Health Psychol. 2015;34(9):966-9. 352 29. Stubbs B, Vancampfort D, Firth J, Solmi M, Siddigi N, Smith L, et al. Association between 353 depression and smoking: A global perspective from 48 low- and middle-income 354 countries. J Psychiatr Res. 2018;103:142-9. 355 30. Siddiqi K, Khan A, Ahmad M, Dogar O, Kanaan M, Newell JN, et al. Action to stop 356 smoking in suspected tuberculosis (ASSIST) in Pakistan: a cluster randomized, 357 controlled trial. Ann Intern Med. 2013;158(9):667-75. 358 31. Siddiqi K, Lin Y, Elsey H, Siddiqi N. TB multimorbidity: a global health challenge 359 demanding urgent attention. International Journal of Tubrculosis & Lung Disease. 360 2021;25(2):87-90. 361 362