

1 **Depressive symptoms are associated with cigarette smoking among tuberculosis patients in**

2 **Botswana**

3

4 Alexandria Jones-Patten MSN, RN. Sue & Bill Gross School of Nursing, University of

5 California Irvine, Irvine, CA, USA;

6 Qiao Wang MPH: Fielding School of Public Health, University of California Los Angeles, Los

7 Angeles, CA, USA;

8 Keneilwe Molebatsi. Department of Psychiatry, Faculty of Medicine, University of Botswana,

9 Gaborone, Botswana;

10 Thomas E. Novotny MD, MPH: School of Public Health, San Diego State University, San

11 Diego, CA, USA;

12 Kamran Siddiqi MBBS, MRCP, MPH, FFPH, PhD. Department of Health Sciences, University

13 of York, York, England;

14 Chawangwa Modongo: Victus Global Botswana Organisation, Gaborone, Botswana;

15 Nicola M. Zetola MD, MPH: Victus Global Botswana Organisation, Gaborone, Botswana;

16 Bontle Mbongwe, PhD: Department of Environmental Health, Faculty of Health Sciences,

17 University of Botswana, Gaborone, Botswana;

18 Sanghyuk S. Shin PhD. Sue & Bill Gross School of Nursing, University of California Irvine,

19 Irvine, CA, USA.

20

21

22

23

24 **Corresponding Author:**

25 Sanghyuk S. Shin, PhD

26 Associate Professor, Sue & Bill Gross School of Nursing

27 Director, UCI Infectious Disease Science Initiative

28 University of California Irvine

29 106F Berk Hall

30 Irvine, CA 92697

31 sshin2@uci.edu

32 Tel: 949-576-8675

33

34

35 **Acknowledgements**

36 We would like to thank study participants for their time and efforts. We would also like to thank

37 University of California, Irvine and Botswana Ministry of Health and Wellness Human Research

38 Development Committee. Dr. Siddiqi was funded by Medical Research Council (UK) from TB

39 Multimorbidity Grant (Ref No: MC_PC_MR/T037806/1).

40

41 **Conflict of Interest Declaration**

42 All authors declare we have no conflict of interest.

43

44

45 **Abstract**

46 Background: Researchers have increasingly recognized the adverse effects of smoking on
47 tuberculosis (TB) outcomes. Smoking may be a maladaptive coping mechanism for depression
48 and anxiety among TB patients; however, this association has not yet been investigated.

49
50 Design/Methods: We conducted a cross-sectional study among newly diagnosed TB patients
51 between January and December 2019 in Gaborone, Botswana, and evaluated factors associated
52 with cigarette smoking. Using the Patient Health Questionnaire-9 and the Zung Self-Rating
53 Anxiety scale, we collected depression and anxiety scores, respectively; scores of ≥ 10 indicate
54 depression and scores of ≥ 36 indicate anxiety. We performed Poisson regression analyses with
55 robust variance to examine whether depression and anxiety were associated with smoking.

56
57 Results: One hundred and eighty participants with TB were enrolled from primary health clinics.
58 Among all enrollees, depression was reported in 46 (27.1%) participants, while anxiety was
59 reported in 60 (44.4%) participants. Overall, 45 (25.0%) participants reported current smoking,
60 and the median number of cigarettes per day was 10. Depressive symptoms were associated with
61 a higher prevalence of smoking (aPR: 1.82; 95% CI = 1.11, 3.01) after adjusting for sex, HIV
62 status, food insecurity, anxiety, and income. The association between anxiety symptoms and
63 cigarette smoking did not reach statistical significance (aPR 1.26; 95% CI: 0.78-2.05).

64
65 Conclusions: We found the association between depressive symptoms and smoking among TB
66 patients in Botswana to be significant and the association between anxiety symptoms and

67 cigarette smoking insignificant. Future studies should further investigate these associations when
68 addressing TB care.

69

70 **Key words (4-6):** Depression, cigarette smoking cessation, tuberculosis, Botswana

71

72

73 **Association between depressive symptoms and cigarette smoking among tuberculosis**

74 **patients in Botswana.**

75

76 Introduction

77 An estimated 1.4 million people died from tuberculosis (TB) globally in 2019; it is one of
78 the top ten global causes of death worldwide [1]. Tobacco smoking is associated with an
79 increased risk of TB infection and TB disease [2]. Research suggests smoking is associated with
80 more extensive lung disease at time of TB diagnosis [3], higher rates of TB infection [4], and
81 higher rates of TB treatment failures [5]. Despite evidence showing these harmful links between
82 smoking and TB, the World Health Organization [6] estimates that 1.3 billion people use tobacco
83 globally, and more than 80% of smokers live in low- and middle-income countries [7]. In
84 Botswana, an upper middle-income country, 18% of adults use tobacco and 14.2% of adults
85 smoke tobacco [8]. Further, depression is reportedly higher in TB patients than the general
86 population [9]. Studies suggest an association between depression and smoking [10-12] as well
87 as between anxiety and smoking [13, 14]; however, to our knowledge, these associations have
88 not been studied in TB-infected populations. With both smoking and depression being more
89 common in people with TB than in the general population [15], their association may be even
90 stronger with potential adverse effects on patient outcomes. This study explores the association

91 between smoking and common mental health issues among TB patients, including depression
92 and anxiety.

93

94 Methods

95 Study Design, Population and Site Description

96 The methods for this study of TB patients have been described in detail elsewhere [16,
97 17]. Briefly, in an effort to explore the association between depression, anxiety and cigarette
98 smoking in TB patients, this cross-sectional study was conducted from January to December,
99 2019 at 12 primary health care facilities in Gaborone, Botswana. Participants were eligible for
100 inclusion if they were at least 18 years of age and older, with a microbiologically confirmed TB
101 diagnosis, and were recruited by our research staff at the time of receiving TB diagnosis. All
102 patients diagnosed with TB at the participating facilities were consecutively enrolled for this
103 study. All TB patients regardless of disease site and drug susceptibility status were eligible.

104

105 Measures

106 The primary outcome variable was smoking status. For analysis of smoking habits,
107 participants were asked, “Are you a current smoker?” Current smoking was self-reported by
108 participants, in which case participants were categorized into two groups: smokers and non-
109 smokers. The independent variables were depression and anxiety. Depressive symptoms were
110 assessed using the Patient Health Questionnaire-9. This nine-item instrument analyzes frequency
111 of depressive symptoms up to two weeks prior to taking the questionnaire, using a 4-point Likert
112 scale. Scores of 10 or higher indicate depressive symptoms [18]. The internal reliability of the
113 Patient Health Questionnaire-9 in the current study was acceptable, with a Cronbach’s alpha of

114 0.74. This instrument has been validated in primary care clinic settings [18]. The Zung Self-
115 Rating Anxiety Scale is a 20-item self-rating instrument which assesses cognitive, autonomic,
116 motor and central nervous system symptoms of anxiety using a 4-point Likert-type scale ranging
117 from ‘a little of the time,’ ‘some of the time,’ ‘good part of the time,’ and ‘most of the time’ [19].
118 Scores of 36 or higher on the Zung scale was indicative of anxiety symptoms. The Zung scale
119 demonstrated acceptable internal reliability with a Cronbach’s alpha of 0.80 in this study.

120

121 We also administered a questionnaire to collect data on age, sex, income, education,
122 employment status, history of alcohol use, and smoking status. Clinical information collected
123 included TB diagnosis, TB symptoms and their duration, HIV status and antiretroviral treatment
124 history, and CD4 counts (from the national HIV database for HIV-infected participants).
125 Rifampin-resistant results from Xpert MTB/RIF were available for some participants.

126

127 Food insecurity was evaluated by the Household Food Insecurity Access Scale. This
128 nine-item questionnaire distinguishes food-insecure households and determines the degree of
129 food insecurity using a Likert-type scale that ranged from ‘never’, ‘rarely (1 or 2 times)’,
130 ‘sometimes (3–10 times)’, and ‘often (more than 10 times)’.

131

132 Data Collection

133 Potential participants were identified first from the lab and clinic register and were
134 approached for recruitment by Research Assistants, who then screened them for eligibility and
135 obtained informed consent. The Research Assistants read consent materials in English and/or
136 Setswana (official languages in Botswana) to patients who were illiterate; all participants verified

137 their consent to participate by signature. Data on socioeconomic, demographic characteristics
138 and clinical information were collected in one-time interviews using the standardized
139 questionnaires described above, prepared in both English and Setswana.

140

141 Participants whose total depressive symptom score corresponded with mild depression
142 (score 5-9) were given a contact sheet for psychiatric services to seek help if symptoms
143 worsened or persisted. Research Assistants helped make appointments for participants scoring a
144 10 or higher on the Patient Health Questionnaire-9 or those presenting with suicidal ideation.
145 This study was approved by the Institutional Review Board of the University of California,
146 Irvine and Botswana Ministry of Health Human Research Development Committee.

147

148 Statistical Analysis

149 We investigated whether depression or anxiety was associated with the primary outcome
150 of interest, smoking status, in people with TB. Participants were categorized into two exposure
151 groups: current smokers and non-smokers. Participants were identified as experiencing
152 depressive and anxiety symptoms if they scored ≥ 10 on Patient Health Questionnaire-9 and ≥ 36
153 on the Zung scale, respectively. To control for potential confounding, participants were
154 categorized as experiencing ‘food security or mild insecurity’ or ‘moderate to severe insecurity
155 [20]. We also dichotomized participants into having “no monthly income” and “some income,”
156 and having “no education” and “some education,” as the variation between these categories was
157 minimal. We examined the bivariate associations of current smoking with age, education, sex,
158 monthly income, HIV status, food insecurity, anxiety, and depression. Poisson regression
159 modeling with robust variance was used to estimate prevalence ratios (PRs) and 95% confidence

160 intervals (95% CI) for bivariate and multivariable analysis with current smoking status specified
161 as the outcome variable [21, 22]. The multivariate Poisson regression model assessed the
162 association between mental health disorders and current smoking status adjusting for sex,
163 income, HIV status, and food insecurity. A sensitivity analysis was also conducted; the analysis
164 was restricted to men only as there were very few women identifying as smokers. We also
165 conducted a sensitivity analysis to assess whether alcohol consumption would confound our
166 results. Data were analyzed using SAS version 9.4 (SAS Institute, Cary, North Carolina, U.S.).

167

168 Results

169 A total of 180 participants were included in the study; the majority were male (64.4%,
170 n=116), and the mean age was 37 years. Among all enrollees, depression was reported in 46
171 (27.1%) participants, while anxiety was reported in 60 (44.4%) participants. One quarter (n=45)
172 of participants identified as smokers. Smokers were more likely to be male (91.1%, n=41), and
173 smoked a median of ten cigarettes per day (67%, n=30). Compared to non-smokers, current
174 smokers were more likely to report depressive symptoms (PHQ-9 ≥ 10 : 40.0% vs. 21.5%) and
175 anxiety symptoms (Zung ≥ 36 : 55.6% vs. 44.4%). Moderate to severe food insecurity was also
176 reported more in smokers than non-smokers (48.9% vs. 34.8%), as was alcohol use (100% vs.
177 25.2%) respectively (Table 1).

178

179 In the unadjusted model, we found female participants were less likely than male
180 participants to be current smokers (cPR: 0.18; 95% CI: 0.07-0.47). Having income (cPR: 1.98;
181 95% CI: 1.05-3.73) and depressive symptoms (cPR:1.89; 95% CI: 1.15-3.10) were more likely to
182 be associated with current smoking (Table 2). After adjusting for HIV co-infection, income,

183 anxiety symptoms, and food insecurity, depressive symptoms (aPR: 1.82; 95% CI: 1.11-3.01)
184 and male sex (aPR: 0.18; 95% CI: 0.07-0.48) remained to be significantly associated with current
185 smoking (Table 2). Depression (aPR: 1.09; 95% CI: 1.03-1.15) and income (aPR: 2.15; 95% CI:
186 1.02-4.53), were each positively associated with smoking intensity. Compared to men, women
187 demonstrated lower smoking intensity (aPR: 0.18; 95% CI: 0.07-0.48). In our sensitivity
188 analyses, we found similar results for male participants only, (aPR: 2.22, 95%CI: 1.41-3.48), and
189 a reduced level of association among alcohol consumers only (aPR: 1.44, 95% CI: 1.00-2.07) for
190 the association between depression and smoking. The association between anxiety symptoms and
191 cigarette smoking did not reach statistical significance (aPR 1.26; 95% CI: 0.78-2.05).

192

193 Discussion

194 In this study, depression was associated with current smoking among TB patients after
195 adjusting for potential confounders. Additionally, we found that men were more likely to engage
196 in smoking than women in this study. Patients with TB have reported both depression [23, 24]
197 and smoking during treatment of TB [3, 25], with both potentially leading to reduced treatment
198 success. In addition to the hypothesis that this relationship could be bidirectional, another
199 hypothesis for the association between depression and smoking is the desire to self-medicate
200 [26]. Nicotine causes the release of dopamine; when smoking stops, depressed mood can occur
201 as a result of nicotine withdrawal [27]. The negative effect of depression may be enhanced
202 among people with TB, as treatment for TB can take a minimum of six to nine months. Our
203 findings suggest that current efforts to help TB patients quit smoking may benefit from the
204 management of multi-morbidity, including mental health and smoking cessation counseling.

205 The findings of this study should be considered with several limitations. First, these data
206 are from a cross-sectional study, leaving the direction of causality between smoking and
207 depression unclear. However, other studies show consistent associations between depression and
208 smoking [28, 29]. Given the high proportion of smoking and depression among TB patients, a
209 more consistent approach would be addressing the multi-morbidities of smoking and depression
210 in TB treatment programs. Additionally, data about recent quit attempts was not collected. It is
211 possible people recently diagnosed with TB could have quit smoking upon learning of their TB
212 diagnosis and therefore classified as non-smokers. Further, because we used a convenience
213 sample in Botswana for this study, our results may not be generalizable to other TB populations.
214 Finally, some data collected in our study were based on self-report (anxiety scores, smoking
215 status), which may lead to misclassification of variables included in our model. However,
216 research staff were trained to minimize this bias via objective interview techniques; for questions
217 related to mental health disorders, participants were able to directly record their answers on the
218 questionnaire.

219

220 Conclusion

221 This study expands our understanding of the association between depression and smoking
222 status among TB patients in Botswana. Additionally, this study adds evidence to the need for
223 specific public health efforts including smoking cessation interventions as part of TB treatment
224 programs. Previous programs have evaluated smoking cessation as part of the core treatment
225 program for people with TB [30], and should be considered in future TB treatment programs.
226 Mental health screenings should also be included in treatment of TB to provide person-centered
227 care among these higher risk patients and further develop treatment plans for patients with TB

228 multimorbidity. Future studies should employ larger sample size in different settings to validate
229 our findings. Additionally, future research should review the course of depression and anxiety
230 and its association with smoking cessation during TB treatment [31].

231

232

233 Statements

234 Informed Consent

235 Informed consent was obtained from all individual participants included in the study.

236

237 Ethical Approval

238 All procedures performed in studies involving human participants were in accordance with the
239 ethical standards of the institutional and national research committee and with the 1964 Helsinki
240 declaration and its later amendments or comparable ethical standards. This study was approved
241 by the UCI IRB and the Botswana Ministry of Health Human Research Development Program.

242

243 Welfare of Animals

244 This article does not contain any studies with animals performed by any of the authors.

245

246

247

248

249

250

251

252 **Table 1. Characteristics of study participants in Gaborone, Botswana, 2019 (n=180)**
 253

Characteristic		Non smoker n/N (%) or median (IQR)	Smoker n/N (%) or median (IQR)
Overall		135/180 (75.0)	45/180 (25.0)
Age (in years)		36.0 (27.0-46.0)	39.0 (30.0-43.0)
Number of cigarettes smoked per day		NA	10.0 (5.0-20.0)
HIV	Negative	60/135 (44.4)	21/45 (46.7)
	Positive	75/135 (55.6)	24/45 (53.3)
Food insecurity	None or mildly insecure	88/135 (65.2)	23/45 (51.1)
	Moderate to severe	47/135 (34.8)	22/45 (48.9)
Depression	No	106/135 (78.5)	27/45 (60.0)
	Yes	29/135 (21.5)	18/45 (40.0)
Anxiety	No	75/135 (55.6)	20/45 (44.4)
	Yes	60/135 (44.4)	25/45 (55.6)
Education	No	12/135 (8.9)	5/45 (11.1)
	Yes	123/135 (91.1)	40/45 (88.9)
Income	No	55/135 (40.7)	10/45 (22.2)
	Yes	80/135 (59.3)	35/45 (77.8)
Gender	Male	75/135 (55.6)	41/45 (91.1)
	Female	60/135 (44.4)	4/45 (8.9)
Alcohol	No	101/135 (74.8)	0
	Yes	34/135 (25.2)	45/45 (100.0)
CD4* (cells/mm ³)	≤ 200	23/75 (30.7)	7/24 (29.2)
	> 200	40/75 (53.3)	15/24 (62.5)
	Missing	12/75 (16.0)	2/24 (8.3)
ART*	Never taken ART	25/75 (33.3)	6/24 (25.0)
	Taking ART	45/75 (60.0)	14/24 (58.3)
	Took ART but stopped	5/75 (6.7)	4/24 (16.7)

254 Abbreviations: IQR=Interquartile range; ART=Antiretroviral therapy

255 *Among HIV-positive patients only (n = 99)

256

257

258 **Table 2. Correlates of current smoking (n=180)**

259

Characteristic		Crude PR (95% CI)	Adjusted PR (95% CI)
HIV	Negative	1.00	1.00
	Positive	0.94 (0.56-1.55)	0.87 (0.54-1.39)
Food insecurity	None or mildly insecure	1.00	1.00
	Moderate to severe	1.54 (0.93-2.54)	1.43 (0.85-2.40)
Depression	No	1.00	1.00
	Yes	1.89 (1.15-3.10)	1.82 (1.11-3.01)
Anxiety	No	1.00	1.00
	Yes	1.40 (0.84-2.33)	1.26 (0.78-2.05)
Education	No	1.00	Not included
	Yes	0.83 (0.38-1.83)	
Income	No	1.00	1.00
	Yes	1.98 (1.05-3.73)	1.67 (0.92-3.04)
Gender	Male	1.00	1.00
	Female	0.18 (0.07-0.47)	0.18 (0.07-0.48)
Age		1.00 (0.99-1.02)	Not included

260 Abbreviations: PR=Prevalence ratio; CI=Confidence interval

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281 **References**

- 282 1. World Health Organization. Tuberculosis 2020 [Available from: [https://www.who.int/news-](https://www.who.int/news-room/fact-sheets/detail/tuberculosis)
283 [room/fact-sheets/detail/tuberculosis](https://www.who.int/news-room/fact-sheets/detail/tuberculosis)].
- 284 2. Lin HH, Ezzati M, Murray M. Tobacco smoke, indoor air pollution and tuberculosis: a
285 systematic review and meta-analysis. *PLoS Med*. 2007;4(1):e20.
- 286 3. Leung CC, Yew WW, Chan CK, Chang KC, Law WS, Lee SN, et al. Smoking adversely
287 affects treatment response, outcome and relapse in tuberculosis. *Eur Respir J*.
288 2015;45(3):738-45.
- 289 4. Alavi-Naini R, Sharifi-Mood B, Metanat M. Association between tuberculosis and smoking.
290 *Int J High Risk Behav Addict*. 2012;1(2):71-4.
- 291 5. Mahishale V, Patil B, Lolly M, Eti A, Khan S. Prevalence of Smoking and Its Impact on
292 Treatment Outcomes in Newly Diagnosed Pulmonary Tuberculosis Patients: A Hospital-
293 Based Prospective Study. *Chonnam Med J*. 2015;51(2):86-90.
- 294 6. World. Health Organization. WHO Report on the global tobacco epidemic 2019: Offer help to
295 quit tobacco use 2019 [Available from: [https://www.who.int/teams/health-](https://www.who.int/teams/health-promotion/tobacco-control/who-report-on-the-global-tobacco-epidemic-2019)
296 [promotion/tobacco-control/who-report-on-the-global-tobacco-epidemic-2019](https://www.who.int/teams/health-promotion/tobacco-control/who-report-on-the-global-tobacco-epidemic-2019)].
- 297 7. World Health Organization. Tobacco 2020 [Available from: [https://www.who.int/health-](https://www.who.int/health-topics/tobacco#tab=tab_2)
298 [topics/tobacco#tab=tab_2](https://www.who.int/health-topics/tobacco#tab=tab_2)].
- 299 8. Republic of Botswana WHOCfDCP. GLOBAL ADULT TOBACCO SURVEY. 2018.
- 300 9. Duko B, Bedaso A, Ayano G. The prevalence of depression among patients with tuberculosis:
301 a systematic review and meta-analysis. *Annals of general psychiatry*. 2020;19(1):30.
- 302 10. Ojo-Fati O, Thomas JL, Vogel RI, Ogedegbe O, Jean-Louis G, Okuyemi KS. Predictors of
303 Adherence to Nicotine Replacement Therapy (Nicotine Patch) Among Homeless Persons
304 Enrolled in a Randomized Controlled Trial Targeting Smoking Cessation. *J Fam Med*.
305 2016;3(7).
- 306 11. Harris T, Winetrobe H, Rhoades H, Wenzel S. The Role of Mental Health and Substance Use
307 in Homeless Adults' Tobacco Use and Cessation Attempts. *J Dual Diagn*. 2019;15(2):76-
308 87.
- 309 12. Weinberger AH, Kashan RS, Shpigel DM, Esan H, Taha F, Lee CJ, et al. Depression and
310 cigarette smoking behavior: A critical review of population-based studies. *Am J Drug*
311 *Alcohol Abuse*. 2017;43(4):416-31.
- 312 13. Stanton R, To QG, Khalesi S, Williams SL, Alley SJ, Thwaite TL, et al. Depression, Anxiety
313 and Stress during COVID-19: Associations with Changes in Physical Activity, Sleep,
314 Tobacco and Alcohol Use in Australian Adults. *Int J Environ Res Public Health*.
315 2020;17(11).
- 316 14. Zvolensky MJ, Bakhshaie J, Norton PJ, Smits JAJ, Buckner JD, Garey L, et al. Visceral
317 sensitivity, anxiety, and smoking among treatment-seeking smokers. *Addict Behav*.
318 2017;75:1-6.
- 319 15. Marshall AM, Barua D, Mitchell A, Keding A, Huque R, Khan A, et al. Smoking prevalence
320 among tuberculosis patients: A cross-sectional study in Bangladesh and Pakistan. *Tob*
321 *Induc Dis*. 2020;18:70.
- 322 16. Molebatsi K, Wang Q, Dima M, Ho-Foster A, Modongo C, Zetola N, et al. Depression and
323 delayed tuberculosis treatment initiation among newly diagnosed patients in Botswana.
324 *Glob Public Health*. 2020:1-11.

- 325 17. Wang Q, Dima M, Ho-Foster A, Molebatsi K, Modongo C, Zetola NM, et al. The association
326 of household food insecurity and HIV infection with common mental disorders among
327 newly diagnosed tuberculosis patients in Botswana. *Public Health Nutr.* 2020;1-9.
- 328 18. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity
329 measure. *J Gen Intern Med.* 2001;16(9):606-13.
- 330 19. Zung WW. A rating instrument for anxiety disorders. *Psychosomatics.* 1971;12(6):371.
- 331 20. Maxwell D, Coates J, Vaitla B, Maxwell D. How Do Different Indicators of Household Food
332 Security Compare? Empirical Evidence from Tigray 2013.
- 333 21. Greenland S. Model-based Estimation of Relative Risks and Other Epidemiologic Measures
334 in Studies of Common Outcomes and in Case-Control Studies. *Am J Epidemiol.*
335 2004;160(4):301-5.
- 336 22. Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J*
337 *Epidemiol.* 2004;159(7):702-6.
- 338 23. Amreen, Nbsp, Rizvi N. Frequency of Depression and Anxiety among Tuberculosis Patients.
339 *Journal of Tuberculosis Research.* 2016;04(04):183-90.
- 340 24. Shrestha P, Subba UK, Brouwer M, Sweetland AC. Depression among TB patients and
341 associated factors in Kathmandu Valley, Nepal. *Glob Ment Health (Camb).* 2020;7:e4.
- 342 25. Altet N, Latorre I, Jiménez - Fuentes MÁ, Maldonado J, Molina I, González - Díaz Y, et al.
343 Assessment of the influence of direct tobacco smoke on infection and active TB
344 management. info:eu-repo/grantAgreement/ISCIII/PI 13/01546.
- 345 26. Fluharty M, Taylor AE, Grabski M, Munafò MR. The Association of Cigarette Smoking
346 With Depression and Anxiety: A Systematic Review. *Nicotine Tob Res.* 2017;19(1):3-13.
- 347 27. Benowitz NL. Pharmacology of nicotine: addiction, smoking-induced disease, and
348 therapeutics. *Annu Rev Pharmacol Toxicol.* 2009;49:57-71.
- 349 28. Ellis EM, Orom H, Giovino GA, Kiviniemi MT. Relations between negative affect and
350 health behaviors by race/ethnicity: Differential effects for symptoms of depression and
351 anxiety. *Health Psychol.* 2015;34(9):966-9.
- 352 29. Stubbs B, Vancampfort D, Firth J, Solmi M, Siddiqi N, Smith L, et al. Association between
353 depression and smoking: A global perspective from 48 low- and middle-income
354 countries. *J Psychiatr Res.* 2018;103:142-9.
- 355 30. Siddiqi K, Khan A, Ahmad M, Dogar O, Kanaan M, Newell JN, et al. Action to stop
356 smoking in suspected tuberculosis (ASSIST) in Pakistan: a cluster randomized,
357 controlled trial. *Ann Intern Med.* 2013;158(9):667-75.
- 358 31. Siddiqi K, Lin Y, Elsey H, Siddiqi N. TB multimorbidity: a global health challenge
359 demanding urgent attention. *International Journal of Tuberculosis & Lung Disease.*
360 2021;25(2):87-90.

361
362