It is made available under a CC-BY-NC-ND 4.0 International license .
DNA-METHYLATION AND MENTAL HEALTH
R

Raffington et al. 1

1. Title page

Title: Socially stratified DNA-methylation profiles are associated with disparities in child and adolescent mental health

Authors: Raffington^{1,2}, L., PhD, Tanksley^{1,2}, P., PhD, Vinnik¹, L., MSc, Sabhlok¹, A., MA,

Patterson³, M.W., PhD, Mallard, T.¹, PhD, Malanchini⁴, M., PhD, Ayorech⁵, Z., PhD, Tucker-

Drob^{1,2}, E.M., PhD, & Harden^{1,2}, K.P., PhD

Affiliations:

¹Department of Psychology, The University of Texas at Austin, Austin, TX, USA.

²Population Research Center, The University of Texas at Austin, Austin, TX, USA

³Department of Psychiatry, University of Colorado, Denver – Anschutz Medical Campus,

Aurora, CO, USA

⁴Department of Biological and Experimental Psychology, Queen Mary University of London, UK.

⁵PROMENTA Research Center, Department of Psychology, University of Oslo, Norway

Corresponding Authors: Laurel Raffington, University of Texas at Austin, 108 E. Dean Keeton Stop A8000, Austin, Texas 78712-1043. E-mail: laurel.raffington@austin.utexas.edu

Kathryn P. Harden, University of Texas at Austin, 108 E. Dean Keeton Stop A8000, Austin,

Texas 78712-1043. E-mail: harden@utexas.edu

These authors contributed equally: Elliot M. Tucker-Drob, Kathryn P. Harden

Manuscript word count (not including title, abstract, acknowledgment, references, tables, and figure legends, max 3000): 2995

It is made available under a CC-BY-NC-ND 4.0 International license.

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 2

Key Points

- 1. **Question**: We examined whether salivary DNA-methylation profiles are socially stratified and associated with child mental health.
- Findings: In this preregistered, cross-sectional observational study of 1,183 children and adolescents, socioeconomic, racial, and ethnic disparities in mental health were associated with salivary DNA-methylation profiles of inflammation and the pace of biological aging.
- 3. **Meaning**: DNA-methylation biomarkers hold promise as tools to quantify the biological impact of socioeconomic inequality and being racially minoritized in a manner that is tied to social disparities in mental health.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 3

2. Abstract

Importance: Economic and racial inequality is linked to disparities in children's mental health. Biomarkers that reflect these social disparities are lacking.

Objective: We examined the hypothesis that salivary DNA-methylation patterns of higher inflammation and faster pace of biological aging are economically, racially and ethnically stratified and are associated with child mental health.

Design: The Texas Twin Project is an on-going, observational, longitudinal study that began in

May 2012. Analyses were preregistered on May 7, 2021, and completed on August 23, 2021.

Setting: The population-based study identified and recruited participants from public school rosters in the greater Austin area.

Participants: Participants in the analytic data set included all participants that agreed to contribute DNA samples and whose samples were assayed by January 2021.

Exposures: Family- and neighborhood-level socioeconomic inequality, racial and ethnic identities (White, Latinx, Black, Asian).

Main Measure(s): Environmental exposures were analyzed in relation to salivary DNAmethylation profiles of higher inflammation (DNAm-CRP) and faster pace of biological aging (DunedinPoAm). Child internalizing problems, attention problems, aggression, rule-breaking, ADHD, oppositional defiant disorder, and conduct disorder were measured using parent-reports and self-reports on abbreviated versions of the Achenbach Child Behavior Checklist and Conners 3. The hypotheses being tested were formulated after data collection of the present data freeze and were pre-registered prior to analyses being conducted.

Results: In a sample of *N*=1,183 8-to-19-year-olds (609 female, age M=13.38y), children's salivary DNA-methylation profiles and psychiatric symptoms differed by socioeconomic

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 4

conditions, race and ethnicity. Children with more parent-reported internalizing symptoms had higher DNAm-CRP (r=0.15, 95% CI=0.05 to 0.25, P=0.004) and DunedinPoAm (r=0.15, CI=0.05 to 0.25, P=0.002), and children with more parent-reported aggression problems had higher DNAm-CRP (r=0.17, CI=0.04 to 0.31, P=0.013). DNAm-CRP partially mediated advantage of higher family socioeconomic status (16% of total effect) and White racial identity (12% of total effect) on reduced internalizing symptoms. DunedinPoAm also partially mediated advantage of White racial identity on internalizing (19% of total effect).

Conclusions and Relevance: Socioeconomic and racial inequality are visible in children's epigenetic profiles of inflammation and the rate of biological aging in a manner that is tied to social disparities in mental health.

Keywords: DNA-methylation; epigenetics; mental health; children; socioeconomic status; racism

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 5

3. Text

Children who experience environmental adversities, including financial scarcity and the discrimination, prejudice, and oppression associated with racial and ethnic marginalization, are at increased risk of experiencing symptoms of psychiatric disorders. Their mental health burden includes both problems with internalizing such as anxiety and depression, as well as behaviors characteristic of externalizing disorders like aggression and hyperactivity^{1–3}. In contrast, White racial identity may confer protective effects on children's mental health, as White children experience the generational legacy of state-sanctioned social power, resources, and favoritism lived by White people, *i.e.* White privilege^{4,5}. We capitalize these terms to highlight that racial and ethnic identities are social constructions that are not based on "innate" biosocial boundaries, but may have biosocial effects through people's lived experiences⁶ (**see Box 1**).

Biomarkers that reflect these social disparities are lacking in child populations. Here we test whether DNA-methylation alterations are sensitive to socioeconomic, racial, and ethnic disparities in mental health. Biomarkers developed and validated in adults that associate social disparities with children's mental health would be a novel tool to (1) improve forecasting of psychiatric illness from childhood to later life, (2) study the early-life social determinants of lifelong mental health, and (3) assess the efficacy of interventions and policies through a new surrogate outcome relevant for later life.

In the current study, we leverage results from *discovery* epigenome-wide studies of adults to create DNA-methylation composite scores that can be used for *prediction* of mental health outcomes and sensitivity to environmental inequality in our independent pediatric sample. Following a preregistered analysis plan (https://osf.io/t3vnp/), two salivary DNA-methylation composite scores and one genetic composite score were selected because their blood-derived

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 6

composites have been associated with psychiatric symptoms or they have been found to be sensitive to social inequality. First, we examined DNA-methylation profiles of a peripheral proxy for systemic low-grade inflammation (*i.e.*, C-reactive protein, CRP; DNAm-CRP⁷), which in blood samples have previously been found to be associated with internalizing and externalizing symptoms in children⁸. While inflammation is an essential component of immunosurveillance and host defense, a chronic low-grade inflammatory state is a pathological feature of a wide range of chronic health conditions that is also known to affect brain and psychological development^{9,10}. Second, we examined genetic profiles of inflammation (*i.e.*, polygenic scores of CRP, PGS-CRP¹¹), because there are genetic correlations of CRP with mood disorders in adults, such as PTSD¹² and depressive symptoms¹³ that may include causal effects of CRP on depression^{14,15}. Third, we previously reported that socioeconomic disadvantage and Latinx compared to White identity is associated with the pace of multi-system biological aging, as indicated by DunedinPoAm¹⁶, in an earlier sub-sample of salivary DNA-methylation data from the Texas Twin Project $(N=600)^{17}$. In the current analysis, we examine data from 1,183 children and adolescents from the Texas Twin Project aged 8-to-19-years.

Box 1. Racial and Ethnic Identity, Racism, and Genetic Ancestry.

Racial and ethnic identity, social and institutional racism, and genetic ancestry are not the same thing^{18,19}. Racial and ethnic groups as they are defined in the U.S. were pseudoscientific inventions used as a political tool benefitting White people to justify the enslavement of Africans and genocide of indigenous Americans^{4,5}. Social and institutional racism includes the generational legacy of state-sanctioned social power, resources, representation, and favoritism lived by White people, *i.e.* White privilege, which comes at the cost of marginalized racial and ethnic groups^{4,5}. Racial and ethnic identity is context-specific both in terms of how people self-identify²⁰ and how they are categorized by others²¹.

Racial and ethnic disparities in child mental health arise through various factors tied to classism and racism, including inequitable access to high-quality childcare, educational resources, healthcare, nutrition, and differences in exposure to toxicants, family stress, and neighborhood threat, among other factors^{22,23}. Importantly, racial and ethnic identity itself is not a form of adversity or risk factor for mental health, whereas the institutional racial discrimination and chronic nature of interpersonal discrimination associated with racial and ethnic marginalization is^{24,25}.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 7

In contrast, genetic ancestry describes patterns of gene frequencies that people have inherited from their genetic ancestors, and people's genetic ancestry does not change across social contexts^{18,26}. The lack of biological "reality" underlying race does not mean that race is unimportant: Racial and ethnic identity remains relevant as it pertains to people's lived experiences, culture, community, social challenges, and opportunity. Thus, these social constructs are not based on "innate" biosocial boundaries, but may have biosocial effects through people's lived experiences⁶.

Because we are integrating genetic, epigenetic, and social levels of analysis, we employ both labels based on biogeographical ancestry (like "predominantly recent European ancestries") as relevant to our genomic measure and socially based labels (like "White") that are more appropriate for population health disparities research²⁷.

Method

Sample

The Texas Twin Project is an ongoing longitudinal study²⁸. Participants in the current study were 1213 children and adolescents that had at least one DNA-methylation sample. 195 participants contributed two DNA-methylation samples (time between repeated samples: M=22 months, SD=6.5, range 3 to 38 months) and 16 samples were assayed in duplicate for reliability analyses (total methylation sample n=1424). After exclusions based on DNA-methylation quality control criteria, data from 1,183 (609 female) children, including 426 monozygotic and 757 dizygotic twins from 611 unique families, aged 8 to 19 years (age M=13.38y, SD=2.99y) was analyzed. Participants self-identified as White only (n=752), Latinx only (n=147), Latinx and White (n=97), Black and potentially another race/ethnicity (Black+, n=120), Asian and potentially another race/ethnicity (but not Latinx or Black; Asian+, n=90), and Indigenous American, Pacific Islander or other (but not Latinx, Black, or Asian; n=7). The University of Texas Institutional Review board granted ethical approval.

Measures

DNA-methylation

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 8

Saliva samples were collected during a laboratory visit using Oragene kits (DNA Genotek, Canada). The Infinium MethylationEPIC BeadChip kit (Illumina, Inc., USA) was used to assess methylation levels at 850,000 methylation sites. **Supplemental Methods** contain information on preprocessing, **Table 1** a description of DNA-methylation profiles calculation, and **Table 2** descriptive statistics.

Genetics

Genetic data was used to calculate a polygenic score for CRP (PGS-CRP; description in **Table 1**). The PGS-CRP is an approximate indicator of an individual's genetic liability for developing high levels of CRP²⁶. All PGS-CRP analyses were restricted to individuals solely of European ancestries in order to reduce the risk of spurious findings due to population stratification²⁶. **Supplemental Methods** contain information on genotyping.

Mental health

Multiple dimensions of child and adolescent mental health were measured using self-reports and parent-reports on abbreviated versions of the Achenbach Child Behavior Checklist^{29,30}, and the *DSM-IV* symptom count scales of the Conners 3³¹. **Table 1** contains a description of mental health composite measure calculation and **Table S1** a full list of items included in each measure.

Socioeconomic context

We measured children's socioeconomic disadvantage at the family and neighborhood levels of analysis (**Table 1**).

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 9

Developmental covariates and tobacco exposure

Body mass index³² (BMI), pubertal status^{33,34}, and tobacco exposure³⁵ have been associated with early life disadvantage as well as differential DNA-methylation patterns^{36–38}. We therefore consider these factors in our analysis as covariates (**Table 1**).

Statistical analyses

Fourteen mental health outcomes were examined in each model, including 7 self-reported and 7 parent-reported aggregate measures of (1) internalizing, (2) attention problems, (3) aggression, (4) rule-breaking, (5) ADHD, (6) oppositional defiant disorder (ODD), and (7) conduct disorder (CD).

We performed multilevel, multivariate regression models fit with FIML in *Mplus* 8.2 software³⁹. All models included a random intercept, representing the family-level intercept of the dependent variable, to correct for non-independence of twins. To account for nesting of repeated measures within individuals, a sandwich correction was applied to the standard errors in all analyses. All models included age, gender, and an age by gender interaction as covariates. We used the Benjamini-Hochberg false discovery rate method (FDR)⁴⁰ to account for multiple testing. We report standardized effect size estimates and nominal *P*-values as significant, when the FDR-corrected *P*-values were below alpha<0.05.

Results

Mental health is socially stratified in children

We observed socioeconomic disparities in parent- and child-reported psychiatric burden. After correcting for seven comparisons, family-level disadvantage was associated with more

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 10

parent-reported internalizing, attention problems, aggression, rule-breaking, ADHD, and ODD symptoms, but not CD symptoms (effect sizes r=0.11 to 0.26; Figure 1 and Supplemental Table S2). Family-level disadvantage was also associated with higher children's self-reported attention problems, aggression, and rule-breaking (effect sizes r=0.16 to 0.24; Figure 1 and

Supplemental Table S3).

Neighbourhood-level disadvantage was associated with higher parent-reported internalizing (r=0.13, CI=0.04 to 0.22, P=0.003) and attention problems (r=0.12, CI=0.03 to 0.20, P=0.008). Neighborhood opportunity was associated with lower parent-reported internalizing, attention problems, aggression, and ODD (effect sizes r=-0.12 to -0.19; **Figure 1** and **Table S2**). Neighbourhood-level disadvantage and less neighbourhood opportunity were both also associated with higher child-reported aggression, rule-breaking, and ODD (effect sizes r=0.13 to 0.16 and r=-0.13 to -0.16, respectively; **Figure 1** and **Table S3**).

We also observed racial/ethnic disparities in parent- and child-reported mental health. Children identifying as Latinx-only or Black+ had higher parent-reported internalizing and attention problems than White-only identifying children (effect sizes b=0.09 to 0.23; **Figure 1** and **Table S2**). Black+ identifying children also showed more parent-reported symptoms of ADHD. Asian+ identifying children had lower parent-reported ADHD, ODD, and CD than White-only children (effect sizes b=-0.06 to -0.11). Child-reported mental health was largely unrelated to their race/ethnicity except that Black+ identifying children self-reported higher rates of aggression than White-only children (b=0.15, CI=0.06 to 0.23, P=0.001).

We then examined the degree to which socioeconomic inequality statistically accounted for racial/ethnic disparities in mental health. Children reporting Black+ and Latinx-only identities lived in substantially more socioeconomically disadvantaged families and

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 11

neighborhoods compared to children reporting White-only identity (effect sizes *b*=0.25 to 0.43; **Table S4**). Family-level disadvantage accounted for differences between Latinx-only and White-only children's parent-reported internalizing, but not attention problems (**Table S2**). Family-level disadvantage accounted for differences between Black+ and White-only children's parent-reported ADHD and child-reported aggression, but not differences in parent-reported internalizing or attention problems (**Table S2 and S3**). After accounting for family-level disadvantage, Latinx-only children self-reported lower rates of CD than White-only children.

Linear regression analysis (not preregistered) indicated that White identity compared to racial/ethnic categories associated with marginalization was associated with less parent-reported internalizing (r=-0.16, 95% CI=-0.23 to -0.08, P<0.001) and fewer attention problems (r=-0.19, CI=-0.27 to -0.11, P<0.001), but not less aggression (r=-0.09, CI=-0.19 to 0.01, P=0.079), rule-breaking (r=-0.05, CI=-0.13 to 0.02, P=0.139), ADHD (r=-0.01, CI=-0.09 to 0.07, P=0.796), ODD (r=0.03, CI=-0.05 to 0.11, P=0.471), or CD (r=0.00, CI=-0.13 to 0.13, P=0.990). Though attenuated, White identity remained a statistically significant predictor of parent-reported internalizing (r=-0.09, CI=-0.16 to -0.02, P=0.012) and attention problems (r=-0.15, CI=-0.38 to -0.19, P<0.001) after accounting for family-level disadvantage (racial/ethnic identity and family-level disadvantage: r=-0.29, CI=-0.38 to -0.19, P<0.001). White identity also remained a statistic (r=-0.09, CI=-0.16 to -0.01, P=0.020) and attention problems (r=-0.15, CI=-0.23 to -0.19, P<0.001). White identity also remained a significant predictor of parent-reported internalizing (r=-0.29, CI=-0.38 to -0.19, P<0.001). White identity also remained a significant predictor of parent-reported internalizing (r=-0.15, CI=-0.23 to -0.06, P=0.001), after accounting for both family- and neighborhood-level socioeconomic disadvantage together (r=-0.34, CI=-0.42 to -0.26, P<0.001).

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 12





Salivary DNA-methylation profiles are socially stratified in children

Analyses of 15 technical replicates (one sample did not pass quality control) suggested moderate-to-good reliability of DNA-methylation profiles residualized for technical artifacts and cell composition (ICC for DNAm-CRP=0.73, DunedinPoAm=0.84; see **Table 1** for descriptive

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 13

statistics). Biometric models using the twin family structure, where the sum of additive genetic factors (*A*) and environmental factors shared by twins living in the same home (*C*) represents a lower bound estimate of reliability (because it does not include measurement error), also suggested good reliability of DNA-methylation profiles (A+C variation for DNAm-CRP= 60.7%, DunedinPoAm= 54.2%, accounting for age and gender).

Higher DNAm-CRP was correlated with higher DunedinPoAm (r=0.89, CI=0.81 to 0.96, P<0.001, accounting for age and gender). This is not surprising, as CRP levels were one of the 18 biomarkers that the pace of aging algorithm was trained on, and 7 CpG sites overlapped across measures¹⁶.

Older children had higher DNAm-CRP (r=0.35, CI=0.26 to 0.44, P<0.001) and DunedinPoAm profiles (r=0.13, CI=0.02 to 0.23, P=0.018). Boys had lower DNAm-CRP (b=-0.26, CI=-0.34 to -0.18, P<0.001) and DunedinPoAm (b=-0.18, CI=-0.27 to -0.10, P<0.001) profiles (b=0.06, CI= -0.02 to 0.14, P=0.143) as compared to girls. All models included age, gender, and an age by gender interaction as covariates.

We observed salivary DNA-methylation profiles that differed by socioeconomic and racial/ethnic inequality. As reported in Raffington et al.⁴¹, children from socioeconomically disadvantaged families, socioeconomically disadvantaged neighborhoods, neighborhoods with less neighborhood opportunity, and Latinx-only or Black+ identity relative to White-only identity exhibited DNA-methylation profiles associated with higher chronic inflammation and a faster pace of biological aging (effect sizes r=0.08 to 0.28; **Table S5**). The analyses of Raffington et al.⁴¹ were preregistered at the same time as the preregistration of the present study. See **Table S6** for effect size estimates between socioeconomic inequality and DNA-methylation profiles reported separately for each racial/ethnic group (not preregistered).

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 14

Salivary DNA-methylation profiles are associated with child and adolescent mental health

We assessed whether salivary DNA-methylation profiles were associated with child mental health. Higher DNAm-CRP was associated with higher parent-reported internalizing (r=0.15, CI=0.05 to 0.25, P=0.004) and aggression (r=0.17, CI=0.04 to 0.31, P=0.013; **Figure 2** and **Table S7**). Faster DunedinPoAm was also associated with higher parent-reported internalizing (r=0.15, CI=0.05 to 0.25, P=0.002).

Since both profiles were associated with parent-reported internalizing, we performed commonality analyses to examine the proportion of overlapping and unique variation explained. DNAm-CRP and DunedinPoAm explained largely unique variation in parent-reported internalizing (DNAm-CRP alone: 2.5%, DunedinPoAm alone: 2.2%, combined: 4.5%).

DNA-methylation profiles were not associated with children's self-reported mental health (**Table S8**). See **Table S9 and S10** for effect size estimates between DNA-methylation profiles and mental health reported separately for each racial/ethnic group (not preregistered).

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH



Figure 2. Associations between A) DNAm-CRP, B) PGS-CRP, and C) DunedinPoAm with parent-reported mental health in children and adolescents. DNA-methylation profiles and mental health values are in standard deviation units. Higher DNAm-CRP values indicate a methylation profile associated with higher chronic inflammation. Higher PGS-CRP values indicate a genetic profile associated with higher chronic inflammation. Higher DunedinPoAm values indicate a methylation profile associated with higher chronic inflammation. Higher DunedinPoAm values indicate a methylation profile associated with faster biological aging. See **Supplemental Table S7** for standardized regression coefficients.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 16

DNA-methylation partially accounts for associations of family-level disadvantage and racial and ethnic marginalization with mental health

We observed that DNAm-CRP and DunedinPoAm partially accounted for associations of family-level disadvantage and racial/ethnic disparities in mental health. DNAm-CRP, but not DunedinPoAm, partially mediated family-level disadvantage on internalizing (indirect effect 16% of total effect), but not on aggression. Both DNAm-CRP and DunedinPoAm partially mediated the impact of being racially minoritized compared to not on internalizing (indirect DNAm-CRP effect 12% of total effect; indirect DunedinPoAm effect 19% of total effect).

BMI statistically accounts for associations of DNA-methylation with mental health

We next examined the role of BMI, pubertal status, and DNAm-smoke in associations of DNA-methylation with mental health (see **Table S7**). Associations of DNAm-CRP and DunedinPoAm with mental health were largely accounted for by controlling for BMI, but not by controlling for pubertal status. Associations of DNAm-CRP with aggression were largely accounted for by controlling for DNAm-smoke.

Within-family analyses indicate family-level stratification of DNA-methylation

We further assessed the extent to which DNA-methylation associations with mental health are robust to complete genetic and family-level environmental control in a bivariate ACE model that used the twin family structure. We found no evidence to suggest that monozygotic twins who differ from their co-twins in DNA-methylation show corresponding differences in their mental health (see **Table S11**). This is consistent with the hypothesis that DNA-methylation

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 17

associations with mental health represents (potentially partially unmeasured) effects of familylevel stratification.

Polygenic scores of CRP are associated with mental health

In an analysis restricted to participants solely of European genetic ancestry, PGS-CRP was not correlated with DNAm-CRP (r=0.04, CI=-0.08 to 0.17, P=0.476) or DunedinPoAm (r=-0.04, CI=-0.18 to 0.10, P=0.563).

Higher PGS-CRP was associated with higher parent-reported internalizing (r=0.14, CI 0.04 to 0.23, P=0.004) and attention problems (r=0.16, CI=0.05 to 0.28, P=0.004; **Figure 1** and **Table S7**) and these associations were unaffected by controlling for BMI, pubertal status, DNAm-smoke, and family-level disadvantage. PGS-CRP was not associated with children's self-reported mental health (**Table S8**).

Commonality analyses showed that PGS-CRP explained largely unique variation in parent-reported internalizing (PGS-CRP alone: 2%) relative to both DNAm-CRP (combined: 4.5%) and DunedinPoAm (combined: 4.2%).

In exploratory preregistered analyses of 364 dizygotic twin pairs, PGS-CRP did not account for differences in internalizing (r=0.07, 95% CI=-0.06 to 0.20, P=0.316) or attention problems (r=0.13, 95% CI=-0.02 to 0.27, P=0.094), although we caution that effect size estimates warrant a repeated examination of this question in a larger sample.

Discussion

We analyzed epigenetic data from 1183 children and adolescents participating in the Texas Twin Project to examine whether salivary DNA-methylation patterns are socially stratified

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 18

and are associated with mental health. We find that salivary DNA-methylation profiles of inflammation (DNAm-CRP), faster pace of biological aging (DunedinPoAm) and parent-reported child and adolescent mental health, consistently differed by socioeconomic inequality and self-reported racial/ethnic identity. Additionally, these salivary DNA-methylation profiles and genetic profiles of inflammation (PGS-CRP) were associated with children's internalizing, aggression, and attention problems. Moreover, salivary DNA-methylation profiles partially accounted for child mental health disparities related to family-level disadvantage, racial, and ethnic identities.

Our findings linking environmental inequality to epigenetic profiles and mental health were most consistent for internalizing symptoms, which is in line with previous research on peripheral inflammation⁴² and a blood-based study on DNA-methylation of inflammation in children⁸. These results add to a growing body of research in animals and humans that links exposure to social adversity, DNA-methylation changes, and gene expression markers associated with inflammation^{10,43,44}. Animal research suggests that inflammation affects synaptogenesis, synaptic survival, and myelination, for instance in neural circuitries subserving executive control, and threat and reward processing that are pertinent to emotional and behavioral regulation⁴⁵. Further, inflammation and DNA-methylation alterations are environmentally-sensitive hallmarks of biological aging^{46–48}. We found that body mass index largely accounted for associations of DNAm-CRP and DunedinPoAm with mental health, which corresponds to new insights casting obesity as an inflammatory disease of accelerated biological aging⁴⁹.

Highlighting the distinctly harmful role of social and institutional racism, we found that even after accounting for the lower rates of socioeconomic disadvantage experienced by White children, White identifying children sustained a lower burden of mental health symptoms. Thus,

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 19

socioeconomic inequality captures some but not all of the pathways through which social and institutional racism impacts children's mental health²⁴. Moreover, our categorization of children's complex racial and ethnic identities that for instance grouped Black and Black-White identifying children together is likely to dilute and underestimate the bio-psychological effects of racial marginalization as the proximity to White privilege and experiences of marginalization are not the same between these identities.

Further research is needed to contextualize the biological and mental health effects of race-based discrimination, prejudice, and oppression experienced by marginalized youth compared to the favoritism and privilege experienced by Whites^{3,6,50}. While our findings are in line with the hypothesis that DNA-methylation alterations are one mechanism for how psychosocial stress is biologically embedded to injure children's psychiatric health, our observational design is, of course, not well-suited to evaluate mechanistic hypotheses. Randomized controlled trials and natural experiments that involve exogenous changes to various aspects of social inequality (*e.g.*, eliminating redlining, child tax credits) are necessary to establish whether children's salivary DNA-methylation profiles change in response to intervention in a way that is tied to mental health.

In contrast to results seen for parent-reported psychiatric symptoms, child-reported mental health was generally not associated with DNA-methylation or PGS-CRP profiles. Previous research in this sample has found that children and parents agree minimally on specific symptoms scales⁵¹. In addition, both children and parent self-reports can be influenced by their level of insight of their specific psychopathological burden, and the degree to which they have internalised narratives associated with being racially minoritized, *i.e.*, internalised racism⁵². We further acknowledge that restricting PGS-CRP analyses to participants solely of European

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 20

genetic ancestry is a major study limitation. Future research should explore whether recent approaches to compute ancestry deconvoluted polygenic scores⁵³ can be applied to PGS-CRP and validated with serum CRP in ancestrally diverse samples.

In conclusion, our findings suggest that salivary DNA-methylation patterns of higher inflammation and faster pace of biological aging are economically and racially stratified and are associated with child mental health. Because saliva can easily be collected in large-scale pediatric epidemiological studies and may indicate emerging health conditions, these DNAmethylation profiles can be employed in research seeking to understand and prevent economic and racial disparities in childhood mental health.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 21

Table 1. Description of study measures.

DNAm-CRP	DNAm-CRP was computed on the basis of an epigenome-wide association study of CRP ⁷ . Using the
	summary statistics of the associations between CpG sites and adult CRP, we created one methylation
	score per person by summing the product of the weight and the individual beta estimate for each
	individual at each of the 218 CpG sites significantly associated ($P < 1.15 \times 10^{-7}$) with CRP.
DunedinPoAm	DunedinPoAm was developed from DNA-methylation analysis of Pace of Aging in the Dunedin
	Study birth cohort. Pace of Aging is a composite phenotype derived from analysis of longitudinal
	change in 18 biomarkers of organ-system integrity measured when Dunedin Study members were all
	26, 32, and 38 years of age 17. Elastic-net regression machine learning analysis was used to fit Pace of
	Aging to Illumina 450k DNA-methylation data generated from blood samples collected when
	participants were aged 38 years. The elastic net regression produced a 46-CpG algorithm. Increments
	of DunedinPoAm correspond to "years" of physiological change occurring per 12-months of
	chronological time. The Dunedin Study mean was 1, <i>i.e.</i> the typical pace of aging among 38-year-olds
	in that birth cohort. Thus, 0.01 increment of DunedinPoAm corresponds to a percentage point increase
	or decrease in an individual's pace of aging relative to the Dunedin birth cohort at midlife.
	DunedinPoAm was calculated based on the published algorithm 10 using code available at
	https://github.com/danbelsky/DunedinPoAm38.
PGS-CRP	PGS-CRP was computed in two steps. First, GWAS summary statistics were adjusted for linkage
	disequilibrium, or LD (i.e., correlation structures in the genome that capture population stratification).
	The preregistered analysis plan proposed using SBayesR ⁵⁴ for LD-adjustment. However, as the GWAS
	summary statistics used to compute PGS-CRP did not meet the data requirements of SBayesR (e.g.,
	effect allele frequency, per SNP sample size), we elected to use PRScs for LD-adjustment instead.
	PRScs is a program that uses Bayesian regression to infer posterior SNP effects using continuous
	shrinkage priors. PRScs has been shown to improve prediction accuracy of PGSs over other widely used
	PGS approaches ⁵⁵ . PRScs requires GWAS summary statistics and an external reference panel of the
	same ancestry as the GWAS. For the summary statistics, we used publicly available data from a GWAS
	of CRP in 148,164 individuals solely of European ancestry ¹¹ . For the reference panel, we used the 1000
	Genomes Project ⁵⁶ European reference panel (phase 3 v5; provided with the software) that was
	restricted to HapMap3 SNPs ⁵⁷ .

Second, we used PLINK v2⁵⁸ to apply the LD-adjusted SNP effects from PRScs in the Texas Twins Project sample. The resulting PGS-CRP is described by the following equation:

$$PGS = \sum_{i}^{m} x_{i} \widehat{\beta}_{i}$$

where *m* is the number of SNPs, $\hat{\beta}$ is the estimated effect of the *i*th SNP and *x*, coded as 0, 1 or 2, is the number of effect alleles of the *i*th SNP. All PGS analyses were restricted to individuals solely of European ancestries in order to reduce the risk of spurious findings due to population stratification. PGS-CRP was residualized for the top five genetic principal components and genotyping batch and then standardized using Z-scores (M=1; SD=0).

Mental Health Self and parent-reported scales of CBCL and Conners were analyzed separately. Based on prior published work in this sample^{51,59,60}, CBCL items were averaged into means scores of internalizing (5 self-reported anxiety items, 6 depression items; 9 parent-reported anxiety items, 6 depression items, 4 somatic complaint items), aggression (13 self-reported items; 11 parent-reported items), rule-breaking (12 self-reported items; 5 parent-reported items), and attention problems (7 self-reported items; 6 parent-reported items). All items were rated on a 3-point scale ranging from 0 – *Not true* to 2 – *Very true or often true.*"

Conners items were coded as summed symptom counts of ADHD (self-reported 10 inattentive items and 11 hyperactive/impulsive items; parent-reported 9 inattentive items and 10 hyperactive/impulsive items; items correspond to 9 symptoms for inattention and 9 symptoms for hyperactivity-impulsivity), oppositional defiant disorder (8 self-reported items; 8 parent-reported; each item corresponds to one symptom), and conduct disorder (9 self-reported items; 12 parent-reported items; each item corresponds to one symptom). All items were rated on a 4-point scale, ranging from 0 - Not at all to 3 - Very often. Responses of 2 ("often") or 3 ("very often") on each item were classified as a symptom count. See **Table S1** for a full list of items included in each measure.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

 socioeconomic disadvantage Neighborhood -level socioeconomic disadvantage Neighborhood -level socioeconomic disadvantage Neighborhood -level socioeconomic disadvantage Socioeconomic disadvantage was computed from tract-level proportions of residents reported as
disadvantageSurvey Module 61, father absence, residential instability (changes in home address), and family receipt of public assistance. These were aggregated to form a composite measure of household-level cumulative socioeconomic disadvantage described in 62, and coded such that higher scores reflect greater disadvantage.Neighborhood -levelThe neighbourhood-level measure was composed from tract-level US Census data according to the method described in 62. Briefly, participant addresses were linked to tract-level data from the US Census Bureau American Community Survey averaged over five years (https://www.census.gov/programs-surveys/acs). A composite score of neighbourhood-level socioeconomic disadvantage was computed from tract-level proportions of residents reported as
 of public assistance. These were aggregated to form a composite measure of household-level cumulative socioeconomic disadvantage described in ⁶², and coded such that higher scores reflect greater disadvantage. Neighborhood -level socioeconomic disadvantage American Community Survey averaged over five years (https://www.census.gov/programs-surveys/acs). A composite score of neighbourhood-level as
Neighborhood -levelThe neighbourhood-level measure was composed from tract-level US Census data according to the method described in 62. Briefly, participant addresses were linked to tract-level data from the US Census Bureau American Community Survey averaged over five years (https://www.census.gov/programs-surveys/acs). A composite score of neighbourhood-level socioeconomic disadvantage was computed from tract-level proportions of residents reported as
NeighborhoodThe neighbourhood-level measure was composed from tract-level US Census data according to the method described in 62. Briefly, participant addresses were linked to tract-level data from the US Census Bureau American Community Survey averaged over five years (https://www.census.gov/programs-surveys/acs). A composite score of neighbourhood-level socioeconomic disadvantage was computed from tract-level proportions of residents reported as
NeighborhoodThe neighbourhood-level measure was composed from tract-level US Census data according to the method described in 62. Briefly, participant addresses were linked to tract-level data from the US Census Bureau American Community Survey averaged over five years (https://www.census.gov/programs-surveys/acs). A composite score of neighbourhood-level socioeconomic disadvantage was computed from tract-level proportions of residents reported as
-level method described in ⁶² . Briefly, participant addresses were linked to tract-level data from the US census Bureau American Community Survey averaged over five years (https://www.census.gov/programs-surveys/acs). A composite score of neighbourhood-level socioeconomic disadvantage was computed from tract-level proportions of residents reported as
socioeconomic disadvantage Census Bureau American Community Survey averaged over five years (https://www.census.gov/programs-surveys/acs). A composite score of neighbourhood-level socioeconomic disadvantage was computed from tract-level proportions of residents reported as
disadvantage (https://www.census.gov/programs-surveys/acs). A composite score of neighbourhood-level socioeconomic disadvantage was computed from tract-level proportions of residents reported as
socioeconomic disadvantage was computed from tract-level proportions of residents reported as
unemployed, living below the federal poverty threshold, having less than 12 years of education, not
being employed in a management position, and single mothers. These were aggregated to form a
neighbourhood-level socioeconomic disadvantage composite measure described in ⁶² , and coded such
that higher scores reflect greater disadvantage.
Neighborhood The neighborhood opportunity measure indexed the intergenerational economic mobility of children
opportunity of low-income parents. It examines average annual household income in 2014-15 of offspring (born
between 1978-1983, who are now in their mid-thirties) of low-income parents (defined as mean pre-
tax income at the household level across five years (1994, 1995, 1998-2000) at the 25th percentile of
the national income distribution, or \$27000/year) within each census tract. Household income was
obtained from federal tax return records between 1989-2015, the 2000 and 2010 Decennial Census
(US Census Bureau, 2000, 2010; https://data2.nhgis.org/main), and 2005-2015 American Community
Surveys (https://www.census.gov/programs-surveys/acs). Census tracts reflect where the child resided
through the age of 23. This data was compiled by and obtained from the Opportunity Atlas
(https://opportunityatlas.org: 53).
Body mass We measured BMI from in-laboratory measurements of height and weight transformed to gender- and
index (BMI) age-normed z-scores according to the method published by the US Centers for Disease Control and
Prevention (https://www.cdc.gov/growthcharts/percentile_data_files.htm).
Pubertal status Pubertal development was measured using children's self-reports on the Pubertal Development Scale
⁶⁴ . The scale assesses the extent of development across five sex-specific domains (for both: height,
body hair growth, skin changes; for girls: onset of menses, breast development; for boys: growth in
body hair, deepening of voice). A total pubertal status score was computed as the average response (1
= "Not vet begun" to $4 =$ "Has finished changing") across all items. Pubertal development was
residualized for age, gender, and an age by gender interaction.
Tobacco We indexed tobacco exposure using a DNA-methylation smoking (DNAm-smoke) score created by
exposure summing the product of the weight and the individual beta estimate for each individual at each CpG
site significantly associated with smoking in the discovery EWAS ³⁸ . Excluding self-reported tobacco
users (n=53) did not significantly alter results.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 23

Sample	п	М	SD
DNAm-CRP ^a	1365	0.03	0.02
DunedinPoAm ^a	1365	1.02	0.06
PGS-CRP ^b	654	0	1
Parent-reported internalizing	1311	3.05	0.39
Parent-reported attention problems	1311	3.27	0.50
Parent-reported aggression	805	3.02	0.46
Parent-reported rule-breaking	1311	3.14	0.70
Parent-reported ADHD	1266	2.39	3.87
Parent-reported oppositional defiant disorder	1266	0.63	1.39
Parent-reported conduct disorder	1266	0.06	0.32
Child-reported internalizing	1255	0.55	0.40
Child -reported attention problems	1255	0.72	0.43
Child -reported aggression	1255	0.39	0.27
Child -reported rule-breaking	1255	0.25	0.24
Child -reported ADHD	1104	4.45	3.99
Child -reported oppositional defiant disorder	1104	1.06	1.27
Child -reported conduct disorder	1104	0.11	0.46
Family-level socioeconomic disadvantage	993	-0.02	0.96
Neighborhood-level socioeconomic disadvantage	1218	-0.02	1
Neighborhood opportunity	950	0.31	0.62
Body mass index	1364	0.4	1.34
Pubertal development	1325	2.61	0.92
Tobacco use (yes/no)	58/631	_	_
DNAm-smoke	1365	-13.62	1.84

Table 2. Descriptive statistics.

^a After exclusion of participants based on DNA-methylation preprocessing (n=44), excluding technical replicates (n=15), and including repeated samples (n= 182). Means of raw scores before residualizing for cell composition, array, slide, and batch. Scores were standardized (mean = 0, SD = 1) for analyses.

^b PGS-CRP only computed for individuals solely of recent European ancestries.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 24

4. References

- 1. Kinge JM, Øverland S, Flatø M, et al. Parental income and mental disorders in children and adolescents: prospective register-based study. *International Journal of Epidemiology*. Published online May 11, 2021:dyab066. doi:10.1093/ije/dyab066
- 2. Peverill M, Dirks MA, Narvaja T, Herts KL, Comer JS, McLaughlin KA. Socioeconomic status and child psychopathology in the United States: A meta-analysis of population-based studies. *Clinical Psychology Review*. 2021;83:101933. doi:10.1016/j.cpr.2020.101933
- 3. Vaid E, Lansing AH. Discrimination, Prejudice, and Oppression and the Development of Psychopathology. In: Benuto LT, Duckworth MP, Masuda A, O'Donohue W, eds. *Prejudice, Stigma, Privilege, and Oppression: A Behavioral Health Handbook*. Springer International Publishing; 2020:235-248. doi:10.1007/978-3-030-35517-3_14
- 4. Davis FJ. *Who Is Black? One Nation's Definition*. Pennsylvania State University Press; 2001. doi:10.2307/4612582
- 5. Kendi IX. *Stamped from the Beginning: The Definitive History of Racist Ideas in America*. Random House; 2017.
- 6. Goosby BJ, Cheadle JE, Mitchell C. Stress-related biosocial mechanisms of discrimination and African American health inequities. *Annual Review of Sociology*. 2018;44(1):319-340. doi:10.1146/annurev-soc-060116-053403
- 7. Ligthart S, Marzi C, Aslibekyan S, et al. DNA methylation signatures of chronic low-grade inflammation are associated with complex diseases. *Genome Biology*. 2016;17(1):255. doi:10.1186/s13059-016-1119-5
- 8. Barker ED, Cecil CAM, Walton E, et al. Inflammation-related epigenetic risk and child and adolescent mental health: A prospective study from pregnancy to middle adolescence. *Dev Psychopathol.* 2018;30(3):1145-1156. doi:10.1017/S0954579418000330
- 9. Danese A, J Lewis S. Psychoneuroimmunology of Early-Life Stress: The Hidden Wounds of Childhood Trauma? *Neuropsychopharmacol*. 2017;42(1):99-114. doi:10.1038/npp.2016.198
- Snyder-Mackler N, Burger JR, Gaydosh L, et al. Social determinants of health and survival in humans and other animals. *Science*. 2020;368(6493):eaax9553. doi:10.1126/science.aax9553
- Ligthart S, Vaez A, Võsa U, et al. Genome Analyses of >200,000 Individuals Identify 58 Loci for Chronic Inflammation and Highlight Pathways that Link Inflammation and Complex Disorders. *The American Journal of Human Genetics*. 2018;103(5):691-706. doi:10.1016/j.ajhg.2018.09.009

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

- Muniz Carvalho C, Wendt FR, Maihofer AX, et al. Dissecting the genetic association of Creactive protein with PTSD, traumatic events, and social support. *Neuropsychopharmacol*. Published online March 16, 2020. doi:10.1038/s41386-020-0655-6
- Kappelmann N, Arloth J, Georgakis MK, et al. Dissecting the Association Between Inflammation, Metabolic Dysregulation, and Specific Depressive Symptoms: A Genetic Correlation and 2-Sample Mendelian Randomization Study. *JAMA Psychiatry*. 2021;78(2):161-170. doi:10.1001/jamapsychiatry.2020.3436
- 14. Dardani C, Yarmolinsky J, Robinson J, et al. Disentangling causal relationships between inflammatory markers and depression: a bidirectional Mendelian randomization analysis. *bioRxiv*. Published online November 25, 2019:712133. doi:10.1101/712133
- 15. Milton DC, Ward J, Ward E, et al. The association between C-reactive protein, mood disorder, and cognitive function in UK Biobank. *European Psychiatry*. 2021;64(1). doi:10.1192/j.eurpsy.2021.6
- Belsky D, Caspi A, Arseneault L, et al. Quantification of the pace of biological aging in humans through a blood test: The DunedinPoAm DNA methylation algorithm. *eLife*. 2020;9. doi:10.1101/2020.02.05.927434
- Raffington L, Belsky DW, Kothari M, Malanchini M, Tucker-Drob EM, Harden KP. Socioeconomic Disadvantage and the Pace of Biological Aging in Children. *Pediatrics*. Published online May 17, 2021:e2020024406. doi:10.1542/peds.2020-024406
- 18. Rutherford A. *How to Argue with a Racist: History, Science, Race, and Reality.* Hachette UK; 2020.
- 19. Yudell M, Roberts D, DeSalle R, Tishkoff S. Taking race out of human genetics. *Science*. 2016;351(6273):564-565. doi:10.1126/science.aac4951
- 20. Pauker K, Meyers C, Sanchez DT, Gaither SE, Young DM. A review of multiracial malleability: Identity, categorization, and shifting racial attitudes. *Soc Personal Psychol Compass*. 2018;12(6):e12392. doi:10.1111/spc3.12392
- 21. Abascal M. Contraction as a Response to Group Threat: Demographic Decline and Whites' Classification of People Who Are Ambiguously White. *American Sociological Review*. Published online March 3, 2020:000312242090512. doi:10.1177/0003122420905127
- 22. Anglin DM, Ereshefsky S, Klaunig MJ, et al. From Womb to Neighborhood: A Racial Analysis of Social Determinants of Psychosis in the United States. *AJP*. Published online May 3, 2021:appi.ajp.2020.2. doi:10.1176/appi.ajp.2020.20071091
- 23. Krieger N. Measures of Racism, Sexism, Heterosexism, and Gender Binarism for Health Equity Research: From Structural Injustice to Embodied Harm—An Ecosocial Analysis. *Annu Rev Public Health*. 2020;41(1):37-62. doi:10.1146/annurev-publhealth-040119-094017

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

- Williams DR, Priest N, Anderson NB. Understanding associations among race, socioeconomic status, and health: Patterns and prospects. *Health Psychology*. 2016;35(4):407-411. doi:10.1037/hea0000242
- 25. Adam EK, Hittner EF, Thomas SE, Villaume SC, Nwafor EE. Racial discrimination and ethnic racial identity in adolescence as modulators of HPA axis activity. *Dev Psychopathol*. 2020;32(5):1669-1684. doi:10.1017/S095457942000111X
- 26. Raffington L, Mallard T, Harden KP. Polygenic Scores in Developmental Psychology: Invite Genetics In, Leave Biodeterminism Behind. *Annu Rev Dev Psychol*. 2020;2(1):389-411. doi:10.1146/annurev-devpsych-051820-123945
- 27. Olson S, Berg K, Bonham V, et al. The use of racial, ethnic, and ancestral categories in human genetics research. *The American Journal of Human Genetics*. 2005;77(4):519-532. doi:10.1086/491747
- 28. Harden KP, Tucker-Drob EM, Tackett JL. The Texas Twin Project. *Twin Research and Human Genetics*. 2013;16(01):385-390. doi:10.1017/thg.2012.97
- 29. Achenbach T. Integrative Guide for the 1991 CBCL/4-18, YSR, and TRF Profiles. Department of Psychiatry, University of Vermont; 1991.
- Lizotte AJ, Chard-Wierschem DJ, Loeber R, Stern SB. A Shortened Child Behavior Checklist for Delinquency Studies. *Journal of Quantitative Criminology*. 1992;8(2):233-245.
- Conners CK, Sitarenios G, Parker JDA, Epstein JN. The Revised Conners' Parent Rating Scale (CPRS-R): Factor Structure, Reliability, and Criterion Validity. J Abnorm Child Psychol. 1998;26(4):257-268. doi:10.1023/A:1022602400621
- 32. Datar A, Chung PJ. Changes in Socioeconomic, Racial/Ethnic, and Sex Disparities in Childhood Obesity at School Entry in the United States. *JAMA Pediatr*. 2015;169(7):696. doi:10.1001/jamapediatrics.2015.0172
- 33. Braithwaite D, Moore DH, Lustig RH, et al. Socioeconomic status in relation to early menarche among black and white girls. *Cancer Causes Control*. 2009;20(5):713-720. doi:10.1007/s10552-008-9284-9
- Sumner JA, Colich NL, Uddin M, Armstrong D, McLaughlin KA. Early experiences of threat, but not deprivation, are associated with accelerated biological aging in children and adolescents. *Biological Psychiatry*. 2019;85(3):268-278. doi:10.1016/j.biopsych.2018.09.008
- 35. US Department of Health and Human Services (USDHHS).; 2012.
- 36. Wahl S, Drong A, Lehne B, et al. Epigenome-wide association study of body mass index, and the adverse outcomes of adiposity. *Nature*. 2017;541(7635):81-86. doi:10.1038/nature20784

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

- 37. Almstrup K, Lindhardt Johansen M, Busch AS, et al. Pubertal development in healthy children is mirrored by DNA methylation patterns in peripheral blood. *Scientific Reports*. 2016;6(1):28657. doi:10.1038/srep28657
- Joehanes R, Just AC, Marioni RE, et al. Epigenetic signatures of cigarette smoking. *Circulation: Cardiovascular Genetics*. 2016;9(5):436-447. doi:10.1161/CIRCGENETICS.116.001506
- 39. Muthén LK, Muthén BO. Mplus User's Guide. Eighth Edi. (Muthén & Muthén, ed.).; 2017.
- 40. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate : A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society*. 1995;57(1):289-300.
- 41. Raffington L, Tanksley P, Sabhlok A, et al. Socially stratified epigenetic profiles are associated with cognitive functioning in children and adolescents. *bioRxiv [Preprint]*. Published online 2021.
- 42. Danese A, Baldwin JR. Hidden Wounds? Inflammatory Links Between Childhood Trauma and Psychopathology. *Annu Rev Psychol*. 2017;68(1):517-544. doi:10.1146/annurev-psych-010416-044208
- 43. Cole SW, Shanahan MJ, Gaydosh L, Harris KM. Population-based RNA profiling in Add Health finds social disparities in inflammatory and antiviral gene regulation to emerge by young adulthood. *Proceedings of the National Academy of Sciences*. 2020;(17):201821367. doi:10.1073/pnas.1821367117
- 44. McDade TW, Ryan CP, Jones MJ, et al. Genome-wide analysis of DNA methylation in relation to socioeconomic status during development and early adulthood. *American Journal of Physical Anthropology*. 2019;169(1):3-11. doi:10.1002/ajpa.23800
- 45. Hostinar CE, Nusslock R, Miller GE. Future Directions in the Study of Early-Life Stress and Physical and Emotional Health: Implications of the Neuroimmune Network Hypothesis. *Journal of Clinical Child & Adolescent Psychology*. 2018;47(1):142-156. doi:10.1080/15374416.2016.1266647
- 46. Emeny RT, Carpenter DO, Lawrence DA. Health disparities: Intracellular consequences of social determinants of health. *Toxicology and Applied Pharmacology*. 2021;416:115444. doi:10.1016/j.taap.2021.115444
- 47. Epel ES. The geroscience agenda: Toxic stress, hormetic stress, and the rate of aging. *Ageing Research Reviews*. 2020;63:101167. doi:10.1016/j.arr.2020.101167
- 48. Peters A, Nawrot TS, Baccarelli AA. Hallmarks of environmental insults. *Cell*. 2021;184(6):1455-1468. doi:10.1016/j.cell.2021.01.043
- 49. Santos AL, Sinha S. Obesity and aging: Molecular mechanisms and therapeutic approaches. *Ageing Research Reviews*. 2021;67:101268. doi:10.1016/j.arr.2021.101268

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

- 50. Card D, Giuliano L. Universal screening increases the representation of low-income and minority students in gifted education. *Proceedings of the National Academy of Sciences*. 2016;113(48):13678-13683. doi:10.1073/pnas.1605043113
- 51. Harden KP, Engelhardt LE, Mann FD, et al. Genetic Associations Between Executive Functions and a General Factor of Psychopathology. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2020;59(6):749-758. doi:10.1016/j.jaac.2019.05.006
- 52. Williams DR. Stress and the Mental Health of Populations of Color: Advancing Our Understanding of Race-related Stressors. *J Health Soc Behav*. 2018;59(4):466-485. doi:10.1177/0022146518814251
- Marnetto D, Pärna K, Läll K, et al. Ancestry deconvolution and partial polygenic score can improve susceptibility predictions in recently admixed individuals. *Nat Commun.* 2020;11(1):1628. doi:10.1038/s41467-020-15464-w
- 54. Lloyd-Jones LR, Zeng J, Sidorenko J, et al. Improved polygenic prediction by Bayesian multiple regression on summary statistics. *Nature Communications*. 2019;10(1):5086. doi:10.1038/s41467-019-12653-0
- Pain O, Glanville KP, Hagenaars SP, et al. Evaluation of polygenic prediction methodology within a reference-standardized framework. Plagnol V, ed. *PLoS Genet*. 2021;17(5):e1009021. doi:10.1371/journal.pgen.1009021
- 56. Auton A, Abecasis GR, Altshuler DM, et al. A global reference for human genetic variation. *Nature*. 2015;526(7571):68-74. doi:10.1038/nature15393
- 57. The International HapMap 3 Consortium. Integrating common and rare genetic variation in diverse human populations. *Nature*. 2010;467(7311):52-58. doi:10.1038/nature09298
- 58. Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*. 2015;4(1):7. doi:10.1186/s13742-015-0047-8
- Harden KP, Patterson MW, Briley DA, et al. Developmental changes in genetic and environmental influences on rule-breaking and aggression: age and pubertal development. *Journal of Child Psychology and Psychiatry*. 2015;56(12):1370-1379. doi:https://doi.org/10.1111/jcpp.12419
- 60. Patterson MW, Mann FD, Grotzinger AD, Tackett JL, Tucker-Drob EM, Harden KP. Genetic and environmental influences on internalizing psychopathology across age and pubertal development. *Developmental Psychology*. 2018;54(10):1928-1939. doi:10.1037/dev0000578
- 61. Service UD of AER. US Household Food Security Survey Module: three-stage design, with screeners. Published online 2012.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

- 62. Engelhardt LE, Church JA, Harden KP, Tucker-Drob EM. Accounting for the shared environment in cognitive abilities and academic achievement with measured socioecological contexts. *Developmental Science*. 2019;22(1):e12699. doi:10.1111/desc.12699
- 63. Chetty R, Friedman JN, Hendren N, Jones MR, Porter SR. *The Opportunity Atlas: Mapping the Childhood Roots of Social Mobility*. National Bureau of Economic Research; 2018. doi:10.3386/w25147
- 64. Petersen AC, Crockett L, Richards M, Boxer A. A self-report measure of pubertal status: Reliability, validity, and initial norms. *Journal of youth and adolescence*, 1988;17(2):117-133.
- 65. Aryee MJ, Jaffe AE, Corrada-Bravo H, et al. Minfi: A flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. *Bioinformatics*. Published online 2014. doi:10.1093/bioinformatics/btu049
- 66. Triche TJ, Weisenberger DJ, Van Den Berg D, Laird PW, Siegmund KD. Low-level processing of Illumina Infinium DNA Methylation BeadArrays. *Nucleic Acids Research*. 2013;41(7):e90-e90. doi:10.1093/nar/gkt090
- 67. Pidsley R, Zotenko E, Peters TJ, et al. Critical evaluation of the Illumina MethylationEPIC BeadChip microarray for whole-genome DNA methylation profiling. *Genome Biology*. 2016;17(1):208. doi:10.1186/s13059-016-1066-1
- 68. Middleton LYM, Dou J, Fisher J, et al. Saliva cell type DNA methylation reference panel for epidemiological studies in children. *Epigenetics*. Published online February 22, 2021:1-17. doi:10.1080/15592294.2021.1890874
- 69. Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics*. 2007;8(1):118-127. doi:10.1093/biostatistics/kxj037
- Anderson CA, Pettersson FH, Clarke GM, Cardon LR, Morris AP, Zondervan KT. Data quality control in genetic case-control association studies. *Nat Protoc*. 2010;5(9):1564-1573. doi:10.1038/nprot.2010.116
- 71. Turner S, Armstrong LL, Bradford Y, et al. Quality Control Procedures for Genome-Wide Association Studies. *Current Protocols in Human Genetics*. 2011;68(1):1.19.1-1.19.18. doi:10.1002/0471142905.hg0119s68
- 72. Roshyara NR, Kirsten H, Horn K, Ahnert P, Scholz M. Impact of pre-imputation SNPfiltering on genotype imputation results. *BMC Genetics*. 2014;15(1):88. doi:10.1186/s12863-014-0088-5

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 30

Acknowledgment Section

Funding: We gratefully acknowledge all participants of the Texas Twin Project. This research was supported by National Institutes of Health (NIH) grants R01HD083613 and R01HD092548. LR was supported by the German Research Foundation (DFG). MWP was supported by a training grant from NIMH, T32MH015442. ZA is supported by a Marie Skłodowska-Curie Fellowship from the European Union (894675). KPH and EMTD are Faculty Research Associates of the Population Research Center at the University of Texas at Austin, which is supported by a NIH grant P2CHD042849. EMTD is a member of the Center on Aging and Population Sciences (CAPS) at The University of Texas at Austin, which is supported by NIH grant P30AG066614. KPH and EMTD were also supported by Jacobs Foundation Research Fellowships. LR and KPH had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Conflicts of interest: Not applicable.

Ethics approval: The University of Texas at Austin Institutional Review board granted ethical approval.

Consent to participate: Informed consent to participate in the study was obtained from all participants and their parent or legal guardian.

Consent for publication: Not applicable.

Availability of data and material: Because of the high potential for deductive identification in this special population from a geographically circumscribed area, and the sensitive nature of information collected, data from the Texas Twin Project are not shared with individuals outside of the research team.

Code availability: Code will be shared by the first author upon request.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 31

Author Contributions: KPH, EMTD, & LR developed the study concept and design. LR, PT,

LV, & TM performed the data analysis under the supervision of EMTD and KPH. LR drafted the manuscript. All authors provided critical revisions to the analysis plan and manuscript and approved the final version of the manuscript for submission.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 32

Supplemental Methods

DNA-methylation preprocessing. DNA extraction and methylation profiling was conducted by Edinburgh Clinical Research Facility (UK). DNA-methylation preprocessing was primarily conducted with the 'minfi' package in R⁶⁵. Within-array normalization was performed to address array background correction, red/green dye bias, and probe type I/II correction, and it has been noted that at least part of the probe type bias is a combination of the first two factors (. Noob preprocessing as implemented by minfi's "preprocessNoob"⁶⁶ is a background correction and dye-bias equalization method that has similar within-array normalization effects on the data as probe type correction methods such as BMIQ (Teschendorff et al., 2013).

In line with our preregistered preprocessing plan, CpG probes with detection p > 0.01 and fewer than 3 beads in more than 1% of the samples and probes in cross-reactive regions were excluded ⁶⁷. None of these failed probes overlapped with the probes used for DNA-methylation scores. 44 samples were excluded because (1) they showed low intensity probes as indicated by the log of average methylation <9 and their detection *p*-value was > 0.01 in >10% of their probes, (2) their self-reported and methylation-estimated sex mismatch, and/or (3) their self-reported and DNA-estimated sex mismatch. Cell composition of immune and epithelial cell types (*i.e.*, CD4+ T-cell, natural killer cells, neutrophilseosinophils, B cells, monocytes, CD8+ T-cell, and granulocytes) were estimated using a newly developed child saliva reference panel implemented in the R package "BeadSorted.Saliva.EPIC" within "ewastools" ⁶⁸. Surrogate variable analysis was used to correct methylation values for batch effects using the "combat" function in the SVA package ⁶⁹.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 33

Genotyping and imputation. DNA samples were genotyped at the University of Edinburgh using the Illumina Infinium PsychArray, which assays ~590,000 single nucleotide polymorphisms (SNPs), insertions-deletions (indels), copy number variants (CNVs), structural variants, and germline variants across the genome. Genetic data was subjected to quality control procedures recommended for chip-based genomic data ^{70,71}. Briefly, samples were excluded on the basis of poor call rate (< 98%) or inconsistent self-reported and biological sex, while variants were excluded if missingness exceeded 2%. As further variant-level filtering has been shown to have a detrimental effect on imputation quality ⁷², quality control thresholds for minor allele frequency (MAF) and Hardy–Weinberg equilibrium (HWE) were applied after phasing and imputation.

Untyped markers were imputed on the Michigan Imputation Server (https://imputationserver.sph.umich.edu). Specifically, genotypes were phased and imputed with Eagle v2.4 and Minimac4 (v1.5.7), respectively, while using the 1000 Genomes Phase 3 v5 reference panel ⁵⁶. To ensure that only high-quality typed and imputed markers were used for analysis, variants were excluded if they had a MAF < 1e-3, a HWE p-value < 1e-6, or an imputation quality score < .90. These procedures produced a final set of 4,703,309 genetic markers to be used in analyses.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 34

Supplemental Tables

- Table S1. CBCL and Conners items included in each mental health summary measure.
- Table S2. Associations between socioeconomic inequality and racial/ethnic identity with parent-reported mental health.
- Table S3. Associations between socioeconomic inequality and racial/ethnic identity with self-reported mental health.
- Table S4. Associations between racial/ethnic identity and socioeconomic inequality.
- Table S5. Associations between socioeconomic inequality and racial/ethnic identity with DNA-methylation profiles.
- Table S6. Associations between socioeconomic inequality with DNA-methylation profiles for each racial/ethnic group.
- Table S7. Associations between DNA-methylation and genetic profiles with parentreported mental health.
- Table S8. Associations between DNA-methylation and genetic profiles with self-reported mental health.
- Table S9. Associations between DNA-methylation with parent-reported mental health for each racial/ethnic group.
- Table S10. Associations between DNA-methylation with self-reported mental health for each racial/ethnic group.
- Table S11. Co-twin-control associations between DNA-methylation and parent-reported mental health.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 35

Measure	#	CBCL self-report question	CBCL parent-report question
Attention Problems	1	I fail to finish things that I start.	Acts too young for his/her age.
		I have trouble concentrating or paying	Can't concentrate, can't pay attention
Attention Problems	2	attention.	for long.
Attention Problems	3	I have trouble sitting still.	Can't sit still, restless, or hyperactive.
Attention Problems	4	I feel confused or in a fog	thoughts
Attention Problems	4 5	I devidencem a lot	Inoughts.
Attention Problems	5	I daydreann a lot.	Deer ask ash work
Attention Problems	0	I act without stopping to unitk.	
Attention Problems	/	I am inattentive and easily distracted.	/ Deliberately harms self or attempts
Internalizing - depression	1	There is very little that I enjoy.	suicide.
Internalizing - depression	2	I feel lonely.	Secretive, keeps things to self.
Internalizing - depression	3	I crv a lot.	Self-conscious or easily embarrassed.
Internalizing - depression	4	I feel overtired without good reason.	Too shy or timid.
Internalizing - depression	5	I don't have much energy.	Unhappy, sad, or depressed.
Internalizing - depression	6	I am unhappy, sad, or depressed.	Withdrawn, doesn't get involved with others.
Internalizing - anxiety	7	I am nervous or tense.	Cries a lot.
Internalizing - anxiety	8	I am too fearful or anxious	Fears going to school
internalizing anxiety	0		Fears he/she might think or do
Internalizing - anxiety	9	I feel too guilty.	something bad.
T . 1	10	I am self-conscious and easily	
Internalizing - anxiety	10	embarrassed.	Feels he/she has to be perfect.
Internalizing - anxiety	11	I worry a lot.	Feels worthless or inferior.
Internalizing - anxiety	12		Too fearful or anxious.
Internalizing - anxiety	13	/	Feels too guilty.
Internalizing - anxiety	14	/	Whining.
Internalizing - anxiety	15	/	Worries.
Internalizing - somatic		,	Aches or pains (not stomach or
complaints	16	/	headaches).
Internalizing - somatic	17	1	Handaahaa
Internalizing comptia	1/	7	Headacnes.
complaints	18	1	Neuson fools sick
Internalizing - somatic	10	1	Ivausea, icers siek.
complaints	19	/	Stomachaches
Rule-Breaking	1	I disobey my parents.	Disobedient at home.
Rule-Breaking	2	I disobey at school	Disobedient at school
Rule Dieuking	2	I don't feel guilty after doing something	Hangs around with others who get in
Rule-Breaking	3	I shouldn't.	trouble.
6	-	I break rules at home, school, or	
Rule-Breaking	4	elsewhere.	Lying or cheating.
6		I hang around with kids who get in	
Rule-Breaking	5	trouble	Steals outside the home.
Rule-Breaking	6	I lie or cheat.	/
Rule-Breaking	7	I run away from home.	/
Rule-Breaking	8	I set fires.	/
Rule-Breaking	9	I steal at home.	/
Rule-Breaking	10	I steal from places other than home.	/
Rule-Breaking	11	I swear or use dirty language.	/

Table S1. CBCL and Conners items included in each mental health summary measure.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Rule-Breaking	12	I cut classes or skip school.	/
Aggression	1	I argue a lot.	Cruel to animals.
			Cruelty, bullying, or meanness to
Aggression	2	I brag.	others.
Aggression	3	I am mean to others.	Demands a lot of attention.
			Destroys things belonging to his/her
Aggression	4	I destroy my own things.	family or others.
Aggression	5	I destroy things belonging to others.	Gets in many fights.
Aggression	6	I get in many fights.	Screams a lot.
Aggression	7	I physically attack people.	Stubborn, sullen, or irritable.
Aggression	8	I scream a lot.	Sudden changes in mood or feelings.
Aggression	9	I am stubborn.	Temper tantrums or hot temper.
Aggression	10	I talk too much.	Threatens people.
Aggression	11	I tease others a lot.	Unusually loud.
Aggression	12	I have a hot temper.	/
Aggression	13	I threaten to hurt people.	/
Measure	#	Conner self-report question	Conner parent-report question
ADHD-			
hyperactive/impulsive	1	I blurt out the first thing that I think of.	Talks too much.
ADHD-			Blurts out answers before the question
hyperactive/impulsive	2	I interrupt other people.	has been completed.
ADHD-	2	T I	
hyperactive/impulsive	3	I am restless.	Acts as if he/she is driven by a motor.
ADHD-	4	I blurt out the answer before the	Has difficulty waiting for his/har turn
	4	Linu or climb even when Lam not	Runs or climbs when he/she is not
hyperactive/impulsive	5	supposed to	supposed to
ADHD-	5	supposed to.	Is noisy and loud when playing or
hyperactive/impulsive	6	I have trouble waiting for my turn.	using free time.
ADHD-			Leaves seat when he/she should stay
hyperactive/impulsive	7	I have too much energy to stay still.	seated.
ADHD-		I like to be on the go rather than being	
hyperactive/impulsive	8	in one place.	Fidgets or squirms in seat.
ADHD-		I get out of my seat when I am not	
hyperactive/impulsive	9	supposed to.	ls restless or overactive.
ADHD-	10		Interrupts others (butts into
	10	I leef like I am driven by a motor.	conversations or games).
hyperactive/impulsive	11	quietly	/
ADHD_inattentive	1	Llose stuff that I need	/ Is forgetful in daily activities
	1	Tiose stuff that Theed.	Avoids or dislikes things that take a lot
ADHD-inattentive	2	I have trouble finishing things.	of effort and are not fun.
		I have trouble keeping myself	Does not seem to listen to what is being
ADHD-inattentive	3	organized.	said to him/her.
		It is hard for me to pay attention to	Doesn't pay attention to details, makes
ADHD-inattentive	4	details.	careless mistakes.
			Does not follow through on instructions
	_		(even when he/she understands and is
ADHD-inattentive	5	I forget stuff.	trying to cooperate).
			Fails to complete schoolwork, chores,
	C	I maleo mistoleon har anni dana	or tasks (even when he/she understands
ADHD-inattentive	0	I make mistakes by accident.	and is trying to cooperate).
ADHD inattentivo	7	what people are saving to me	nas trouble organizing tasks or
	/	what people are saying to me.	מכנו אונובא.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

		I don't like doing things that make me	Has trouble keeping his/her mind on
ADHD-inattentive	8	think hard.	work or play for long.
			Loses things (like schoolwork, pencils,
ADHD-inattentive	9	I have trouble following instructions.	books, tools, or toys).
		I have trouble keeping my mind on	
ADHD-inattentive	10	what I am doing.	/
		I steal from other people (by mugging	
Conduct Disorder	1	or purse snatching).	Skips class.
		I tell lies to get out of doing things or to	Has forced someone into sexual
Conduct Disorder	2	get stuff.	activity.
Conduct Disorder	3	I start fights with other people.	Bullies, threatens, or scares others.
			Uses a weapon (like a bat, brick,
Conduct Disorder	4	I am mean to animals.	broken bottle, knife or gun).
		I steal important things when no one is	
Conduct Disorder	5	watching.	Starts fights with others on purpose.
		I use a weapon (like a bat, brick,	
		broken glass, knife, or gun) to scare or	
Conduct Disorder	6	hurt people.	Physically hurts people.
			Lies to get things or to avoid having to
Conduct Disorder	7	I break into houses, buildings, or cars,	do something.
		8-,8-,	Runs away from home for at least one
Conduct Disorder	8	I do things to hurt people	night
	Ũ	I go out at night even when I am	Has set fires with the intention of
Conduct Disorder	9	supposed to be at home	causing damage
Conduct Disorder		supposed to be at nonie.	Has broken into someone else's house
Conduct Disorder	10	/	huilding or car
Conduct Disorder	10	,	Goes out at night even though it breaks
Conduct Disorder	11	/	the rules
Conduct Disorder	11	1	Steals while confronting a person (like
Conduct Disorder	12	/	mugging purse snatching)
Oppositional Defiant	12	I do what my parents or other adults	mugging, purse snatching).
Disorder	1	ask me to do	Loses temper
Oppositional Defiant	1	ask life to do.	Blames others for his/her mistakes or
Disordor	2	I try to approve other people	mishahavior
Oppositional Defiant	2	Tuy to annoy other people.	misochavior.
Disordor	3	I argue with adults	Is anorry or resentful
Oppositional Defient	5	l'algue with adults.	is angly of resentrui.
Disorder	4	I blome others for things I do wrong	Tries to get even with people
Oppositional Defient	4	I blame bulers for unings I do wrong.	Thes to get even with people.
Disorder	5	Lloss my temper	Annous other needle on numero
Oppositional Defient	5	Tiose my temper.	Annoys other people on purpose.
Disorder	6	I am apply approved by others	is initiable and easily annoyed by
Disorder Organitienal Definet	0	I am easily annoyed by others.	Ouriers.
Disorder	7	Deemle melte me enemy	Actively refuses to do what adults tell
Disorder	/	When I act used at accurate I act to a	mm/ner to do.
Oppositional Defiant	0	when I get mad at someone, I get even	A 11 1 1
Disorder	8	with them.	Argues with adults.

	Int	ternalizi	ing	Atten	tion Pro	blems	A	ggressio	on	Ru	le-Breal	king	-	ADHD			ODD			CD	
									No furth	ner cova	riates ^a										
	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р
Family-level disadvantage	0.25	0.16, 0.33	0.000	0.16	0.07, 0.25	0.001	0.26	0.14, 0.39	0.000	0.11	0.04, 0.19	0.004	0.18	0.08, 0.27	0.000	0.14	0.04, 0.24	0.005	0.16	-0.02, 0.34	0.088
Neighborhood- level disadvantage	0.13	0.04, 0.22	0.003	0.12	0.03, 0.20	0.008	0.07	-0.04, 0.19	0.219	0.04	-0.04, 0.11	0.364	0.04	-0.04, 0.12	0.286	0.04	-0.05, 0.14	0.365	0.16	-0.03, 0.34	0.097
Neighborhood opportunity	-0.17	-0.25, -0.09	0.000	-0.19	-0.28, -0.10	0.000	-0.12	-0.22, -0.02	0.016	-0.01	-0.09, 0.07	0.826	-0.09	-0.18, 0.01	0.068	-0.18	-0.31, -0.05	0.008	-0.23	-0.42, -0.03	0.022
Latinx	0.09	0.01, 0.17	0.021	0.18	0.09, 0.27	0.000	0.03	-0.08, 0.12	0.689	0.08	-0.08, 0.23	0.352	-0.05	-0.13, 0.02	0.168	-0.06	-0.13, 0.00	0.060	-0.04	-0.08, 0.00	0.047
Latinx-White	0.06	-0.03, 0.14	0.218	0.05	-0.03, 0.14	0.213	0.13	-0.05, 0.22	0.232	0.02	-0.18, 0.21	0.854	0.05	-0.03, 0.13	0.250	0.03	-0.07, 0.13	0.570	0.07	-0.05, 0.18	0.254
Black+	0.23	0.15, 0.30	0.000	0.15	0.06, 0.23	0.001	0.16	0.01, 0.24	0.027	0.10	-0.06, 0.26	0.204	0.12	0.03, 0.22	0.009	0.07	-0.03, 0.18	0.173	0.06	-0.09, 0.20	0.399
Asian+	0.00	-0.07, 0.07	0.946	0.06	-0.03, 0.14	0.173	0.00	-0.09, 0.09	0.995	0.04	-0.14, 0.22	0.647	-0.08	-0.13, -0.04	0.001	-0.11	-0.14, -0.08	0.000	-0.06	-0.08, -0.03	0.001
							С	ontrollin	ng for fai	mily-lev	vel disad	vantage									
	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р
Family-level disadvantage	0.17	0.07, 0.26	0.000	0.09	-0.02, 0.19	0.105	0.24	0.11, 0.37	0.000	0.13	0.03, 0.22	0.008	0.16	0.05, 0.27	0.004	0.16	0.05, 0.26	0.004	0.14	-0.01, 0.29	0.063
Latinx	0.07	-0.02, 0.16	0.120	0.16	0.06, 0.26	0.002	-0.02	-0.15, 0.11	0.774	-0.01	-0.11, 0.10	0.909	-0.09	-0.18, 0.01	0.078	-0.11	-0.20, -0.03	0.007	-0.10	-0.17, -0.03	0.005
Latinx-White	0.06	-0.04, 0.15	0.239	0.05	-0.04, 0.15	0.283	0.13	-0.03, 0.28	0.109	-0.02	-0.10, 0.07	0.692	0.02	-0.08, 0.12	0.685	0.03	-0.09, 0.14	0.682	0.07	-0.08, 0.22	0.360
Black+	0.16	0.06, 0.25	0.002	0.14	0.03, 0.26	0.017	0.09	-0.06, 0.23	0.250	-0.01	-0.12, 0.10	0.844	0.06	-0.05, 0.16	0.286	-0.02	-0.13, 0.09	0.732	0.09	-0.12, 0.30	0.410
Asian+	-0.03	-0.11, 0.04	0.433	0.08	-0.02, 0.18	0.133	0.02	-0.1, 0.13	0.773	0.04	-0.06, 0.14	0.421	-0.07	-0.13, -0.01	0.016	-0.10	-0.14, -0.06	0.000	-0.05	-0.11, 0.00	0.075

	Table S2.	Associations	between	socioeconor	nic inequ	ality and	l racial/ethnic	identity	with	parent-re	ported n	nental h	ealth
--	-----------	--------------	---------	-------------	-----------	-----------	-----------------	----------	------	-----------	----------	----------	-------

Standardized regression coefficients (*b*) and 95% confidence intervals (CIs) and uncorrected *p*-value calculated by regressing parent-reported mental health on socioeconomic inequality and racial/ethnic identity (with and without controlling for family-level disadvantage). *P*-values, where FDR corrected *p*-values < 0.05, are marked in bold. White-only identity is reference group. ^aAll models included covariate adjustment for age, gender, and an age by gender interaction. *Note:* ADHD=attention-deficit hyperactivity disorder; ODD=oppositional defiant disorder; CD= conduct disorder.

	Interr	nalizing		Atten	tion Pro	blems	Aggre	ession		Rule-	Breakin	g	ADH	D		ODD			CD		
									No furtl	ner cova	riates ^a										
	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р
Family-level disadvantage	0.07	-0.04, 0.17	0.237	0.16	0.06, 0.26	0.001	0.24	0.13, 0.36	0.000	0.16	0.04, 0.28	0.010	0.14	0.02, 0.26	0.024	0.08	-0.03, 0.19	0.142	-0.01	-0.07, 0.06	0.778
Neighborhood- level disadvantage	0.08	-0.01, 0.17	0.085	0.07	-0.02, 0.17	0.117	0.13	0.03, 0.22	0.008	0.15	0.04, 0.26	0.009	0.09	0.00, 0.19	0.046	0.16	0.04, 0.28	0.009	0.04	-0.05, 0.13	0.409
Neighborhood opportunity	-0.06	-0.16, 0.05	0.280	-0.08	-0.18, 0.03	0.171	-0.15	-0.27, -0.03	0.013	-0.16	-0.29, -0.03	0.017	-0.09	-0.21, 0.04	0.165	-0.13	-0.22, -0.04	0.007	-0.11	-0.21, -0.01	0.039
Latinx	-0.03	-0.12, 0.07	0.578	-0.01	-0.10, 0.08	0.844	0.01	-0.08, 0.09	0.891	0.05	-0.05, 0.16	0.324	0.04	-0.07, 0.15	0.494	-0.01	-0.11, 0.08	0.766	-0.05	-0.11, 0.00	0.045
Latinx-White	0.10	0.00, 0.19	0.050	0.09	-0.01, 0.19	0.092	-0.03	-0.12, 0.06	0.518	0.03	-0.07, 0.12	0.599	0.01	-0.10, 0.11	0.863	-0.03	-0.11, 0.04	0.417	-0.01	-0.08, 0.06	0.763
Black+	0.00	-0.09, 0.08	0.988	0.07	-0.02, 0.15	0.111	0.15	0.06, 0.23	0.001	0.08	-0.01, 0.18	0.087	0.07	-0.02, 0.15	0.125	0.05	-0.04, 0.14	0.240	-0.01	-0.07, 0.05	0.742
Asian+	-0.04	-0.12, 0.03	0.284	-0.06	-0.14, 0.02	0.174	-0.03	-0.12, 0.06	0.499	-0.01	-0.10, 0.09	0.913	-0.08	-0.16, 0.00	0.038	-0.03	-0.11, 0.05	0.435	0.00	-0.06, 0.06	0.983
							C	ontrollir	ng for fa	mily-lev	vel disad	lvantage	•								
	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р	b	CI	р
Family-level disadvantage	0.10	-0.02, 0.23	0.106	0.18	0.06, 0.29	0.003	0.21	0.08, 0.33	0.001	0.02	-0.10, 0.15	0.022	0.11	0.24, -0.03	0.122	0.08	-0.04, 0.21	0.187	0.03	-0.05, 0.10	0.508
Latinx	-0.06	-0.17, 0.05	0.286	-0.06	-0.17, 0.05	0.253	-0.02	-0.12, 0.09	0.762	0.01	-0.11, 0.13	0.740	0.04	0.18, -0.09	0.538	-0.02	-0.12, 0.09	0.791	-0.08	-0.13, -0.03	0.001
Latinx-White	0.10	-0.01, 0.21	0.077	0.07	-0.06, 0.19	0.302	-0.11	-0.22, 0.00	0.055	-0.03	-0.15, 0.10	0.895	-0.03	0.10, -0.16	0.684	-0.03	-0.10, 0.04	0.480	0.02	-0.07, 0.11	0.726
Black+	-0.07	-0.19, 0.05	0.264	-0.02	-0.12, 0.08	0.732	0.06	-0.05, 0.16	0.311	-0.06	-0.18, 0.06	0.658	0.02	0.13, -0.10	0.775	-0.03	-0.14, 0.08	0.616	-0.05	-0.12, 0.02	0.172
Asian+	-0.04	-0.13, 0.05	0.387	-0.07	-0.16, 0.02	0.138	-0.08	-0.17, 0.02	0.122	0.16	0.02, 0.29	0.315	-0.10	0.00, -0.18	0.040	-0.06	-0.14, 0.02	0.129	-0.04	-0.08, 0.01	0.096

Table S3. Association	ns between	socioeconomi	c inequality	y and raci	al/ethnic ide	entity with	self-reported	mental health.
						•	-	

Standardized regression coefficients (*b*) and 95% confidence intervals (CIs) and uncorrected *p*-value calculated by regressing parent-reported mental health on socioeconomic inequality and racial/ethnic identity (with and without controlling for family-level disadvantage). *P*-values, where FDR corrected *p*-values < 0.05, are marked in bold. White-only identity is reference group. ^aAll models included covariate adjustment for age, gender, and an age by gender interaction. *Note:* ADHD=attention-deficit hyperactivity disorder; ODD=oppositional defiant disorder; CD= conduct disorder.

	Family	-level disadvanta	ıge	Neighb	orhood-level dis	advantage	Neighb	orhood opportu	inity
	b	95% CI	р	b	95% CI	р	b	95% CI	p
Latinx	0.25	0.16 - 0.35	<0.001	0.43	0.33-0.52	<0.001	-0.29	-0.41	<0.001
								0.17	
Latinx-White	0.02	-0.09 - 0.12	0.763	0.04	-0.06-0.15	0.444	0.03	-0.11-0.16	0.700
Black+	0.43	0.35 - 0.52	<0.001	0.33	0.25 - 0.41	<0.001	-0.28	-0.38– -	<0.001
								0.18	
Asian+	-0.10	-0.190.02	0.017	-0.06	-0.130.01	0.037	0.09	0.01 - 0.17	0.040

Table S4. Associations between racial/ethnic identity and socioeconomic inequality.

Standardized regression coefficients (*b*), 95% confidence intervals (CIs), and uncorrected *p*-values calculated by regressing socioeconomic measures on racial/ethnic identity. White-only identity is reference group. *P*-values, where FDR corrected *p*-values < 0.05, are marked in bold.

	DNAm-(CRP		DunedinP	oAm	
		No furth	er covariates ^a			
	b	95% CI	р	b	95% CI	р
Family-level disadvantage	0.22	0.12-0.31	<0.001	0.28	0.18-0.37	<0.001
Neighborhood-level disadvantage	0.25	0.17 - 0.33	<0.001	0.24	0.15 - 0.34	<0.001
Neighborhood opportunity	-0.12	-0.210.03	0.012	-0.19	-0.300.08	0.001
Latinx	0.15	0.07 - 0.24	0.001	0.18	0.10-0.27	<0.001
Latinx-White	0.10	-0.02 - 0.22	0.108	0.09	-0.01-0.19	0.089
Black+	0.08	0.02 - 0.15	0.012	0.19	0.11 - 0.28	<0.001
Asian+	-0.01	-0.09-0.07	0.776	0.11	-0.01-0.22	0.058
		Control	ling for BMI			
	b	95% CI	р	b	95% CI	р
BMI	0.33	0.24 - 0.42	<0.001	0.30	0.21 - 0.40	<0.001
Family-level disadvantage	0.11	0.01 - 0.21	0.024	0.19	0.08 - 0.29	<0.001
Neighborhood-level disadvantage	0.15	0.07 - 0.23	<0.001	0.14	0.05 - 0.24	0.004
Neighborhood opportunity	-0.05	-0.15-0.05	0.314	-0.09	-0.20 - 0.01	0.088
Latinx	0.10	0.01 - 0.18	0.029	0.13	0.04 - 0.22	0.004
Latinx-White	0.07	-0.03-0.17	0.180	0.06	-0.03 - 0.14	0.213
Black+	0.01	-0.07 - 0.07	0.975	0.11	0.02 - 0.19	0.011
Asian+	0.01	-0.07-0.09	0.857	0.13	0.02 - 0.24	0.025
		Controlli	ng for puberty			
	b	95% CI	р	b	95% CI	р
Puberty	0.13	0.02 - 0.24	0.020	0.05	-0.07 - 0.18	0.402
Family-level disadvantage	0.21	0.12 - 0.31	<0.001	0.28	0.18-0.39	<0.001
Neighborhood-level disadvantage	0.25	0.16-0.34	<0.001	0.25	0.11 - 0.34	<0.001
Neighborhood opportunity	-0.12	-0.220.01	0.027	-0.17	-0.280.06	0.003
Latinx	0.15	0.06-0.24	0.001	0.18	0.09 - 0.27	<0.001
Latinx-White	0.14	0.01 - 0.26	0.033	0.11	0.01 - 0.21	0.035
Black+	0.09	0.03 - 0.16	0.007	0.19	0.11 - 0.27	<0.001
Asian+	-0.01	-0.09 - 0.07	0.801	0.11	-0.01 - 0.22	0.059
		Controlling	for DNAm-smoke			
	b	95% CI	p	b	95% CI	p
DNAm-smoke	0.40	0.28-0.51	<0.001	0.24	0.11 - 0.37	<0.001
Family-level disadvantage	0.21	0.11 - 0.30	<0.001	0.28	0.18-0.38	<0.001

Table S5. Associations between socioeconomic inequality and racial/ethnic identity with DNA-methylation profiles.

DNA-METHYLATION AND MENTAL HEALTH

Neighborhood-level disadvantage	0.22	0.14 - 0.30	<0.001	0.23	0.14 - 0.33	<0.001
Neighborhood opportunity	-0.11	-0.210.02	0.022	-0.17	-0.270.06	0.003
Latinx	0.11	0.03 - 0.20	0.011	0.16	0.07 - 0.25	0.001
Latinx-White	0.12	0.01 - 0.23	0.048	0.11	0.01 - 0.20	0.028
Black+	0.14	0.07 - 0.20	<0.001	0.22	0.14 - 0.31	<0.001
Asian+	-0.02	-0.09 - 0.06	0.677	0.11	-0.01 - 0.22	0.067
		Controlling for fai	mily-level disadv	antage		
	b	95% CI	р	b	95% CI	р
Family-level disadvantage	0.19	0.07 - 0.30	0.002	0.24	0.13 - 0.36	<0.001
Latinx	0.10	0.01 - 0.20	0.036	0.12	0.02 - 0.22	0.019
Latinx-White	0.12	0.00 - 0.23	0.050	0.10	0.01 - 0.20	0.047
Black+	0.01	-0.06 - 0.09	0.739	0.09	-0.01 - 0.18	0.057

Standardized regression coefficients (*b*), 95% confidence intervals (CIs), and uncorrected p-values calculated by regressing DNA-methylation measures on socioeconomic measures and racial/ethnic identity with and without controlling for normed BMI z-scores, puberty (residualized for age within each gender), and DNA-methylation profiles of smoking (DNAm-smoke), separately. *P*-values, where FDR corrected *p*-values < 0.05, are marked in bold. ^aAll models included covariate adjustment for child's age, gender, and an age by gender interaction. Methylation scores were residualized for technical covariates (array, slide, batch, cell composition).

		DNAm-CRP	-		DunedinPoAm	
			White			
	b	95% CI	р	b	95% CI	р
Family-level disadvantage	0.11	-0.02-0.24	0.098	0.23	0.08-0.37	0.002
Neighborhood-level disadvantage	0.23	0.11-0.35	< 0.001	0.21	0.06-0.37	0.007
Neighborhood opportunity	-0.07	-0.19-0.06	0.312	-0.12	-0.3-0.06	0.182
			Latinx			
	b	95% CI	p	b	95% CI	р
Family-level disadvantage	0.33	0.07-0.59	0.014	0.118	-0.18-0.42	0.439
Neighborhood-level disadvantage	0.11	-0.12-0.33	0.352	0.01	-0.27-0.29	0.925
Neighborhood opportunity	-0.18	-0.43-0.07	0.160	-0.18	-0.51-0.15	0.278
			Latinx-White			
	b	95% CI	p	b	95% CI	р
Family-level disadvantage	0.53	0.13-0.93	0.010	0.29	-0.07-0.64	0.118
Neighborhood-level disadvantage	0.19	-0.1-0.47	0.197	0.11	-0.16-0.38	0.436
Neighborhood opportunity	-0.01	-0.20-0.18	0.945	-0.12	-0.33-0.08	0.232
			Black +			
	b	95% CI	p	b	95% CI	р
Family-level disadvantage	0.06	-0.22-0.34	0.678	0.19	-0.13-0.51	0.252
Neighborhood-level disadvantage	0.49	0.22 - 0.75	< 0.001	0.47	0.14-0.79	0.005
Neighborhood opportunity	-0.24	-0.440.04	0.018	-0.25	-0.52-0.02	0.065
			Asian +			
	b	95% CI	р	b	95% CI	р
Family-level disadvantage	0.08	-0.18-0.34	0.562	0.07	-0.26-0.41	0.668
Neighborhood-level disadvantage	0.07	-0.2-0.33	0.629	-0.10	-0.45-0.25	0.590
Neighborhood opportunity	0.25	-0.36 -0.86	0.428	-0.02	-0.85-0.8	0.959

Fable S6. Associations between socioeconomic ine	qualit	y with DNA-methy	ylation	profiles for eac	h racial/ethnic g	group.
---	--------	------------------	---------	------------------	-------------------	--------

Standardized regression coefficients (*b*) and 95% confidence intervals (CIs) and uncorrected *p*-value calculated by regressing DNAmethylation measures on socioeconomic measures, separately, within each racial/ethnic group. All models included covariate adjustment for age, gender, and an age by gender interaction. Methylation scores were residualized for technical covariates (for methylation: array, slide, batch, cell composition).

DNA-METHYLATION AND MENTAL HEALTH

	Inter	nalizing		Atten	tion Pro	blems	Aggre	ession	8	Rule-	Breakin	g	ADH	D		ODD			CD		
									No furtl	ner cova	riates ^a	-									
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	0.15	0.05, 0.25	0.004	0.10	-0.01, 0.21	0.083	0.17	0.04, 0.31	0.013	0.06	-0.05, 0.16	0.270	0.09	-0.01, 0.19	0.084	0.08	-0.04, 0.19	0.194	0.16	0.02, 0.29	0.026
DunedinPoAm	0.15	0.05, 0.25	0.002	0.10	0.00, 0.20	0.058	0.11	-0.02, 0.25	0.094	0.00	-0.07, 0.08	0.948	0.11	0.00, 0.21	0.042	0.05	-0.08, 0.17	0.459	0.05	-0.09, 0.46	0.301
PGS-CRP	0.14	0.04, 0.23	0.004	0.16	0.05, 0.28	0.004	0.11	-0.03, 0.23	0.115	0.09	0.00, 0.17	0.059	0.02	-0.09, 0.12	0.766	0.05	-0.06, 0.15	0.386	0.01	-0.18, 0.19	0.953
									Contro	lling for	BMI										
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
BMI	0.13	0.04, 0.22	0.004	-0.02	-0.12, 0.07	0.669	0.14	0.02, 0.26	0.020	0.05	-0.03, 0.14	0.201	0.09	-0.01, 0.2	0.076	0.03	-0.08, 0.15	0.589	0.09	-0.1, 0.29	0.342
DNAm-CRP	0.08	-0.03, 0.19	0.158	0.10	-0.02, 0.23	0.112	0.10	-0.05, 0.25	0.182	0.03	-0.09, 0.15	0.627	0.04	-0.08, 0.16	0.471	0.06	-0.08, 0.19	0.405	0.11	-0.02, 0.23	0.110
DunedinPoAm	0.10	-0.01, 0.2	0.069	0.10	-0.01, 0.22	0.085	0.05	-0.1, 0.19	0.526	-0.03	-0.13, 0.07	0.583	0.08	-0.04, 0.19	0.179	0.03	-0.1, 0.15	0.668	0.15	-0.1, 0.4	0.240
PGS-CRP	0.12	0.03, 0.22	0.009	0.17	0.06, 0.28	0.003	0.09	-0.04, 0.22	0.182	0.08	-0.01, 0.17	0.075	0.01	-0.1, 0.11	0.934	0.05	-0.06, 0.15	0.383	0.00	-0.19, 0.18	0.974
								Con	trolling	for pube	ertal stat	us									
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
Puberty	-0.04	-0.14, 0.06	0.435	-0.14	-0.25, -0.03	0.013	-0.01	-0.15, 0.12	0.865	-0.03	-0.13, 0.08	0.598	-0.03	-0.13, 0.07	0.568	-0.04	-0.17, 0.08	0.496	-0.16	-0.43, 0.11	0.245
DNAm-CRP	0.16	0.05, 0.26	0.004	0.12	0.01, 0.24	0.034	0.18	0.03, 0.32	0.016	0.06	-0.07, 0.19	0.356	0.10	-0.01, 0.2	0.076	0.08	-0.04, 0.2	0.175	0.19	0.02, 0.35	0.025
DunedinPoAm	0.15	0.05, 0.25	0.002	0.11	0.01, 0.21	0.040	0.11	-0.02, 0.25	0.098	0.01	-0.09, 0.1	0.920	0.11	0.01, 0.22	0.041	0.05	-0.07, 0.17	0.439	0.20	-0.09, 0.48	0.180
PGS-CRP	0.14	0.04, 0.23	0.004	0.17	0.06, 0.28	0.003	0.10	-0.03, 0.23	0.116	0.09	0, 0.18	0.051	0.02	-0.09, 0.12	0.775	0.05	-0.06, 0.15	0.374	0.01	-0.18, 0.19	0.949
								Con	trolling	for DNA	Am-smo	ke									
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-smoke	-0.07	-0.2, 0.06	0.301	0.05	-0.09, 0.19	0.500	-0.10	-0.26, 0.06	0.217	-0.03	-0.15, 0.1	0.674	-0.05	-0.18, 0.08	0.454	0.06	-0.07, 0.2	0.354	-0.02	-0.2, 0.17	0.850
DNAm-CRP	0.19	0.06, 0.31	0.004	0.08	-0.07, 0.22	0.299	0.11	-0.15, 0.37	0.400	0.08	-0.05, 0.21	0.245	0.12	-0.01, 0.24	0.069	0.05	-0.09, 0.18	0.517	0.16	0.03, 0.3	0.021
DunedinPoAm	0.16	0.06, 0.26	0.002	0.08	-0.02, 0.19	0.127	0.13	-0.01, 0.27	0.076	0.00	-0.09, 0.1	0.926	0.12	0.01, 0.23	0.031	0.03	-0.1, 0.15	0.665	0.19	-0.08, 0.45	0.177

Table S7. Associations between DNA-methylation and genetic profiles with parent-reported mental health.

PGS-CRP	0.14	0.04, 0.23	0.004	0.16	0.05, 0.27	0.005	0.11	-0.02, 0.24	0.100	0.09	0, 0.11	0.057	0.02	-0.09, 0.12	0.753	0.04	-0.06, 0.15	0.411	0.01	-0.18, 0.19	0.958
Controlling for l	Family-	Level Di	isadvant	age																	
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
Family-level disadvantage	0.21	0.13, 0.3	0.000	0.14	0.04, 0.23	0.005	0.21	0.1, 0.32	0.000	0.10	0.02, 0.18	0.017	0.15	0.06, 0.25	0.001	0.13	0.04, 0.22	0.007	0.13	-0.04, 0.29	0.136
DNAm-CRP	0.09	-0.01, 0.19	0.078	0.06	-0.05, 0.17	0.301	0.12	-0.03, 0.26	0.111	0.03	-0.08, 0.14	0.593	0.05	-0.06, 0.15	0.375	0.04	-0.07, 0.15	0.472	0.12	0, 0.25	0.056
DunedinPoAm	0.08	-0.02, 0.18	0.108	0.06	-0.05, 0.16	0.289	0.05	-0.09, 0.18	0.497	-0.03	-0.13, 0.06	0.490	0.06	-0.05, 0.17	0.258	0.01	-0.11, 0.12	0.935	0.15	-0.11, 0.41	0.251
PGS-CRP	0.13	0.04, 0.21	0.005	0.16	0.05, 0.27	0.004	0.10	-0.03, 0.22	0.135	0.08	0, 0.17	0.062	0.01	-0.09, 0.12	0.813	0.04	-0.06, 0.15	0.406	0.00	-0.18, 0.19	0.962

Standardized regression coefficients (*r*) and 95% confidence intervals (CIs) and uncorrected *p*-value calculated by regressing parent-reported mental health on DNA-methylation measures and PGS-CRP with and without controlling for normed BMI z-scores, puberty (residualized for age within each gender), DNA-methylation profiles of smoking (DNAm-smoke), and family-level disadvantage separately. *P*-values, where FDR corrected *p*-values < 0.05, are marked in bold. PGS analyses were restricted to participants of European ancestries as indicated by genetic ancestry PCs that are comparable to the GWAS discovery sample. ^aAll models included covariate adjustment for age, gender, and an age by gender interaction. Methylation scores and PGS-CRP were residualized for technical covariates (for methylation: array, slide, batch, cell composition; for PGS-CRP: genetic ancestry PCs). *Note:* ADHD=attention-deficit hyperactivity disorder; ODD=oppositional defiant disorder; CD= conduct disorder.

DNA-METHYLATION AND MENTAL HEALTH

	In	ternaliz	ing	Atten	tion Pro	oblems	A	ggressi	on	Ru	le-Brea	king		ADHD			ODD			CD	
									No furt	ner cova	riates ^a										
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	0.00	-0.12, 0.12	0.999	0.05	-0.07, 0.18	0.386	0.10	-0.03, 0.22	0.116	0.16	0.03, 0.29	0.018	0.05	-0.08, 0.19	0.427	0.00	-0.11, 0.12	0.952	0.04	-0.05, 0.13	0.386
DunedinPoAm	-0.11	-0.22, 0.00	0.060	0.00	-0.05, 0.05	0.976	0.03	-0.09, 0.14	0.667	-0.01	-0.13, 0.10	0.848	-0.03	-0.14, 0.07	0.562	0.03	-0.08, 0.15	0.564	0.01	-0.09, 0.11	0.783
PGS-CRP	0.01	-0.10, 0.12	0.885	0.01	-0.10, 0.12	0.833	-0.03	-0.13, 0.07	0.597	0.00	-0.11, 0.11	0.980	-0.04	-0.16, 0.08	0.497	-0.02	-0.13, 0.08	0.690	-0.02	-0.09, 0.06	0.692
									Contro	lling for	BMI										
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
BMI	0.06	-0.04, 0.16	0.244	0.04	-0.06, 0.14	0.443	0.10	0.00, 0.2	0.050	-0.02	-0.12, 0.08	0.670	-0.01	-0.11, 0.09	0.799	0.15	0.05, 0.26	0.005	-0.03	-0.13, 0.08	0.631
DNAm-CRP	-0.02	-0.16, 0.11	0.741	0.04	-0.09, 0.17	0.555	0.06	-0.08, 0.19	0.393	0.18	0.03, 0.32	0.016	0.06	-0.08, 0.21	0.399	-0.05	-0.17, 0.07	0.421	0.05	-0.05, 0.15	0.305
DunedinPoAm	-0.14	-0.27, -0.02	0.025	-0.02	-0.13, 0.1	0.777	-0.02	-0.14, 0.11	0.769	-0.02	-0.15, 0.10	0.731	-0.04	-0.15, 0.08	0.545	-0.01	-0.13, 0.12	0.940	0.02	-0.11, 0.15	0.760
PGS-CRP	0.00	-0.11, 0.11	0.963	0.01	-0.1, 0.12	0.895	-0.03	-0.13, 0.07	0.534	0.00	-0.11, 0.11	0.985	-0.04	-0.14, 0.08	0.494	-0.03	-0.13, 0.08	0.598	-0.02	-0.09, 0.06	0.706
								(Controll	ing for I	Puberty										
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
Puberty	0.15	0.04, 0.26	0.008	0.15	0.04, 0.26	0.008	0.15	0.04, 0.26	0.010	0.17	0.05, 0.3	0.008	0.10	-0.02, 0.22	0.106	0.18	0.07, 0.29	0.001	0.08	0, 0.16	0.047
DNAm-CRP	-0.03	-0.15, 0.10	0.686	0.03	-0.1, 0.15	0.651	0.07	-0.05, 0.2	0.242	0.13	0, 0.26	0.049	0.04	-0.1, 0.18	0.566	-0.02	-0.14, 0.1	0.716	0.03	-0.07, 0.12	0.555
DunedinPoAm	-0.12	-0.23, 0.00	0.042	-0.01	-0.11, 0.1	0.910	0.02	-0.1, 0.13	0.776	-0.02	-0.13, 0.09	0.718	-0.03	-0.14, 0.07	0.517	0.03	-0.09, 0.15	0.614	0.01	-0.09, 0.11	0.826
PGS-CRP	0.00	-0.1, 0.11	0.941	0.01	-0.1, 0.11	0.890	-0.03	-0.13, 0.07	0.518	-0.01	-0.11, 0.1	0.919	-0.04	-0.16, 0.07	0.463	-0.03	-0.14, 0.08	0.590	-0.02	-0.09, 0.06	0.636
								Con	trolling	for DNA	Am-smo	ke									
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-smoke	-0.08	-0.23, 0.06	0.264	-0.07	-0.21, 0.07	0.334	-0.07	-0.23, 0.08	0.339	-0.05	-0.21, 0.11	0.565	-0.52	-0.25, 0.04	0.153	-0.05	-0.21, 0.1	0.515	-0.07	-0.17, 0.07	0.335
DNAm-CRP	0.04	-0.12, 0.2	0.623	0.09	-0.06, 0.25	0.250	0.14	-0.02, 0.29	0.083	0.18	0.02, 0.34	0.032	0.43	-0.06, 0.27	0.196	0.01	-0.14, 0.17	0.857	0.07	-0.04, 0.12	0.235
DunedinPoAm	-0.10	-0.22, 0.02	0.111	0.01	-0.1, 0.13	0.821	0.03	-0.09, 0.16	0.601	-0.02	-0.14, 0.1	0.754	-0.01	-0.13, 0.1	0.825	0.04	-0.08, 0.17	0.511	0.03	-0.09, 0.14	0.653

Table S8. Associations between DNA-methylation and genetic profiles with self-reported mental health.

PGS-CRP	0.01	-0.1, 0.12	0.841	0.01	-0.1, 0.12	0.830	-0.03	-0.13, 0.08	0.616	0.00	-0.11, 0.11	0.994	-0.04	-0.16, 0.08	0.521	-0.02	-0.13, 0.09	0.720	-0.01	-0.09, 0.06	0.730
							Co	ontrolling	g for Fa	mily-Le	evel Disa	dvantag	je								
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
Family-level disadvantage	0.07	-0.04, 0.18	0.216	0.16	0.06, 0.26	0.002	0.22	0.12, 0.33	0.000	0.13	0.02, 0.24	0.026	0.13	0.02, 0.24	0.023	0.11	-0.01, 0.23	0.063	-0.01	-0.07, 0.05	0.665
DNAm-CRP	-0.02	-0.15, 0.11	0.803	0.02	-0.11, 0.14	0.814	0.04	-0.08, 0.17	0.510	0.13	-0.01, 0.26	0.064	0.02	-0.11, 0.16	0.751	-0.02	-0.14, 0.1	0.735	0.04	-0.05, 0.13	0.344
DunedinPoAm	-0.14	-0.26, -0.01	0.031	-0.05	-0.16, 0.07	0.432	-0.04	-0.16, 0.08	0.488	-0.06	-0.18, 0.07	0.385	-0.07	-0.18, 0.04	0.197	0.01	-0.11, 0.13	0.894	0.02	-0.09, 0.12	0.768
PGS-CRP	0.01	-0.11, 0.12	0.914	0.01	-0.1, 0.12	0.885	-0.03	-0.13, 0.07	0.570	0.00	-0.11, 0.11	0.989	-0.04	-0.16, 0.08	0.491	-0.02	-0.12, 0.08	0.709	-0.02	-0.09, 0.06	0.695

Standardized regression coefficients (*r*) and 95% confidence intervals (CIs) and uncorrected *p*-value calculated by regressing parent-reported mental health on DNA-methylation measures and PGS-CRP with and without controlling for normed BMI z-scores, puberty (residualized for age within each gender), DNA-methylation profiles of smoking (DNAm-smoke), and family-level disadvantage separately. *P*-values, where FDR corrected *p*-values < 0.05, are marked in bold. PGS analyses were restricted to participants of European ancestries as indicated by genetic ancestry PCs that are comparable to the GWAS discovery sample. ^aAll models included covariate adjustment for age, gender, and an age by gender interaction. Methylation scores and PGS-CRP were residualized for technical covariates (for methylation: array, slide, batch, cell composition; for PGS-CRP: genetic ancestry PCs). *Note:* ADHD=attention-deficit hyperactivity disorder; ODD=oppositional defiant disorder; CD= conduct disorder.

	In	ternaliz	ing	Atten	tion Pro	blems	A	Aggressio	n	Ru	le-Break	king		ADHD			ODD			CD	
										White											
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	0.22	0.09, 0.34	0.001	0.08	-0.06, 0.22	0.235	0.28	0.09, 0.46	0.003	0.09	-0.05, 0.22	0.211	0.15	0.01, 0.29	0.038	0.10	-0.04, 0.24	0.156	0.26	-0.87, 1.39	0.652
DunedinPoAm	0.18	0.04, 0.31	0.010	0.12	-0.02, 0.27	0.090	0.10	-0.1, 0.3	0.320	0.06	-0.07, 0.19	0.372	0.11	-0.04, 0.25	0.157	0.01	-0.14, 0.15	0.945	-0.08	-1.07, 0.9	0.869
										Latinx											
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	0.21	-0.02, 0.44	0.069	-0.03	-0.37, 0.3	0.841	0.26	-0.05, 0.58	0.104	0.08	-0.33, 0.49	0.700	0.10	-0.11, 0.31	0.339	0.03	-0.15, 0.21	0.737	-0.08	-0.26, 0.11	0.405
DunedinPoAm	0.28	0.07, 0.49	0.009	0.00	-0.34, 0.34	0.980	0.47	0.09, 0.84	0.014	-0.02	-0.26, 0.21	0.853	0.21	0.02, 0.4	0.034	0.02	-0.12, 0.17	0.742	0.06	-0.13, 0.26	0.521
									Lat	inx-Whi	te										
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	-0.27	-0.5, - 0.04	0.020	-0.01	-0.32, 0.3	0.935	-0.18	-0.5, 0.13	0.252	0.07	-0.14, 0.28	0.508	-0.02	-0.36, 0.33	0.917	0.19	-0.24, 0.61	0.387	0.10	-0.16, 0.36	0.438
DunedinPoAm	-0.07	-0.29, 0.15	0.523	0.05	-0.26, 0.36	0.756	-0.07	-0.4, 0.26	0.681	-0.11	-0.35, 0.13	0.356	0.09	-0.18, 0.35	0.525	0.12	-0.18, 0.42	0.442	0.04	-0.17, 0.25	0.693
]	Black +											
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	0.17	-0.44, 0.77	0.585	0.49	-0.3, 1.27	0.223	0.12	-0.5, 0.74	0.701	-0.05	-0.66, 0.56	0.872	-0.21	-0.62, 0.2	0.321	-0.30	-0.92, 0.32	0.342	0.52	-0.23, 1.28	0.175
DunedinPoAm	-0.04	-0.4, 0.31	0.813	-0.07	-0.36, 0.22	0.645	0.04	-0.39, 0.48	0.852	-0.15	-0.44, 0.15	0.326	0.14	-0.23, 0.52	0.459	0.20	-0.32, 0.73	0.450	0.68	0.13, 1.23	0.016
									1	Asian +											
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	-0.14	-0.42, 0.13	0.308	-0.17	-0.65, 0.31	0.484	0.04	-0.27, 0.36	0.791	-0.25	-0.61, 0.12	0.186	0.04	-0.16, 0.25	0.696	-0.08	-0.37, 0.22	0.614	-0.11	-0.38, 0.17	0.439
DunedinPoAm	-0.28	-0.63, 0.08	0.130	-0.26	-0.75, 0.23	0.294	-0.29	-0.67, 0.1	0.149	-0.24	-0.61, 0.13	0.200	-0.08	-0.34, 0.18	0.541	-0.07	-0.35, 0.22	0.648	0.55	0.3, 0.8	0.000

Table S9. Associations between DNA-methylation with parent-reported mental health for each racial/ethnic group.

Standardized regression coefficients (*r*) and 95% confidence intervals (CIs) and uncorrected *p*-value calculated by regressing parent-reported mental health on DNA-methylation measures within each racial/ethnic group. ^aAll models included covariate adjustment for age, gender, and an age by gender interaction. Methylation scores were residualized for technical covariates (array, slide, batch, cell composition). *Note:* ADHD=attention-deficit hyperactivity disorder; ODD=oppositional defiant disorder; CD= conduct disorder.

	In	ternalizi	ing	Atten	tion Pro	blems	A	ggressio	n	Ru	le-Break	king		ADHD			ODD			CD	
										White											
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	0.07	-0.1, 0.23	0.432	0.08	-0.07, 0.23	0.278	0.073	-0.08, 0.22	0.344	0.16	0.01, 0.32	0.038	0.12	-0.04, 0.28	0.137	0.00	-0.15, 0.14	0.956	0.08	0, 0.16	0.058
DunedinPoAm	-0.12	-0.29, 0.05	0.161	0.05	-0.1, 0.19	0.522	-0.01	-0.16, 0.15	0.938	0.05	-0.1, 0.2	0.497	-0.01	-0.16, 0.14	0.892	0.02	-0.15, 0.2	0.795	0.10	-0.01, 0.22	0.073
										Latinx											
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	-0.40	-0.76, -0.04	0.030	-0.08	-0.38, 0.22	0.587	0.05	-0.37, 0.46	0.823	-0.05	-0.38, 0.29	0.790	-0.24	-0.66, 0.18	0.269	-0.23	-0.52, 0.06	0.115	-0.14	-0.9, 0.61	0.714
DunedinPoAm	-0.03	-0.29, 0.24	0.853	0.02	-0.25, 0.29	0.903	0.14	-0.11, 0.38	0.266	-0.06	-0.26, 0.14	0.548	-0.07	-0.33, 0.2	0.616	0.05	-0.2, 0.3	0.705	-0.40	-1.11, 0.3	0.260
									Lat	inx-Whi	te										
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	-0.30	-0.67, 0.07	0.110	0.03	-0.27, 0.33	0.845	0.07	-0.34, 0.48	0.740	-0.05	-0.35, 0.26	0.768	-0.14	-0.56, 0.28	0.511	0.31	-0.17, 0.78	0.207	-0.41	-0.83, 0.01	0.058
DunedinPoAm	-0.02	-0.28, 0.24	0.889	0.03	-0.24, 0.3	0.839	0.14	-0.11, 0.39	0.262	-0.01	-0.26, 0.14	0.533	-0.03	-0.31, 0.25	0.828	0.08	-0.14, 0.3	0.463	-0.22	-0.94, 0.51	0.559
]	Black +											
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	0.16	-0.39, 0.7	0.575	-0.28	-0.95, 0.39	0.411	0.33	-0.35, 1.01	0.344	0.29	-0.3, 0.89	0.330	-0.04	-0.8, 0.71	0.917	0.27	-0.16, 0.69	0.218	1.01	-2.43, 4.46	0.564
DunedinPoAm	-0.12	-0.41, 0.16	0.393	-0.28	-0.64, 0.08	0.125	0.18	-0.18, 0.53	0.325	-0.07	-0.35, 0.22	0.637	-0.27	-0.58, 0.04	0.089	0.28	-0.06, 0.62	0.104	-0.05	-2.92, 2.82	0.972
									1	Asian +											
	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р	r	CI	р
DNAm-CRP	0.13	-0.24, 0.49	0.496	0.14	-0.24, 0.51	0.474	0.09	-0.32, 0.49	0.664	0.08	-0.33, 0.49	0.714	-0.21	-0.47, 0.04	0.103	-0.19	-0.53, 0.14	0.255	-0.11	-0.55, 0.33	0.619
DunedinPoAm	-0.20	-0.62, 0.22	0.355	-0.10	-0.56, 0.37	0.689	-0.28	-0.77, 0.2	0.252	-0.25	-0.66, 0.17	0.251	-0.27	-0.56, 0.03	0.073	-0.40	-0.72, -0.08	0.014	-0.28	-0.78, 0.22	0.269

Table S10. Associations between DNA-methylation with self-reported mental health for each racial/ethnic group.

Standardized regression coefficients (*r*) and 95% confidence intervals (CIs) and uncorrected *p*-value calculated by regressing self-reported mental health on DNA-methylation measures within each racial/ethnic group. ^aAll models included covariate adjustment for age, gender, and an age by gender interaction. Methylation scores were residualized for technical covariates (array, slide, batch, cell composition). *Note:* ADHD=attention-deficit hyperactivity disorder; ODD=oppositional defiant disorder; CD= conduct disorder.

It is made available under a CC-BY-NC-ND 4.0 International license .

DNA-METHYLATION AND MENTAL HEALTH

Raffington et al. 50

	DNAm-	CRP		Dunedi	nPoAm	
	rA	CI	р	rA	CI	р
Internalizing	0.28	-0.10, 0.66	0.146	0.30	-0.19, 0.78	0.226
Attention problems	0.23	-0.24, 0.70	0.339	0.21	-0.38, 0.80	0.488
Aggression	0.30	0.02, 0.58	0.037	0.15	-0.23, 0.53	0.431
Rule-breaking	0.19	-0.03, 0.41	0.086	0.12	-0.14, 0.38	0.375
ADHD	0.05	-0.09-0.18	0.505	0.077	-0.08, 0.24	0.340
ODD	0.08	-0.10, 0.26	0.370	0.03	-0.30, 0.36	0.854
CD	0.26	-0.50, 1.02	0.502	0.40	-0.43, 1.22	0.348
	rC	CI	р	rC	CI	р
Internalizing	-1	-1, -1	< 0.001	-0.48	-5.72, 4.75	0.857
Attention problems	-1	-1, -1	< 0.001	0.18	-2.70, 3.06	0.904
Aggression	1	1, 1	< 0.001	1	1, 1	< 0.001
Rule-breaking	-0.96	-1.71, -0.22	0.011	1	1, 1	< 0.001
ADHD	-0.96	-0.96, -0.96	< 0.001	1	1, 1	< 0.001
ODD	-0.78	-3.34, 1.78	0.553	1	1, 1	< 0.001
CD	-1	-1, -1	< 0.001	-1	-1, -1	< 0.001
	rE	CI	р	rE	CI	р
Internalizing	-0.07	-0.19, 0.05	0.248	-0.03	-0.16, 0.10	0.671
Attention problems	-0.04	-0.17, 0.10	0.591	-0.05	-0.18, 0.08	0.470
Aggression	-0.12	-0.27, 0.03	0.117	-0.14	-0.28, 0.01	0.062
Rule-breaking	-0.09	-0.22, 0.03	0.145	-0.11	-0.23, 0.01	0.075
ADHD	0.08	-0.05, 0.21	0.221	0.09	-0.02, 0.20	0.106
ODD	0.02	-0.09, 0.12	0.766	-0.04	-0.13, 0.06	0.424
CD	0.04	-0.07, 0.15	0.436	0.01	-0.09, 0.10	0.915

Table S11. Co-twin-control associations between DNA-methylation and parent-reported
mental health.

Regression coefficients, 95% confidence intervals (CIs), and *p*-value calculated in a bivariate biometric model that decomposed the association between DNA-methylation and cognition into components representing additive genetic factors (*A*), environmental factors shared by twins living in the same home (*C*), and environmental factors unique to each twin (*E*). *r*A is the correlation between the *A* components of variation in DNA-methylation and cognition, which reflects the extent to which genetic variation in DNAm accounts for differences in cognitive functioning. *r*C is the correlation between the *C* components of variation in DNAm accounts for differences in cognition, which reflects the extent to which reflects the extent to which shared environmental variation in DNAm accounts for differences in cognitive functioning. *r*E is the correlation between the *E* components of variation in DNA-methylation and cognition in DNAm accounts for differences in cognitive functioning. *r*E is the extent to which shared environmental variation in DNAm accounts for differences in cognitive functioning. *r*E is the correlation between the *E* components of variation in DNA-methylation and cognition, which reflects the extent to which identical twins who differ from their co-twins in DNAm show corresponding differences in their cognitive functioning. *Note:* ADHD=attention-deficit hyperactivity disorder; ODD=oppositional defiant disorder; CD= conduct disorder.