It is made available under a CC-BY-NC-ND 4.0 International license .

New insights from GWAS on longitudinal and cross-sectional BMI and related phenotypes in admixed children with Native American and European ancestries

Lucas Vicuña<sup>1,2</sup>, Esteban Barrientos<sup>1,2</sup>, Tomás Norambuena<sup>1,2</sup>, Danilo Alvares<sup>3</sup>, Juan Cristobal Gana<sup>4</sup>, Valeria Leiva<sup>3</sup>, Veronica Mericq<sup>5</sup>, Cristian Meza<sup>6</sup>, Ana Pereira<sup>7</sup>, José L. Santos<sup>8</sup>, Susana Eyheramendy<sup>1,2\*</sup>

 Faculty of Engineering and Sciences, Universidad Adolfo Ibáñez, Santiago, Chile.
 Instituto Milenio de Investigación Sobre los Fundamentos de los Datos (IMFD), Santiago, Chile.

3 Department of Statistics, Pontificia Universidad Católica de Chile, Santiago, Chile.
4 Department of Pediatric Gastroenterology and Nutrition. Division of Pediatrics.
School of Medicine. Pontificia Universidad Católica de Chile, Santiago, Chile
5 Institute of Maternal and Child Research, Faculty of Medicine, University of Chile, Santiago, Chile.

**6** INGEMAT-CIMFAV, Faculty of Engineering, Universidad de Valparaiso, Valparaiso, Chile.

7 Institute of Nutrition and Food Technology, University of Chile, Santiago, Chile.

8 Department of Nutrition, Diabetes and Metabolism, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile.

\* Corresponding author: susana.eyheramendy@uai.cl

### Abstract

Body-mass index (BMI) is a well-known marker of adiposity across all ages. The genetic architecture of BMI has been thoroughly studied among adults. In contrast, there are a few genome-wide association studies (GWAS) on children. Further, GWAS on children have been performed almost exclusively in Europeans at single ages. We aimed to better understand the genetic architecture of BMI trajectory across ages and how BMI is affected by Native American genetic ancestry. We performed cross-sectional and longitudinal GWAS for BMI-related traits on 904 admixed Chilean children with mostly European and Mapuche Native American genetic ancestry. We focused on BMI and two traits that occur at the minimum of the childhood BMI growth trajectory, namely, age at adiposity rebound (Age-AR) and BMI at adiposity rebound (BMI-AR). We found several variants in the immune gene HLA-DQB3 that are strongly associated with BMI at ages 1.5-2.5 years old, but not at other ages. We also identified a variant in the sex-determining gene DMRT1 significantly associated with Age-AR ( $P = 9.8 \times 10^{-9}$ ). Further, BMI was significantly higher in Mapuche than in European children at all ages between 5.5 and 16.5 years old, but not before. Finally, Age-AR was significantly lower (P = 0.013) by 1.64 years in the Mapuche children compared with Europeans.

# Introduction

Childhood obesity is a major public health problem across the world. One well-studied indicator of obesity is the body mass index (BMI), whose alterations in children have been associated with risk of adult obesity, as well as related diseases, including type 2

diabetes 1 and cardiometabolic diseases 2. Even though several environmental causes explain an important part of childhood obesity, there is still a lack of knowledge about the genetic factors underlying susceptibility to this disease.

BMI follows a nonlinear trajectory over time during childhood. BMI trajectory is characterized by three main phases: i) a rapid increase with an adiposity peak (AP) at  $\sim 9$  month of age; ii) a decline reaching its lowest value at  $\sim 5.5$  years, also called the adiposity rebound (AR); and iii) a subsequent peak at early adulthood [3].

The heritability of adult obesity has been estimated to be 40 - 70% [34]. Approximately 22.4% of adult BMI heritability is explained by 941 SNPs, according to a genome-wide association study (GWAS) meta-analysis performed in  $\sim 700,000$ individuals of European ancestry 5. Such studies have been mostly performed in adults, but for most of these loci it is unknown whether or not and to which extent they also affect BMI in children. Some cross-sectional GWA studies on European populations have shown that several loci influence BMI in adulthood and childhood, whereas other loci seem to act only during adulthood or childhood. For instance, a GWAS found variants in the FTO and MC4R genes significantly associated with BMI in children (< 6 vears old) as well as in adults, but a locus in MAF only associated in adults 6. A GWAS meta-analysis found significant associations with FTO, MC4R, TMEM18, SDCCAG8 and TNKS/MSRA loci, but associations for the later gene were limited to children and adolescents 7. Another GWAS meta-analysis identified 15 loci significantly associated with BMI (they considered a single BMI value at the oldest age between 2 and 10 years). Among them, 12 were previously associated with adult BMI or childhood obesity, suggesting that the 3 remaining loci, located near ELP3, RAB27B and ADAM23 genes, act only during childhood 8. A further GWAS meta-analysis identified 18 SNPs significantly associated with BMI in paediatric cohorts from diverse ancestries pooled together, although most of these loci only reached the genome-wide significance threshold in Europeans when same-ancestry cohorts were analyzed separately. Most of these loci had been previously associated with adult BMI, childhood BMI or childhood obesity 9.

A few GWAS on European children have identified loci affecting BMI in distinct phases of child growth. A recent GWAS meta-analysis identified 4 loci significantly associated with childhood BMI-related traits: one LEPR/LEPROT gene variant associated with BMI-AP; one FTO variant and one TFAP2B variant associated with Age-AR; and a GNPDA2 variant associated with BMI-AR. Among them, the TFAP2B. GNPDA2 and FTO variants are also associated with adult BMI, whereas the LEPR/LEPROT variant only associates with a BMI-related trait in the infant phase. The authors suggested that adult BMI-related variants start influencing BMI by the time of AR [3]. A candidate gene study on a European multi-cohort found that a FTOlocus is positively associated with BMI from 5.5 years onward, but is inversely associated below age 2.5 years old. They also confirmed this findings using a longitudinal linear mixed model 13. Warrington et al 2015 performed a longitudinal GWAS to identify associations between genotypes and BMI trajectories across childhood [14]. They found 4 loci previously associated with BMI in adults or children as well as a novel locus in the FAM120AOS gene significantly associated with BMI at 8 years of age. To our knowledge, Warrington et al 2015 is the only longitudinal GWAS on childhood BMI.

As mentioned before, the vast majority of these GWA studies have been performed in Europeans, but we do not know to what extent these loci affect BMI in populations with other continental ancestries, in particular those with Native American ancestry. This is relevant, as obesity and obesity-related disorders are markedly affected by genetic ancestry 34. For instance, Latinos, which usually have mixed European, Native American and African ancestries, are more susceptible to lipid-related disorders than any other US group, in part due to their Native-American genetic heritage 15. 10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

36

37

38

39

40

41

42

43

44

45

46

47

49

50

51

52

53

54

55

It is made available under a CC-BY-NC-ND 4.0 International license .

The aims of this study are twofold: i) to estimate how much does Native American ancestry affect childhood BMI trajectory, Age-AR as well as BMI-AR; and ii) to identify genetic variants associated with these traits.

# Materials and methods

### Sample collection and genotyping.

We analyzed individuals of the "Growth and Obesity Chilean Cohort Study" (GOCS; [22]), who were genotyped using the Infinium <sup>®</sup> Multhi-Ethnic Global BeadChip (Illumina), as described in [21]. After quality controls were applied, we obtained a filtered data set of 904 individuals and 774, 433 autosomal SNPs. Further details can be found in [21].

### Weight and height measurements.

Weight and height measurements were taken since 2006 at Instituto de Nutrición y Tecnología de los Alimentos (INTA), Santiago, Chile, on children/adolescents in barefoot and light clothes by a trained dietitian following standardized protocols; ICC for all measurements was  $\geq 0.75$  [41]. Weight and height were measured twice at each individual visit, and the average was considered as the final value. Before and after puberty, weight and height were measured once per year; during puberty they were measured every 6 months. Weight was measured with a portable electronic scale (Seca 770), precision of 0.1 kg, and height was measured with a portable stadiometer (Harpenden 603) to the nearest 0.1 cm. BMI (kg/m<sup>2</sup>) and Z-scores (for < 18 years old) were calculated based on the World Health Organization 2007 growth references. Weight and height data prior to 2006 were retrieved from medical records.

### Local Ancestry Estimation.

We used RFMIX [48] to infer the local ancestry of each SNP allele from our Chilean sample. We used reference populations from the 1000 Genomes Project [49], namely, Yoruba (YRI, n=108) for the African ancestry, Utah Residents with Northern and Western European Ancestry (CEU, n=36) for the European ancestry, and Peruvian (PEL) individuals with > 95% Native American ancestry (n=29). We excluded individuals with > 5% SNP missing rate. We inferred the gametic phase of individuals with the Beagle 5 software [24], using the HapMap37 human genome build 37 recombination map. To obtain the local ancestry, we used the --forward-backward parameter, as recommended in the manual. Local ancestry estimation identified three continental ancestries: European, African and Native American.

### **Global Ancestry Estimation.**

Global ancestry proportions of Chilean children were estimated with Admixture 1.3 23, using K=4 ancestral populations. This K value was chosen because it was able to distinguish the two main Native American subcomponents of Chileans, namely, Mapuche and Aymara 21. In order to distinguish Peruvian (PEL) individuals with > 95% Native American ancestry, we used K=3 ancestral populations and the reference populations mentioned in the previous section. Global ancestry estimation identified four proportion of ancestries corresponding to European, African, Aymara and Mapuche. We used the Mapuche proportion ancestry in the GWAS to identify its effect on the phenotypes.

57

58

61

63

65

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

qq

It is made available under a CC-BY-NC-ND 4.0 International license .

Variant and gene annotations.

Variant annotations [GRCh37 (hg19) assembly] were retrieved with the web tool Variant 101 Effect Predictor (VEP) from Ensembl 25. Upstream and downstream variants were 102 defined as those located 10 Kb upstream or downstream of the gene, respectively. 103 Intergenic variants were defined as those located > 100 Kb upstream or downstream of 104 the closest gene. We used HaploReg v4.1 26 to identify the genes located closest to 105 associated intergenic variants. Reported GWAS associations were retrieved from the 106 NHGRI GWAS Catalog 27. When more than one variant in a gene has been 107 associated with the same phenotype, we reported the strongest association. 108

### Derivation of Age-AR and BMI-AR

Before modeling BMI trajectory over time, we applied some filters on the individuals. We excluded 51 individuals who had no sufficient measurements between 2 and 10 years old. We also excluded 156 individuals with all their measurements taken after 4.5 years old. We obtained a final dataset of 696 individuals with measurements between 2 and 10 years old. To estimate Age-AR, we implemented a longitudinal statistical model of BMI (in logarithm scale) on this filtered data. The model equation is:

$$\log(BMI)_{ij} = (\beta_1 S_i + \beta_0) + (\beta_2 S_i + \beta_3) Age_{ij} + (\beta_4 S_i + \beta_5) Age_{ij}^2 + (\beta_6 S_i + \beta_7) Age_{ij}^3 + \beta_{8i} + \beta_{9i} Age_{ij} + \beta_{10i} Age_{ij}^2 + \beta_{11i} Age_{ij}^3 + \epsilon_{ij}$$

$$j = 1, 2, \dots, n_i \text{ and } i = 1, \dots, 696.$$
(1)

where  $n_i$  corresponds to the number of Age/BMI measurements for individual i,  $S_i$ represents the gender of the individual (0 female, 1 male). The parameters  $\beta_k$  for k = 1, ..., 7 are fixed effects whereas the parameters  $\beta_{ki}$  for k = 8, ..., 11 are random effects for each individual i (i.e.  $\beta_{ki} \sim N(0, \sigma_k^2)$ ). The errors  $\{\epsilon_{ij}\}$  are assumed independent across the different individuals i but dependent between observations for the same individual (index j), and are normally distributed with variance  $\sigma^2$ .

The predicted trajectory for individual i can then be written as

$$\log(\hat{BMI})_{ij} = (\hat{\beta}_1 S_i + \hat{\beta}_0) + (\hat{\beta}_2 S_i + \hat{\beta}_3) Age_{ij} + (\hat{\beta}_4 S_i + \hat{\beta}_5) Age_{ij}^2 + (\hat{\beta}_6 S_i + \hat{\beta}_7) Age_{ij}^3 + \hat{\beta}_{8i} + \hat{\beta}_{9i} Age_{ij} + \hat{\beta}_{10i} Age_{ij}^2 + \hat{\beta}_{11i} Age_{ij}^3$$
$$j = 1, 2, \dots, n_i \text{ and } i = 1, \dots, 696.$$
(2)

where  $\hat{}$  (hat) represents the estimators of the parameters. From this trajectory, the Age 123 of Adiposity rebound (Age-AR) and the BMI at Age-AR are obtained by finding the 124 Age at which the minimum BMI is found. 125

A local minimum of BMI is found by solving the following equation:

$$3(\hat{\beta}_6 S_i + \hat{\beta}_7 + \hat{\beta}_{11i})x_i^2 + 2(\hat{\beta}_4 S_i + \hat{\beta}_5 + \hat{\beta}_{10i})x_i + (\hat{\beta}_2 S_i + \hat{\beta}_3 + \hat{\beta}_{9i}) = 0$$
(3)

which has as its solution:

$$Age-AR_{i} = \frac{1}{3(\hat{\beta}_{6}S_{i} + \hat{\beta}_{7} + \hat{\beta}_{11i})} (-(\hat{\beta}_{4}S_{i} + \hat{\beta}_{5} + \hat{\beta}_{10i}) + \pm 2\sqrt{(\hat{\beta}_{4}S_{i} + \hat{\beta}_{5} + \hat{\beta}_{10i})^{2} - 3(\hat{\beta}_{6}S_{i} + \hat{\beta}_{7} + \hat{\beta}_{11i})(\hat{\beta}_{2}S_{i} + \hat{\beta}_{3} + \hat{\beta}_{9i})} )$$
(4)

100

109

122

126

It is made available under a CC-BY-NC-ND 4.0 International license .

where the minimum obtained above needs to satisfy:

$$6(\hat{\beta}_6 S_i + \hat{\beta}_7 + \hat{\beta}_{11i}) \text{Age-AR}_i + 2(\hat{\beta}_4 S_i + \hat{\beta}_5 + \hat{\beta}_{10i}) > 0$$
(5)

To identify BMI-AR, we replace the Age term from equation (1) by the Age-AR value obtained from equation (3).

#### Cross-sectional GWAS on BMI.

For the cross-sectional GWAS, we analyzed individuals between 1.5 and 16.5 years old, 132 across 16 strata of 1 year interval each (S1 Table). BMI was expressed as  $kg/m^2$  and 133 age in years. Individuals with no weight or height measurements in a particular stratum 134 were removed. When > 1 measurement was taken in the same individual and time 135 interval, we took the mean of the BMI measurements. To get a normal distribution of 136 BMI values, we applied Box-Cox transformations ( $\lambda$  values for each stratum are shown 137 in S1 Table). These transformed values were standardized to get mean = 0 and 138 standard deviation = 1, obtaining zBMI scores [13] [14]. These zBMI scores were used 139 to test for genome-wide associations. We applied the following linear regression model: 140

$$zBMI_i = \beta_0 + \beta_1 GT_{ij} + \beta_2 S_i + \beta_3 GA_i + \beta_4 LA_{ij} + \epsilon_i \tag{6}$$

where  $\beta_1$  represents the effect of the additive genotype (GT) of SNP j;  $\beta_2$  represents 141 the effect of gender (S);  $\beta_3$  represents the effect of the global Mapuche Native American 142 ancestry proportion (GA); and  $\beta_4$  represents the effect of local Native American 143 ancestry, where  $LA_i$  takes the values 0, 1 or 2 depending on the number of Native 144 American alleles. Of note, while our global ancestry analysis is able to separately 145 estimate the Mapuche and Aymara ancestral Native American subcomponents of 146 Chileans, the local ancestry analysis only considers a general Native American 147 population. 148

### GWAS on longitudinal BMI data.

To estimate longitudinal genotype-phenotype associations, we implemented a linear mixed model for BMI. We adjusted for gender, age, global ancestry, and local ancestry. We assumed an additive inheritance model. The model equation is as follows:

$$\log(BMI)_{ij} = (\beta_1 S_i + \beta_0) + (\beta_2 S_i + \beta_3) Age_{ij} + \beta_4 Age_{ij}^2 + \beta_5 Age_{ij}^3 + \beta_6 GA_i + \beta_7 LA_i + \beta_8 GT_i + \beta_{9i} + \beta_{10i} Age_{ij} + \beta_{11i} Age_{ij}^2 + \epsilon_{ij}$$
(7)  

$$j = 1, 2, \dots, m_i \text{ and } i = 1, \dots, 901.$$

where  $m_i$  corresponds to the number of Age/BMI measurements for individual  $i, S_i$  represents the gender of the individual (0 female, 1 male).

### GWAS on Age-AR and BMI-AR.

Similarly as with BMI, we obtained standardized Age-AR and BMI-AR values by applying Box-Cox transformations followed by standardization. We obtained estimates of  $\lambda = 0.841$  for Age-AR and  $\lambda = -0.759$  for BMI-AR. We performed genome-wide associations for standardized Age-AR, using the following linear regression model: 159

128

129

130

131

149

153

154

It is made available under a CC-BY-NC-ND 4.0 International license .

$$zAge-AR_{i} = (\beta_{1}S_{i} + \beta_{0}) + (\beta_{2}S_{i} + \beta_{3})zBMI-AR_{i} + (\beta_{4}S_{i} + \beta_{5})zBMI-AR_{i}^{2} + (\beta_{6}S_{i} + \beta_{7})zBMI-AR_{i}^{3} + \beta_{8}GA_{i} + (\beta_{9}S_{i} + \beta_{10})GT_{i} + \beta_{11}LA_{i} + \epsilon_{i}$$

$$(8)$$

where  $\beta_9$  represents the interaction effect between gender and genotype;  $(\beta_2 S_i + \beta_3), (\beta_4 S_i + \beta_5), (\beta_6 S_i + \beta_7) =$  interaction effect between sex and a third degree polynomial of zBMI-AR. For the GWAS on BMI-AR, we used a similar model: 160

$$zBMI-AR_{i} = (\beta_{1}S_{i} + \beta_{0}) + (\beta_{2}S_{i} + \beta_{3})zAge-AR_{i} + (\beta_{4}S_{i} + \beta_{5})zAge-AR_{i}^{2} + (\beta_{6}S_{i} + \beta_{7})zAge-AR_{i}^{3} + \beta_{8}GA_{i} + (\beta_{9}S_{i} + \beta_{10})GT_{i} + \beta_{11}LA_{i} + \epsilon_{i}$$

$$(9)$$

### Results

### Estimation of Age-AR and BMI-AR

We first characterized BMI trajectories in 904 children from the "Growth and Obesity" 165 Chilean Cohort Study" (GOCS) [22], from age 2 years to age 10 years (Fig 1). We 166 implemented longitudinal mixed models (see details in Methods) in boys and girls 167 pooled together as well as separately. Fig 1 shows the expected BMI trajectories for the 168 three analyses. We observed a strong fit between observed and expected curves, as 169 revealed by a Pearson's correlation coefficient of  $\rho = 0.97$ . Using the same longitudinal 170 model, we estimated BMI-AR and Age-AR (see Methods). The mean Age-AR was 4.5 171 (SD = 1.5) years in pooled children, 4.35 (SD = 1.35) years in boys and 4.55 (SD =172 1.71) years in girls. The mean BMI-AR was 16.3 (SD = 1.4) kg/m<sup>2</sup> in pooled children, 173 16.4 (SD = 1.29) kg/m<sup>2</sup> in boys and 16.2 (SD = 1.42) kg/m<sup>2</sup> in girls. 174

### Effect of global ancestry on BMI, Age-AR and BMI-AR.

The children of this cohort have on average 52.1% European, 43.8% Mapuche Native American, 2.6% Aymara Native American, and 1.5% African global ancestry proportions [21] (S1 Fig). To quantify how much Mapuche and European ancestries affect BMI at each age stratum, we applied linear regressions with BMI as the dependent variable and global Mapuche ancestry as the independent variable. We estimated BMI values for hypothetical individuals with 100% Mapuche ancestry and with 100% European ancestry, hereafter referred to as "Mapuche" and "Europeans". When compared with Europeans, Mapuche individuals have significantly higher BMI at all age strata between 5.5 and 16.5 years old, but not before 5.5 years old (Fig 2A). Indeed, intercept *P*-values were almost all < 0.01 except at 7.5-8.5 years old (P < 0.05; Fig 2A and S1 Table). Further, BMI differences between Mapuche and European individuals tend to increase with age (Fig 2A). Similarly, global ancestry had a mild but significant effect in the whole longitudinal BMI trajectory (P = 0.045; effect size = 0.094).

We explored whether individual Mapuche and European global ancestries have an effect on Age-AR and BMI-AR. By regressing Age-AR onto Mapuche global ancestry, our model predicted that Age-AR is significantly lower by 1.64 years in a Mapuche child (mean = 3.53 years) than in a European child (mean = 5.17 years; P=0.013; Fig 2B). On the other hand, BMI-AR was 16.9 and  $15.9 \text{ kg/m}^2$  in a Mapuche and a European child, respectively (Fig 2C). However, this difference was not significant.

### Identification of variants associated with BMI.

We performed GWAS on BMI in each age stratum by performing linear regression models (see Methods for details; S1 Table shows the number of individuals in each

163

164

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

6/36

It is made available under a CC-BY-NC-ND 4.0 International license .



Fig 1. Estimation of Age-AR and BMI-AR. Fitted BMI  $(kg/m^2)$  curves over time (years) in the whole cohort as well as in boys and girls considered separately. Vertical and horizontal dotted lines pinpoint the observed mean Age-AR and mean BMI-AR in boys and girls.

stratum). Besides gender and age, we adjusted for the global ancestry proportion of each individual to account for population substructure as well as for local ancestry to adjust for the ancestry of each haplotype (i.e. SNP). S3 Fig S4 Fig S5 Fig S6 Fig S7 Fig S8 Fig S9 Fig S10 Fig S11 Fig S12 Fig S13 Fig S14 Fig S15 Fig S16 Fig S17 Fig show the corresponding Manhattan plots. We found a significant association for the rs75964957 variant at 4.5 - 5.5 years old stratum ( $P < 3.5 \times 10^{-8}$ ; S1 Table). The second strongest association ( $P < 1.6 \times 10^{-7}$ ) was achieved by a genetic signal at chromosome 6 harboring several variants in high linkage disequilibrium, as revealed by a clear peak in the Manhattan plot of Fig 3A. This genetic signal was apparent in the 1.5 - 2.5 years old stratum, but not at other strata (Fig 3B).

The strongest variants of this peak, namely, rs9275582, rs9275593 and rs9275595, map a promoter flanking region of the *HLA-DQB3* pseudogene, which is part of the human leukocyte antigen (HLA) region (see Discussion). S2 Table shows all variants from the HLA peak achieving nominal genome-wide significance ( $P < 1 \times 10^{-6}$ ), with their corresponding annotations. We identified additional association peaks at chromosomes 4 and 20 (S18 Fig). The strongest peak variant at chromosome 4, rs12501266, maps the gene SORCS2, whereas the strongest peak variant at chromosome 20, rs474169, maps the gene SNAP25-AS1. Of note, rs474169 is also associated with BMI at age stratum 10.5 – 11.5 (S1 Table). In contrast to the HLA peak, peak variants at chromosomes 4 and 20 show associations across several strata during childhood and/or adolescence (S18 Fig).

In order to identify variants associated with the whole BMI trajectory during body growth, we performed a longitudinal GWAS on BMI (see Methods for details). The strongest association was achieved by rs35266519 ( $P = 2.9 \times 10^{-7}$ ), which is a missense variant of the the *GSDMB* gene. This gene has been GWAS-associated with

198

It is made available under a CC-BY-NC-ND 4.0 International license .



Fig 2. Effect of global Mapuche ancestry on BMI trajectory, Age-AR and BMI-AR. A: BMI  $(kg/m^2)$  estimated at each age stratum for predicted individuals with 100% Mapuche (red line) and 100% European global ancestry (blue line). B: Linear model of Age-AR with Mapuche ancestry as covariate. C: Linear model of BMI-AR with Mapuche ancestry as covariate.

obesity-related traits (see Discussion). S19 Fig shows the corresponding Manhattan plot, whereas S3 Table shows the associations achieving nominal genome-wide significance  $(P < 1 \times 10^{-6})$ , as well as their annotations.

### Identification of variants associated with Age-AR and BMI-AR.

To identify variants associated with Age-AR, we implemented a regression model described in the Methods section. We detected one variant significantly associated with Age-AR ( $P < 9.8 \times 10^{-9}$ ; S2 Table and S20 Fig). This variant, namely rs445398, maps an intron of the *DMRT1* gene. *DMRT1* is a key gene involved in sex differentiation (see Discussion). In addition, we identified 33 variants achieving nominal significance ( $P < 1 \times 10^{-6}$ ; S2 Table).

We also performed a GWAS for BMI-AR (see Methods for details). We detected 8 variants achieving nominal significance (S3 Table). S21 Fig shows the Manhattan plot. 234

228

229

230

231

232

223

224

It is made available under a CC-BY-NC-ND 4.0 International license .

SNP ID-Allele	Location	Consequence	Gene	ß-GT	P-GT	Age stratum	Frequency
rs75964957-G	1:239032385	intergenic	-	1.56	4.7E-08	4.5-5.5	0.0077
rs269511-A	5:114688671	intergenic	-	-0.27	2.2E-07	1.5-2.5	0.4646
rs9275582-T	6:32770853	promoter flanking region	HLA-DQB3	0.30	2.5E-07	1.5-2.5	0.2865
rs9275593-A	6:32771630	promoter flanking region	HLA-DQB3	0.30	2.5E-07	1.5-2.5	0.2865
rs9275595-C	6:32772142	promoter flanking region	HLA-DQB3	0.30	2.5E-07	1.5-2.5	0.2865
rs7134291-A	12:14128322	intron	GRIN2B	-0.33	3.0E-07	8.5-9.5	0.1726
rs7896870-T	10:127827684	intron	ADAM12	0.25	4.0E-07	0.5-1.5	0.4906
rs474169-T	20:10081800	intron/NCT	SNAP25-AS1	-0.29	4.8E-07	10.5-11.5	0.3031
rs1495271-C	15:101962723	intron	PCSK6	0.26	6.0E-07	4.5-5.5	0.4060
rs11244839-G	10:127834046	intron	ADAM12	0.25	6.7E-07	0.5-1.5	0.4751
rs13257360-G	8:6117252	upstream	RP11-124B13.1	0.28	7.2E-07	14.5-15.5	0.3335
rs10150218-A	14:73174338	intron	DPF3	0.35	8.3E-07	5.5-6.5	0.4629
rs77468591-T	10:12937704	downstream	CCDC3	-1.84	9.2E-07	4.5-5.5	0.0039

Table 1. Top cross-sectional associations for BMI. Shown is the SNP rs ID with the associated allele, physical location, SO consequence/feature type, gene, effect size of the genotype ( $\beta_{GT}$ ), association *P*-value of the genotype ( $P_{GT}$ ), age stratum and frequency of associated allele. NCT: noncoding transcript variant.

SNP ID-Allele	Location	Consequence	Gene	ß-GTxSex	P-GTxSex	Frequency
rs445398-C	9:954336	intron	DMRT1	5.81	9.8E-09	0.165
rs73952010-G	18:37721370	downstream	RPL17P45	-5.45	7.3E-08	0.007
rs78106261-G	6:110655312	intron	METTL24	-5.13	3.8E-07	0.008
rs73139959-T	3:13815160	intergenic	-	-4.97	8.4E-07	0.003
rs969092-G	12:42234778	intron/NCT	RP11-630C16.2	4.96	8.9E-07	0.238
rs11066997-T	12:114661524	intergenic	-	4.94	9.9E-07	0.358

Table 2. Strongest associations for Age-AR. Shown is the SNP rs ID with the associated allele, physical position in the chromosome, consequence type, gene, effect size of the interaction between genotype and sex ( $\beta_{\text{GTxSex}}$ ) with the corresponding association *P*-value ( $P_{\text{GTxSex}}$ ), and the frequency of the associated allele. NCT: noncoding transcript variant.

# Discussion

In this study, we investigated the genetic architecture of BMI-related traits on admixed children with mainly European and Mapuche Native American ancestry. There are a few studies that have analyzed the genetic architecture of longitudinal growth traits during childhood and adolescence **6 7 8 9 14**, and most of them have been performed in European populations. To our knowledge, Vicuña *et al.*, 2021 **21** is the only study that has quantified the relationship between Native American genetic ancestry and growth traits. The authors showed that the pubertal age where maximum height growth occurs (peak height velocity), is significantly older by 0.73 years in Europeans than in Mapuche adolescents on average. We are the first to quantify how Native American ancestry affects childhood growth traits. One of our most important findings is the observation that increases in Mapuche ancestry associate with significant increases in BMI between 5.5 and 16.5 years old as well as with BMI trajectory between 2 - 10 years old. Moreover, the difference in BMI between Mapuche and Europeans

It is made available under a CC-BY-NC-ND 4.0 International license .



Fig 3. Genome-wide associations with BMI. A: GWAS at age stratum 1.5 - 2.5. B: Association peak of chromosome 6 across age strata. The region shown has physical coordinates 6: 32676320 - 32683820 and was centered at the variant with the strongest association.

increases steadily after 5.5 years old. This makes sense, since it is known that the contribution of heritable genetic variation to BMI increases from childhood (4 years) to young adulthood (19 years)  $\boxed{11}$ . We did not find significant differences at age stratum 4.5 - 5.5 years old, likely due to a lack of statistical power derived from the sample size of our cohort. Another important finding was the significantly lower Age-AR in the Mapuche compared with Europeans on average. A higher adiposity in the Mapuche during childhood could explain in part why this population is particularly susceptible to developing metabolic disorders such as insulin resistance, obesity, cholesterol gallstones and metabolic syndrome during adulthood, since all of these disorders are strongly associated with elevated lipid levels  $\boxed{20}$ . Moreover, Mapuche ancestry among Chileans is distinctly associated with heart diseases, hypertension and diabetes mellitus  $\boxed{19}$ . Further, the prevalence of type II diabetes and obesity among the Mapuche increased significantly following the change from a rural to an urban life style  $\boxed{33}$ , suggesting that

genotype-environment interactions may lead to a higher genetic susceptibility to developing cardiometabolic diseases.

It is unknown whether the effect of Mapuche genetic ancestry on BMI-related traits was originated by random genetic drift in ancient Native American or European populations, by adaptation to selective pressures (see 16) or by a combination of adaptation, drift and/or other evolutionary forces. It is also possible that environmental factors could partially contribute to such differences. However, the children of our cohort belong to the same urban district in Santiago and to the same middle-lower socioeconomic group. Therefore, we do not expect high variation in environmental factors such as nutrition, exposure to pollutants or incidence of pathogenic diseases. However, it is worth noting that in this same cohort, mothers of boys with high Mapuche ancestry ( $\geq 3$  Mapuche last names: 4.8% of total boys) have a lower educational level than mothers of boys with low Mapuche ancestry (0 Mapuche last names: 80% of boys) 35, which in turn could potentially have an effect on their son's BMI.

Our GWAS on BMI detected a significant association for the intergenic variant rs75964957 at age stratum 4.5 - 5.5 years. This was despite the sample size at this stratum (n = 708). The second stronger association was for the rs9275582, rs9275593 and rs9275595 variants, which map a promoter flanking region of the HLA-DQB3 unprocessed pseudogene. Noteworthy, rs9275595 has been previously associated with BMI in adults  $(P = 6 \times 10^{-6}; 0.016 \text{ kg/m}^2 \text{ increase}; 28)$ . These three variants constitute the top of an association peak that was observed at age stratum 1.5 - 2.5years, but not at other age strata. The HLA-DQB3 gene belongs to the human leucocyte antigen (HLA) gene cluster, which harbors hundreds of genes that are fundamental for immune function. HLA genes encode proteins of the major histocompatibility complex, which present antigenic peptides to immune cells, in order to distinguish between "self" and "non-self" agents [29]. It is unknown why these HLA-DQB3 variants have such a distinct effect on BMI at 1.5 - 2.5 years old, but the observation that they map a promoter flanking region suggests that they affect early BMI by regulating the expression of HLA-DQB3 or additional HLA genes in high linkage disequilibrium with HLA-DQB3. Future studies will be needed to validate this finding. It is worth noting that the HLA region shows high variability among ethnic groups  $\overline{37}$  and that the HLA-DR/DQ region is the major determinant of susceptibility to childhood type 1 diabetes 38. The strongest association of the longitudinal GWAS on BMI was for rs35266519, a missense variant of the GSDMB gene. Among other phenotypes, GSDMB variants have been GWAS-associated with type 1 diabetes (P  $< 6 \times 10^{-13}$ ) [39] and subcutaneous adipose tissue ( $P < 3 \times 10^{-8}$ ) [40], two traits that are strongly related with obesity.

We identified a variant of the DMRT1 gene significantly associated with Age-AR. Remarkably, this association was captured using a sample size of n = 696. DMRT1 is a hallmark gene involved in sex differentiation by maintaining the fates of testes or ovaries in adult mammals 30. Indeed, deletion or inactivation of DMRT1 in humans causes XY male-to-female sex reversal 31. Thus, it is possible that genetic variation in this gene could affect endocrine mechanisms of infant growth. Moreover, this effect on infant growth may lead to an altered pubertal timing, since it has been observed that Mapuche boys have a higher incidence of precocious puberty 35.

We could not replicate our findings in an independent cohort, due to the lack of longitudinal paediatric growth cohorts with enough longitudinal BMI measurements and relatively high Native American ancestry. To our knowledge, the Salvador-SCAALA cohort from Brazil 32 and the COIPIS cohort from Mexico 10 are the only paediatric cohorts with genome-wide genotype information and Native American admixture. However, these cohorts present severe limitations for replication. In the

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

It is made available under a CC-BY-NC-ND 4.0 International license .

Salvador-SCAALA cohort, the mean Native American global ancestry is too low (6%), in contrast to high mean African and European ancestries (51% and 46%, respectively; 32). Also, in this cohort only a single BMI measurement per individual was taken in a broad age window (4 – 11 years old). Regarding the COIPIS cohort, while the level of Native American ancestry is appropriate (36%; 10), it has a similar problem, namely, that only a single BMI measurement per individual was taken in children aged 3 – 18 years old.

## Conclusion

Our study contributes to better understand the biological mechanisms underlying BMI and Age-AR differences between Native Americans and Europeans. Also, it highlights the importance of performing serial cross-sectional GWAS for complex longitudinal traits like BMI trajectory, since longitudinal studies alone might not be well-suited to capture effects that occur at short time intervals. Finally, our results have medical relevance, since they could be useful to identify genetic risk factors for infant obesity based on the individual's proportion of Native American genetic ancestry.

# **Consent to Participate**

Informed consents from participants were obtained from parents or guardians. Children agreed to participate when they turned 7 years old. This study was approved by the Scientific Ethics Committees of Instituto de Nutrición y Tecnología en Alimentos (INTA) and Pontificia Universidad Católica de Chile.

# Acknowledgments

This work was supported by the Fondo Nacional de Ciencia y Tecnología (FONDECYT) 335 [1200146 to S.E., L.V. and E.B.; 1190346 to V.M.; 1150416 and 1150486 to JL.S., 336 1190801 to C.M. ]. S.E., T.N. and L.V. were additionally supported by the Instituto 337 Milenio de Investigación Sobre los Fundamentos de los Datos (IMFD). S.E. conceived 338 the project. S.E. and L.V. designed experiments. E.B., L.V., T.N., C.M., D.A. and V.L. 330 analyzed the data. A.P. and V.M. collected phenotype data. S.E., JL. S. and JC. G. 340 raised funds for the genotyping data. L.V., S.E., A.P., E.B. wrote the manuscript. All 341 authors critically reviewed and accepted the final version. 342

# References

- Can, B. & Can, B. Change in overweight from childhood to early adulthood and risk of type 2 diabetes. N Engl J Med 378, 2537 (2018).
- 2. Sovio, U. *et al.* How do changes in body mass index in infancy and childhood associate with cardiometabolic profile in adulthood? findings from the northern finland birth cohort 1966 study. *Int J Obes (Lond)* **38**, 53–9 (2014).
- Couto Alves, A. et al. Gwas on longitudinal growth traits reveals different genetic factors influencing infant, child, and adult bmi. Sci Adv 5, eaaw3095 (2019).
- Llewellyn, C. H., Trzaskowski, M., van Jaarsveld, C. H. M., Plomin, R. & Wardle, J. Satiety mechanisms in genetic risk of obesity. JAMA Pediatr 168, 338–344 (2014).

343

346

347

348

321

329

It is made available under a CC-BY-NC-ND 4.0 International license .

5.	Yengo, L. <i>et al.</i> Meta-analysis of genome-wide association studies for height and body mass index in $\sim$ 700000 individuals of european ancestry. <i>Hum Mol Genet</i> <b>27</b> , 3641–3649 (2018).	354 355 356
6.	Meyre, D. <i>et al.</i> Genome-wide association study for early-onset and morbid adult obesity identifies three new risk loci in european populations. <i>Nat Genet</i> <b>41</b> , 157–9 (2009).	357 358 359

- 7. Scherag, A. *et al.* Two new loci for body-weight regulation identified in a joint analysis of genome-wide association studies for early-onset extreme obesity in french and german study groups. *PLoS Genet* **6**, e1000916 (2010).
- Felix, J. F. *et al.* Genome-wide association analysis identifies three new susceptibility loci for childhood body mass index. *Hum Mol Genet* 25, 389–403 (2016).
- Bradfield, J. P. et al. A trans-ancestral meta-analysis of genome-wide association studies reveals loci associated with childhood obesity. Hum Mol Genet 28, 3327–3338 (2019).
- 10. Costa-Urrutia, P. *et al.* Genome-wide association study of body mass index and body fat in mexican-mestizo children. *Genes (Basel)* **10** (2019). 370
- 11. Dubois, L. *et al.* Genetic and environmental contributions to weight, height, and bmi from birth to 19 years of age: an international study of over 12,000 twin pairs. *PLoS One* **7**, e30153 (2012).
- 12. Jacob, R. *et al.* The role of eating behavior traits in mediating genetic susceptibility to obesity. *Am J Clin Nutr* **108**, 445–452 (2018).
- 13. Sovio, U. *et al.* Association between common variation at the fto locus and changes in body mass index from infancy to late childhood: the complex nature of genetic association through growth and development. *PLoS Genet* **7**, e1001307 (2011).
- 14. Warrington, N. M. *et al.* A genome-wide association study of body mass index across early life and childhood. *Int J Epidemiol* 44, 700–12 (2015).
- 15. Ko, A. *et al.* Amerindian-specific regions under positive selection harbour new lipid variants in latinos. *Nat Commun* **5**, 3983 (2014).
- 16. NEEL, J. V. Diabetes mellitus: a "thrifty" genotype rendered detrimental by "progress"? Am J Hum Genet 14, 353–62 (1962).
- 17. Amorim, C. E. *et al.* Genetic signature of natural selection in first americans. *Proc Natl Acad Sci U S A* **114**, 2195–2199 (2017).
- Eyheramendy, S., Martinez, F. I., Manevy, F., Vial, C. & Repetto, G. M. Genetic structure characterization of chileans reflects historical immigration patterns. *Nat Commun* 6, 6472 (2015).
- 19. Lorenzo Bermejo, J. *et al.* Subtypes of native american ancestry and leading causes of death: Mapuche ancestry-specific associations with gallbladder cancer risk in chile. *PLoS Genet* **13**, e1006756 (2017).
- Miquel, J. Salud y enfermedad en nuestras poblaciones originarias. *Rev Chil Cardiol* 31, 129–133 (2012).

360

361

362

363

364

365

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

391

392

393

394

It is made available under a CC-BY-NC-ND 4.0 International license .

21.	Vicuña, L. <i>et al.</i> Novel loci and mapuche genetic ancestry are associated with pubertal growth traits in chilean boys. <i>Hum Genet</i> (2021).	396 397
22.	Pereira, A. <i>et al.</i> Breast bud detection: a validation study in the chilean growth obesity cohort study. <i>BMC Womens Health</i> <b>14</b> , 96 (2014).	398 399
23.	Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated individuals. <i>Genome Res</i> <b>19</b> , 1655–64 (2009).	400 401
24.	Browning, S. R. & Browning, B. L. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. <i>Am J Hum Genet</i> <b>81</b> , 1084–97 (2007).	402 403 404
25.	McLaren, W. <i>et al.</i> The ensembl variant effect predictor. <i>Genome Biol</i> <b>17</b> , 122 (2016).	405 406
26.	Ward, L. D. & Kellis, M. Haploreg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. <i>Nucleic Acids Res</i> <b>40</b> , D930–4 (2012).	407 408 409
27.	Welter, D. <i>et al.</i> The nhgri gwas catalog, a curated resource of snp-trait associations. <i>Nucleic Acids Res</i> <b>42</b> , D1001–6 (2014).	410 411
28.	Locke, A. E. <i>et al.</i> Genetic studies of body mass index yield new insights for obesity biology. <i>Nature</i> <b>518</b> , 197–206 (2015).	412 413
29.	Lokki, ML. & Paakkanen, R. The complexity and diversity of major histocompatibility complex challenge disease association studies. <i>HLA</i> <b>93</b> , 3–15 (2019).	414 415 416
30.	Huang, S., Ye, L. & Chen, H. Sex determination and maintenance: the role of dmrt1 and foxl2. <i>Asian J Androl</i> <b>19</b> , 619–624 (2017).	417 418
31.	Raymond, C. S. <i>et al.</i> A region of human chromosome 9p required for testis development contains two genes related to known sexual regulators. <i>Hum Mol Genet</i> <b>8</b> , 989–96 (1999).	419 420 421
32.	Scliar, M. O. <i>et al.</i> Admixture/fine-mapping in brazilians reveals a west african associated potential regulatory variant (rs114066381) with a strong female-specific effect on body mass and fat mass indexes. <i>Int J Obes (Lond)</i> <b>45</b> , 1017–1029 (2021).	422 423 424 425
33.	Pérez-Bravo, F. <i>et al.</i> Prevalence of type 2 diabetes and obesity in rural mapuche population from chile. <i>Nutrition</i> <b>17</b> , 236–8 (2001).	426 427
34.	Stryjecki, C., Alyass, A. & Meyre, D. Ethnic and population differences in the genetic predisposition to human obesity. <i>Obes Rev</i> <b>19</b> , 62–80 (2018).	428 429
35.	Fernández, M., Pereira, A., Corvalán, C. & Mericq, V. Precocious pubertal events in children: ethnic disparities. <i>J Endocrinol Invest</i> <b>42</b> , 385–395 (2019).	430 431
36.	Vicuña, L. <i>et al.</i> Novel loci and mapuche genetic ancestry are associated with pubertal growth traits in chilean boys. <i>Hum Genet</i> (2021).	432 433
37.	Degenhardt, F. <i>et al.</i> Construction and benchmarking of a multi-ethnic reference panel for the imputation of hla class i and ii alleles. <i>Hum Mol Genet</i> <b>28</b> , 2078–2092 (2019).	434 435 436

It is made available under a CC-BY-NC-ND 4.0 International license .

38.	Ilonen, J. <i>et al.</i> Genetic susceptibility to type 1 diabetes in childhood - estimation of hla class ii associated disease risk and class ii effect in various phases of islet autoimmunity. <i>Pediatr Diabetes</i> <b>17 Suppl 22</b> , 8–16 (2016).	437 438 439
39.	Barrett, J. C. <i>et al.</i> Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. <i>Nat Genet</i> <b>41</b> , 703–7 (2009).	440 441
40.	Chu, A. Y. <i>et al.</i> Multiethnic genome-wide meta-analysis of ectopic fat depots identifies loci associated with adipocyte development and differentiation. <i>Nat Genet</i> <b>49</b> , 125–130 (2017).	442 443 444
41.	Kain, J. <i>et al.</i> [evolution of the nutritional status of chilean children from preschool to school age: anthropometric results according to the source of the data]. <i>Arch Latinoam Nutr</i> <b>60</b> , 155–9 (2010).	445 446 447
42.	Helgeland, Ø. <i>et al.</i> Genome-wide association study reveals dynamic role of genetic variation in infant and early childhood growth. <i>Nat Commun</i> <b>10</b> , 4448 (2019).	448 449 450
43.	Berkey, C. S. & Reed, R. B. A model for describing normal and abnormal growth in early childhood. <i>Hum Biol</i> <b>59</b> , 973–87 (1987).	451 452
44.	Stergiakouli, E. <i>et al.</i> Genome-wide association study of height-adjusted bmi in childhood identifies functional variant in adcy3. <i>Obesity (Silver Spring)</i> <b>22</b> , 2252–9 (2014).	453 454 455
45.	Cole, T. J. Weight/heightp compared to weight/height2 for assessing adiposity in childhood: influence of age and bone age on p during puberty. Ann Hum Biol 13, 433–51 (1986).	456 457 458
46.	Cole, T. J. & Green, P. J. Smoothing reference centile curves: the lms method and penalized likelihood. <i>Stat Med</i> <b>11</b> , 1305–19 (1992).	459 460
47.	Tobar, H. E. <i>et al.</i> Identification and functional analysis of missense mutations in the lecithin cholesterol acyltransferase gene in a chilean patient with hypoalphalipoproteinemia. <i>Lipids Health Dis</i> <b>18</b> , 132 (2019).	461 462 463
48.	Maples, B. K., Gravel, S., Kenny, E. E. & Bustamante, C. D. Rfmix: a discriminative modeling approach for rapid and robust local-ancestry inference. <i>Am J Hum Genet</i> <b>93</b> , 278–88 (2013).	464 465 466
49.	1000 Genomes Project Consortium <i>et al.</i> A global reference for human genetic variation. <i>Nature</i> <b>526</b> , 68–74 (2015).	467 468
50.	Chang, C. C. <i>et al.</i> Second-generation plink: rising to the challenge of larger and richer datasets. <i>Gigascience</i> 4, 7 (2015).	469 470
51.	Delaneau, O., Zagury, JF. & Marchini, J. Improved whole-chromosome phasing for disease and population genetic studies. <i>Nat Methods</i> <b>10</b> , 5–6 (2013).	471 472