

1 **Exoskeleton-assisted walking improves pulmonary and exercise performances**
2 **more than conventional exercise program in individuals with spinal cord injury:**
3 **a randomised controlled study**

4 **Running head:** Exoskeleton-assisted walking in SCI

5
6 **Authors:**

7 Xiao-Na Xiang,^{1,2,3,†} (0000-0001-8659-1080) Li-Ming Zhang,^{1,†} Hui-Yan Zong,¹ Yi
8 Ou,¹ Xi Yu,^{1,2} Yan Liu,^{1,2} Hong-Ying Jiang,¹ Hong Cheng,⁴ Hong-Chen He,^{1,2,3,*} (0000-
9 0002-3016-022X) and Cheng-Qi He^{1,2,3,*}

10 †These authors contributed equally to this work.

11
12 **Setting:**

13 Department of Rehabilitation Medicine, West China Hospital of Sichuan University.

14
15 **Affiliations:**

16 ¹ Department of Rehabilitation Medicine, West China Hospital, Sichuan University,
17 Chengdu, 610041 Sichuan, PR China

18 ² School of Rehabilitation Sciences, West China School of Medicine, Sichuan
19 University, Chengdu, 610041 Sichuan, PR China

20 ³ Key Laboratory of Rehabilitation Medicine in Sichuan Province, West China
21 Hospital, Sichuan University, Chengdu, 610041 Sichuan, PR China

22
23 **Corresponding Author:**

24 Hong-Chen He, PhD, Rehabilitation Medicine Centre, West China Hospital, Sichuan
25 University, Chengdu, 610041 Sichuan, PR China. E-mail: hxkfhhc@126.com.

26 Cheng-qi He, PhD, Rehabilitation Medicine Centre, West China Hospital, Sichuan
27 University, Chengdu, 610041 Sichuan, PR China. E-mail: hxkfhcq2015@126.com.

28

29 ABSTRACT

30 **Question:** In people with spinal cord injury, does exoskeleton-assisted walking training
31 improve pulmonary ventilation function, motor function and related body structure,
32 walking, and activities of daily life equally comparing those with conventional exercise
33 program? **Design:** Randomised controlled trial with concealed allocation, assessor
34 blinding and intention-to-treat analysis. **Intervention:** Both groups undertook 16
35 sessions of 50-60min training (4 days/week, 4 weeks). Participants in the experimental
36 group received EAW trainings using AIDER system, which assisted standing, walking,
37 and climbing the stairs. The control group received a conventional exercise program
38 which combined aerobic, resistance, flexibility and walking training. **Outcome**
39 **measures:** The primary outcome was the pulmonary function test. The secondary
40 outcomes included: a 6-minute walk test with Borg scale (0 to 10) rating of exertion,
41 10-metre walk test, basic activities of daily living, trunk control test, lower extremity
42 motor score, muscle tone of lower limb, bone mineral density, and distal femoral
43 cartilage at baseline and upon completion of treatment. **Results:** After 4 weeks of
44 trainings, the experimental group improved more on the forced vital capacity (MD 0.53,
45 95%CI 0.01 to 1.06), predicted FVC% (MD 19.59, 95%CI 6.63 to 32.54) and forced
46 expiratory volume in 1s (MD 0.61, 95%CI 0.15 to 1.07); BADL (MD 19.75, 95%CI
47 10.88 to 28.62); and distal femoral cartilage than the control group. Participants
48 completed 6-minute walk test with median 17.3 meters while wearing the exoskeleton.
49 There was no difference in trunk control test, lower extremity motor score, muscle tone,
50 bone mineral density and adverse event. **Conclusions:** In people with lower thoracic
51 neurological level of spinal cord injury, exoskeleton-assisted walking training instead
52 of a conventional excise program has potential benefits to facilitate pulmonary
53 ventilation function, walking, basic activities of daily living and thickness of cartilage.
54 **Trial registration:** ChiCTR2000034623
55 **Keywords:** Spinal cord injury; Exoskeleton; Exercise; Pulmonary function test; Walk

56

57 Introduction

58 Spinal cord injury (SCI) is a worldwide life-disrupting pathological condition with

59 estimated 17,810 injuries occurred in the United States in 2017,^{1,2} and 3.5 per million
60 in the United Kingdom each year.³ Restoring functions (neuromusculoskeletal and
61 respiratory function, etc.), activities of daily living (walk, grasp, lift, etc.), quality of
62 life and life expectancy post-SCI remain to be medical challenges.⁴ Despite great
63 progress in treatments, it is still difficult to achieve the recovery of motor that leads to
64 immobility.⁵ Due to accelerated aging, lifestyle factors, and decreased mobility, even
65 incomplete SCI lesions as low as L4 should be considered as risk factors for
66 cardiopulmonary disability.⁶ A longitudinal decline was reported in forced vital capacity
67 (FVC) and forced expiratory volume in 1 second (FEV₁),⁷ that indicates persons with
68 SCI are more likely to suffer pulmonary capacity deficit.⁸

69 Exercise and Sports Science Australia recommended people with SCI to undertake
70 a combined exercise program which contained moderate aerobic exercise
71 (>30min, >5d/week), moderate strength training and flexibility training (>2d/week) for
72 maintaining respiratory function, quality of life and functional independence.⁹⁻¹²
73 However, the improvements are limited for people with SCI due to lower extremity
74 motor lesions and training types.¹³ Exercise of upper body limits their maximal exercise
75 capacity and puts them at a disadvantage compared with leg exercise.¹⁴ Hence, a new
76 rehabilitation therapy consisting leg exercise is needed for improving the function and
77 independence of people with SCI.

78 Exoskeleton-assisted walking (EAW) refers to a robotic suit worn on the body
79 enabling a person with paralysis to stand and walk,¹⁵ that has been confirmed to help
80 individuals with thoracic and lumbar SCI to walk safely.¹⁶⁻¹⁸ A quantity of studies^{19,20}
81 has manifested the EAW can be used a moderate-intensity level of exercise according
82 to the heart rate, oxygen demand, and rate of perceived exertion. Nevertheless, these
83 clinical trials were designed as observational study, and the changes of pulmonary
84 ventilation function followed by EAW have not been reported yet. It is unclear that
85 whether better pulmonary caused by EAW training performed better exercise capacity
86 in walking. There is lack of evidence that related to the effect of EAW training on
87 respiratory function, quality of life and functional independence for people with SCI.
88 Further, Further, the relation between musculoskeletal function, pulmonary function,

89 and walk ability are not yet completely understood.

90 Therefore, the research question for this randomised trial was:

91 In people with SCI, do EAW training provide equivalent benefits on the pulmonary
92 ventilation and musculoskeletal function, and the functional independence to those
93 obtained in a conventional exercise program?

94

95 **METHODS**

96 **Design**

97 This was a single-blinded, randomized controlled efficacy trial with 2 parallel
98 groups and intention-to-treat analysis. The study protocol has been registered at Chinese
99 Clinical Trial Registry (ChiCTR2000034623) and approved by the medical ethics
100 committee of West China Hospital of Sichuan University. All patients were informed
101 of the procedure, the use of their data and images for research. They understood the
102 purposes and provided written informed consent according to the 1964 Declaration of
103 Helsinki prior to their participation.

104 The individuals were randomly divided into one of two groups in a 1:1 ratio by
105 simple randomization method, using a computer-generated simple random table. The
106 sequences were preserved using closed envelop method by one researcher who did not
107 participant in the trainings and assessments. Clinical data except walking parameters
108 were measured at baseline and post-training by two clinical researchers who were
109 blinded and did not know the group to which each participant belonged. Walking
110 parameters were assessed by two researchers who participated in an EAW program or
111 a conventional program. Clinical data was recorded after averaging.

112 **Participants**

113 From July 2020 to March 2021, we prospectively enrolled all individuals aged
114 between 15 and 65 years with a diagnosed SCI between T4 and L1 at least 1 month.
115 Participants were recruited from inpatients in 3 units of Rehabilitation centre, West
116 China Hospital, Sichuan University. In addition, the eligible individuals met the
117 following inclusion criteria: (1) American spinal injuries association impairment scale
118 (AIS)²¹ classified with A, B or C, (2) the height was between 1.50 meters and 1.85

119 meters and (3) stopped smoking for over 6 months. Exclusion criteria were: (1)
120 spasticity of any lower extremity muscle scored over 2 according to the Modified
121 Ashworth Scale,²² (2) unstable fracture, (3) had EAW training experience before, (4)
122 hypertension, (5) severe osteoporosis (bone mineral density t-score< -3.5), (6) any
123 respiratory or other neurological diseases, (7) overweight (> 100 Kg).

124 **Interventions**

125 All participants were received 16 sessions of exercise training for 50 to 60 minutes
126 per session, 1 session per day, 4 sessions per week for 4 weeks. Exercise intensity was
127 requested to reach 60% to 70% maximal heart rate (HR, $HR_{max}=220 - age$) that is
128 checked with the values of a heart rate sensor (Polar H10, POLAR® China). All
129 participants were scheduled to perform occupational therapy, endurance training
130 cycling using upper limb, and biofeedback therapy once per day. Medications and
131 rehabilitation nursing were ordered based on the medical condition.

132 *Experimental group*

133 The AIDER (AssItive DEvice for paRalyzed patient) powered robotic exoskeleton
134 (generation IV, Buffalo Robot Technology Co. Ltd, Chengdu, China) was used for the
135 EAW training. A mobile phone application (Android) as an additional treatment system
136 switched different training modes (from sit to stand, walk, climb, etc.). Body-weight-
137 supported AIDER system was applied at the beginning of this program to protect users.
138 All subjects were individually fitted to the robotic exoskeleton according to pelvic
139 width, thigh length, and shank length. Training session included sitting, standing,
140 walking, climbing stairs and slope with maximal assistance-walking mode under
141 monitoring of experimented therapists. A progressive EAW program was designed for
142 participants that shown in Table 1.

143 *Control group*

144 Individuals in the control group undertook a conventional exercise program that
145 included strength training using dumbbell between 5 kg and 20 kg, dynamic balance
146 training in sitting or standing position, flexibility training, and walking training with
147 brace. The time proportion of each training depended on the situations of individuals.

148 **Outcome measures**

149 Outcome measures were collected and analysed at the baseline and end of 16-
150 session intervention period.

151 *Primary outcome*

152 Pulmonary ventilation function test was completed with a computerized
153 spirometer (Vyntus™ SPIRO PC Spirometer, Vyair Medical Inc., Mettawa, US) based
154 on the standardized procedures as the American Thoracic Society²³ described. To
155 determine pulmonary ventilation function, participants performed test seating in the
156 wheelchair and were forbidden to disclose their intervention assignment to the assessor.
157 The test was performed with the participants wearing a nose clip. If the participant
158 coughed or made a mistake, the numerical values were not recorded. Three repeated
159 maneuvers were performed, separated by a five-minute rest and the best result was
160 recorded automatically. The test consisted of the assessments of FVC, predicted FVC%,
161 FEV₁, forced expiratory flow (FEF_{25/50/75}), peak expiratory flow (PEF), and maximal
162 voluntary ventilation (MVV).²⁴

163 *Secondary outcomes*

164 Walking parameters were assessed by the 6MWT (6-minute walk test) for distance
165 and 10MWT (10-metre walk test) for speed. Tests were performed in door in accordance
166 with the guidelines of the American Thoracic Society²⁵. The participant's heart rate, and
167 the rate of perceived exertion (RPE) based on the Borg scale²⁶ during the 6MWT were
168 recorded. Participants in EAW group were allowed to wear the exoskeleton, while those
169 in control group using the knee-ankle-foot orthoses if they had one. Modified Barthel
170 index (MBI) were reported to demonstrate the ability of basic activities of daily living
171 (BADL).²⁷

172 Moreover, trunk control test (TCT)²⁸ and lower extremity motor score (LEMS)²⁹
173 were reported to clarify the reasons of ventilation and walking improvement, and the
174 recovery of muscle strength. The thickness of distal femoral cartilage that consisted of
175 medial condyle, intercondylar area, and lateral condyle measured by musculoskeletal
176 ultrasound were compared. The value of Wards triangle bone mineral density (BMD)
177 evaluated by dual-energy X-Ray absorptiometry, and muscle tone of lower limb
178 estimated by modified Ashworth scale (MAS)³⁰ were recorded. The rate of adverse

179 events was used for safety indicator.

180 **Data Analysis**

181 As for the sample size, a pilot study³¹ was performed for determining the effect
182 size (1.03975) of primary outcome (FVC). It was calculated that at least 16 participants
183 were required for each group according to a significance level of 5% and study power
184 of 80%. The final sample size was 40 participants when considered the rate of dropping
185 out as 20%.

186 Analyses were completed according to intention-to-treat principle. Data with
187 normally distributed were recorded as means (SDs), others were described as median
188 (IQR) where necessary. The differences between groups were reported as means (95%
189 CI). Pearson correlation test or Spearman rank correlation test was performed to discuss
190 the relation between the results of 6MWT, TCT and ventilation parameters with
191 statistically significant difference between groups.

192

193 **RESULTS**

194 **Flow of participants**

195 A total of 134 individuals with SCI were screened from July 2020 to March 2021,
196 of which 94 were excluded as per exclusion criteria (n=81) or declining to commit the
197 full participation (n=13). 40 eligible individuals were randomized to either EAW group
198 (n=20) or control group (n=20). Two groups were comparable on the baseline
199 characteristics (Table 2). Figure 1 represents the Consolidated Standards of Reporting
200 Trials (CONSORT) diagram.

201 **Compliance with the study protocol**

202 After pre-intervention assessment, one individual in the EAW group, and one in
203 the control group withdrew the study and did not accept the final assessment because
204 of own desire to be discharged from hospital.

205 **Effect of intervention on pulmonary ventilation function**

206 EAW group showed improvements in pulmonary ventilation function by the end
207 of 16-session intervention period. EAW training was estimated to be more favourable
208 than conventional exercise program for several measures of pulmonary ventilation

209 function, including FVC, FVC%_{predicted}, and FEV₁. The results for FEF₂₅₋₇₅, PEF, and
210 MVV supported that EAW training might be equivalent to conventional exercise
211 program. The estimates and their 95% confidence intervals are presented in Table 3.

212 **Effect of intervention on walking parameters and BADL**

213 Of the 21 participants who completed the final 6MWT and 10MWT, 2 were in the
214 control group, 19 in the EAW group while wearing the exoskeleton in door. The distance
215 and speed in the EAW group were 15.9 (5.0) meters and 0.049 (0.164) m/s, respectively.
216 PRE and heart rate were 3.47 (1.47) and 114.5 (18.0) bpm during the 6MWT.

217 Both groups showed statistical improvement in the BADL after interventions. The
218 improvement was estimated to be 19.75 greater in the EAW group (95% CI 10.88 to
219 28.62) than in the control group. These better results on the MBI were achieved with a
220 mean difference of 28.8 (SD 15.0) within the EAW group and 14.0 (SD 11.5) within the
221 control group by the end of trainings. These results are presented in Table 4.

222 **Effect of intervention on motor function**

223 Both groups showed statistical improvement in the TCT by the end of trainings,
224 while the improvement of the EAW training might be equivalent to conventional
225 exercise program. Additionally, the control group showed improvement in LEMS by
226 the end of trainings. Nonetheless, the results of LEMS and MAS did not have estimates
227 that clearly favoured one intervention over the other. These results are presented in
228 Table 4, with individual-participant data in Table 6 on the eAddenda.

229 **Effect of intervention on bone and cartilage**

230 EAW training was estimated to be more favourable than conventional exercise
231 program for protecting distal femoral cartilage. The effect on BMD was unclear (MD
232 0.07 g/cm³, 95% CI -0.21 to 0.06). The confidence intervals around these effect
233 estimates confirmed the benefit (Table 5).

234 The exercise training in both groups was well tolerated and there were 3 adverse
235 events in the EAW group. Adverse events consisted of fall (n=1) and heel indentation
236 (n=2), that disappeared within 24 hours and caused no injury to participants. Individual
237 participant data for all outcomes are presented in Table 6 on the eAddenda.

238

239 DISCUSSION

240 For individuals with T4 to L1 SCI, EAW training instead of conventional exercise
241 program improved the pulmonary ventilation function, trunk control, and BADL.
242 Moreover, it promoted the walking ability, and protected the cartilage. However, it is
243 estimated that neither were effective for motor function recovery and bone loss
244 prevention. It is essential to consider whether the benefits from EAW training are robust
245 (as reflected in the confidence intervals) to warrant recommending it over conventional
246 exercise program.

247 The estimate of the differences between the groups on the predicted FVC% and
248 FEV₁ (19.6% and 0.61L) were more beneficial than the minimal clinically important
249 difference. Because the minimal clinically important difference for predicted FVC%
250 and FEV₁ are 2-6% by distribution-based method,³² and 0.1L according to previous
251 study by the anchor-based method.³³ Moreover, the estimate of the effects on predicted
252 FVC% from post- to pre- intervention in EAW group and control group and were 14.4%
253 and -0.6%, respectively. And effects on changes of FEV₁ were 0.53L and -0.02L for
254 EAW trainings and conventional exercise program.

255 Both groups were required equal intensity and amounts of time training in this
256 study. Therefore, the effects on pulmonary ventilation function might be considered as
257 a ‘potential’ bonus, that was meaningful for people with SCI. Upper limb exercise are
258 limited for improving a maximal external power output, peak oxygen consumption,
259 heart rate, total peripheral resistance, and responses for cardiac stroke volume.¹⁴
260 Another underlying mechanism might be the potential of EAW to recruit trunk
261 muscles.^{34,35} Alamro et al.³⁶ and Guan et al.³⁷ have demonstrated that overground
262 walking by exoskeleton elicits greater activation of trunk muscles compared to
263 treadmill walking, even after controlling for the use of hand-held assistive devices. The
264 benefits for pulmonary ventilation function are consistent with other research,^{20,38-40}
265 although it reported a higher VO_{2 peak} among the incomplete SCI individuals after EAW
266 training.

267 Robotic exoskeleton was achievable to enhance the walking capacity. The greater
268 distance on the 6MWT was achieved with a higher predicted FVC% (r=0.54) and less

269 exertion ($r=-0.61$). The improvement of walking performance was not caused by the
270 recovery of lower limb motor function. All participants were able to perform walking
271 while wearing the exoskeleton, that was similar to McIntosh et al.⁴¹ and Sale et al.⁴²
272 Longer training duration may result in longer distance. Benson et al.⁴³ found the
273 minimal distance of 6MWT after 10-week trainings was 91 meters which was more
274 than 5 times than our average distance.

275 Additionally, we found that individuals with lower neurologic injury level (T11 to
276 L1) had similar mean improvement (15.6 meters) than others (16.0 meters). The age or
277 neurologic injury level did not perform as an associated factor for walking distance.
278 Participants who younger than 40 years-old completed 15.5 meters during 6WMT, with
279 that of 17.3 meters for older participants. This was inconsistent with previous study⁴⁴
280 that reported walking speed and performances were significantly associated with injury
281 level. The inconsistency caused by product update according to the result of Guanziroli
282 et al.⁴⁵

283 Beyond this, the application of EAW and product update led to the differences
284 between groups on walking and going stair, that two issues of MBI. Hence, participants
285 in the EAW group were accessible to gain the scores on these two aspects with
286 exoskeleton. However, besides EAW and conventional exercise program, occupational
287 therapy is also benefit for the ability of ADL.⁴⁶ Thus, the possible effect of two
288 technologies on BADL is overestimated and needs to be considered.

289 In addition, the effects on bone and cartilage were further explored. There was an
290 increase of BDM in the EAW group, although the estimate was unclear and the values
291 before intervention were different. The effect on EAW on BMD of Wards triangle was
292 consistent with the study of Karelis et al.⁴⁷, which reported an improvement with
293 clinical meaning measured by peripheral quantitative computed tomography on tibia.
294 Different training prescriptions lead to different effects on joint, although the effect on
295 cartilage was merely reported in EAW training. Optional duration (less than 1 hour) is
296 helpful for increasing the thickness of distal femoral cartilage according to Yilmaz et
297 al.,⁴⁸ while higher intensity is harmful for cartilage.⁴⁹ Therefore, the EAW had potential
298 benefit for protecting the cartilage for SCI and the training prescription in this study

299 was suitable.

300 This study was limited to the number of training session, although our feasibility
301 study has proved this training period realized the application of EAW and was most
302 achievable in inpatient rehabilitation in the health care system. Although we tried to
303 avoid the detection bias, it was inevitable during the 6MWT. In this study, the range of
304 age is 40 and the used equation underestimates the HR_{max} in older adults.⁵⁰ Moreover,
305 the validity of equation has not been established in a study sample that included an
306 adequate number of individuals with SCI. This might have resulted in misestimation of
307 the training intensity. Further, gait parameters should be recorded and compared while
308 walking without the exoskeleton.

309 In summary, the study successfully manifested that EAW training has the potential
310 to improve performance in pulmonary ventilation function, trunk control, and BADL
311 among individuals with SCI at least equivalently with conventional exercise program.
312 Additionally, it promoted the walking ability, and protected the cartilage better than
313 conventional exercise program. However, it is estimated that neither were effective for
314 motor function recovery and bone loss prevention.

315

316 **What was already known on this topic:** Pulmonary ventilation and motor function
317 are most frequently affected by spinal cord injury. Recovery of motor function is hard
318 to achieve, while progression in pulmonary ventilation function helps to improve motor
319 performance and activities of daily livings. Conventional exercise programs help to
320 maintain pulmonary ventilation function and activities of daily livings. Exoskeleton-
321 assisted walking is a new technique for gain walking ability among people with spinal
322 cord injury.

323 **What this study adds:** In people with spinal cord injury, exoskeleton-assisted walking
324 training improve performance in pulmonary ventilation function, trunk control, and
325 basic activities of daily livings among individuals with SCI equivalently with
326 conventional exercise program. Other benefits of the exoskeleton-assisted walking
327 were better improvement in walking ability and protection for joint cartilage. However,
328 neither were effective for motor function recovery and bone loss prevention.

329

330 **Ethics approval and consent to participate:** This study received approval from the
331 Ethics Committee of West China Hospital of Sichuan University and registered at the
332 Chinese Clinical Trial Registry with the following identifier: ChiCTR2000034623.
333 Participants gave written informed consent before starting data collection.

334

335 **Acknowledgments:** The authors are grateful for the assistance from Professor Hong
336 Cheng. Additionally, we also thank the participants who were involved in the project.

337

338 **Conflict of interest:** The authors report no conflicts of interest.

339

340 **Source of support:** This research study was supported by the National Key R&D
341 Program of China [grant numbers 2017YFB1302305].

342

343 **Availability of data and materials:** The dataset used in the study is available from the
344 corresponding author on reasonable request.

345

346 **Correspondence:** Hong-Chen He, Rehabilitation Medicine Centre, West China
347 Hospital, Sichuan University, Chengdu, Sichuan, PR China. E-mail: hxfhch@126.com;
348 Cheng-qi He, Rehabilitation Medicine Centre, West China Hospital, Sichuan University,
349 Chengdu, Sichuan, PR China. E-mail: hxfhchq2015@126.com.

350

351

352 **References**

353 1. Jain NB, Ayers GD, Peterson EN, et al. Traumatic spinal cord injury in the United
354 States, 1993-2012. *JAMA*. 2015;313(22):2236-2243.

355 2. Ge L, Arul K, Ikpeze T, Baldwin A, Nickels JL, Mesfin A. Traumatic and
356 Nontraumatic Spinal Cord Injuries. *World Neurosurg*. 2018;111:e142-e148.

- 357 3. Bao FP, Zhang HG, Zhu SM. Anesthetic considerations for patients with acute
358 cervical spinal cord injury. *Neural Regen Res.* 2017;12(3):499-504.
- 359 4. Zhai HW, Gong ZK, Sun J, et al. Ganglioside with nerve growth factor for the
360 recovery of extremity function following spinal cord injury and somatosensory evoked
361 potential. *Eur Rev Med Pharmacol Sci.* 2015;19(12):2282-2286.
- 362 5. Lee BB, Cripps RA, Fitzharris M, Wing PC. The global map for traumatic spinal
363 cord injury epidemiology: update 2011, global incidence rate. *Spinal Cord.*
364 2014;52(2):110-116.
- 365 6. Legg Ditterline BE, Aslan SC, Randall DC, Harkema SJ, Castillo C, Ovechkin AV.
366 Effects of Respiratory Training on Heart Rate Variability and Baroreflex Sensitivity in
367 Individuals With Chronic Spinal Cord Injury. *Arch Phys Med Rehabil.* 2018;99(3):423-
368 432.
- 369 7. Stolzmann KL, Gagnon DR, Brown R, Tun CG, Garshick E. Longitudinal change
370 in FEV1 and FVC in chronic spinal cord injury. *American journal of respiratory and*
371 *critical care medicine.* 2008;177(7):781-786.
- 372 8. Malas FÜ, Köseoğlu F, Kara M, et al. Diaphragm ultrasonography and pulmonary
373 function tests in patients with spinal cord injury. *Spinal cord.* 2019;57(8):679-683.
- 374 9. Tweedy SM, Beckman EM, Geraghty TJ, et al. Exercise and sports science
375 Australia (ESSA) position statement on exercise and spinal cord injury. *J Sci Med Sport.*
376 2017;20(2):108-115.
- 377 10. Gaffurini P BL, Calza S, Calabretto C, Orizio C, Gobbo M. Energy metabolism
378 during activity-promoting video games practice in subjects with spinal cord injury:

- 379 evidences for health promotion. *Eur J Phys Rehabil Med*. 2013;49((1)):23-29.
- 380 11. Neville BT, Murray D, Rosen KB, Bryson CA, Collins JP, Guccione AA. Effects
381 of Performance-Based Training on Gait and Balance in Individuals With Incomplete
382 Spinal Cord Injury. *Arch Phys Med Rehabil*. 2019;100(10):1888-1893.
- 383 12. Liu H, Li J, Du L, et al. Short-term effects of core stability training on the balance
384 and ambulation function of individuals with chronic spinal cord injury: a pilot
385 randomized controlled trial. *Minerva Med*. 2019;110(3):216-223.
- 386 13. Battikha M, Sa L, Porter A, Taylor JA. Relationship between pulmonary function
387 and exercise capacity in individuals with spinal cord injury. *Am J Phys Med Rehabil*.
388 2014;93(5):413-421.
- 389 14. Theisen D. Cardiovascular determinants of exercise capacity in the Paralympic
390 athlete with spinal cord injury. *Exp Physiol*. 2012;97(3):319-324.
- 391 15. Geigle PR, Kallins M. Exoskeleton-Assisted Walking for People With Spinal Cord
392 Injury. *Archives of physical medicine and rehabilitation*. 2017;98(7):1493-1495.
- 393 16. Kozlowski AJ, Bryce TN, Dijkers MP. Time and Effort Required by Persons with
394 Spinal Cord Injury to Learn to Use a Powered Exoskeleton for Assisted Walking. *Top*
395 *Spinal Cord Inj Rehabil*. 2015;21(2):110-121.
- 396 17. Birch N, Graham J, Priestley T, et al. Results of the first interim analysis of the
397 RAPPER II trial in patients with spinal cord injury: ambulation and functional exercise
398 programs in the REX powered walking aid. *J Neuroeng Rehabil*. 2017;14(1):60.
- 399 18. Xiang XN, Ding MF, Zong HY, et al. The safety and feasibility of a new
400 rehabilitation robotic exoskeleton for assisting individuals with lower extremity motor

- 401 complete lesions following spinal cord injury (SCI): an observational study. *Spinal*
402 *Cord*. 2020;58(7):787-794.
- 403 19. Asselin P, Knezevic S, Kornfeld S, et al. Heart rate and oxygen demand of powered
404 exoskeleton-assisted walking in persons with paraplegia. *J Rehabil Res Dev*.
405 2015;52(2):147-158.
- 406 20. Escalona MJ, Brosseau R, Vermette M, et al. Cardiorespiratory demand and rate of
407 perceived exertion during overground walking with a robotic exoskeleton in long-term
408 manual wheelchair users with chronic spinal cord injury: A cross-sectional study. *Ann*
409 *Phys Rehabil Med*. 2018;61(4):215-223.
- 410 21. Roberts TT, Leonard GR, Cepela DJ. Classifications In Brief: American Spinal
411 Injury Association (ASIA) Impairment Scale. *Clin Orthop Relat Res*.
412 2017;475(5):1499-1504.
- 413 22. Meseguer-Henarejos AB, Sanchez-Meca J, Lopez-Pina JA, Carles-Hernandez R.
414 Inter- and intra-rater reliability of the Modified Ashworth Scale: a systematic review
415 and meta-analysis. *Eur J Phys Rehabil Med*. 2018;54(4):576-590.
- 416 23. Graham BL, Steenbruggen I, Miller MR, et al. Standardization of Spirometry 2019
417 Update. An Official American Thoracic Society and European Respiratory Society
418 Technical Statement. *Am J Respir Crit Care Med*. 2019;200(8):e70-e88.
- 419 24. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur*
420 *Respir J*. 2005;26(2):319-338.
- 421 25. ATS statement: guidelines for the six-minute walk test. *American journal of*
422 *respiratory and critical care medicine*. 2002;166(1):111-117.

- 423 26. Holmlund T, Ekblom-Bak E, Franzen E, Hultling C, Wahman K. Intensity of
424 physical activity as a percentage of peak oxygen uptake, heart rate and Borg RPE in
425 motor-complete para- and tetraplegia. *PLoS One*. 2019;14(12):e0222542.
- 426 27. Furlan JC, Noonan V, Singh A, Fehlings MG. Assessment of disability in patients
427 with acute traumatic spinal cord injury: a systematic review of the literature. *Journal of*
428 *neurotrauma*. 2011;28(8):1413-1430.
- 429 28. Quinzaños J, Villa AR, Flores AA, Pérez R. Proposal and validation of a clinical
430 trunk control test in individuals with spinal cord injury. *Spinal cord*. 2014;52(6):449-
431 454.
- 432 29. Wilson JR, Jaja BNR, Kwon BK, et al. Natural History, Predictors of Outcome, and
433 Effects of Treatment in Thoracic Spinal Cord Injury: A Multi-Center Cohort Study from
434 the North American Clinical Trials Network. *J Neurotrauma*. 2018;35(21):2554-2560.
- 435 30. Akpinar P, Atici A, Ozkan FU, et al. Reliability of the Modified Ashworth Scale
436 and Modified Tardieu Scale in patients with spinal cord injuries. *Spinal Cord*.
437 2017;55(10):944-949.
- 438 31. Xiao-na X, Hui-yan Z, Yi O, et al. *Journal of NeuroEngineering and Rehabilitation*.
439 2021.
- 440 32. du Bois RM, Weycker D, Albera C, et al. Forced vital capacity in patients with
441 idiopathic pulmonary fibrosis: test properties and minimal clinically important
442 difference. *Am J Respir Crit Care Med*. 2011;184(12):1382-1389.
- 443 33. Gompelmann D, Kontogianni K, Schuhmann M, Eberhardt R, Heussel CP, Herth
444 FJ. The minimal important difference for target lobe volume reduction after endoscopic

- 445 valve therapy. *Int J Chron Obstruct Pulmon Dis*. 2018;13:465-472.
- 446 34. Triolo RJ, Boggs L, Miller ME, Nemunaitis G, Nagy J, Bailey SN. Implanted
447 electrical stimulation of the trunk for seated postural stability and function after cervical
448 spinal cord injury: a single case study. *Archives of physical medicine and rehabilitation*.
449 2009;90(2):340-347.
- 450 35. Hart N, Laffont I, de la Sota AP, et al. Respiratory effects of combined truncal and
451 abdominal support in patients with spinal cord injury. *Archives of physical medicine*
452 *and rehabilitation*. 2005;86(7):1447-1451.
- 453 36. Alamro RA, Chisholm AE, Williams AMM, Carpenter MG, Lam T. Overground
454 walking with a robotic exoskeleton elicits trunk muscle activity in people with high-
455 thoracic motor-complete spinal cord injury. *J Neuroeng Rehabil*. 2018;15(1):109.
- 456 37. Guan X, Kuai S, Ji L, Wang R, Ji R. Trunk muscle activity patterns and motion
457 patterns of patients with motor complete spinal cord injury at T8 and T10 walking with
458 different un-powered exoskeletons. *J Spinal Cord Med*. 2017;40(4):463-470.
- 459 38. Evans N, Hartigan C, Kandilakis C, Pharo E, Clesson I. Acute Cardiorespiratory
460 and Metabolic Responses During Exoskeleton-Assisted Walking Overground Among
461 Persons with Chronic Spinal Cord Injury. *Top Spinal Cord Inj Rehabil*. 2015;21(2):122-
462 132.
- 463 39. Jang YC, Park HK, Han JY, Choi IS, Song MK. Cardiopulmonary function after
464 robotic exoskeleton-assisted over-ground walking training of a patient with an
465 incomplete spinal cord injury: Case report. *Medicine (Baltimore)*. 2019;98(50):e18286.
- 466 40. Gorman PH, Scott W, VanHiel L, Tansey KE, Sweatman WM, Geigle PR.

- 467 Comparison of peak oxygen consumption response to aquatic and robotic therapy in
468 individuals with chronic motor incomplete spinal cord injury: a randomized controlled
469 trial. *Spinal Cord*. 2019;57(6):471-481.
- 470 41. McIntosh K, Charbonneau R, Bensaada Y, Bhatiya U, Ho C. The Safety and
471 Feasibility of Exoskeletal-Assisted Walking in Acute Rehabilitation After Spinal Cord
472 Injury. *Arch Phys Med Rehabil*. 2020;101(1):113-120.
- 473 42. Sale P, Russo EF, Scarton A, Calabro RS, Masiero S, Filoni S. Training for mobility
474 with exoskeleton robot in spinal cord injury patients: a pilot study. *Eur J Phys Rehabil*
475 *Med*. 2018;54(5):745-751.
- 476 43. Benson I, Hart K, Tussler D, van Middendorp JJ. Lower-limb exoskeletons for
477 individuals with chronic spinal cord injury: findings from a feasibility study. *Clin*
478 *Rehabil*. 2016;30(1):73-84.
- 479 44. Louie DR, Eng JJ, Lam T, Spinal Cord Injury Research Evidence Research T. Gait
480 speed using powered robotic exoskeletons after spinal cord injury: a systematic review
481 and correlational study. *J Neuroeng Rehabil*. 2015;12:82.
- 482 45. Guanziroli E, Cazzaniga M, Colombo L, Basilico S, Legnani G, Molteni F.
483 Assistive powered exoskeleton for complete spinal cord injury: correlations between
484 walking ability and exoskeleton control. *Eur J Phys Rehabil Med*. 2019;55(2):209-216.
- 485 46. Caizhong X, Chunlei S, Beibei L, Zhiqing D, Qinneng D, Tong W. The application
486 of somatosensory evoked potentials in spinal cord injury rehabilitation.
487 *NeuroRehabilitation*. 2014;35(4):835-840.
- 488 47. Karelis AD, Carvalho LP, Castillo MJ, Gagnon DH, Aubertin-Leheudre M. Effect

489 on body composition and bone mineral density of walking with a robotic exoskeleton
490 in adults with chronic spinal cord injury. *J Rehabil Med*. 2017;49(1):84-87.

491 48. Yilmaz B, Demir Y, Ozyoruk E, Kesikburun S, Guzelkucuk U. The effect of knee
492 joint loading and immobilization on the femoral cartilage thickness in paraplegics.
493 *Spinal Cord*. 2016;54(4):283-286.

494 49. Teichtahl AJ, Wang Y, Heritier S, et al. The interaction between physical activity
495 and amount of baseline knee cartilage. *Rheumatology (Oxford)*. 2016;55(7):1277-1284.

496 50. Tanaka H, Monahan KD, Seals DR. Age-predicted maximal heart rate revisited.
497 *Journal of the American College of Cardiology*. 2001;37(1):153-156.

498




Figure 1. Design and flow of participants through the trial.

Table 1
Description of the EAW program in the experimental group

Exercise	Session
From sit to stand and standing training using body-weight-supported AIDER system, controlled by therapist.	1
Standing training and shifting of the body centre of gravity using body-weight-supported AIDER system, controlled by patient.	1
Walking training using body-weight-supported AIDER system, controlled by patient.	1
From sit to stand, standing training and shifting of the body centre of gravity using AIDER system, controlled by patient.	2
Walking training (walk straight) using AIDER system, controlled by patient. Turning around with the help of therapists.	1
Walking training (walk straight and turn around) using AIDER system, controlled by patient.	6
Climb the slope using AIDER system, controlled by therapist.	1
Climb the slope using AIDER system, controlled by patient.	1
Take the stairs using AIDER system, controlled by therapist.	1
Take the stairs using AIDER system, controlled by patient.	1

EAW = exoskeleton-assisted walking, AIDER = Assltive DEvice for paRalyzed patient

Table 2
Baseline characteristics of participants

Characteristic	EAW (n=20)	Con (n=20)
Demographic		
Age (<i>yr</i>), mean (SD)	36.6 (11.7)	37.7 (12.6)
Height (<i>cm</i>), mean (SD)	166.2 (6.4)	170.0 (6.2)
Weight (<i>kg</i>), mean (SD)	56.1 (9.1)	64.0 (10.0)
Gender, No. (%)		
Female	4 (20.0)	3 (15.0)
Male	16 (80.0)	17 (75.0)
Clinical		
NLI, No. (%)		
T4-T10	12 (60.0)	12 (60.0)
T11-L1	8 (40.0)	8 (40.0)
AIS, No. (%)		
A	15 (75.0)	10 (50.0)
B	3 (15.0)	2 (10.0)
C	2 (10.0)	8 (40.0)
DOI (<i>month</i>), median (IQR)	2.0 (4.5)	2.0 (0.5)

EAW = exoskeleton-assisted walking group, Con = control group, NLI = Neurological level of injury, AIS = American spinal injuries association impairment scale, DOI = duration of injury

Table 3

Mean (SD) of groups, mean (SD) difference within groups, and mean (95% CI) difference between groups for the pulmonary ventilation function

Outcome	EAW (n=20)		Con (n=20)		Difference within groups (Post minus pre)		Difference between groups
	Pre	Post	Pre	Post	EAW	Con	EAW minus Con
FVC (L)	3.0 (0.9)	3.4 (0.9)	2.9 (0.6)	2.8 (0.8)	0.40 (0.62)	-0.04 (0.48)	0.53 (0.01 to 1.06)
Predicted FVC%	72.4 (16.8)	86.9 (22.6)	67.9 (14.1)	67.3 (17.5)	14.41 (15.89)	-0.62 (11.25)	19.59 (6.30 to 32.54)
FEV ₁ (L)	2.6 (0.8)	3.1 (0.8)	2.5 (0.6)	2.5 (0.7)	0.53 (0.50)	-0.25 (0.38)	0.61 (0.15 to 1.07)
FEF ₇₅ (L/s)	5.0 (1.6)	6.4 (1.4)	5.5 (1.6)	5.7 (1.7)	1.36 (1.19)	0.21 (1.15)	0.61 (-0.38 to 1.59)
FEF ₅₀ (L/s)	3.6 (1.1)	4.3 (1.4)	3.6 (1.0)	3.5 (1.2)	0.67 (1.14)	-0.09 (1.04)	0.79 (-0.23 to 1.61)
FEF ₂₅ (L/s)	1.5 (0.7)	1.9 (1.5)	1.4 (0.9)	1.5 (1.1)	0.40 (0.79)	0.68 (0.82)	0.40 (-0.28 to 1.07)
PEF (L/s)	5.6 (1.5)	6.7 (1.4)	5.8 (1.7)	5.9 (1.9)	1.04 (1.40)	0.11 (1.59)	0.78 (-0.28 to 1.84)
MVV (L)	99.6 (31.1)	111.3 (28.9)	102.1 (30.0)	97.7 (27.0)	11.65 (13.88)	-4.42 (20.72)	13.59 (-4.33 to 31.51)

EAW = exoskeleton-assisted walking group, Con = control group, FVC = Forced vital capacity, FEV₁ = Forced expiratory volume in 1 second, FEF = Forced expiratory flow, PEF = Peak expiratory flow, MVV = Maximal voluntary ventilation

Table 4

Mean (SD) or median (IQR) of groups, mean (SD) difference within groups, and mean (95% CI) difference between groups for motor function, the distal femoral cartilage and bone mineral density (BMD)

Outcome	EAW (n=20)		Con (n=20)		Difference within groups (Post minus pre)		Difference between groups
	Pre	Post	Pre	Post	EAW	Con	EAW minus Con
TCT	54.4 (23.6) [†]	74.0 (35.8) [†]	40.6 (19.6)	61.0 (26.0)	25.0 (35.0) [†]	19.6 (19.2)	N/A
MAS, n (%)							
Grade 0	18 (90)	18 (90)	18 (90)	18 (90)	N/A	N/A	N/A
Grade 1	1 (5)	1 (5)	2 (10)	1 (5)	N/A	N/A	N/A
Grade 1+	1 (5)	1 (5)	0 (0)	1 (5)	N/A	N/A	N/A
LEMS [†]	1.0 (10.8)	1.5 (7.8)	0 (7.5)	5 (13.8)	0 (0)	0.5 (4.5)	N/A
Left medial condyle (mm)	1.49 (0.51)	1.80 (0.31)	1.59 (0.54)	1.46 (0.52)	0.31 (0.50)	-0.13 (0.35)	0.34 (0.06 to 0.61)
Left intercondylar area (mm)	2.02 (0.25)	2.31 (0.45)	1.94 (0.50)	1.85 (0.44)	0.29 (0.41)	-0.09 (0.45)	0.47 (0.18 to 0.75)
Left lateral condyle (mm)	1.99 (0.38)	2.13 (0.30)	1.91 (0.44)	1.87 (0.43)	0.14 (0.35)	-0.04 (0.28)	0.26 (0.02 to 0.50)
Right medial condyle (mm)	1.52 (0.40)	1.78 (0.28)	1.55 (0.42)	1.52 (0.39)	0.26 (0.34)	-0.03 (0.38)	0.26 (0.04 to 0.47)
Right intercondylar area (mm)	2.04 (0.48)	2.30 (0.39)	1.85 (0.50)	1.94 (0.48)	0.26 (0.34)	-0.09 (0.40)	0.32 (0.03 to 0.60)
Right lateral condyle (mm)	1.97 (0.43)	2.20 (0.28)	2.05 (0.54)	1.90 (0.51)	0.24 (0.45)	-0.15 (0.39)	0.31 (0.04 to 0.60)
BMD (g/cm ³)	0.78 (0.14)	0.81 (0.20)	0.90 (0.19)	0.89 (0.22)	0.04 (0.09)	-0.02 (0.13)	0.07 (-0.21 to 0.06)

EAW = exoskeleton-assisted walking group, Con = control group, TCT = trunk control test, MAS = modified Ashworth scale, LEMS = lower extremity motor score, BMD= bone mineral density, [†]= median (IQR)

Table 5

Mean (SD) of groups, mean (SD) difference within groups, and mean (95% CI) difference between groups for the basic activities of daily living (BADL)

Outcome	EAW (n=20)		Con (n=20)		Difference within groups (Post minus pre)		Difference between groups
	Pre	Post	Pre	Post	EAW	Con	EAW minus Con
MBI	34.8 (14.7)	65.0 (17.5)	29.8 (14.7)	42.5 (15.0)	28.8 (15.0)	14.0 (11.5)	19.8 (10.9 to 28.6)

EAW = exoskeleton-assisted walking group, Con = control group, MBI = modified Barthel index

Table 6

Outcomes of correlation of physical function and fitness

Characteristic	Correlation coefficient	p-value
FVC – distance	0.122	0.620
Predicted FVC% – distance	0.540	0.017
FEV ₁ – distance	0.075	0.761
TCT – distance	0.427	0.068
PRE – distance	-0.605	0.006
FVC – TCT	-0.062	0.801
Predicted FVC% – TCT	0.210	0.387
FEV ₁ – TCT	0.044	0.858
RPE – TCT	-0.394	0.095
FVC – RPE	-0.295	0.220
Predicted FVC% – RPE	-0.596	0.007
FEV ₁ – RPE	-0.290	0.228

FVC = Forced vital capacity, FEV₁ = Forced expiratory volume in 1 second, TCT = trunk control test, RPE = rate of perceived exertion