Safety Monitoring of mRNA Vaccines Administered During the Initial 6 Months of the U.S. COVID-19 Vaccination Program: Reports to Vaccine Adverse Events Reporting System (VAERS) and v-safe

Hannah G. Rosenblum, MD^{1,2}; Julianne M. Gee, MPH¹; Ruiling Liu, PhD¹; Paige L. Marquez, MSPH¹; Bicheng Zhang, MS¹; Penelope Strid, MPH¹, Winston E. Abara, MD¹; Michael M. McNeil, MD, MPH¹; Tanya R. Myers, PhD¹; Anne M. Hause, PhD, MSPH¹; John R. Su, MD, PhD, MPH¹; Bethany Baer, MD³; David Menschik, MD, MPH³; Lauri E. Markowitz, MD¹; Tom T. Shimabukuro, MD, MPH, MBA¹, David K. Shay, MD, MPH¹

¹CDC COVID-19 Response Team, Centers for Disease Control and Prevention, Atlanta, Georgia

²Epidemic Intelligence Service, Centers for Disease Control and Prevention, Atlanta, Georgia

³Food and Drug Administration, Silver Spring, Maryland

Corresponding author: Julianne Gee, eocevent416@cdc.gov

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention (CDC) or the Food and Drug Administration (FDA). Mention of a product or company name is for identification purposes only and does not constitute endorsement by the CDC or the FDA.

Acknowledgements

We wish to acknowledge the following contributors: CDC: Amelia Jazwa, Tara Johnson, Charles Licata, Stacey Martin, Florence Lee, Carla Black; FDA: Jane Baumblatt, Deborah Thompson, Kerry Welsh, Narayan Nair, Kosal Nguon (Commonwealth Informatics); v-safe participants; Oracle v-safe development team.

Funding: No external sources of funding were used. CDC received nonfinancial technical support to develop and maintain the v-safe infrastructure from Oracle.

Abstract

Background: In December 2020, two mRNA-based COVID-19 vaccines were authorized for use in the United States. Vaccine safety was monitored using the Vaccine Adverse Event Reporting System (VAERS), a passive surveillance system, and v-safe, an active surveillance system.

Methods: VAERS and v-safe data during December 14, 2020—June 14, 2021 were analyzed. VAERS reports were categorized as non-serious, serious, or death; reporting rates were calculated. Rates of reported deaths were compared to expected mortality rates by age. Proportions of v-safe participants reporting local and systemic reactions or health impacts the week following doses 1 and 2 were determined.

Findings: During the analytic period, 298,792,852 doses of mRNA vaccines were administered in the United States. VAERS processed 340,522 reports; 92·1% were non-serious; 6·6%, serious, non-death; and 1·3%, death. Over half of 7,914,583 v-safe participants self-reported local and systemic reactogenicity, more frequently after dose 2. Injection-site pain, fatigue, and headache were commonly reported during days 0–7 following vaccination. Reactogenicity was reported most frequently one day after vaccination; most reactions were mild. More reports of being unable to work or do normal activities occurred after dose 2 ($32 \cdot 1\%$) than dose 1 ($11 \cdot 9\%$); <1% of participants reported seeking medical care after vaccination. Rates of deaths reported to VAERS were lower than expected background rates by age group.

Interpretation: Safety data from >298 million doses of mRNA COVID-19 vaccine administered in the first 6 months of the U.S. vaccination program show the majority of reported adverse events were mild and short in duration.

Introduction

In December 2020, two mRNA coronavirus disease 2019 (COVID-19) vaccines (BNT162b2; Pfizer-BioNTech and mRNA-1273; Moderna) were granted Emergency Use Authorization (EUA) by the U.S. Food and Drug Administration (FDA) as 2-dose series and recommended for use by the Advisory Committee on Immunization Practices (ACIP).^{1,2} In clinical trials, both mRNA COVID-19 vaccines showed acceptable safety profiles;^{3,4} the most frequently reported local and systemic symptoms were injection site pain, fatigue, and headache. Reactogenicity was more frequently reported following dose 2, and more common among participants aged <65 years.³⁻⁵

Post-authorization safety monitoring better characterizes the safety profiles of mRNA-based COVID-19 vaccines in larger and more heterogeneous populations.⁶ Phased administration of COVID-19 vaccines in the United States began with healthcare workers and long-term care facility residents and expanded to the general population by Spring 2021; however, implementation plans varied by state.⁷ The major sources of initial U.S. safety data were the Vaccine Adverse Event Reporting System (VAERS), a spontaneous, passive reporting system,⁸ and v-safe,⁹ a new active monitoring system. Reports from these systems have been issued through websites, publications, and presentations to advisory committees.¹⁰⁻¹⁴ We review VAERS and v-safe data during the first 6 months of the U.S. vaccination program, when >298 million doses of mRNA COVID-19 vaccines were administered.

Methods

VAERS

VAERS is a national spontaneous reporting system for detecting potential adverse events (AEs) for authorized or licensed U.S. vaccines⁸ co-administered by the Centers for Disease Control and Prevention (CDC) and FDA. VAERS accepts reports from healthcare providers, manufacturers, and the public. Reports include information about the vaccinated person, type of vaccine administered, and AEs experienced. We included all VAERS reports for mRNA vaccines submitted and processed from

3

December 14, 2020 through June 14, 2021, inclusive of any interval from vaccination to event report.¹⁵ Processed reports were those quality checked and coded using the Medical Dictionary for Regulatory Activities (MedDRA) terminology.⁸ Each VAERS report may be assigned more than one MedDRA Preferred Term (PT); PTs do not necessarily indicate a medically confirmed diagnosis, and include signs and symptoms of illness and results of diagnostic tests.

Based on the Code of Federal Regulations,¹⁶ VAERS reports were classified as 'serious' if any of the following were documented: hospitalization, prolongation of hospitalization, permanent disability, life-threatening illness, congenital anomaly or birth defect, or death. Adverse events of special interest were selected for enhanced COVID-19 vaccine safety monitoring based on biologic plausibility, previous vaccine safety experience, and theoretical concerns related to COVID-19.¹⁷ Death certificates and autopsy reports were requested for death reports. CDC physicians reviewed VAERS reports and available death certificates for each decedent to form an impression about cause of death. Impressions were assigned to one of the following categories: one of the 15 most common *International Classification of Disease, Tenth Revision (ICD-10)* diagnostic categories reported on U.S. death certificates¹⁸; COVID-19-related; other (i.e., impression was not included in a pre-specified categories); or unknown/unclear if a likely cause could not be determined.

V-safe

V-safe is a voluntary smartphone-based system that uses text messaging and secure web-based surveys to actively monitor COVID-19 vaccine safety for common local injection site and systemic reactions.⁹ V-safe participants receive text messages that link to web-based health check-in surveys following vaccination, initially daily (days 0–7), then at longer intervals post vaccination. The system resets to the initial survey frequency after entry of another dose. We analyzed survey reports from days 0–7 for reactogenicity, severity⁹ (mild, moderate, severe), and health impact (i.e., unable to perform normal daily

4

activities, unable to work, and/or received care from a medical professional). Participants who reported receiving medical care were contacted and VAERS reports were completed, if clinically indicated.

Data analyses

We conducted descriptive analyses of available VAERS and v-safe data from December 14, 2020 through June 14, 2021 following dose 1 and dose 2 of BNT162b2 and mRNA-1273 vaccines among individuals aged \geq 16 years. For VAERS, bivariate analyses included sex, age groups, race/ethnicity, serious AEs, and vaccine type/manufacturer administered and for death reports, time from vaccination to reported death (i.e., onset interval) and cause of death. Reporting rates to VAERS were calculated for AEs using the total number of doses of mRNA vaccine administered during the 6-month period. COVID-19 vaccine administration data were provided through CDC's COVID-19 Data Tracker.¹⁹ Reporting rates for deaths within 7 days following vaccination and 42 days following vaccination were calculated per vaccinated persons using persons who received at least one dose as the estimation of persons vaccinated and compared to expected background mortality rates.²⁰

Empirical Bayesian (EB) data mining was used to detect disproportional reporting of post-vaccine outcomes by vaccine among all VAERS serious and non-serious reports received by June 14, 2021.²¹ This statistical method calculates observed to expected MedDRA PT pairings by comparing a specific vaccine-MedDRA PT pair to all vaccine-PT pairs in VAERS, adjusting for age, sex, and year received.²² These ratios are ranked by the lower 5% bound of the EB geometric mean confidence interval (EB05) and a standard alert threshold of EB05 >2 was used. An EB05 >2 represents a high degree of confidence that a vaccine-PT pair was reported at least twice as frequently as expected.

V-safe participants who responded to at least one health check-in survey during day 0–7 after vaccination were included in analyses. Descriptive statistics were calculated for participants' characteristics (sex, age,

race/ethnicity), reaction (type and severity) and health impact by manufacturer, dose number, and number of days following vaccination.

SAS software, version 9.4 (SAS Institute; Cary, NC, USA) was used for analyses. Both VAERS and vsafe conduct surveillance as a public health function and are exempt from institutional review board review. Activities were reviewed by the CDC and conducted in accordance with applicable federal law and CDC policy (See: 45 C.F.R. part 46.102(l)(2), 21 C.F.R. part 56; 42 U.S.C. §241(d); 5 U.S.C. §552a; 44 U.S.C. §3501 et seq.).

Results

During December 14, 2020–June 14, 2021, a total of 298,792,852 doses of mRNA COVID-19 vaccines were administered in the United States: 167,177,332 were BNT162b2 and 131,639,515 were mRNA-1273 (Supplemental Table 1). A greater proportion of vaccines were administered to females (53·2%) compared with males (45·8%). The median age at vaccination was 50 years (inter-quartile range [IQR]: 33–65) for BNT162b2 and 56 years (IQR: 39–68) for mRNA-1273, respectively. Non-Hispanic White persons accounted for 38·4% of vaccine recipients; however, race/ethnicity was unknown for 34·9% of all vaccine recipients.

VAERS

During the analytic period, VAERS received and processed a total of 340,522 reports: 164,669 following BNT162b2 and 175,816 following mRNA-1273 vaccination (Table 1). Of these reports, 92·1% were classified as non-serious, 6·6% were serious, not resulting in a death (non-death), and 1·3% were deaths. Seventy-two percent of reports were among females, and 45·3% of reports were among those aged 18–49 years; median age was 50 years (IQR: 36–64). Fifty percent of those reporting race/ethnicity identified as non-Hispanic White; for 22·1%, race/ethnicity was unknown. The most common MedDRA PTs among

6

non-serious reports were headache (20.4%), fatigue (16.6%), pyrexia (16.3%), chills (15.7%), and pain (15.2%). The most common MedDRA PTs among serious reports were dyspnea (15.4%), death (14.1%), pyrexia (11.0%), fatigue (9.7%), and headache (9.5%). The reporting rate to VAERS was 1,049 non-serious reports per million doses, and 90 serious reports per million doses (Table 2). Among the pre-specified AESIs, reporting rates ranged from 0.1 narcolepsy reports per million doses administered to 32 COVID-19 disease reports per million doses administered.

There were 4,496 reports of death to VAERS (Table 3). After review, 24 reports were excluded because of miscoding of death or duplicate reporting. Of the 4,472 reports of deaths analyzed, 2,087 (46·7%) were reported following BNT162b2 and 2,385 (53·3%) following mRNA-1273. Females accounted for 42·6% of reported deaths; the median age was 76 years (IQR: 66–86). More than 80% of deaths were reported among individuals aged 60 years or older (deaths reported per million doses administered by age group: 60–69 years, 14·4; 70–79 years, 28·5; 80–89 years, 75·4; ≥90 years, 2·1). 18·3% of decedents were identified as long-term care facility residents. Time interval to death following vaccination was available for 4,119 reports (92·1%); median time interval was 10·0 days (range: 0—161 days). The greatest number of death reports occurred on day 1 (10·5%) and day 2 (7·0%) following vaccination (Supplemental Figure 1). Compared to expected background rates of death from all causes per million vaccinated persons²⁰, deaths reported to VAERS following mRNA vaccination were consistently 15-30 times less frequent within 7 days of vaccination, and 50 times less frequent within 42 days of vaccination, by age (Table 4).

Death certificates or autopsy reports were available for clinical review for 808 (18·1%) reports of deaths analyzed. Among these, causes of death were most commonly diseases of the heart (46·5%) and COVID-19 disease (12·6%) (Supplemental Table 2, Supplemental Table 3). Among the 3,664 reports of death without a death certificate or autopsy, causes of death were most commonly unknown/unclear (54·1%), diseases of the heart (17·0%), and COVID-19 disease (8·7%). Causes of death by age among reports with death certificate or autopsy reports available are shown by age (Supplemental Table 4).

No adverse health outcome alerts were identified in EB data mining. However, five mRNA COVID-19 administrative error alerts (e.g. 'product temperature excursion issue') with disproportionality (EB05>2) were identified during the surveillance period.

v-safe

During the analytic period, 7,914,583 mRNA COVID-19 vaccine recipients enrolled in v-safe and completed at least one post-vaccination health survey during days 0–7 (Table 5). The median age of vsafe participants was 50 years (IQR: 36-63), 62.9% were female, and 59.4% identified as non-Hispanic White. A total of 6,775,515 participants completed at least one survey during day 0-7 after dose 1 (3,455,778 following BNT162b2; 3,319,737 following mRNA-1273). Of these participants, 68-6% reported a local injection site reaction and 52.7% reported a systemic reaction. Of the 5,674,420 participants who completed surveys after dose 2, a greater percentage reported an injection site reaction (71.7%) and/or a systemic reaction (70.8%) (Table 6). Local injection site reactions were reported more frequently after mRNA-1273 (dose 1: 73.3%; dose 2: 78.4%) than after BNT162b2 (dose 1: 64.0%; dose 2: 65.3%). A similar pattern was found for systemic reactions after mRNA-1273 (dose 1: 54.3%; dose 2: 75.8%) versus BNT162b2 (dose 1: 51.3%; dose 2: 66.1%). The most frequently reported events after dose 1 of either mRNA vaccine included injection site pain ($66 \cdot 2\%$), fatigue ($33 \cdot 9\%$), and headache (27.0%); these were also more frequent after dose 2: injection site pain (68.6\%), fatigue (55.7%), headache (46.2%). Differences in proportions of reactogenicity by dose number were similar after stratifying by age (<65 vs. ≥65 years) and sex. More reactogenicity was reported among younger participants aged <65 years and by females. (Supplemental Table 5).

Local and systemic reactions by manufacturer, dose, days after vaccination, and severity are shown in Figure 1. The majority of reported symptoms were mild. Participants reported moderate and severe

reactogenicity most commonly on day 1 after dose 2 of either vaccine. The proportion of participants who reported symptoms was greatest on day 1 and then decreased subsequently. The highest proportion of participants reported severe symptoms on day 1 following dose 2 of mRNA-1273 (Supplemental Table 7). On all other days, proportions of participants reporting severe symptoms did not exceed 3.0% for any individual symptom (Supplemental Tables 6 and 7).

Reported health impact was greater following dose 2 of either vaccine (32.1%) compared with dose 1 (11.9%) and after mRNA-1273 of either dose compared with BNT162b2 (Table 6). After dose 2 of BNT162b2, 20.5% were unable to do normal activities, and 12.3% were unable to work. After dose 2 of mRNA-1273, 32.8% were unable to do normal activities, and 20.0% were unable to work. Less than 1.0% reported receiving medical care after receiving either dose of either vaccine. Fewer participants reported an emergency room visit (dose 1: 0.1%; dose 2: 0.2%) or hospitalization (dose 1: 0.03%; dose 2: 0.04%).

When stratified by sex, females reported a health impact more frequently than males, peaking on day 1 after vaccination (Supplemental Figure 2). Following dose 2 of mRNA-1273 vaccine, 41·4% of females reported an inability to perform normal activities in the day 1 survey, and 23·5% an inability to work. Among males receiving dose 2 of mRNA-1273 on the day 1 survey, 25·6% were unable to perform normal activity and 16·9% were unable to work (Supplemental Table 8).

Discussion

In this analysis of the first six months of VAERS and v-safe data, when over 298 million doses of mRNA vaccines were administered, we report reactogenicity following mRNA COVID-19 vaccination, and reported adverse events, including deaths. Reactogenicity was similar to what was reported from clinical trials and from early post-authorization monitoring.^{3-5,10,11} In both VAERS and v-safe, local injection site and systemic reactions were commonly reported. V-safe participants more frequently reported transient

reactions following mRNA-1273 compared with BNT162b2, and more frequently following dose 2 of either vaccine. Females and individuals aged <65 years reported AEs and reactions more frequently. Analysis of deaths reported to VAERS demonstrated lower than expected reported mortality rates compared with background mortality rates.

Safety monitoring of COVID-19 vaccines has been the most comprehensive in recent U.S. history. COVID-19 vaccine safety has been monitored with established systems, including the Vaccine Safety Datalink (VSD)²³ and VAERS, and a new system, v-safe, developed specifically for COVID-19 vaccine safety monitoring. During the study period, all COVID-19 vaccines were administered under EUAs, which require vaccine providers to report serious safety events that follow vaccination to VAERS. Heightened public awareness of the COVID-19 vaccination program, outreach and education to healthcare providers and hospitals about COVID-19 EUA AE reporting requirements, and adherence to EUA reporting requirements by providers and health systems, are likely contributing to the high volume of VAERS reports following mRNA vaccination. Data from U.S. safety monitoring systems have been reviewed regularly by the ACIP COVID-19 Vaccine Safety Technical (VaST) work group²⁴ and at public ACIP meetings.²⁵ Specific safety evaluations have covered serious AEs including anaphylaxis,¹⁴ thrombosis with thrombocytopenia syndrome (TTS),²⁶ myocarditis,²⁷ and Guillain-Barré Syndrome (GBS).²⁸ ACIP has assessed the benefit-risk balance of each of the currently licensed or authorized U.S. COVID-19 vaccines; these evaluations have not prompted any changes in U.S. COVID-19 vaccine recommendations.^{27,28}

Reactogenicity findings from VAERS and v-safe were similar to those from a large study of reactogenicity conducted in the United Kingdom.²⁹ The observed patterns may be explained in part by host characteristics known to influence reactogenicity, including age, sex, and the presence of underlying medical conditions.³⁰ Females have more vigorous antibody responses³¹ to certain vaccines and also tend to report more severe local and systemic reactions to influenza vaccine.³² Females may also be more

likely than males to respond to surveys.^{33,34} Younger persons may be more comfortable with smartphonebased surveys and more likely to respond to surveys, generally.^{35,36}

Using v-safe data, we were able to assess the effects of mRNA vaccination on daily life activities for the first time in adults. Effects on daily activities were most frequently reported on day 1 after vaccination. Reports about the health impact measures used in v-safe, while self-assessed and subjective, correlate with reports about reactogenicity: more health impact was reported by females than males, by participants aged <65 years compared with older participants, after dose 2 compared with dose 1, and by those who received mRNA-1273 versus BNT162b2. Reports of seeking medical care after mRNA vaccine were rare; v-safe did not ask which symptoms prompted the participant to seek medical care. Reactogenicity and its associated health effects, even if transient, may deter some from seeking vaccination. Surveys found that nearly half of unvaccinated adults aged <50 years expressed concern about missing work due to vaccine side effects and that employees provided time off were more likely to get vaccinated;^{37,38} employer policies that accommodate this might increase vaccination coverage.³⁹

Patterns of VAERS reports following mRNA COVID-19 vaccination were similar to reports following receipt of other vaccines routinely administered to adults; most reported events were non-serious.^{40,43} EB data mining findings did not suggest any unexpected vaccine safety problems based on pre-specified statistical thresholds. Serious AEs have been detected following receipt of COVID-19 vaccines during U.S. safety monitoring and reviewed in detail in other reports.²⁶⁻²⁸ Reports of anaphylaxis and myocarditis following mRNA vaccination prompted clinical guidance and management recommendations.^{27, 44-47} TTS²⁶ and GBS²⁸ have been associated with Janssen COVID-19 vaccination (Ad26.COV2.S) but not with mRNA vaccination to date.

In our review and analysis of death reports to VAERS following mRNA vaccination, we found no unusual patterns in cause of death among the large number of death reports received. Reporting rates of

death to VAERS among vaccinated persons in 7- and 42-day periods following vaccination were lower than expected background mortality rates. Under the COVID-19 vaccine EUA regulations, healthcare providers are required to report deaths and life-threatening potential adverse events that follow COVID-19 vaccinations to VAERS regardless of their potential association with vaccination. These requirements make comparing the number of reported deaths to VAERS for COVID-19 vaccines to reported deaths following other adult vaccines⁴⁸ difficult. Older individuals and those in long-term care facilities with the highest baseline mortality risk were initially prioritized for COVID-19 vaccination while most U.S. immunization programs target pediatric and adolescent populations.⁴⁹ The concentrated reporting of deaths on the first few days following vaccination follows patterns similar to those observed for other adult vaccinations.⁴⁸ This pattern may represent reporting bias, as the likelihood to report a serious AE may increase when it occurs in close temporal proximity to vaccination.

There are clear limitations in any review of preliminary mortality data. A comparison to national mortality data,^{20,50} suggests that certain causes of death such as accidents, suicides, or cancer, are less likely to be reported to VAERS. The predominance of heart disease as a cause of death reports to VAERS warrants continued monitoring and assessment, but may be driven by non-specific causes, like cardiac arrest, that might be chosen as a terminal event if no immediate explanation is available. Death certificate or autopsy reports were available only for a small proportion of deaths reported to VAERS. Finally, VAERS is designed as an early warning system to detect potential safety signals⁸ and VAERS data alone generally cannot establish causal relationships between vaccination and AEs. Other studies and surveillance systems, including the VSD, are needed to better understand patterns of mortality following mRNA COVID-19 vaccination.^{23, 51}

This study has several strengths, including the large population under surveillance and comprehensive capture of national data from two complementary surveillance systems. Because the U.S. government purchased all COVID-19 doses and collected administration data, we were able to calculate VAERS

reporting rates using the number of mRNA vaccine doses administered.⁵² In contrast, past VAERS analyses for other vaccines could only use doses distributed, which are always greater than doses administered as denominators, underestimating vaccine adverse event reporting rates. Data concerning how reactogenicity following mRNA vaccination affects daily activities and work during the week following vaccination from v-safe provide information has not been previously described. Limitations of this analysis include a limitation shared by all analyses using VAERS data: it is a passive reporting system subject to underreporting and variable or incomplete reporting.⁸ While VAERS death reports were individually reviewed by CDC physicians and follow-up is ongoing to obtain additional and missing records, other serious AE reports were not individually reviewed. Additionally, VAERS reports require interpretation to determine if AE reports meet clinical case definitions.⁵³ EB data mining has multiple limitations,^{21,22} including that the absence of a disproportionality alert for an event does not rule out a possible corresponding adverse event. A new concern with disproportionality scores, which are adjusted by year to control for time-dependent confounders, is that during the study period most VAERS reports were for COVID-19 vaccinations. As all VAERS reports are used for vaccine-event comparisons in EB data mining, potential associations with mRNA COVID-19 vaccines plausibly could be missed. A limitation of v-safe is the need for smartphone access. Because a subset of all vaccine recipients chose or were able to participate in v-safe, the results likely are not generalizable to the entire vaccinated U.S population. In addition, participants in v-safe may be lost to follow up as there is no requirement for continuous enrollment. Finally, this report only included v-safe responses received during the first week post-vaccination.

During the first six months of the U.S. COVID-19 vaccination program, more than 50% of the eligible population received at least one vaccine dose.¹⁹ VAERS and v-safe data from this period demonstrate a post-authorization safety profile for mRNA COVID-19 vaccines that is generally consistent with pre-authorization trials^{3,4} and early post-authorization surveillance reports.^{10,11} Serious AEs have been identified following mRNA vaccinations; however, based on the most current information, these events

are rare. Vaccines are the most effective tool to prevent serious COVID-19 disease outcomes⁵⁴ and the benefits of immunization in preventing serious morbidity and mortality clearly favor vaccination.²⁶⁻²⁸ VAERS and v-safe, two complementary surveillance systems, will continue to provide data needed to inform immunization policy makers, medical and immunization providers, and the public about the safety of COVID-19 vaccination.

References

1. Oliver SE, Gargano JW, Marin M, et al. The Advisory Committee on Immunization Practices' Interim Recommendation for Use of Pfizer-BioNTech COVID-19 Vaccine - United States, December 2020. *MMWR Morbidity and mortality weekly report 2020*; **69**(50): 1922-4.

2. Oliver SE, Gargano JW, Marin M, et al. The Advisory Committee on Immunization Practices' Interim Recommendation for Use of Moderna COVID-19 Vaccine - United States, December 2020. *MMWR Morbidity and mortality weekly report* 2021; **69**(5152): 1653-6.

3. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *The New England Journal of Medicine* 2020; **383**(27): 2603-15.

4. Baden LR, El Sahly HM, Essink B, et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *The New England Journal of Medicine* 2021; **384**(5): 403-16.

5. Anderson EJ, Rouphael NG, Widge AT, et al. Safety and Immunogenicity of SARS-CoV-2 mRNA-1273 Vaccine in Older Adults. *New England Journal of Medicine* 2020; **383**(25): 2427-38.

 Chen RT, Davis RL, Rhodes PH. Special methodological issues in pharmacoepidemiology studies of vaccine safety. In: Strom BL, editor. Pharmacoepidemiology. 4th John Wiley & Sons; Sussex: 2005

7. Dooling K, McClung N, Chamberland M, et al. The Advisory Committee on Immunization Practices' Interim Recommendation for Allocating Initial Supplies of COVID-19 Vaccine - United States, 2020. *MMWR Morbidity and mortality weekly report* 2020; **69**(49): 1857-9.

8. Shimabukuro TT, Nguyen M, Martin D, DeStefano F. Safety monitoring in the Vaccine Adverse Event Reporting System (VAERS). *Vaccine* 2015; **33**(36): 4398-405.

9. <u>https://www.cdc.gov/coronavirus/2019-ncov/vaccines/safety/vsafe.html</u>

10. Gee J, Marquez P, Su J, et al. First Month of COVID-19 Vaccine Safety Monitoring - United States, December 14, 2020-January 13, 2021. *MMWR Morbidity and mortality weekly report* 2021; **70**(8): 283-8.

11. Chapin-Bardales J, Gee J, Myers T. Reactogenicity Following Receipt of mRNA-Based COVID-19 Vaccines. *JAMA* 2021; **325**(21): 2201-2.

12. Shimabukuro TT, Cole M, Su JR. Reports of Anaphylaxis After Receipt of mRNA COVID-19 Vaccines in the US—December 14, 2020-January 18, 2021. *JAMA* 2021; **325**(11): 1101-2.

13. Shay DK, Gee J, Su JR, et al. Safety Monitoring of the Janssen (Johnson & Johnson) COVID-19 Vaccine - United States, March-April 2021. *MMWR Morbidity and mortality weekly report* 2021; **70**(18): 680-4.

14. Allergic Reactions Including Anaphylaxis After Receipt of the First Dose of Pfizer-BioNTech COVID-19 Vaccine - United States, December 14-23, 2020. *MMWR Morbidity and mortality weekly report* 2021; **70**(2): 46-51.

15. <u>https://www.cdc.gov/vaccinesafety/ensuringsafety/monitoring/vaers/index.html</u>

16. Code of Federal Regulations Title 21

https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/cfrsearch.cfm?fr

17. <u>https://www.cdc.gov/vaccinesafety/pdf/VAERS-v2-SOP.pdf</u>

18. https://www.cdc.gov/nchs/nvss/leading-causes-of-death.htm

19. https://covid.cdc.gov/covid-data-tracker/#datatracker-home

20. Abara WE, Gee J, Mu Y, et al. Expected Rates of Select Adverse Events following Immunization for COVID-19 Vaccine Safety Monitoring. *medRxiv* 2021: 2021.08.31.21262919.

21. Szarfman A, Machado SG, O'Neill RT. Use of screening algorithms and computer systems to efficiently signal higher-than-expected combinations of drugs and events in the US FDA's spontaneous reports database. Drug Saf 2002; **25**(6): 381-92.

22. Martin, D, Menschik D, Bryant-Genevier M, Ball R. Data mining for prospective early detection of safety signals in the Vaccine Adverse Event Reporting System (VAERS): a case study of febrile seizures after a 2010-2011 seasonal influenza virus vaccine. *Drug Saf.* 2013; Jul:**36**(7):547-56.

23. <u>https://www.cdc.gov/vaccinesafety/ensuringsafety/monitoring/vsd/index.html</u>

24. <u>https://www.cdc.gov/vaccines/acip/work-groups-vast/index.html</u>

25. https://www.cdc.gov/vaccines/acip/meetings/index.html

26. MacNeil JR, Su JR, Broder KR, et al. Updated Recommendations from the Advisory Committee on Immunization Practices for Use of the Janssen (Johnson & Johnson) COVID-19 Vaccine After Reports of Thrombosis with Thrombocytopenia Syndrome Among Vaccine Recipients - United States, April 2021. *MMWR Morbidity and mortality weekly report* 2021; **70**(17): 651-6.

27. Gargano JW, Wallace M, Hadler SC, et al. Use of mRNA COVID-19 Vaccine After Reports of Myocarditis Among Vaccine Recipients: Update from the Advisory Committee on Immunization Practices - United States, June 2021. *MMWR Morbidity and mortality weekly report* 2021; **70**(27): 977-82.

28. Rosenblum HG, Hadler SC, Moulia D et al. Use of COVID-19 Vaccines After Reports of Adverse Events Among Adult Recipients of Janssen (Johnson & Johnson) and mRNA COVID-19 Vaccines (Pfizer-BioNTech and Moderna): Update from the Advisory Committee on Immunization Practices – United States, July 2021. *MMWR Morbidity and mortality weekly report* 2021: **70**(32); 1094-1099.

29. Menni C, Klaser K, May A, et al. Vaccine side-effects and SARS-CoV-2 infection after vaccination in users of the COVID Symptom Study app in the UK: a prospective observational study. *Lancet Infect Dis.* 2021; **21**(7):939-949

30. Hervé C, Laupèze B, Del Giudice G, Didierlaurent AM, Tavares Da Silva F. The how's and what's of vaccine reactogenicity. *NPJ Vaccines* 2019;**4**: 39.

31. Zimmermann P, Curtis N. Factors That Influence the Immune Response to Vaccination. *Clin Microbiol Rev* 2019; **32**(2).

32. Klein SL, Jedlicka A, Pekosz A. The Xs and Y of immune responses to viral vaccines. *Lancet Infect Dis* 2010; **10**(5): 338-49.

33. Pathiravasan CH, Zhang Y, Trinquart L, et al. Adherence of Mobile App-Based Surveys and Comparison With Traditional Surveys: eCohort Study. *J Med Internet Res* 2021; **23**(1): e24773.

34. Guo X, Vittinghoff E, Olgin JE, Marcus GM, Pletcher MJ. Volunteer Participation in the Health eHeart Study: A Comparison with the US Population. *Sci Rep* 2017; **7**(1): 1956.

35. Millar MM, Elena JW, Gallicchio L, et al. The feasibility of web surveys for obtaining patientreported outcomes from cancer survivors: a randomized experiment comparing survey modes and brochure enclosures. *BMC Med Res Methodol* 2019; **19**(1): 208.

36. Messer BL, Dillman DA. Surveying the General Public over the Internet Using Address-Based Sampling and Mail Contact Procedures. *Public Opinion Quarterly* 2011; **75**(3): 429-57.

37. <u>https://www.kff.org/coronavirus-covid-19/poll-finding/kff-covid-19-vaccine-monitor-april-2021/</u>

38. <u>https://www.kff.org/coronavirus-covid-19/poll-finding/kff-covid-19-vaccine-monitor-june-2021/</u>

39. <u>https://www.whitehouse.gov/briefing-room/statements-releases/2021/04/21/fact-sheet-president-biden-to-call-on-all-employers-to-provide-paid-time-off-for-employees-to-get-vaccinated-after-meeting-goal-of-200-million-shots-in-the-first-100-days/</u>

40. Miller ER, Lewis P, Shimabukuro TT, et al. Post-licensure safety surveillance of zoster vaccine live (Zostavax®) in the United States, Vaccine Adverse Event Reporting System (VAERS), 2006-2015. *Human vaccines & immunotherapeutics* 2018; **14**(8): 1963-9

41. Moro PL, Haber P, McNeil MM. Challenges in evaluating post-licensure vaccine safety: observations from the Centers for Disease Control and Prevention. *Expert Review of Vaccines* 2019; **18**(10): 1091-101

42. Haber P, Moro PL, Ng C, et al. Safety review of tetanus toxoid, reduced diphtheria toxoid, acellular pertussis vaccines (Tdap) in adults aged ≥65 years, Vaccine Adverse Event reporting System (VAERS), United States, September 2010-December 2018. *Vaccine* 2020; **38**(6): 1476-80

43. Woo EJ, Moro PL. Postmarketing safety surveillance of quadrivalent recombinant influenza vaccine: Reports to the vaccine adverse event reporting system. *Vaccine* 2021; **39**(13): 1812-7.

44. <u>https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-</u> anaphylaxis.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fvaccines%2Fcovid-19%2Finfo-by-product%2Fpfizer%2Fanaphylaxis-management.html

45. Abu Mouch S, Roguin A, Hellou E, et al. Myocarditis following COVID-19 mRNA vaccination. Vaccine 2021.

46 Marshall M, Ferguson ID, Lewis P, et al. Symptomatic Acute Myocarditis in Seven Adolescents Following Pfizer-BioNTech COVID-19 Vaccination. *Pediatrics* 2021 Jun 4;e2021052478. *Online ahead* of print

47. <u>https://www.cdc.gov/vaccines/covid-19/clinical-considerations/myocarditis.html</u>

48. Moro PL, Arana J, Cano M, Lewis P, Shimabukuro TT. Deaths Reported to the Vaccine Adverse Event Reporting System, United States, 1997-2013. Clin Infect Dis 2015; 61(6): 980-7.

49. <u>https://wonder.cdc.gov/vaers.html</u>

50. Murphy SL, Xu J, Kochanek KD, Arias E, Tejada-Vera B. Deaths: Final Data for 2018. Natl Vital Stat Rep 2021; **69**(13): 1-83.

51. <u>https://www.cdc.gov/vaccinesafety/pdf/SCK_COVID_Vaccine_Mortality-508.pdf</u>

- 52. https://www.cdc.gov/coronavirus/2019-ncov/vaccines/distributing/first-doses.html
- 53. <u>https://vaers.hhs.gov/data/dataguide.html</u>
- 54. <u>https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/fully-vaccinated-people.html</u>

Table 1: Characteristics of reports received and processed by Vaccine Adverse Events Reporting System (VAERS) for mRNA COVID-19 vaccines—December 14, 2020–June 14, 2021

	Both mRNA vaccines [†] $(n-240.522)$	BNT162b2 vaccine	mRNA-1273 vaccine
Donorta	(11=340,322)	(II=104,009)	(II=1/3,810)
Non serious adverse event reports	212 400 (02 1)	150 486 (01 4)	162 077 (02 7)
Serious reports including death	27 023 (7.9)	14 183 (8.6)	102,977(92,77) 12 830(7,3)
Serious non-death adverse event reports	22,527 (6.6)	12,078,(7,3)	10,448 (5.9)
Death	4 496 (1.3)	2,078(7.5)	2 301 (1.4)
Sev	4,490 (1.3)	2,105 (1.3)	2,391 (1.4)
Female	246 085 (72 3)	116 597 (70.9)	120 475 (72 6)
Mala	240,085 (72.3)	45 157 (27 4)	129,475(73.0)
Unknown	6126(1.8)	(27.4)	(24.3)
	0,120 (1.8)	2,923 (1.8)	5,201 (1.8)
16 17	6 874 (2 0)	2 292 (2 0)	2 501 (2 0)
10-17	(0,874)(2.0)	76385(2.0)	3,391(2.0) 77 773 (44.2)
50 64	84 949 (24.9)	40.367 (24.5)	11,115 (11 ,2) 14 572 (25.4)
50-04 65 7 <i>1</i>	49 755 (14.6)	20.048(12.2)	20702(16.0)
75 94	(1+0)	20,040(12.2) 0.021(5.5)	12,702(10.9)
>85	7595(2.2)	3,021(3.3) 3,564(2.2)	12,32(7.1) 4,027(2.3)
≥05 Unknown	15 760 (4.6)	12,001,(7,2)	2,750 (2,1)
Pace/Ethnicity	15,700 (4.0)	12,001 (7.5)	5,759 (2.1)
Hispanic/Latino	23 480 (6.9)	11 217 (6.8)	12 260 (7.0)
Non Hispanic	23,480 (0.9)	11,217 (0.8)	12,200 (7.0)
White	160 877 (40.0)	72 208 (44 6)	06 460 (54 0)
Plack	109,877(49.9) 10,446(2,1)	5 104 (2 1)	5 342 (2 0)
Asian	10,440 (3.1)	5,104(5.1) 5,028(2,1)	5,542 (3.0)
Asian American Indian or Alaska Nativa	10,172(5.0) 1,414(0,4)	5,058 (5.1)	3,131(2.9) 700 (0,5)
Native Heweijen or Other Pacific Islander	1,414(0.4)	200(0.1)	799(0.3)
Multiple means	441(0.1)	209(0.1)	232(0.1)
Other mass	3,342(1.0)	1,378(1.0)	$1,904(1\cdot1)$ 876(0,5)
Uniter races	2 502 (0.8)	1 422 (0, 0)	870 (0.5)
Unknown race	2,393 (0.8)	1,422 (0.9)	1,171 (0.7)
White	29 797 (9 5)	15 407 (0.4)	12 280 (7.6)
Plast	20,707 (0.3)	15,497(9.4)	15,289 (7.0)
Asian	$4,109(1\cdot 2)$	$2,324(1\cdot 3)$	1,002(1.0) 1,020(0.6)
Asian American Indian or Alaska Nativa	2,455 (0.7)	1,590(0.9)	1,059(0.0)
Native Heneiian or Other Pacific Islander	105 (0.02)	56 (0.02)	40 (0.02)
Multiple reces	500 (0.2)	30 (0.03)	280 (0.2)
Other mass	4700(1.4)	301(0.2) 2.838(1.7)	289(0.2) 1 870(1.1)
Unknown race and ethnicity	4,709 (1.4)	42,330 (1.7)	22 000 (18 8)
Signs or symptoms most frequently reported non	75,554 (22.1)	42,320 (23.7)	52,999 (18.8)
serious*			
Headache	64,064 (20.4)	30,907 (20.5)	33,154 (20.3)
Fatigue	52,048 (16.6)	24,805 (16.5)	27,241 (16.7)
Pyrexia	51,023 (16.3)	22,185 (14.7)	28,837 (17.7)
Chills	49,234 (15.7)	21,638 (14.4)	27,595 (16.9)
Pain	47,745 (15.2)	21,506 (14.3)	26,238 (16.1)
Nausea	37,333 (11.9)	18,066 (12.0)	19,267 (11.8)
Dizziness	37,257 (11.9)	20,307 (13.5)	16,950 (10.4)
Pain in extremity	31,753 (10.1)	14,098 (9.4)	17,653 (10.8)
Injection site pain	28,949 (9.2)	10,462 (7.0)	18,487 (11.3)
Injection site erythema	22,351 (7.1)	2,991 (2.0)	19,360 (11.9)
Signs or symptoms most frequently reported, serious*			
Dyspnea	4,175 (15.4)	2,210 (15.6)	1,965 (15.3)
Death [‡]	3,802 (14.1)	1,753 (12.4)	2,039 (15.9)
Pyrexia	2,986 (11.0)	1,469 (10.4)	1,517 (11.8)
Fatigue	2,608 (9.7)	1,395 (9.8)	1,213 (9.4)
Headache	2,567 (9.5)	1,360 (9.6)	1,207 (9.4)
Chest pain	2,300 (8.5)	1,310 (9.2)	990 (7.7)
Nausea	2,228 (8.2)	1,160 (8.2)	1,068 (8.3)
Pain	2,222 (8.2)	1,195 (8.4)	1,027 (8.0)
Asthenia	2,194 (8.1)	1,084 (7.6)	1,110 (8.6)
Dizziness	2,069 (7.7)	1,111 (7.8)	958 (7.5)

Data are n (%).

*Symptoms refers to MedDRA preferred terms (PTs) and are ordered by most frequently reported for both vaccines. MedDRA PTs are not mutually exclusive.

[†]Total includes reports without a vaccine manufacturer listed.

[‡]Not all reports of death were coded with the MedDRA PT of 'death'

Table 2: Frequency and reporting rates of adverse events of special interest reported to Vaccine Adverse Event Reporting System (VAERS) by recipients of mRNA COVID-19 vaccine—December 14, 2020–June 14, 2021

	Both mRNA	vaccines	BNT162b2 vaco	cine	mRNA-1273 vaccine		
	n	Reports per million doses administered [*]	n	Reports per million doses administered [†]	n	Reports per million doses administered [‡]	
Non-serious adverse event reports	313,499	1,049.2	150,486	900.2	162,977	1,238.1	
Serious reports, including death	27,023	90.4	14,183	84.8	12,839	97.5	
Serious, non-death adverse event reports	22,527	75.4	12,078	72.2	10,448	79.4	
Reports $^{\$}$ of adverse events of special interest **							
COVID-19	9,344	31.3	7,184	43.0	2,160	16.4	
Coagulopathy ^{††}	4,320	14.5	2,343	14.0	1,977	15.0	
Seizure	2,733	9.1	1,478	8.8	1,255	9.5	
Stroke ^{‡‡}	1,937	6.5	981	5.9	955	7.3	
Bells' Palsy	1,918	6.4	1,057	6.3	861	6.5	
Anaphylaxis	1,639	5.5	972	5.8	667	5.1	
Myopericarditis	1,307	4.4	813	4.9	494	3.8	
Acute Myocardial Infarction	1,118	3.7	610	3.6	508	3.9	
Appendicitis	383	1.3	258	1.5	125	1.0	
Guillain-Barré Syndrome	293	$1 \cdot 0$	154	0.9	139	1.1	
Multisystem Inflammatory Syndrome in Adults	119	0.4	60	0.4	59	0.4	
Transverse Myelitis	98	0.3	55	0.3	43	0.3	
Narcolepsy	21	0.1	12	0.1	9	0.1	

*298,792,852 doses of mRNA vaccine were administered in the study period.

[†]167,177,332 doses of BNT162b2 vaccine were administered in the study period.

[‡]131,639,515 doses of mRNA-1273 vaccine were administered in the study period.

[§]These represent reports, not confirmed by case definition.

**Reported death is an adverse event of special interest but counts appear in following tables. Events are not mutually exclusive. ††Coagulopathy is an aggregate term capturing three specific adverse events: 1) thrombocytopenia, 2) deep venous

thrombosis/pulmonary embolism, and 3) disseminated intravascular coagulopathy.

^{‡‡}No vaccine manufacturer was provided for one report of stroke.

Table 3: Frequency and reporting rates of death reported to Vaccine Adverse Event Reporting System (VAERS) by recipients of mRNA COVID-19 vaccine, by sex and age group—December 14, 2020–June 14, 2021

	Both mRNA vaccines		BNT162b2 vac	cine	mRNA-1273 vaccine			
	(n=4,472*)		(n=2,087)		(n=2,385)			
	n (%)	Reports per million doses administered [†]	n (%)	Reports per million doses administered [†]	n (%)	Reports per million doses administered [†]		
Sex								
Female	1,906 (42.6)	12.2	918 (44.0)	10.6	988 (41.4)	14.2		
Male	2,486 (55.6)	18.5	1,117 (53.5)	15.1	1,369 (57.4)	22.6		
Unknown [‡]	80 (1.8)		52 (2.5)		28 (1.2)	••-		
Age (years)								
16–17	6 (0.1)	1.1	6 (0.3)	1.1				
18–29	51 (1.1)	1.3	27 (1.3)	1.1	24 (1.0)	1.6		
30–39	94 (2.1)	2.4	50 (2.4)	2.2	44 (1.8)	2.8		
40–49	151 (3.4)	3.8	74 (3.5)	3.2	77 (3.2)	4.6		
50–59	328 (7.3)	6.9	132 (6.3)	5.0	196 (8-2)	9.3		
60–69	765 (17.1)	14.43	354 (17.0)	13.0	411 (17·2)	16.0		
70–79	1,118 (25.0)	28.5	497 (23.8)	25.9	621 (26.0)	31.0		
80–89	1,128 (25.2)	75.4	529 (25.3)	72.1	599 (25.1)	78.6		
≥90	637 (14·2)	207.7	302 (14.5)	188.1	335 (14.0)	229.3		
Unknown [‡]	194 (4.3)		116 (5.6)		78 (3.3)			

*Of 4,496 deaths, 24 were excluded as they could not be confirmed or were duplicate reports upon review.

[†]Doses of vaccine administered in the study period were used for denominators in each age group; does not include doses administered in Texas as data for Texas were reported to CDC in aggregate

[‡]Reporting rates not shown for unknown categories due to less reliable dose denominators

Table 4: Expected rates of death per million vaccinated persons* and reported deaths to VAERS per million vaccinated persons within 7 and 42 days of vaccination among mRNA COVID-19 vaccine recipients, by age— December 14, 2020–June 14, 2021

	Expected	death rates	Rates based on reported deaths in VAERS						
	All-caus per million vac	se deaths cinated persons	Both mRN per million vac	A vaccines ccinated persons	BNT162 per million vac	b2 vaccine ccinated persons	mRNA-12 per million vac	273 vaccine ccinated persons	
	Within 7 days of vaccination	Within 42 days of vaccination	Within 7 daysWithin 42 daysWithin 42 daysof vaccinationof vaccinationof vaccination		Within 7 days of vaccination	Within 42 days of vaccination	Within 7 days of vaccination	Within 42 days of vaccination	
Age (years) ^{\dagger}									
16–24‡	14.2	85.1	0.9	1.7	0.8	1.6	$1 \cdot 0$	1.9	
25–34	25.5	152.7	1.6	2.8	1.5	2.6	1.8	3.2	
35–44	37.4	224.5	2.5	4.6	2.3	3.8	2.9	5.7	
45–54	77.0	461.7	5.2	8.8	4.3	7.0	6.4	11.1	
55–64	169.8	1,018.6	9.0	16.6	6.7	13.5	11.8	20.2	
65–74	343.2	2,059.4	17.9	34.4	16.5	30.6	19.3	38.0	
75–84	857-2	5,143.0	30.0	66.1	26.8	59.1	33.0	72.9	
≥85	2,601.4	15,608.3	105.5	236.6	95.9	212.8	115.4	261.3	
Total	165.6	993.3	11.7	23.6	9.2	18.5	15.0	30.5	

*Expected rates of deaths were derived from Abara WE, Gee J, Mu Y, Deloray M, Ye T, Shay DK, et al. Expected Rates of Select Adverse Events following Immunization for COVID-19 Vaccine Safety Monitoring. medRxiv. 2021.

[†]Age grouping were selected for comparison to the expected rates age groups

[‡] The expected rates are among individuals aged 15-24

	Both mRNA vaccines	BNT162b2 vaccin	e	mRNA-1273 vaccine		
Characteristics	(- 7.014.592)	Dose 1	Dose 2	Dose 1	Dose 2	
	(n = 7,914,583)	n=3,455,778	n=2,920,526	n= 3,319,737	n=2,753,894	
Sex						
Female	4,975,209 (62.9)	2,150,068 (62.2)	1,861,599 (63.7)	2,073,542 (62.5)	1,779,200 (64.6)	
Male	2,860,738 (36.1)	1,272,011 (36.8)	1,032,941 (35.4)	1,210,622 (36.5)	947,612 (34.4)	
Other	8,872 (0.1)	4,027 (0.1)	3,464 (0.1)	3,443 (0.1)	2,947 (0.1)	
Unknown	69,764 (0.9)	29,672 (0.9)	22,522 (0.8)	32,130 (1.0)	24,135 (0.9)	
Age (years)						
16–17	73,347 (0.9)	63,865 (1.8)	38,530 (1.3)	946 (0.03)	473 (0.02)	
18–49	3,791,839 (47.9)	1,726,465 (50.0)	1,431,627 (49.0)	1,505,760 (45.4)	1,219,210 (44.3)	
50–59	1,500,981 (19.0)	653,799 (18.9)	574,422 (19.7)	627,214 (18.9)	531,200 (19.3)	
60–64	739,381 (9.3)	315,404 (9.1)	279,350 (9.6)	316,768 (9.5)	270,831 (9.8)	
65–74	1,344,721 (17.0)	516,227 (14.9)	452,928 (15.5)	643,663 (19.4)	557,279 (20.2)	
≥75	464,314 (5.9)	180,018 (5.2)	143,669 (4.9)	225,386 (6.8)	174,901 (6.4)	
Race/Ethnicity						
Hispanic	782,301 (9.9)	346,197 (10.0)	288,263 (9.9)	316,460 (9.5)	256,185 (9.3)	
Non-Hispanic						
White	4,701,715 (59.4)	2,059,560 (59.6)	1,896,823 (64.9)	1,979,056 (59.6)	1,830,413 (66.5)	
Black	443,938 (5.6)	202,598 (5.9)	176,164 (6.0)	178,981 (5.4)	153,667 (5.6)	
Asian	467,932 (5.9)	215,713 (6.2)	196,173 (6.7)	154,498 (4.7)	138,793 (5.0)	
American Indian or Alaska Native	27,899 (0.4)	11,161 (0.3)	9,194 (0.3)	13,486 (0.4)	11,410 (0.4)	
Native Hawaiian or Other Pacific Islander	19,393 (0.2)	8,500 (0.2)	7,373 (0.3)	7,689 (0.2)	6,664 (0.2)	
Multiple races	110,326 (1.4)	50,954 (1.5)	46,129 (1.6)	41,977 (1.3)	38,772 (1.4)	
Other races	42,230 (0.5)	19,252 (0.6)	16,757 (0.6)	15,885 (0.5)	13,880 (0.5)	
Unknown race	23,420 (0.3)	10,249 (0.3)	9,090 (0.3)	9,502 (0.3)	8,270 (0.3)	
Unknown ethnicity*						
White	115,766 (1.5)	48,084 (1.4)	38,674 (1.3)	52,143 (1.6)	42,070 (1.5)	
Black	26,865 (0.3)	11,602 (0.3)	8,570 (0.3)	11,993 (0.4)	8,406 (0.3)	
Asian	33,146 (0.4)	14,134 (0.4)	11,844 (0.4)	11,356 (0.3)	9,153 (0.3)	
American Indian or Alaska Native	3,142 (0.04)	1,206 (0.03)	848 (0.03)	1,582 (0.05)	1,151 (0.04)	
Native Hawaiian or Other Pacific Islander	1,945 (0.02)	815 (0.02)	659 (0.02)	800 (0.02)	613 (0.02)	
Multiple races	6,370 (0.1)	2,902 (0.1)	2,408 (0.1)	2,478 (0.1)	2,041 (0.1)	
Other races	13,148 (0.2)	5,681 (0.2)	4,528 (0.2)	5,414 (0.2)	4,263 (0.2)	
Unknown race and ethnicity*	129,647 (1.6)	56,481 (1.6)	45,410 (1.6)	54,969 (1.7)	44,340 (1.6)	
Unavailable [†]	965,400 (12.2)	390,689 (11.3)	161,619 (5.5)	461,468 (13.9)	183,803 (6.7)	
Pregnant at time of vaccination	86,801 (1.1)	39,884 (1.2)	39,163 (1.3)	25,255 (0.8)	25,428 (0.9)	
Pregnancy test positive after vaccination	27,370 (0.3)	1,548 (0.04)	11,677 (0.4)	4,009 (0.1)	10,199 (0.4)	

Table 5: Demographic characteristics of v-safe participants reporting receipt of mRNA COVID-19 vaccine and completing at least one health survey 0-7 days after vaccination—December 14, 2020–June 14, 2021

Data are n (%) unless otherwise stated.

*Unknown indicates that v-safe participants selected unknown or preferred not to say.

[†]Unavailable refers to information that was not collected or missing in v-safe.

Table 6: Reported local and systemic reactions*, and reported health impact following mRNA COVID-19 vaccines reported days 0–7 after vaccination to v-safe, by manufacturer and dose—December 14, 2020 – June 14, 2021

	Both mRNA vaccines		BNT162b2 vaccine		mRNA-1273 vaccine	
	Dose 1	Dose 2	Dose 1	Dose 2	Dose 1	Dose 2
	(n=6,775,515)	(n=5,674,420)	(n=3,455,778)	(n=2,920,526)	(n=3,319,737)	(n=2,753,894)
Any injection site reaction	4,644,989 (68.6)	4,068,447 (71.7)	2,212,051 (64.0)	1,908,124 (65.3)	2,432,938 (73.3)	2,160,323 (78.4)
Injection site pain	4,488,402 (66.2)	3,890,848 (68.6)	2,140,843 (61.9)	1,835,398 (62.8)	2,347,559 (70.7)	2,055,450 (74.6)
Swelling	703,790 (10.4)	976,946 (17·2)	246,230 (7.1)	309,718 (10.6)	457,560 (13.8)	667,228 (24.2)
Redness	353,788 (5.2)	640,739 (11.3)	116,108 (3.4)	167,127 (5.7)	237,680 (7.2)	473,612 (17.2)
Itching	376,076 (5.6)	605,633 (10.7)	145,596 (4.2)	191,132 (6.5)	230,480 (6.9)	414,501 (15.1)
Any systemic reaction	3,573,429 (52.7)	4,018,920 (70.8)	1,771,509 (51.3)	1,931,643 (66.1)	1,801,920 (54.3)	2,087,277 (75.8)
Fatigue	2,295,205 (33.9)	3,158,299 (55.7)	1,127,904 (32.6)	1,475,646 (50.5)	1,167,301 (35.2)	1,682,653 (61.1)
Headache	1,831,471 (27.0)	2,623,721 (46.2)	893,992 (25.9)	1,189,444 (40.7)	937,479 (28.2)	1,434,277 (52.1)
Myalgia	1,423,336 (21.0)	2,478,170 (43.7)	653,821 (18.9)	1,085,365 (37.2)	769,515 (23.2)	1,392,805 (50.6)
Chills	631,546 (9.3)	1,680,185 (29.6)	263,617 (7.6)	642,856 (22.0)	367,929 (11.1)	1,037,329 (37.7)
Fever	642,092 (9.5)	1,679,577 (29.6)	274,650 (7.9)	656,454 (22.5)	367,442 (11.1)	1,023,123 (37.2)
Joint pain	642,006 (9.5)	1,440,927 (25.4)	285,812 (8.3)	591,877 (20.3)	356,194 (10.7)	849,050 (30.8)
Nausea	562,273 (8.3)	901,103 (15.9)	267,160 (7.7)	384,525 (13.2)	295,113 (8.9)	516,578 (18.8)
Diarrhea	383,576 (5.7)	419,044 (7.4)	190,542 (5.5)	198,618 (6.8)	193,034 (5.8)	220,426 (8.0)
Abdominal pain	233,511 (3.4)	359,107 (6.3)	113,872 (3.3)	158,251 (5.4)	119,639 (3.6)	200,856 (7.3)
Rash	85,766 (1.3)	99,878 (1.8)	41,565 (1.2)	42,662 (1.5)	44,201 (1.3)	57,216 (2.1)
Vomiting	55,710 (0.8)	91,727 (1.6)	25,336 (0.7)	36,761 (1.3)	30,374 (0.9)	54,966 (2.0)
With reported health impact*	808,963 (11.9)	1,821,421 (32.1)	361,834 (10.5)	740,529 (25.4)	447,129 (13.5)	1,080,892 (39.2)
Unable to do normal activity	658,330 (9.7)	1,501,679 (26.5)	290,207 (8.4)	598,584 (20.5)	368,123 (11.1)	903,095 (32.8)
Unable to work	305,709 (4.5)	911,366 (16-1)	135,063 (3.9)	360,411 (12.3)	170,646 (5.1)	550,955 (20.0)
Reported medical care	56,647 (0.8)	53,077 (0.9)	27,358 (0.8)	25,568 (0.9)	29,289 (0.9)	27,509 (1.0)
Telehealth	19,562 (0.3)	19,770 (0.3)	9,318 (0.3)	9,238 (0.3)	10,244 (0.3)	10,532 (0.4)
Clinic	18,671 (0.3)	16,793 (0.3)	9,109 (0.3)	8,487 (0.3)	9,562 (0.3)	8,306 (0.3)
Emergency visit	9,907 (0.1)	8,907 (0.2)	5,087 (0.1)	4,494 (0.2)	4,820 (0.1)	4,413 (0.2)
Hospitalization	1,896 (0.03)	2,053 (0.04)	915 (0.03)	1,001 (0.03)	981 (0.03)	1,052 (0.04)

Data are n (%) unless otherwise stated.

*Reports of local and systemic reactions, and reports of health impact are not mutually exclusive.



Figure 1: Local and systemic reactions* to mRNA COVID-19 vaccine reported in v-safe, by manufacturer, dose, days after vaccination, and severity[†]

*Top five reactions determined by reported frequency after dose 2 of both mRNA COVID-19 vaccines in v-safe, excluding fever because it was not rated mild, moderate, or severe. [†]Mild was defined as "noticeable symptoms but they aren't a problem", moderate was defined as "symptoms that limit normal activities, and severe symptoms "make normal daily activities difficult or impossible"

Characteristics	Both mRNA vaccines [†]	BNT162b2 vaccine [†]	mRNA-1273 vaccine [†]	
	(n=298,792,852)	(n=167,177,332)	(n=131,639,515)	
Sex				
Female	155,969,573 (53-2)	86,507,992 (53.5)	69,461,582 (52.8)	
Male	134,373,958 (45.8)	73,768,602 (45.6)	60,605,356 (46.1)	
Unknown	2,868,979 (1.0)	1,452,344 (0.9)	1,416,634 (1.1)	
Age (years)				
16–17*	5,506,763 (1.8)	5,365,855 (3.2)	140,908 (0.1)	
18–49	126,288,626 (42.3)	74,999,327 (44.9)	51,289,299 (39.0)	
50-64	79,207,752 (26.5)	43,595,972 (26.1)	35,611,780 (27.1)	
65–74	51,699,307 (17.3)	25,402,217 (15.2)	26,297,090 (20.0)	
75–84	27,731,181 (9.3)	13,555,128 (8.1)	14,176,053 (10.8)	
≥85	8,359,223 (2.8)	4,248,648 (2.5)	4,110,575 (3.1)	
Unknown	23,995 (0.01)	10,185 (0.01)	13,810 (0.01)	
Race/Ethnicity				
Hispanic	31,599,632 (10.8)	17,964,345 (11.1)	13,635,287 (10.4)	
Non-Hispanic				
White	112,698,875 (38.4)	61,996,607 (38.3)	50,702,268 (38.6)	
Asian	11,789,429 (4.0)	7,258,033 (4.5)	4,531,396 (3.4)	
Black	16,848,436 (5.7)	9,665,586 (6.0)	7,182,849 (5.5)	
American Indian or Alaska Native	1,738,938 (0.6)	842,263 (0.5)	896,674 (0.7)	
Native Hawaiian or Other Pacific Islander	508,285 (0.2)	295,634 (0.2)	212,651 (0.2)	
Multiple races	8,856,800 (3.0)	5,037,828 (3.1)	3,818,972 (2.9)	
Other races	6,949,404 (2.4)	4,161,353 (2.6)	2,788,051 (2.1)	
Unknown race and ethnicity	102,227,532 (34.9)	54,511,493 (33.7)	47,716,039 (36.3)	

Supplemental Table 1: mRNA COVID-19 vaccine doses adminis	stered in the United States	s—December 14, 2020–3	June 14, 2021

Data are n (%) unless otherwise stated.

*mRNA-1273 vaccine was not authorized for individuals <18 years during this period, reported mRNA-1273 doses are either from clinical trials or were administered or reported in error [†]Totals reflect the number of doses in age categories. Missing doses for sex and race/ethnicity are due to certain jurisdictions that report data in aggregate.

Supplemental Table 2: Most common causes of death among reports received and processed by Vaccine Adverse Event Reporting System (VAERS) following mRNA COVID-19 vaccination (n=4,472)—December 14, 2020–June 14, 2021

	Death or autopsy certificate available			No death certificate or autopsy available				
ICD-10 Major Group	Both mRNA vaccines	BNT162b2 vaccine	mRNA-1273 vaccine	Both mRNA vaccines	BNT162b2 vaccine	mRNA-1273 vaccine		
	(n=808)	(n=401)	(n=407)	(n=3,664)	(n=1,686)	(n=1,978)		
All reported deaths								
Diseases of the heart	376 (46.5)	161 (40.1)	215 (52.8)	622 (17.0)	296 (17.6)	326 (16.5)		
COVID-19 disease	102 (12.6)	62 (15.5)	40 (9.8)	317 (8.7)	178 (10.6)	139 (7.0)		
Other	68 (8.4)	38 (9.5)	30 (7.4)	141 (3.8)	68 (4.0)	73 (3.7)		
Cerebrovascular diseases	53 (6.6)	28 (7.0)	25 (6.1)	207 (5.6)	101 (6.0)	106 (5.4)		
Dementia	41 (5.1)	20 (5.0)	21 (5.2)	9 (0.2)	3 (0.2)	6 (0.3)		
Chronic lower respiratory diseases	28 (3.5)	17 (4.2)	11 (2.7)	29 (0.8)	10 (0.6)	19 (1.0)		
Malignant neoplasms	27 (3.3)	15 (3.7)	12 (2.9)	68 (1.9)	42 (2.5)	26 (1.3)		
Unknown/unclear	27 (3.3)	9 (2·2)	18 (4.4)	1,984 (54-1)	844 (50.1)	1,140 (57.6)		
Septicemia	23 (2.8)	12 (3.0)	11 (2.7)	72 (2.0)	47 (2.8)	25 (1.3)		
Influenza and pneumonia	22 (2.7)	18 (4.5)	4 (1.0)	113 (3.1)	52 (3.1)	61 (3.1)		
Accidents/unintentional injuries	11 (1.4)	3 (0.7)	8 (2.0)	22 (0.6)	8 (0.5)	14 (0.7)		
Renal disease	8 (1.0)	5 (1.2)	3 (0.7)	25 (0.7)	7 (0.4)	18 (0.9)		
Hematologic disease, other than malignancy	7 (0.9)	5 (1.2)	2 (0.5)	19 (0.5)	9 (0.5)	10 (0.5)		
Pneumonitis due to solids and liquids	6 (0.7)	3 (0.7)	3 (0.7)	8 (0.2)	5 (0.3)	3 (0.2)		
Diabetes mellitus	4 (0.5)	1 (0.2)	3 (0.7)	6 (0.2)	4 (0.2)	2 (0.1)		
Chronic liver disease and cirrhosis	4 (0.5)	3 (0.7)	1 (0.2)	7 (0.2)	4 (0.2)	3 (0.2)		
Intentional self-harm	1 (0.1)	1 (0.2)	0 (0.0)	15 (0.4)	8 (0.5)	7 (0.4)		

Data are n (%) unless otherwise stated.

Supplemental Table 3: Causes and impressions of death among reported deaths to Vaccine Adverse Event Reporting System (VAERS) December 14, 2020–June 14, 2021 following mRNA vaccination

ICD-10 Major Group and impression	All reports of death	Reported deaths with death certificate or autopsy
All reported deaths, n	4,472	808
Diseases of the heart	998 (22.3)	376 (46.5)
Aortic dissection, aneurysm or aortitis	13 (0.3)	7 (0.9)
Arrhythmia	42 (0.9)	18 (2.2)
Atherosclerotic cardiovascular or hypertensive cardiovascular disease	129 (2.9)	98 (12.1)
Cardiac arrest	321 (7.2)	79 (9.8)
Cardiomyopathy or hypertrophy	17 (0.4)	12 (1.5)
Heart failure	104 (2.3)	47 (5.8)
Myocardial infarction	247 (5.5)	87 (10.8)
Myocarditis	4 (0.1)	0 (0.0)
Pulmonary embolism	84 (1.9)	17 (2.1)
Other cardiac cause	37 (0.8)	11 (1.4)
COVID-19 disease	419 (9.4)	102 (12.6)
Cerebrovascular diseases	260 (5.8)	53 (6.6)
Other	209 (4.7)	68 (8.4)
Disseminated herpes zoster	2 (0.04)	1 (0.1)
Drug overdose/intoxication	7 (0.2)	5 (0.6)
Failure to thrive	9 (0.2)	9 (1.1)
$Gastrointestinal^*$	36 (0.8)	8 (1.0)
Hemorrhage/Hemorrhagic shock	4 (0.1)	2 (0.2)
Metabolic derangement	4 (0.1)	1 (0.1)
Multiorgan failure	28 (0.6)	7 (0.9)
Natural	2 (0.04)	2 (0.2)
Neurologic [†]	28 (0.6)	6 (0.7)
Obesity	2 (0.04)	2 (0.2)
Respiratory failure	63 (1.4)	22 (2.7)
Vaccine related [‡]	4 (0.1)	3 (0.4)
Influenza and pneumonia	135 (3.0)	22 (2.7)
Malignant neoplasms	95 (2.1)	27 (3.3)
Septicemia	95 (2.1)	23 (2.8)
Chronic lower respiratory diseases	57 (1.3)	28 (3.5)
Dementia	50 (1.1)	41 (5.1)
Accidents/unintentional injuries	33 (0.7)	11 (1.4)
Renal disease	33 (0.7)	8 (1.0)
Hematologic disease, other than malignancy	26 (0.6)	7 (0.9)
Intentional self-harm	16 (0.4)	1 (0.1)
Pneumonitis due to solids and liquids	14 (0.3)	6 (0.7)
Chronic liver disease and cirrhosis	11 (0.2)	4 (0.5)
Diabetes mellitus	10 (0.2)	4 (0.5)
Unknown/unclear	2,011 (45.0)	27 (3.3)

Data are n (%) unless otherwise stated.

*Gastrointestinal includes gastrointestinal bleeding, bowel obstruction/perforation, mesenteric ischemia, pancreatitis.

[†]Neurologic includes amyotrophic lateral sclerosis, encephalopathy, hydrocephalus, Guillain-Barré syndrome, seizure.

[‡]Vaccine related includes systemic inflammatory response syndrome from vaccine reaction, anaphylaxis post-COVID-19 vaccination

Supplemental Table 4: Causes of death among reports of death with death certificate or autopsy available to Vaccine Adverse Event Reporting System (VAERS) December 14, 2020–June 14, 2021 following mRNA vaccination, by age

	All	10	5-24	25	5-34	35-	-44	4	5-54	55-	-64	65	-74	75	5-84	85	i+
ICD-10 Major Group	808	r	1=4	n	=5	n=	24	n	=37	n=	101	n=	171	n=	202	n=2	264
		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Diseases of the heart	376	1	25.0	2	40.0	10	41.7	21	56.8	50	49.5	98	57.3	91	45.0	103	39.0
COVID-19 disease	102	1	25.0	1	20.0	2	8.3	2	5.4	11	10.9	19	11.1	32	15.8	34	12.9
Cerebrovascular diseases	53	0	0	0	0	0	0	4	10.8	8	7.9	8	4.7	13	6.4	20	7.6
Other	68	0	0	1	20.0	5	20.8	2	5.4	10	9.9	11	6.4	15	7.4	24	9.1
Influenza and pneumonia	22	0	0	0	0	2	8.3	0	0	2	2.0	4	2.3	7	3.5	7	2.7
Malignant neoplasms	27	0	0	0	0	0	0	2	5.4	5	5.0	2	1.2	8	4.0	10	3.8
Septicemia	23	0	0	0	0	0	0	1	2.7	2	2.0	2	1.2	6	3.0	12	4.5
Chronic lower respiratory diseases	28	0	0	0	0	0	0	0	0	2	2.0	11	6.4	4	2.0	11	4.2
Dementia, including Alzheimer's, Parkinson's disease	41	0	0	0	0	0	0	0	0	1	1.0	1	0.6	7	3.5	32	12.1
Accidents/unintentional injuries	11	0	0	0	0	1	4.2	0	0	2	2.0	2	1.2	3	1.5	3	1.1
Renal disease, including nephritis and chronic disease	8	0	0	0	0	0	0	1	2.7	0	0	3	1.8	3	1.5	1	0.4
Hematologic disease, other than malignancy	7	0	0	0	0	0	0	1	2.7	3	3.0	0	0	1	0.5	2	0.8
Intentional self-harm	1	1	25.0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Pneumonitis due to solids and liquids	6	0	0	0	0	0	0	0	0	1	1.0	0	0	3	1.5	2	0.8
Chronic liver disease and cirrhosis	4	0	0	1	20.0	0	0	0	0	1	1.0	1	0.6	1	0.5	0	0
Diabetes mellitus	4	0	0	0	0	1	4.2	0	0	0	0	2	1.2	1	0.5	0	0
Unknown/unclear	27	1	25.0	0	0	3	12.5	3	8.1	3	3.0	7	4.1	7	3.5	3	1.1

Supplemental Table 5: Local and systemic reactions*	* 0–7 days after vaccination by sex, ag	ge, and dose number, reported in v-saf	e—December 14, 2020–
June 14, 2021			

	Female		Male		<65 years		≥65 years		
	Dose 1	Dose 2 Dose 1		Dose 2	Dose 1	Dose 2	Dose 1	Dose 2	
	(n=4,223,610)	(n=3,640,799)	(n=2,482,633)	(n=1,980,553)	(n=5,210,221)	(n=4,345,643)	(n=1,565,294)	(n=1,328,777)	
Any injection site reaction	3,095,194 (73.3)	2,792,488 (76.7)	1,498,108 (60.3)	1,235,278 (62.4)	3,835,618 (73.6)	3,290,206 (75.7)	809,371 (51.7)	778,241 (58.6)	
Injection site pain	2,989,733 (70.8)	2,666,734 (73.3)	1,448,440 (58.3)	1,184,914 (59.8)	3,728,795 (71.6)	3,179,024 (73.2)	759,607 (48.5)	711,824 (53.6)	
Swelling	539,793 (12.8)	771,962 (21.2)	154,980 (6-2)	194,033 (9.8)	604,868 (11.6)	812,126 (18.7)	98,922 (6.3)	164,820 (12.4)	
Redness	283,345 (6.7)	529,175 (14.5)	66,134 (2.7)	104,933 (5.3)	295,413 (5.7)	512,516 (11.8)	58,375 (3.7)	128,223 (9.6)	
Itching	299,407 (7.1)	504,016 (13.8)	72,095 (2.9)	95,356 (4.8)	309,607 (5.9)	466,319 (10.7)	66,469 (4-2)	139,314 (10.5)	
Any systemic reaction	2,444,362 (57.9)	2,752,592 (75.6)	1,088,296 (43.8)	1,226,561 (61.9)	2,972,931 (57.1)	3,237,621 (74.5)	600,498 (38-4)	781,299 (58.8)	
Fatigue	1,624,531 (38.5)	2,221,361 (61.0)	643,206 (25.9)	904,556 (45.7)	1,941,979 (37.3)	2,588,541 (59.6)	353,226 (22.6)	569,758 (42.9)	
Headache	1,349,155 (31.9)	1,906,337 (52.4)	460,786 (18.6)	690,138 (34.8)	1,595,091 (30.6)	2,226,046 (51-2)	236,380 (15.1)	397,675 (29.9)	
Myalgia	954,469 (22.6)	1,724,474 (47-4)	450,562 (18.1)	726,994 (36.7)	1,219,190 (23.4)	2,085,722 (48.0)	204,146 (13.0)	392,448 (29.5)	
Chills	451,583 (10.7)	1,202,364 (33.0)	172,283 (6.9)	459,577 (23.2)	542,285 (10.4)	1,426,710 (32.8)	89,261 (5.7)	253,475 (19.1)	
Fever	446,178 (10.6)	1,182,201 (32.5)	187,713 (7.6)	478,912 (24.2)	565,804 (10.9)	1,449,504 (33.4)	76,288 (4.9)	230,073 (17.3)	
Joint pain	444,630 (10.5)	1,023,525 (28.1)	188,846 (7.6)	400,963 (20.2)	539,196 (10.3)	1,214,624 (28.0)	102,810 (6.6)	226,303 (17.0)	
Nausea	447,766 (10.6)	728,730 (20.0)	106,872 (4.3)	161,455 (8.2)	500,782 (9.6)	794,450 (18.3)	61,491 (3.9)	106,653 (8.0)	
Diarrhea	272,890 (6.5)	313,252 (8.6)	106,079 (4.3)	101,107 (5.1)	323,773 (6-2)	352,077 (8.1)	59,803 (3.8)	66,967 (5.0)	
Abdominal pain	179,210 (4.2)	283,422 (7.8)	50,991 (2.1)	71,115 (3.6)	203,575 (3.9)	316,165 (7.3)	29,936 (1.9)	42,942 (3.2)	
Rash	65,498 (1.6)	79,092 (2.2)	19,193 (0.8)	19,735 (1.0)	70,985 (1.4)	79,913 (1.8)	14,781 (0.9)	19,965 (1.5)	
Vomiting	43,998 (1.0)	75,650 (2.1)	10,936 (0.4)	14,915 (0.8)	49,483 (0.9)	81,733 (1.9)	6,227 (0.4)	9,994 (0.8)	

Data are n (%) unless otherwise stated.

*Reports of local and systemic reactions are not mutually exclusive.

Supplemental Table 6: Most common local and systemic reactions^{*} to mRNA COVID-19 vaccine reported in v-safe, by dose and severity,[†] 0-7 days after vaccination with BNT162b2 vaccine

		Dose 1			Dose 2					
	Day	All, n	Severe	Moderate	Mild	All, n	Severe	Moderate	Mild	
Injection site pain	0	2,272,335	2,533 (0.1)	80,358 (3.5)	599,511 (26.4)	1,766,510	4,359 (0.2)	94,156 (5.3)	503,779 (28.5)	
	1	2,545,271	18,827 (0.7)	334,755 (13.2)	1,289,293 (50.7)	2,027,330	48,810 (2.4)	453,726 (22.4)	885,434 (43.7)	
	2	2,545,434	4,356 (0.2)	62,838 (2.5)	565,455 (22.2)	2,116,614	10,391 (0.5)	126,741 (6.0)	680,408 (32.1)	
	3	2,507,344	2,119 (0.1)	26,602 (1.1)	216,785 (8.6)	2,067,908	3,332 (0.2)	43,037 (2.1)	336,222 (16.3)	
	4	2,436,977	1,420 (0.1)	16,710 (0.7)	102,077 (4.2)	2,028,926	1,820 (0.1)	20,548 (1.0)	149,408 (7.4)	
	5	2,332,032	1,138 (0.05)	11,965 (0.5)	60,401 (2.6)	2,000,426	1,272 (0.1)	12,719 (0.6)	73,486 (3.7)	
	6	2,249,409	909 (0.04)	8,901 (0.4)	40,597 (1.8)	2,000,472	1,106 (0.1)	11,008 (0.6)	46,707 (2.3)	
	7	2,198,611	768 (0.03)	7,783 (0.4)	32,967 (1.5)	2,067,201	1,557 (0.1)	14,159 (0.7)	44,322 (2.1)	
	0	2,272,335	7,280 (0.3)	79,232 (3.5)	193,192 (8.5)	1,766,510	11,293 (0.6)	98,802 (5.6)	185,051 (10.5)	
	1	2,545,271	35,734 (1.4)	229,606 (9.0)	326,015 (12.8)	2,027,330	135,581 (6.7)	523,998 (25.8)	373,601 (18.4)	
	2	2,545,434	16,936 (0.7)	114,562 (4.5)	211,046 (8.3)	2,116,614	39,668 (1.9)	217,269 (10.3)	308,126 (14.6)	
Fatigue	3	2,507,344	10,636 (0.4)	74,341 (3.0)	145,114 (5.8)	2,067,908	15,361 (0.7)	104,673 (5.1)	193,174 (9.3)	
Faugue	4	2,436,977	8,275 (0.3)	58,170 (2.4)	109,266 (4.5)	2,028,926	10,280 (0.5)	69,886 (3.4)	133,603 (6.6)	
	5	2,332,032	7,062 (0.3)	49,739 (2.1)	88,721 (3.8)	2,000,426	8,089 (0.4)	55,840 (2.8)	103,919 (5.2)	
	6	2,249,409	6,428 (0.3)	44,044 (2.0)	76,633 (3.4)	2,000,472	7,200 (0.4)	48,388 (2.4)	86,907 (4.3)	
	7	2,198,611	6,027 (0.3)	40,428 (1.8)	67,168 (3.1)	2,067,201	7,528 (0.4)	46,669 (2.3)	78,361 (3.8)	
	0	2,272,335	3,394 (0.1)	42,501 (1.9)	167,985 (7.4)	1,766,510	5,217 (0.3)	52,759 (3.0)	144,892 (8.2)	
	1	2,545,271	20,011 (0.8)	129,629 (5.1)	265,970 (10.4)	2,027,330	82,393 (4.1)	333,605 (16.5)	381,368 (18.8)	
	2	2,545,434	10,458 (0.4)	69,347 (2.7)	162,658 (6.4)	2,116,614	24,063 (1.1)	134,054 (6.3)	249,895 (11.8)	
	3	2,507,344	6,670 (0.3)	46,850 (1.9)	110,115 (4.4)	2,067,908	10,356 (0.5)	68,461 (3.3)	148,990 (7.2)	
Headache	4	2,436,977	5,552 (0.2)	38,319 (1.6)	85,635 (3.5)	2,028,926	7,238 (0.4)	47,550 (2.3)	103,204 (5.1)	
	5	2,332,032	4,911 (0.2)	34,379 (1.5)	72,831 (3.1)	2,000,426	6,154 (0.3)	40,322 (2.0)	82,191 (4.1)	
	6	2,249,409	4,733 (0.2)	31,540 (1.4)	64,890 (2.9)	2,000,472	5,467 (0.3)	35,177 (1.8)	69,168 (3.5)	
	7	2,198,611	4,381 (0.2)	29,475 (1.3)	58,752 (2.7)	2,067,201	5,372 (0.3)	34,057 (1.6)	63,628 (3.1)	
	0	2,272,335	1,999 (0.1)	29,601 (1.3)	96,095 (4.2)	1,766,510	4,001 (0.2)	38,960 (2.2)	75,790 (4.3)	
	1	2,545,271	18,440 (0.7)	136,939 (5.4)	219,125 (8.6)	2,027,330	101,801 (5.0)	408,637 (20.2)	293,241 (14.5)	
	2	2,545,434	7,441 (0.3)	56,954 (2.2)	112,788 (4.4)	2,116,614	23,521 (1.1)	140,700 (6.6)	209,074 (9.9)	
	3	2,507,344	4,200 (0.2)	33,605 (1.3)	65,696 (2.6)	2,067,908	6,925 (0.3)	54,206 (2.6)	100,982 (4.9)	
Myalgia	4	2,436,977	3,255 (0.1)	25,814 (1.1)	46,369 (1.9)	2,028,926	4,146 (0.2)	32,786 (1.6)	60,489 (3.0)	
	5	2,332,032	2,831 (0.1)	22,598 (1.0)	37,598 (1.6)	2,000,426	3,239 (0.2)	25,326 (1.3)	44,242 (2.2)	
	6	2,249,409	2,543 (0.1)	20,904 (0.9)	33,016 (1.5)	2,000,472	2,973 (0.1)	22,422 (1.1)	36,522 (1.8)	
	7	2,198,611	2,504 (0.1)	19,474 (0.9)	30,222 (1.4)	2,067,201	3,379 (0.2)	23,046 (1.1)	33,563 (1.6)	
	0	2,272,335	879 (0.04)	8,246 (0.4)	34,000 (1.5)	1,766,510	2,091 (0.1)	14,428 (0.8)	38,195 (2.2)	
	1	2,545,271	8,558 (0.3)	45,518 (1.8)	78,033 (3.1)	2,027,330	62,884 (3.1)	210,579 (10.4)	207,218 (10.2)	
	2	2,545,434	3,371 (0.1)	18,659 (0.7)	36,412 (1.4)	2,116,614	11,744 (0.6)	51,490 (2.4)	76,276 (3.6)	
~	3	2,507,344	1,462 (0.1)	9,241 (0.4)	19,569 (0.8)	2,067,908	2,582 (0.1)	13,423 (0.6)	25,421 (1.2)	
Chills	4	2,436,977	1,051 (0.04)	6,915 (0.3)	13,967 (0.6)	2,028,926	1,336 (0.1)	7,424 (0.4)	14,223 (0.7)	
	5	2,332,032	863 (0.04)	5,531 (0.2)	11,284 (0.5)	2,000,426	955 (0.05)	5,423 (0.3)	10,583 (0.5)	
	6	2,249,409	779 (0.03)	5,048 (0.2)	9,932 (0.4)	2,000,472	851 (0.04)	4,763 (0.2)	9,029 (0.5)	
	7	2,198,611	752 (0.03)	4,645 (0.2)	8,889 (0.4)	2,067,201	1,222 (0.1)	5,515 (0.3)	9,039 (0.4)	
	0	2,272,335	1,069 (0.05)	11,375 (0.5)	24,689 (1.1)	1,766,510	2,396 (0.1)	18,677 (1.1)	25,699 (1.5)	
	1	2,545,271	9,676 (0.4)	61,691 (2.4)	69,532 (2.7)	2,027,330	55,446 (2.7)	225,949 (11.2)	137,601 (6.8)	
	2	2,545,434	4,608 (0.2)	31,238 (1.2)	44,072 (1.7)	2,116,614	14,386 (0.7)	80,490 (3.8)	90,461 (4.3)	
.	3	2,507,344	2,675 (0.1)	19,912 (0.8)	29,313 (1.2)	2,067,908	4,624 (0.2)	32,971 (1.6)	46,467 (2.2)	
Joint pain	4	2,436,977	2,165 (0.1)	15,923 (0.7)	22,386 (0.9)	2,028,926	2,882 (0.1)	20,861 (1.0)	29,916 (1.5)	
	5	2,332,032	1,999 (0.1)	13,922 (0.6)	18,869 (0.8)	2,000,426	2,341 (0.1)	16,528 (0.8)	23,366 (1.2)	
	6	2,249,409	1,773 (0.1)	13,018 (0.6)	16,874 (0.8)	2,000,472	2,138 (0.1)	15,046 (0.8)	19,649 (1.0)	
	7	2,198,611	1,686 (0.1)	12,245 (0.6)	15,605 (0.7)	2,067,201	2,462 (0.1)	15,782 (0.8)	18,678 (0.9)	

Data are n (%) unless otherwise stated.

*Top five reactions determined by reported frequency after dose 2 of both mRNA COVID-19 vaccines in v-safe, excluding fever because it was not rated mild/moderate/severe. Symptoms are not mutually exclusive. [†]Mild was defined as "noticeable symptoms but they aren't a problem", moderate was defined as "symptoms that limit normal activities, and severe symptoms", and severe symptoms "make normal daily activities difficult or impossible".

Supplemental Table 7: Most common local and systemic reactions* to mRNA COVID-19 vaccine reported in
v-safe, by dose and severity, [†] 0-7 days after vaccination with mRNA-1273 vaccine

		Dose 1				Dose 2					
	Day	All, n	Severe	Moderate	Mild	All, n	Severe	Moderate	Mild		
Injection site pain	0	2,112,380	4,971 (0.2)	113,992 (5.4)	595,108 (28.2)	1,656,723	11,174 (0.7)	155,045 (9.4)	535,260 (32.3)		
	1	2,424,231	49,225 (2.0)	512,076 (21.1)	1,214,808 (50.1)	1,937,029	131,379 (6.8)	735,284 (38.0)	713,164 (36.8)		
	2	2,474,399	13,723 (0.6)	152,289 (6.2)	932,770 (37.7)	2,035,773	26,351 (1.3)	268,371 (13.2)	890,618 (43.7)		
	3	2,459,431	4,778 (0.2)	47,625 (1.9)	370,863 (15.1)	1,993,354	6,469 (0.3)	78,053 (3.9)	525,158 (26.3)		
	4	2,390,709	2,881 (0.1)	25,370 (1.1)	148,288 (6.2)	1,960,829	3,393 (0.2)	33,315 (1.7)	239,358 (12.2)		
	5	2,285,185	2,087 (0.1)	17,079 (0.7)	77,452 (3.4)	1,939,300	2,493 (0.1)	19,783 (1.0)	109,652 (5.7)		
	6	2,196,757	1,512 (0.1)	12,520 (0.6)	50,245 (2.3)	1,949,754	2,496 (0.1)	18,002 (0.9)	61,677 (3.2)		
	7	2,157,101	1,595 (0.1)	13,301 (0.6)	45,669 (2.1)	2,019,370	3,443 (0.2)	22,300 (1.1)	49,711 (2.5)		
	0	2,112,380	7,221 (0.3)	74,690 (3.5)	170,757 (8.1)	1,656,723	13,938 (0.8)	104,699 (6.3)	173,749 (10.5)		
	1	2,424,231	54,659 (2.3)	275,510 (11.4)	315,454 (13.0)	1,937,029	240,342 (12.4)	706,424 (36.5)	330,245 (17.0)		
	2	2,474,399	23,894 (1.0)	143,988 (5.8)	228,253 (9.2)	2,035,773	60,995 (3.0)	292,539 (14.4)	337,135 (16.6)		
	3	2,459,431	11,711 (0.5)	78,023 (3.2)	147,930 (6.0)	1,993,354	19,031 (1.0)	126,765 (6.4)	213,889 (10.7)		
Fatigue	4	2,390,709	8.430 (0.4)	57.900 (2.4)	107.495 (4.5)	1.960.829	11.806 (0.6)	80.578 (4.1)	146,796 (7.5)		
	5	2,285,185	7.252 (0.3)	48.602 (2.1)	86,144 (3.8)	1.939.300	9,238 (0.5)	61,449 (3.2)	110.778 (5.7)		
	6	2,196,757	6.486 (0.3)	43.439 (2.0)	73.380 (3.3)	1.949.754	8.051 (0.4)	52,161 (2.7)	91,466 (4.7)		
	7	2.157.101	6.309 (0.3)	41.022 (1.9)	65.817 (3.1)	2.019.370	8.702 (0.4)	50.008 (2.5)	80.798 (4.0)		
	0	2 112 380	3 475 (0.2)	40.930 (1.9)	153 086 (7.2)	1 656 723	6 812 (0.4)	58 638 (3.5)	143 161 (8.6)		
	1	2,112,300	33 272 (1.4)	164 590 (6.8)	273 687 (11.3)	1,030,729	154 933 (8.0)	497 171 (25.7)	411 266 (21.2)		
	2	2,424,231	15 614 (0.6)	91.066 (3.7)	186 713 (7.5)	2 035 773	39 797 (2.0)	195 074 (9.6)	311 816 (15:3)		
	2	2,459,431	7 782 (0.3)	50 386 (2.0)	114 639 (4.7)	1 993 354	13,752 (0.7)	85 153 (4.3)	177 740 (8.9)		
Headache	1	2,159,151	5 818 (0.2)	38,567 (1.6)	86 140 (3.6)	1,950,829	8 903 (0.5)	56 974 (2,9)	120 773 (6.2)		
	5	2,390,709	5,316 (0.2)	34,309 (1.5)	72 (82 (3.2)	1,900,829	7 429 (0.4)	46 575 (2.4)	03 323 (4.8)		
	5	2,205,105	4 732 (0.2)	31 891 (1.5)	63 922 (2.9)	1,939,300	6 576 (0.3)	40,092 (2.1)	77 243 (4.0)		
	7	2,150,757	4,732 (0.2)	30,705 (1.4)	59 396 (2.8)	2 019 370	6 705 (0.3)	37,634 (1.9)	68 333 (3.4)		
	0	2,112,380	3,011,(0,1)	35,703 (1.4)	95.048 (4.5)	1 656 723	7,757 (0.5)	54,566 (3.3)	82 160 (5.0)		
	1	2,112,380	28.050 (1.6)	107 871 (8.2)	95,048 (4·5)	1,030,723	108 088 (10 2)	54,500 (5·5)	32,100 (3.0)		
	1	2,424,231	38,950 (1.6)	197,871 (8.2)	226,700 (9·4)	1,937,029	198,988 (10-3)	610,812(31.5)	292,422 (15.1)		
	2	2,474,399	13,531 (0·5) 5 264 (0 2)	$80,102(3\cdot 5)$	150,896 (0.1)	2,035,775	39,990 (2.0)	203,082 (10.0)	251,594 (12.4)		
Myalgia	3	2,439,431	3,204 (0.2)	36,080(1.0)	74,047 (S·0)	1,995,534	9,143 (0·3)	$33,042(3\cdot3)$	(5,072,(2,2))		
	4	2,390,709	3,627 (0.2)	20,050 (1.1)	47,845 (2.0)	1,960,829	5,041 (0.3)	36,805 (1.9)	65,072 (3·3)		
	5	2,285,185	2,989 (0.1)	22,955 (1.0)	37,471 (1.6)	1,939,300	3,796 (0.2)	27,445 (1.4)	45,771 (2.4)		
	6	2,196,757	2,667 (0.1)	21,040 (1.0)	33,060 (1.5)	1,949,754	3,592 (0.2)	24,073 (1-2)	37,073 (1.9)		
	/	2,157,101	2,731 (0.1)	20,799 (1.0)	31,659 (1.5)	2,019,370	4,415 (0.2)	25,026 (1.2)	33,652 (1.7)		
	0	2,112,380	1,395 (0.1)	10,125 (0.5)	35,178 (1.7)	1,656,723	4,685 (0.3)	23,091 (1.4)	45,194 (2.7)		
	1	2,424,231	23,553 (1.0)	84,997 (3.5)	101,682 (4.2)	1,937,029	137,685 (7-1)	402,336 (20.8)	291,400 (15.0)		
	2	2,474,399	7,601 (0.3)	33,238 (1.3)	52,246 (2.1)	2,035,773	23,939 (1.2)	90,569 (4.5)	113,067 (5.6)		
Chills	3	2,459,431	2,358 (0.1)	11,569 (0.5)	21,723 (0.9)	1,993,354	4,164 (0.2)	18,349 (0.9)	31,919 (1.6)		
	4	2,390,709	1,374 (0.1)	7,398 (0.3)	14,470 (0.6)	1,960,829	2,100 (0.1)	9,271 (0.5)	16,372 (0.8)		
	5	2,285,185	1,022 (0.04)	5,986 (0.3)	11,645 (0.5)	1,939,300	1,547 (0.1)	6,581 (0.3)	11,521 (0.6)		
	6	2,196,757	838 (0.04)	5,231 (0.2)	10,167 (0.5)	1,949,754	1,500 (0.1)	6,158 (0.3)	9,622 (0.5)		
	7	2,157,101	891 (0.04)	5,094 (0.2)	9,317 (0.4)	2,019,370	2,376 (0.1)	7,405 (0.4)	9,915 (0.5)		
	0	2,112,380	1,448 (0.1)	13,228 (0.6)	24,666 (1.2)	1,656,723	4,387 (0.3)	26,725 (1.6)	29,864 (1.8)		
	1	2,424,231	20,384 (0.8)	92,808 (3.8)	79,034 (3.3)	1,937,029	115,152 (5.9)	375,004 (19.4)	163,832 (8.5)		
	2	2,474,399	8,102 (0.3)	46,926 (1.9)	57,163 (2.3)	2,035,773	25,592 (1.3)	123,386 (6.1)	116,891 (5.7)		
Joint pain	3	2,459,431	3,509 (0.1)	23,226 (0.9)	33,006 (1.3)	1,993,354	6,277 (0.3)	41,550 (2.1)	55,855 (2.8)		
	4	2,390,709	2,400 (0.1)	16,852 (0.7)	23,553 (1.0)	1,960,829	3,601 (0.2)	24,570 (1.3)	34,118 (1.7)		
	5	2,285,185	2,010 (0.1)	14,447 (0.6)	19,143 (0.8)	1,939,300	2,831 (0.1)	18,829 (1.0)	25,132 (1.3)		
	6	2,196,757	1,821 (0.1)	13,250 (0.6)	16,821 (0.8)	1,949,754	2,582 (0.1)	16,587 (0.9)	21,152 (1.1)		
	7	2,157,101	1,874 (0.1)	13,157 (0.6)	16,230 (0.8)	2,019,370	3,240 (0.2)	17,390 (0.9)	19,800 (1.0)		

Data are n (%) unless otherwise stated.

*Top five reactions determined by reported frequency after dose 2 of both mRNA COVID-19 vaccines in v-safe, excluding fever because it was not rated

mild/moderate/severe. Symptoms are not mutually exclusive. [†]Mild was defined as "noticeable symptoms but they aren't a problem", moderate was defined as "symptoms that limit normal activities, and severe symptoms", and severe symptoms "make normal daily activities difficult or impossible".

Supplemental Table 8: Reported health impact^{*} 0-7 days after vaccination by mRNA COVID-19 vaccine manufacturer, dose, and sex reported in v-safe—December 14, 2020–June 14, 2021

	BNT162b2 va	accine					mRNA-1273 vaccine						
	Dose 1 (n=3,4	455,778)		Dose 2 (n=2,920,526)			Dose 1 (n=3,319,737)			Dose 2 (n=2,753,894)			
Sex/day	Unable to do normal activity	Unable to work	Reported medical care	Unable to do normal activity	Unable to work	Reported medical care	Unable to do normal activity	Unable to work	Reported medical care	Unable to do normal activity	Unable to work	Reported medical care	
Female													
Day 0	29,039 (2.1)	11,480 (0.8)	2,486 (0.2)	38,593 (3.5)	18,411 (1.7)	1,567 (0.1)	30,334 (2.4)	10,888 (0.8)	2,099 (0.2)	48,485 (4.6)	22,274 (2.1)	1,471 (0.1)	
Day 1	105,123 (6.6)	41,398 (2.6)	2,810 (0.2)	324,559 (24.9)	178,887 (13·7)	3,276 (0.3)	151,710 (10.0)	60,161 (4.0)	3,370 (0.2)	522,192 (41·4)	296,178 (23.5)	4,470 (0.4)	
Day 2	53,456 (3.4)	22,016 (1.4)	2,912 (0.2)	121,302 (8.9)	67,628 (5.0)	3,335 (0.2)	74,193 (4.8)	31,936 (2.1)	3,432 (0.2)	188,421 (14·2)	107,761 (8-1)	3,767 (0.3)	
Day 3	35,994 (2.3)	13,399 (0.9)	3,387 (0.2)	55,917 (4.2)	24,334 (1.8)	3,695 (0.3)	40,305 (2.6)	15,496 (1.0)	3,345 (0.2)	73,844 (5.7)	32,292 (2.5)	4,052 (0.3)	
Day 4	28,877 (1.9)	10,799 (0.7)	3,572 (0.2)	36,450 (2.8)	14,154 (1.1)	3,377 (0.3)	29,880 (2.0)	10,773 (0.7)	3,351 (0.2)	43,833 (3.4)	16,702 (1.3)	3,541 (0.3)	
Day 5	24,765 (1.7)	9,468 (0.6)	3,548 (0.2)	29,069 (2.3)	10,747 (0.8)	3,125 (0.2)	25,056 (1.7)	9,512 (0.7)	3,467 (0.2)	32,958 (2.6)	12,195 (1.0)	3,065 (0.2)	
Day 6	22,401 (1.6)	9,187 (0.7)	3,621 (0.3)	25,167 (2.0)	9,669 (0.8)	3,157 (0.2)	22,502 (1.6)	9,271 (0.7)	3,733 (0.3)	28,146 (2.2)	10,840 (0.9)	3,133 (0.2)	
Day 7	20,820 (1.5)	8,801 (0.6)	3,811 (0.3)	24,955 (1.9)	10,060 (0.8)	3,419 (0.3)	21,804 (1.6)	9,242 (0.7)	4,483 (0.3)	28,538 (2.2)	12,066 (0.9)	3,272 (0.3)	
Male													
Day 0	8,905 (1.0)	5,711 (0.7)	569 (0.1)	11,137 (1.7)	8,208 (1.3)	380 (0.1)	8,954 (1.1)	5,339 (0.7)	479 (0.1)	13,450 (2.3)	8,955 (1.5)	337 (0.1)	
Day 1	30,240 (3.2)	16,781 (1.8)	656 (0.1)	93,820 (13.3)	66,375 (9.4)	820 (0.1)	46,535 (5.3)	24,313 (2.8)	955 (0.1)	167,957 (25·6)	110,868 (16.9)	1,104 (0.2)	
Day 2	13,698 (1.5)	7,846 (0.8)	767 (0.1)	29,528 (4.0)	21,766 (3.0)	768 (0.1)	21,696 (2.4)	12,307 (1.4)	836 (0.1)	47,601 (6.9)	33,333 (4.9)	785 (0.1)	
Day 3	8,925 (1.0)	4,669 (0.5)	827 (0.1)	12,163 (1.7)	7,101 (1.0)	788 (0.1)	10,625 (1.2)	5,218 (0.6)	865 (0.1)	15,542 (2.3)	8,550 (1.3)	784 (0.1)	
Day 4	7,267 (0.8)	3,667 (0.4)	967 (0.1)	7,978 (1.1)	4,250 (0.6)	843 (0.1)	7,670 (0.9)	3,801 (0.4)	867 (0.1)	9,428 (1.4)	4,613 (0.7)	754 (0.1)	
Day 5	6,180 (0.7)	3,207 (0.4)	981 (0.1)	6,319 (0.9)	3,224 (0.5)	901 (0.1)	6,516 (0.8)	3,376 (0.4)	932 (0.1)	7,156 (1.1)	3,425 (0.5)	785 (0.1)	
Day 6	5,696 (0.7)	3,019 (0.4)	1,022 (0.1)	5,790 (0.8)	2,902 (0.4)	868 (0.1)	5,829 (0.7)	3,107 (0.4)	1,035 (0.1)	6,433 (1.0)	3,146 (0.5)	793 (0.1)	
Day 7	5,324 (0.7)	2,937 (0.4)	1,050 (0.1)	5,873 (0.8)	3,147 (0.4)	975 (0.1)	5,443 (0.7)	3,047 (0.4)	1,094 (0.1)	6,651 (0.9)	3,547 (0.5)	886 (0.1)	

Data are n (%) unless otherwise stated.

*Reports of health impacts are not mutually exclusive.

[†]Percent corresponds to number of respondents by sex and day.



Supplemental Figure 1: Number of reports of death per day following vaccination, by manufacturer, to Vaccine Adverse Event Reporting System (VAERS)—December 14, 2020–June 14, 2021

Days since last dose*

*x-axis reports through 161 days since last dose.





Top left: Female participants reporting health impact after receiving BNT162b2 vaccine. *Top right:* Female participants reporting health impact after receiving mRNA-1273 vaccine. *Bottom left:* Male participants reporting health impact after receiving mRNA-1273 vaccine.